

$\chi_{c1}(1P)$ 

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the Review on “ $\psi(2S)$  and  $\chi_c$  branching ratios” before the  $\chi_{c0}(1P)$  Listings.

### $\chi_{c1}(1P)$ MASS

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3510.67 ± 0.05</b>		<b>OUR AVERAGE</b>	Error includes scale factor of 1.2.		
3508.4 ± 1.9 ± 0.7		460	<sup>1</sup> AAIJ	17BB LHCB	$p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3510.71 ± 0.04 ± 0.09		4.8k	<sup>2</sup> AAIJ	17BI LHCB	$\chi_{c1} \rightarrow J/\psi\mu^+\mu^-$
3510.30 ± 0.14 ± 0.16			ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
3510.719 ± 0.051 ± 0.019			ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
3509.4 ± 0.9			BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3510.60 ± 0.087 ± 0.019		513	<sup>3</sup> ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3511.3 ± 0.4 ± 0.4		30	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
3512.3 ± 0.3 ± 4.0			<sup>4</sup> GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 ± 1.7		91	<sup>5</sup> LEMOIGNE	82 GOLI	$185\pi^-\text{Be} \rightarrow \gamma\mu^+\mu^-A$
3510.4 ± 0.6			OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3510.1 ± 1.1		254	<sup>6</sup> HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3509 ± 11		21	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3507 ± 3			<sup>6</sup> BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3505.0 ± 4 ± 4			<sup>6,7</sup> TANENBAUM	78 MRK1	$e^+e^-$
3513 ± 7		367	<sup>6</sup> BIDDICK	77 CNTR	$\psi(2S) \rightarrow \gamma X$
• • •			We do not use the following data for averages, fits, limits, etc. • • •		
3500 ± 10		40	TANENBAUM	75 MRK1	Hadrons $\gamma$

<sup>1</sup> From a fit of the  $\phi\phi$  invariant mass with the width of  $\chi_{c1}(1P)$  fixed to the PDG 16 value.

<sup>2</sup> AAIJ 17BI reports also  $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$  MeV.

<sup>3</sup> Recalculated by ANDREOTTI 05A, using the value of  $\psi(2S)$  mass from AULCHENKO 03.

<sup>4</sup> Using mass of  $\psi(2S) = 3686.0$  MeV.

<sup>5</sup>  $J/\psi(1S)$  mass constrained to 3097 MeV.

<sup>6</sup> Mass value shifted by us by amount appropriate for  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.

<sup>7</sup> From a simultaneous fit to radiative and hadronic decay channels.

### $\chi_{c1}(1P)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.84 ± 0.04</b>			<b>OUR FIT</b>		
<b>0.88 ± 0.05</b>			<b>OUR AVERAGE</b>		
1.39 <sup>+0.40</sup> <sub>-0.38</sub> <sup>+0.26</sup> <sub>-0.77</sub>			ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
0.876 ± 0.045 ± 0.026			ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
0.87 ± 0.11 ± 0.08		513	<sup>1</sup> ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<1.3	95	BAGLIN	86B	SPEC	$\bar{p}p \rightarrow e^+e^-X$
<3.8	90	GAISER	86	CBAL	$\psi(2S) \rightarrow \gamma X$

<sup>1</sup> Recalculated by ANDREOTTI 05A. **$\chi_{c1}(1P)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
<b>Hadronic decays</b>		
$\Gamma_1$	$3(\pi^+\pi^-)$	$(5.8 \pm 1.4) \times 10^{-3}$ S=1.2
$\Gamma_2$	$2(\pi^+\pi^-)$	$(7.6 \pm 2.6) \times 10^{-3}$
$\Gamma_3$	$\pi^+\pi^-\pi^0\pi^0$	$(1.19 \pm 0.15) \%$
$\Gamma_4$	$\rho^+\pi^-\pi^0 + \text{c.c.}$	$(1.45 \pm 0.24) \%$
$\Gamma_5$	$\rho^0\pi^+\pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$
$\Gamma_6$	$4\pi^0$	$(5.4 \pm 0.8) \times 10^{-4}$
$\Gamma_7$	$\pi^+\pi^-K^+K^-$	$(4.5 \pm 1.0) \times 10^{-3}$
$\Gamma_8$	$K^+K^-\pi^0\pi^0$	$(1.12 \pm 0.27) \times 10^{-3}$
$\Gamma_9$	$K^+K^-\pi^+\pi^-\pi^0$	$(1.15 \pm 0.13) \%$
$\Gamma_{10}$	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(7.5 \pm 0.8) \times 10^{-3}$
$\Gamma_{11}$	$K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(8.6 \pm 1.4) \times 10^{-3}$
$\Gamma_{12}$	$\rho^-K^+\bar{K}^0 + \text{c.c.}$	$(5.0 \pm 1.2) \times 10^{-3}$
$\Gamma_{13}$	$K^*(892)^0\bar{K}^0\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(2.3 \pm 0.6) \times 10^{-3}$
$\Gamma_{14}$	$K^+K^-\eta\pi^0$	$(1.12 \pm 0.34) \times 10^{-3}$
$\Gamma_{15}$	$\pi^+\pi^-K_S^0K_S^0$	$(6.9 \pm 2.9) \times 10^{-4}$
$\Gamma_{16}$	$K^+K^-\eta$	$(3.2 \pm 1.0) \times 10^{-4}$
$\Gamma_{17}$	$\bar{K}^0K^+\pi^- + \text{c.c.}$	$(7.0 \pm 0.6) \times 10^{-3}$
$\Gamma_{18}$	$K^*(892)^0\bar{K}^0 + \text{c.c.}$	$(10 \pm 4) \times 10^{-4}$
$\Gamma_{19}$	$K^*(892)^+K^- + \text{c.c.}$	$(1.4 \pm 0.6) \times 10^{-3}$
$\Gamma_{20}$	$K_J^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K^+\pi^- + \text{c.c.}$	$< 8 \times 10^{-4}$ CL=90%
$\Gamma_{21}$	$K_J^*(1430)^+K^- + \text{c.c.} \rightarrow$ $K_S^0K^+\pi^- + \text{c.c.}$	$< 2.1 \times 10^{-3}$ CL=90%
$\Gamma_{22}$	$K^+K^-\pi^0$	$(1.81 \pm 0.24) \times 10^{-3}$
$\Gamma_{23}$	$\eta\pi^+\pi^-$	$(4.62 \pm 0.23) \times 10^{-3}$
$\Gamma_{24}$	$a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(3.2 \pm 0.4) \times 10^{-3}$ S=2.2
$\Gamma_{25}$	$a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(1.76 \pm 0.24) \times 10^{-4}$
$\Gamma_{26}$	$a_2(1700)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(4.6 \pm 0.7) \times 10^{-5}$
$\Gamma_{27}$	$f_2(1270)\eta \rightarrow \eta\pi^+\pi^-$	$(3.5 \pm 0.6) \times 10^{-4}$
$\Gamma_{28}$	$f_4(2050)\eta \rightarrow \eta\pi^+\pi^-$	$(2.5 \pm 0.9) \times 10^{-5}$
$\Gamma_{29}$	$\pi_1(1400)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 5 \times 10^{-5}$ CL=90%
$\Gamma_{30}$	$\pi_1(1600)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 1.5 \times 10^{-5}$ CL=90%
$\Gamma_{31}$	$\pi_1(2015)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 8 \times 10^{-6}$ CL=90%

$\Gamma_{32}$	$f_2(1270)\eta$	$( 6.7 \pm 1.1 ) \times 10^{-4}$	
$\Gamma_{33}$	$\pi^+\pi^-\eta'$	$( 2.2 \pm 0.4 ) \times 10^{-3}$	
$\Gamma_{34}$	$K^+K^-\eta'(958)$	$( 8.8 \pm 0.9 ) \times 10^{-4}$	
$\Gamma_{35}$	$K_0^*(1430)^+K^- + \text{c.c.}$	$( 6.4 \begin{smallmatrix} +2.2 \\ -2.8 \end{smallmatrix} ) \times 10^{-4}$	
$\Gamma_{36}$	$f_0(980)\eta'(958)$	$( 1.6 \begin{smallmatrix} +1.4 \\ -0.7 \end{smallmatrix} ) \times 10^{-4}$	
$\Gamma_{37}$	$f_0(1710)\eta'(958)$	$( 7 \begin{smallmatrix} +7 \\ -5 \end{smallmatrix} ) \times 10^{-5}$	
$\Gamma_{38}$	$f_2'(1525)\eta'(958)$	$( 9 \pm 6 ) \times 10^{-5}$	
$\Gamma_{39}$	$\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-$	$( 3.5 \pm 0.9 ) \times 10^{-7}$	
$\Gamma_{40}$	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$( 3.2 \pm 2.1 ) \times 10^{-3}$	
$\Gamma_{41}$	$K^*(892)^0 \bar{K}^*(892)^0$	$( 1.4 \pm 0.4 ) \times 10^{-3}$	
$\Gamma_{42}$	$K^+K^-K_S^0K_S^0$	$< 4 \times 10^{-4}$	CL=90%
$\Gamma_{43}$	$K_S^0K_S^0K_S^0K_S^0$	$( 3.5 \pm 1.0 ) \times 10^{-5}$	
$\Gamma_{44}$	$K^+K^-K^+K^-$	$( 5.4 \pm 1.1 ) \times 10^{-4}$	
$\Gamma_{45}$	$K^+K^-\phi$	$( 4.1 \pm 1.5 ) \times 10^{-4}$	
$\Gamma_{46}$	$\bar{K}^0K^+\pi^-\phi + \text{c.c.}$	$( 3.3 \pm 0.5 ) \times 10^{-3}$	
$\Gamma_{47}$	$K^+K^-\pi^0\phi$	$( 1.62 \pm 0.30 ) \times 10^{-3}$	
$\Gamma_{48}$	$\phi\pi^+\pi^-\pi^0$	$( 7.5 \pm 1.0 ) \times 10^{-4}$	
$\Gamma_{49}$	$\omega\omega$	$( 5.7 \pm 0.7 ) \times 10^{-4}$	
$\Gamma_{50}$	$\omega K^+K^-$	$( 7.8 \pm 0.9 ) \times 10^{-4}$	
$\Gamma_{51}$	$\omega\phi$	$( 2.7 \pm 0.4 ) \times 10^{-5}$	
$\Gamma_{52}$	$\phi\phi$	$( 4.2 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{53}$	$\phi\phi\eta$	$( 3.0 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{54}$	$\rho\bar{\rho}$	$( 7.60 \pm 0.34 ) \times 10^{-5}$	
$\Gamma_{55}$	$\rho\bar{\rho}\pi^0$	$( 1.55 \pm 0.18 ) \times 10^{-4}$	
$\Gamma_{56}$	$\rho\bar{\rho}\eta$	$( 1.45 \pm 0.25 ) \times 10^{-4}$	
$\Gamma_{57}$	$\rho\bar{\rho}\omega$	$( 2.12 \pm 0.31 ) \times 10^{-4}$	
$\Gamma_{58}$	$\rho\bar{\rho}\phi$	$< 1.7 \times 10^{-5}$	CL=90%
$\Gamma_{59}$	$\rho\bar{\rho}\pi^+\pi^-$	$( 5.0 \pm 1.9 ) \times 10^{-4}$	
$\Gamma_{60}$	$\rho\bar{\rho}\pi^0\pi^0$	$< 5 \times 10^{-4}$	CL=90%
$\Gamma_{61}$	$\rho\bar{\rho}K^+K^- \text{ (non-resonant)}$	$( 1.27 \pm 0.22 ) \times 10^{-4}$	
$\Gamma_{62}$	$\rho\bar{\rho}K_S^0K_S^0$	$< 4.5 \times 10^{-4}$	CL=90%
$\Gamma_{63}$	$\rho\bar{n}\pi^-$	$( 3.8 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{64}$	$\bar{\rho}n\pi^+$	$( 3.9 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{65}$	$\rho\bar{n}\pi^-\pi^0$	$( 1.03 \pm 0.12 ) \times 10^{-3}$	
$\Gamma_{66}$	$\bar{\rho}n\pi^+\pi^0$	$( 1.01 \pm 0.12 ) \times 10^{-3}$	
$\Gamma_{67}$	$\Lambda\bar{\Lambda}$	$( 1.27 \pm 0.08 ) \times 10^{-4}$	
$\Gamma_{68}$	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$( 2.9 \pm 0.5 ) \times 10^{-4}$	
$\Gamma_{69}$	$\Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)}$	$( 2.5 \pm 0.6 ) \times 10^{-4}$	
$\Gamma_{70}$	$\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%
$\Gamma_{71}$	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90%
$\Gamma_{72}$	$K^+\bar{\rho}\Lambda + \text{c.c.}$	$( 4.2 \pm 0.4 ) \times 10^{-4}$	S=1.2
$\Gamma_{73}$	$nK_S^0\bar{\Lambda} + \text{c.c.}$	$( 1.66 \pm 0.17 ) \times 10^{-4}$	

$\Gamma_{74}$	$K^*(892)^+ \bar{p} \Lambda + \text{c.c.}$	$(4.9 \pm 0.7) \times 10^{-4}$	
$\Gamma_{75}$	$K^+ \bar{p} \Lambda(1520) + \text{c.c.}$	$(1.7 \pm 0.4) \times 10^{-4}$	
$\Gamma_{76}$	$\Lambda(1520) \bar{\Lambda}(1520)$	$< 9 \times 10^{-5}$	CL=90%
$\Gamma_{77}$	$\Sigma^0 \bar{\Sigma}^0$	$(4.2 \pm 0.6) \times 10^{-5}$	
$\Gamma_{78}$	$\Sigma^+ \bar{p} K_S^0 + \text{c.c.}$	$(1.53 \pm 0.12) \times 10^{-4}$	
$\Gamma_{79}$	$\Sigma^0 \bar{p} K^+ + \text{c.c.}$	$(1.46 \pm 0.10) \times 10^{-4}$	
$\Gamma_{80}$	$\Sigma^+ \bar{\Sigma}^-$	$(3.6 \pm 0.7) \times 10^{-5}$	
$\Gamma_{81}$	$\Sigma^- \bar{\Sigma}^+$	$(5.7 \pm 1.5) \times 10^{-5}$	
$\Gamma_{82}$	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$< 9 \times 10^{-5}$	CL=90%
$\Gamma_{83}$	$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$< 5 \times 10^{-5}$	CL=90%
$\Gamma_{84}$	$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(1.35 \pm 0.24) \times 10^{-4}$	
$\Gamma_{85}$	$\Xi^0 \bar{\Xi}^0$	$< 6 \times 10^{-5}$	CL=90%
$\Gamma_{86}$	$\Xi^- \bar{\Xi}^+$	$(8.0 \pm 2.1) \times 10^{-5}$	
$\Gamma_{87}$	$\pi^+ \pi^- + K^+ K^-$	$< 2.1 \times 10^{-3}$	
$\Gamma_{88}$	$K_S^0 K_S^0$	$< 6 \times 10^{-5}$	CL=90%
$\Gamma_{89}$	$\eta_c \pi^+ \pi^-$	$< 3.2 \times 10^{-3}$	CL=90%

### Radiative decays

$\Gamma_{90}$	$\gamma J/\psi(1S)$	$(34.3 \pm 1.0) \%$	
$\Gamma_{91}$	$\gamma \rho^0$	$(2.16 \pm 0.17) \times 10^{-4}$	
$\Gamma_{92}$	$\gamma \omega$	$(6.8 \pm 0.8) \times 10^{-5}$	
$\Gamma_{93}$	$\gamma \phi$	$(2.4 \pm 0.5) \times 10^{-5}$	
$\Gamma_{94}$	$\gamma \gamma$	$< 6.3 \times 10^{-6}$	CL=90%
$\Gamma_{95}$	$e^+ e^- J/\psi(1S)$	$(3.46 \pm 0.22) \times 10^{-3}$	
$\Gamma_{96}$	$\mu^+ \mu^- J/\psi(1S)$	$(2.33 \pm 0.29) \times 10^{-4}$	

### CONSTRAINED FIT INFORMATION

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a  $\chi^2 = 379.8$  for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

$x_{44}$	3				
$x_{54}$	4	2			
$x_{67}$	11	4	5		
$x_{90}$	23	9	2	29	
$\Gamma$	-12	-5	-63	-15	-41
	$x_{17}$	$x_{44}$	$x_{54}$	$x_{67}$	$x_{90}$

$\chi_{c1}(1P)$  PARTIAL WIDTHS $\chi_{c1}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$  $\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}}$   $\Gamma_{54} \Gamma_{90} / \Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>21.9 ± 0.8 OUR FIT</b>			
<b>21.4 ± 0.9 OUR AVERAGE</b>			
21.5 ± 0.5 ± 0.8	<sup>1</sup> ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
21.4 ± 1.5 ± 2.2	<sup>1,2</sup> ARMSTRONG 92	E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
19.9 <sup>+4.4</sup> <sub>-4.0</sub>	<sup>1</sup> BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+ e^- X$

<sup>1</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .<sup>2</sup> Recalculated by ANDREOTTI 05A. $\chi_{c1}(1P)$  BRANCHING RATIOS

## HADRONIC DECAYS

 $\Gamma(3(\pi^+ \pi^-)) / \Gamma_{\text{total}}$   $\Gamma_1 / \Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.8 ± 1.4 OUR EVALUATION</b>	Error includes scale factor of 1.2. Treating systematic error as correlated.		

**5.8 ± 1.1 OUR AVERAGE**

5.4 ± 0.7 ± 0.9	<sup>1</sup> BAI 99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
16.0 ± 5.9 ± 0.8	<sup>1</sup> TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$ . $\Gamma(2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$   $\Gamma_2 / \Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>7.6 ± 2.6 OUR EVALUATION</b>	Treating systematic error as correlated.		

**8 ± 4 OUR AVERAGE** Error includes scale factor of 1.5.

4.6 ± 2.1 ± 2.6	<sup>1</sup> BAI 99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
12.5 ± 4.2 ± 0.6	<sup>1</sup> TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$ . $\Gamma(\pi^+ \pi^- \pi^0 \pi^0) / \Gamma_{\text{total}}$   $\Gamma_3 / \Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.19 ± 0.15 ± 0.03</b>	604.7	<sup>1</sup> HE 08B	CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $1.28 \pm 0.06 \pm 0.15 \pm 0.08 \%$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- \pi^0 \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.45±0.24±0.04</b>	712.3	<sup>1,2</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $1.56 \pm 0.13 \pm 0.22 \pm 0.10$  % from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Calculated by us. We have added the values from HE 08B for  $\rho^+\pi^-\pi^0$  and  $\rho^-\pi^+\pi^0$  decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

$$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.9±3.5</b>	<sup>1</sup> TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the  $\psi(2S)$  decay.

$$\Gamma(4\pi^0)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.4±0.8±0.1</b>	608	<sup>1</sup> ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> ABLIKIM 11A reports  $(0.57 \pm 0.03 \pm 0.08) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.5±1.0 OUR EVALUATION</b>	Treating systematic error as correlated.		
<b>4.5±0.9 OUR AVERAGE</b>			

4.2±0.4±0.9 <sup>1</sup> BAI 99B BES  $\psi(2S) \rightarrow \gamma\chi_{c1}$

7.3±3.0±0.4 <sup>1</sup> TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ .

$$\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.12±0.27±0.03</b>	45.1	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $(0.12 \pm 0.02 \pm 0.02 \pm 0.01) \times 10^{-2}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.46±0.12±1.29</b>	12k	<sup>1</sup> ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$ .

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{10} / \Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.52±0.11±0.79</b>	5.1k	<sup>1</sup> ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$ .

$$\Gamma(K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{11} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.86±0.13±0.02</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.92 \pm 0.09 \pm 0.11 \pm 0.06$  % from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\rho^- K^+ \bar{K}^0 + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.50±0.12±0.01</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.54 \pm 0.11 \pm 0.07 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^- K^+ \bar{K}^0 + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{13} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.23±0.06±0.01</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.25 \pm 0.06 \pm 0.03 \pm 0.02$  % from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+ K^- \eta \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{14} / \Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.112±0.034±0.003</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.12 \pm 0.03 \pm 0.02 \pm 0.01$  % from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+ \pi^- K_S^0 K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{15} / \Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.9±2.9±0.2</b>	19.8±7.7	<sup>1</sup> ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.67 \pm 0.26 \pm 0.11) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.2 \pm 1.0 \pm 0.1</math></b>	<sup>1</sup> ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ATHAR 07 reports  $(0.34 \pm 0.10 \pm 0.04) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\overline{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>
<b><math>7.0 \pm 0.6</math> OUR FIT</b>	

 $\Gamma(K^*(892)^0 \overline{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.98 \pm 0.37 \pm 0.02</math></b>	22	<sup>1</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> ABLIKIM 06R reports  $(1.1 \pm 0.4 \pm 0.1) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \overline{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.43 \pm 0.65 \pm 0.03</math></b>	27	<sup>1</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> ABLIKIM 06R reports  $(1.6 \pm 0.7 \pm 0.2) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_J^*(1430)^0 \overline{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 8 \times 10^{-4}</math></b>	90	<sup>1</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> ABLIKIM 06R reports  $< 0.9 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^0 \overline{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

 $\Gamma(K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 2.1 \times 10^{-3}</math></b>	90	<sup>1</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> ABLIKIM 06R reports  $< 2.4 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .



$$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}} \qquad \Gamma_{22}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.81 ± 0.24 ± 0.04</b>	<sup>1</sup> ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ATHAR 07 reports  $(1.95 \pm 0.16 \pm 0.23) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}} \qquad \Gamma_{23}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.62 ± 0.23 OUR AVERAGE</b>				
4.58 ± 0.23 ± 0.11		<sup>1,2</sup> ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
4.7 ± 0.5 ± 0.1		<sup>3</sup> ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
5.3 ± 0.9 ± 0.1	222	<sup>4</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(4.67 \pm 0.03 \pm 0.23 \pm 0.16) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> ATHAR 07 reports  $(5.0 \pm 0.3 \pm 0.5) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> ABLIKIM 06R reports  $(5.9 \pm 0.7 \pm 0.8) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}} \qquad \Gamma_{24}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.2 ± 0.4 OUR AVERAGE</b>				Error includes scale factor of 2.2.
3.33 ± 0.19 ± 0.08		<sup>1,2</sup> ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
1.79 ± 0.63 ± 0.04	58	<sup>3</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(3.40 \pm 0.03 \pm 0.19 \pm 0.11) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> ABLIKIM 06R reports  $(2.0 \pm 0.5 \pm 0.5) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$**   **$\Gamma_{25} / \Gamma$** 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.176 \pm 0.023 \pm 0.004</math></b>	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(0.18 \pm 0.01 \pm 0.02 \pm 0.01) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 **$\Gamma(a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$**   **$\Gamma_{26} / \Gamma$** 

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>4.6 \pm 0.7 \pm 0.1</math></b>	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(4.7 \pm 0.4 \pm 0.6 \pm 0.2) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 **$\Gamma(f_2(1270)\eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$**   **$\Gamma_{27} / \Gamma$** 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.5 \pm 0.6 \pm 0.1</math></b>	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(0.36 \pm 0.01 \pm 0.06 \pm 0.01) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 **$\Gamma(f_4(2050)\eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$**   **$\Gamma_{28} / \Gamma$** 

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.5 \pm 0.9 \pm 0.1</math></b>	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(2.6 \pm 0.4 \pm 0.8 \pm 0.1) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_4(2050)\eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 **$\Gamma(\pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$**   **$\Gamma_{29} / \Gamma$** 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt; 5 \times 10^{-5}</math></b>	90	1,2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $< 4.6 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

$$\Gamma(\pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{30} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.5 \times 10^{-5}$	90	1,2 ABLIKIM 17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $< 1.5 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

$$\Gamma(\pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{31} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-6}$	90	1,2 ABLIKIM 17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $< 8 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

$$\Gamma(f_2(1270)\eta) / \Gamma_{\text{total}} \quad \Gamma_{32} / \Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.67 \pm 0.11</math> OUR AVERAGE</b>				
$0.63 \pm 0.11 \pm 0.02$		1,2 ABLIKIM 17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
$2.7 \pm 0.8 \pm 0.1$	53	<sup>3</sup> ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> ABLIKIM 17K reports  $(6.4 \pm 1.1) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> From an amplitude analysis using an isobar model.

<sup>3</sup> ABLIKIM 06R reports  $(3.0 \pm 0.7 \pm 0.5) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^+ \pi^- \eta') / \Gamma_{\text{total}} \quad \Gamma_{33} / \Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>2.2 \pm 0.4 \pm 0.1</math></b>	<sup>1</sup> ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ATHAR 07 reports  $(2.4 \pm 0.4 \pm 0.3) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- \eta') / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(K^+ K^- \eta'(958)) / \Gamma_{\text{total}} \quad \Gamma_{34} / \Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>8.75 \pm 0.87</math></b>	310	<sup>1</sup> ABLIKIM 14J	BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

<sup>1</sup> Derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.2 \pm 0.4)\%$ . Uncertainty includes both statistical and systematic contributions combined in quadrature.

$\Gamma(K_0^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$6.41 \pm 0.57^{+2.09}_{-2.71}$	<sup>1</sup> ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

 $\Gamma(f_0(980)\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$1.65 \pm 0.47^{+1.32}_{-0.56}$	<sup>1</sup> ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

 $\Gamma(f_0(1710)\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$0.71 \pm 0.22^{+0.68}_{-0.48}$	<sup>1</sup> ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

 $\Gamma(f_2'(1525)\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$0.92 \pm 0.23^{+0.55}_{-0.51}$	<sup>1</sup> ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

 $\Gamma(\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$0.35 \pm 0.09$		ABLIKIM	18D	BES3 $\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6                      90                      <sup>1</sup> ABLIKIM                      11D                      BES3                       $\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 11D reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 6.0 \times 10^{-7}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$32 \pm 21$	<sup>1</sup> TANENBAUM	78	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the  $\psi(2S)$  decay.

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.44 \pm 0.36 \pm 0.03$	$28.4 \pm 5.5$	<sup>1,2</sup> ABLIKIM	04H	BES $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 04H reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.40 \pm 0.27 \pm 0.22) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Assumes  $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$ .

$$\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}} \quad \Gamma_{42}/\Gamma$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

$<4 \times 10^{-4}$	90	$3.2 \pm 2.4$	<sup>1</sup> ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$
---------------------	----	---------------	----------------------	----------	-----------------------------------------

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 4.2 \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

$$\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}} \quad \Gamma_{43}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.35 \pm 0.10 \pm 0.01$	22	<sup>1</sup> ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$
--------------------------	----	----------------------	-----------	--------------------------------------

<sup>1</sup> Using  $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ . ABLIKIM 19AA reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (3.4 \pm 0.9 \pm 0.3) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value..

$$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{44}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID
--------------------------	-------------

$0.54 \pm 0.11$ OUR FIT	
-------------------------	--

$$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}} \quad \Gamma_{45}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.41 \pm 0.15 \pm 0.01$	17	<sup>1</sup> ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
--------------------------	----	----------------------	----------	-----------------------------------------

<sup>1</sup> ABLIKIM 06T reports  $(0.46 \pm 0.16 \pm 0.06) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{46}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$3.27 \pm 0.28 \pm 0.46$	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
--------------------------	---------	----------	-----------------------------------------

$$\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}} \quad \Gamma_{47}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.62 \pm 0.12 \pm 0.28$	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
--------------------------	---------	----------	-----------------------------------------

$$\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{48}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.75 \pm 0.06 \pm 0.08$	373	<sup>1</sup> ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$
--------------------------	-----	----------------------	----------	-------------------------------------------------------------

<sup>1</sup> Using  $1.06 \times 10^8 \psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$ .

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.7±0.7±0.1</b>	597	<sup>1</sup> ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons

<sup>1</sup> ABLIKIM 11K reports  $(6.0 \pm 0.3 \pm 0.7) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.78±0.04±0.08</b>	628	<sup>1</sup> ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$ .

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.27±0.04±0.01</b>	105	<sup>1</sup> ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.21±0.06±0.01      15      <sup>2,3</sup> ABLIKIM      11K BES3       $\psi(2S) \rightarrow \gamma$  hadrons

<sup>1</sup> ABLIKIM 19J reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  =  $(2.67 \pm 0.31 \pm 0.27) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 11K reports  $(0.22 \pm 0.06 \pm 0.02) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Superseded by ABLIKIM 19J.

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$   $\Gamma_{52}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.2±0.5±0.1</b>	366	<sup>1</sup> ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons

<sup>1</sup> ABLIKIM 11K reports  $(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$   $\Gamma_{53}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.0±0.4±0.2</b>	83.6	<sup>1</sup> ABLIKIM	20B BES3	$\psi(2S) \rightarrow \gamma\phi\phi\eta$

<sup>1</sup> ABLIKIM 20B reports  $(2.96 \pm 0.43 \pm 0.22) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ .

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$   
VALUE (units  $10^{-4}$ )      DOCUMENT ID  
**0.760±0.034 OUR FIT**

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$   
VALUE (units  $10^{-3}$ )      DOCUMENT ID      TECN      COMMENT  
**0.155±0.018 OUR AVERAGE**

0.163±0.019±0.004      <sup>1</sup> ONYISI      10      CLE3       $\psi(2S) \rightarrow \gamma p\bar{p}X$   
0.112±0.047±0.003      <sup>2</sup> ATHAR      07      CLEO       $\psi(2S) \rightarrow \gamma h^+ h^- h^0$   
<sup>1</sup> ONYISI 10 reports  $(1.75 \pm 0.16 \pm 0.13 \pm 0.11) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
<sup>2</sup> ATHAR 07 reports  $(1.2 \pm 0.5 \pm 0.1) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$   
VALUE (units  $10^{-3}$ )      CL%      DOCUMENT ID      TECN      COMMENT  
**0.145±0.024±0.004**      <sup>1</sup> ONYISI      10      CLE3       $\psi(2S) \rightarrow \gamma p\bar{p}X$

• • • We do not use the following data for averages, fits, limits, etc. • • •  
<0.15      90      <sup>2</sup> ATHAR      07      CLEO       $\psi(2S) \rightarrow \gamma h^+ h^- h^0$   
<sup>1</sup> ONYISI 10 reports  $(1.56 \pm 0.22 \pm 0.14 \pm 0.10) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
<sup>2</sup> ATHAR 07 reports  $< 0.16 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$   
VALUE (units  $10^{-3}$ )      DOCUMENT ID      TECN      COMMENT  
**0.212±0.030±0.005**      <sup>1</sup> ONYISI      10      CLE3       $\psi(2S) \rightarrow \gamma p\bar{p}X$

<sup>1</sup> ONYISI 10 reports  $(2.28 \pm 0.28 \pm 0.16 \pm 0.14) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma$   
VALUE      CL%      DOCUMENT ID      TECN      COMMENT  
**<1.7 × 10<sup>-5</sup>**      90      <sup>1</sup> ABLIKIM      11F      BES3       $\psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$

<sup>1</sup> ABLIKIM 11F reports  $< 1.82 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

$\Gamma(\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

**0.50±0.19 OUR EVALUATION** Treating systematic error as correlated.**0.50±0.19 OUR AVERAGE**

0.46±0.12±0.15

<sup>1</sup> BAI 99B BES  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 

1.08±0.77±0.05

<sup>1</sup> TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma\chi_{c1}$ <sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ . $\Gamma(\rho\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**<5 × 10<sup>-4</sup>** 90 <sup>1</sup> HE 08B CLEO  $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ <sup>1</sup> HE 08B reports  $< 0.05 \times 10^{-2}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ . $\Gamma(\rho\bar{p}K^+K^-(\text{non-resonant}))/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.27±0.22±0.03** 82 ± 9 <sup>1</sup> ABLIKIM 11F BES3  $\psi(2S) \rightarrow \gamma\rho\bar{p}K^+K^-$ <sup>1</sup> ABLIKIM 11F reports  $(1.35 \pm 0.15 \pm 0.19) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{p}K^+K^-(\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\rho\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

**<4.5** 90 <sup>1</sup> ABLIKIM 06D BES2  $\psi(2S) \rightarrow \gamma\chi_{c1}$ <sup>1</sup> Using  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.1 \pm 0.6)\%$ . $\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.8±0.5±0.1** 1412 <sup>1</sup> ABLIKIM 12J BES3  $\psi(2S) \rightarrow \gamma\rho\bar{n}\pi^-$ <sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  =  $(0.37 \pm 0.02 \pm 0.04) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.9±0.5±0.1** 1625 <sup>1</sup> ABLIKIM 12J BES3  $\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$ <sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  =  $(0.38 \pm 0.02 \pm 0.04) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.



$\Gamma(\rho\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.3±1.1±0.2</b>	1082	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\rho\bar{n}\pi^-\pi^0$

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (1.00 \pm 0.05 \pm 0.10) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.1±1.1±0.2</b>	1261	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.98 \pm 0.05 \pm 0.10) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID
<b>1.27±0.08 OUR FIT</b>	

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>29±5±1</b>		105	<sup>1</sup> ABLIKIM	12i BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<150      90      <sup>2</sup> ABLIKIM      06D BES2       $\psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> ABLIKIM 12i reports  $(31.1 \pm 3.4 \pm 3.9) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) (9.1 \pm 0.6)\%$ .

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25±6±1</b>	13	<sup>1</sup> ABLIKIM	12i BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

<sup>1</sup> ABLIKIM 12i reports  $(26.2 \pm 5.5 \pm 3.3) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.3 × 10<sup>-4</sup></b>	90	<sup>1</sup> ABLIKIM	12i BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

<sup>1</sup> ABLIKIM 12i reports  $< 14 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

$\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;13</b>	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^-\bar{\Lambda}\pi^+$
<sup>1</sup> ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .				

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.2 \pm 0.4</math> OUR AVERAGE</b>	Error includes scale factor of 1.2.			
$9.2^{+2.8}_{-2.4} \pm 0.4$	24	<sup>1</sup> LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
$4.2 \pm 0.4 \pm 0.1$	3k	<sup>2,3</sup> ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$
$3.1 \pm 0.9 \pm 0.1$		<sup>4</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
<sup>1</sup> LU 19 reports $(9.15^{+2.63}_{-2.25} \pm 0.86) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(1P) K^+)]$ assuming $B(B^+ \rightarrow \chi_{c1}(1P) K^+) = (4.79 \pm 0.23) \times 10^{-4}$ , which we rescale to our best value $B(B^+ \rightarrow \chi_{c1}(1P) K^+) = (4.74 \pm 0.22) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>2</sup> ABLIKIM 13D reports $(4.5 \pm 0.2 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>3</sup> Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$ .				
<sup>4</sup> ATHAR 07 reports $(3.3 \pm 0.9 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{73}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.66 \pm 0.12 \pm 0.12</math></b>	399	<sup>1</sup> ABLIKIM	21AV BES3	$\psi(2S) \rightarrow \gamma n K_S^0 \bar{\Lambda} + \text{c.c.}$
<sup>1</sup> ABLIKIM 21AV reports $(1.66 \pm 0.12 \pm 0.12) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow n K_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0975 \pm 0.0024$ . Also uses $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = (63.9 \pm 0.5)\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$ .				

 $\Gamma(K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{74}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.9 \pm 0.7 \pm 0.1</math></b>	328	<sup>1</sup> ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+} \bar{p}\Lambda$
<sup>1</sup> ABLIKIM 19AU reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $= (4.8 \pm 0.5 \pm 0.4) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(K^+\bar{p}\Lambda(1520)+\text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.71 \pm 0.44 \pm 0.04</math></b>	$48 \pm 10$	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

<sup>1</sup> ABLIKIM 11F reports  $(1.81 \pm 0.38 \pm 0.28) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda(1520)+\text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 9 \times 10^{-5}</math></b>	90	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

<sup>1</sup> ABLIKIM 11F reports  $< 1.00 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.2 \pm 0.6 \pm 0.1</math></b>		103	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 6$	90		<sup>2</sup> ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
$< 4$	90	$3.8 \pm 2.5$	<sup>3</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.41 \pm 0.05 \pm 0.03) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 13H reports  $< 0.62 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

<sup>3</sup> NAIK 08 reports  $< 0.44 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$   $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.6 \pm 0.6 \pm 0.1</math></b>		59	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 8$	90		<sup>2</sup> ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
$< 6$	90	$4.3 \pm 2.3$	<sup>3</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.35 \pm 0.06 \pm 0.02) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 13H reports  $< 0.87 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

<sup>3</sup> NAIK 08 reports  $< 0.65 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

### $\Gamma(\Sigma^- \bar{\Sigma}^+) / \Gamma_{\text{total}}$ $\Gamma_{81} / \Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5.7 \pm 1.4 \pm 0.6</math></b>	214	<sup>1</sup> ABLIKIM	20i	BES3 $\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

<sup>1</sup> ABLIKIM 20i reports  $(5.7 \pm 1.4 \pm 0.6) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ .

### $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-) / \Gamma_{\text{total}}$ $\Gamma_{82} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 9 \times 10^{-5}</math></b>	90	<sup>1</sup> ABLIKIM	12i	BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 12i reports  $< 10 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

### $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+) / \Gamma_{\text{total}}$ $\Gamma_{83} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 5 \times 10^{-5}</math></b>	90	<sup>1</sup> ABLIKIM	12i	BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 12i reports  $< 5.7 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

### $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.}) / \Gamma_{\text{total}}$ $\Gamma_{84} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.35 \pm 0.24 \pm 0.03</math></b>	49	<sup>1</sup> ABLIKIM	15i	BES3 $\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$

<sup>1</sup> ABLIKIM 15i reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$   $= (1.32 \pm 0.20 \pm 0.12) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\Xi^0 \bar{\Xi}^0) / \Gamma_{\text{total}}$ $\Gamma_{85} / \Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 6 \times 10^{-5}</math></b>	90	$1.7 \pm 2.4$	<sup>1</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$

<sup>1</sup> NAIK 08 reports  $< 0.60 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \bar{\Xi}^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

### $\Gamma(\Xi^- \bar{\Xi}^+) / \Gamma_{\text{total}}$ $\Gamma_{86} / \Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.80 \pm 0.21 \pm 0.02</math></b>	$16.4 \pm 4.3$		<sup>1</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^+ \bar{\Xi}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.4                      90                      <sup>2</sup> ABLIKIM      06D    BES2     $\psi(2S) \rightarrow \gamma\chi_{c1}$   
<sup>1</sup> NAIK 08 reports  $(0.86 \pm 0.22 \pm 0.08) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
<sup>2</sup> Using  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) (9.1 \pm 0.6)\%$ .

**$[\Gamma(\pi^+\pi^-) + \Gamma(K^+K^-)]/\Gamma_{\text{total}}$                        $\Gamma_{87}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 21 × 10 <sup>-4</sup>		<sup>1</sup> FELDMAN    77	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 38 × 10<sup>-4</sup>                      90                      <sup>1</sup> BRANDELIK    79B    DASP     $\psi(2S) \rightarrow \gamma\chi_{c1}$   
<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the  $\psi(2S)$  decay.

**$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$                        $\Gamma_{88}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6 × 10 <sup>-5</sup>	90	<sup>1</sup> ABLIKIM      050	BES2	$\psi(2S) \rightarrow \chi_{c1}\gamma$

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  <  $0.6 \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

**$\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$                        $\Gamma_{89}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2 × 10 <sup>-3</sup>	90	<sup>1,2</sup> ABLIKIM    13B	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.4 × 10<sup>-3</sup>                      90                      <sup>1,3</sup> ABLIKIM      13B    BES3     $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$   
<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$ .  
<sup>2</sup> Using the  $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$  decays.  
<sup>3</sup> Using the  $\eta_c \rightarrow K^+ K^- \pi^0$  decays.

————— **RADIATIVE DECAYS** —————

**$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$                        $\Gamma_{90}/\Gamma$**

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>34.3 ± 1.0</b>	<b>OUR FIT</b>			

• • • We do not use the following data for averages, fits, limits, etc. • • •

34.75 ± 0.11 ± 1.70    1.9M                      <sup>1</sup> ABLIKIM      17U    BES3     $e^+e^- \rightarrow \gamma X$   
37.9 ± 0.8 ± 2.1                      <sup>2</sup> ADAM          05A    CLEO     $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$   
<sup>1</sup> Not independent from  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))$  and the product  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))$  also measured in ABLIKIM 17U.  
<sup>2</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\gamma J/\psi)$  from ADAM 05A and  $B(\psi(2S) \rightarrow \gamma\chi_{c1})$  from ATHAR 04.

### $\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ $\Gamma_{91}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	-------------	--------------------	-------------	----------------

#### **216 ± 17 OUR AVERAGE**

215 ± 22 ± 5	432 ± 25	<sup>1</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
217 ± 24 ± 5	186 ± 15	<sup>2</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$

<sup>1</sup> ABLIKIM 11E reports  $(228 \pm 13 \pm 22) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> BENNETT 08A reports  $(243 \pm 19 \pm 22) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ $\Gamma_{92}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	-------------	--------------------	-------------	----------------

#### **68 ± 8 OUR AVERAGE**

66 ± 9 ± 2	136 ± 14	<sup>1</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
74 ± 17 ± 2	39 ± 7	<sup>2</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$

<sup>1</sup> ABLIKIM 11E reports  $(69.7 \pm 7.2 \pm 6.6) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> BENNETT 08A reports  $(83 \pm 15 \pm 12) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ $\Gamma_{93}/\Gamma$

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	------------	-------------	--------------------	-------------	----------------

<b>24 ± 5 ± 1</b>		43 ± 9	<sup>1</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
-------------------	--	--------	----------------------	----------	-----------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 23$	90	5.2 ± 3.1	<sup>2</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
--------	----	-----------	----------------------	----------	-----------------------------------------

<sup>1</sup> ABLIKIM 11E reports  $(25.8 \pm 5.2 \pm 2.3) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> BENNETT 08A reports  $< 26 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

### $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{94}/\Gamma$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	--------------------	-------------	----------------

$< 6.3 \times 10^{-6}$	90	ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$
------------------------	----	---------	-----------	------------------------------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.5 \times 10^{-5}$	90	ECKLUND	08A	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1} \rightarrow 3\gamma$
$< 150 \times 10^{-5}$	90	<sup>1</sup> YAMADA	77	DASP	$e^+e^- \rightarrow 3\gamma$

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the  $\psi(2S)$  decay.

**$\Gamma(e^+e^- J/\psi(1S))/\Gamma_{\text{total}}$   $\Gamma_{95}/\Gamma$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.65 \pm 0.23 \pm 0.09$	1.9k	<sup>1,2</sup> ABLIKIM	17i	BES3	$\psi(2S) \rightarrow \gamma e^+e^- J/\psi$
--------------------------	------	------------------------	-----	------	---------------------------------------------

<sup>1</sup> ABLIKIM 17i reports  $(3.73 \pm 0.09 \pm 0.25) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow e^+e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Not independent from other measurements reported by ABLIKIM 17i

**$\Gamma(e^+e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$   $\Gamma_{95}/\Gamma_{90}$**

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	-------------	--------------------	-------------	----------------

<b><math>10.1 \pm 0.3 \pm 0.5</math></b>	1.9k	<sup>1</sup> ABLIKIM	17i	BES3	$\psi(2S) \rightarrow e^+e^- \gamma J/\psi$
------------------------------------------	------	----------------------	-----	------	---------------------------------------------

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

**$\Gamma(\mu^+\mu^- J/\psi(1S))/\Gamma(e^+e^- J/\psi(1S))$   $\Gamma_{96}/\Gamma_{95}$**

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	-------------	--------------------	-------------	----------------

<b><math>6.73 \pm 0.51 \pm 0.50</math></b>	222	ABLIKIM	19Z	BES3	$\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma(\mu^+\mu^- J/\psi)$
--------------------------------------------	-----	---------	-----	------	----------------------------------------------------------------------------

**$\chi_{c1}(1P)$  CROSS-PARTICLE BRANCHING RATIOS**

**$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$   $\Gamma_{54}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$**

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	--------------------	-------------	----------------

**$2.14 \pm 0.10$  OUR FIT**

<b><math>1.1 \pm 1.0</math></b>	<sup>1</sup> BAI	98i	BES	$\psi(2S) \rightarrow \gamma \chi_{c1} \rightarrow \gamma p\bar{p}$
---------------------------------	------------------	-----	-----	---------------------------------------------------------------------

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c1} \rightarrow p\bar{p})$  reported in BAI 98i is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.7 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

**$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{\psi(2S)}$**

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------------------------------	-------------	--------------------	-------------	----------------

**$12.4 \pm 0.8$  OUR FIT**

**$12.3 \pm 0.9$  OUR AVERAGE** Error includes scale factor of 1.2.

$12.8 \pm 0.6 \pm 0.6$	528	ABLIKIM	21L	BES3	$\psi(2S) \rightarrow \gamma p\pi^- \bar{p}\pi^+$
$10.5 \pm 1.6 \pm 0.6$	46	<sup>1</sup> NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.2±1.0±0.9      136    2,3 ABLIKIM      13H BES3     $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c1} \rightarrow \Lambda \bar{\Lambda}) = (11.6 \pm 1.8 \pm 0.7 \pm 0.7) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$ .

<sup>2</sup> Superseded by ABLIKIM 21L

<sup>3</sup> Calculated by us. ABLIKIM 13H reports  $B(\chi_{c1} \rightarrow \Lambda \bar{\Lambda}) = (12.2 \pm 1.1 \pm 1.1) \times 10^{-5}$  from a measurement of  $B(\chi_{c1} \rightarrow \Lambda \bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma \chi_{c1})$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.2 \pm 0.4)\%$ .

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda \bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{J/\psi(1S)\pi^+\pi^-} \quad \frac{\Gamma_{67}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}{\Gamma_{163}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10 <sup>-5</sup> )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>3.58±0.22 OUR FIT</b>				

<b>7.1</b>	<b>+2.8 -2.4 ±1.3</b>	9.0 <sup>+3.5</sup> -3.1	<sup>1</sup> BAI	03E BES	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$
------------	---------------------------	-----------------------------	------------------	---------	-----------------------------------------------------

<sup>1</sup> BAI 03E reports [  $B(\chi_{c1} \rightarrow \Lambda \bar{\Lambda}) B(\psi(2S) \rightarrow \gamma \chi_{c1}) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) ] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p \bar{p})] = (1.33_{-0.46}^{+0.52} \pm 0.25)\%$ . We calculate from this measurement the presented value using  $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$  and  $B(J/\psi \rightarrow p \bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$ .

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}}}{J/\psi(1S)} \quad \frac{\Gamma_{90}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}{\Gamma_{163}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10 <sup>-2</sup> )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>3.34 ±0.06 OUR FIT</b>				

**3.24 ±0.16 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.

3.518±0.010±0.120	143k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
3.442±0.010±0.132	1.9M	ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
2.81 ±0.05 ±0.23	13k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
2.56 ±0.12 ±0.20		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
2.78 ±0.30		<sup>2</sup> OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \chi_{c1}$
2.2 ±0.5		<sup>3</sup> BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c1}$
2.9 ±0.5		<sup>3</sup> BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma \chi_{c1}$
5.0 ±1.5		<sup>4</sup> BIDDICK	77 CNTR	$e^+ e^- \rightarrow \gamma X$
2.8 ±0.9		<sup>2</sup> WHITAKER	76 MRK1	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.377±0.009±0.183	142k	<sup>5</sup> ABLIKIM	120 BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3.56 ±0.03 ±0.12	24.9k	<sup>6</sup> MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3.44 ±0.06 ±0.13	3.7k	<sup>7</sup> ADAM	05A CLEO	Repl. by MENDEZ 08

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$ .

<sup>2</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

<sup>3</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .


<sup>4</sup> Assumes isotropic gamma distribution.

<sup>5</sup> Superseded by ABLIKIM 17N.

<sup>6</sup> Not independent from other measurements of MENDEZ 08.

<sup>7</sup> Not independent from other values reported by ADAM 05A.





$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}} \text{ (units } 10^{-2}\text{)}$$

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{90}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)} = \Gamma_{90}/\Gamma \times \Gamma_{163}^{\psi(2S)}/(\Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + \Gamma_{14}^{\psi(2S)} + 0.343\Gamma_{163}^{\psi(2S)} + 0.190\Gamma_{164}^{\psi(2S)})$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.43±0.10 OUR FIT</b>				

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 5.70±0.04±0.15    24.9k    <sup>1</sup> MENDEZ    08    CLEO     $\psi(2S) \rightarrow \gamma \chi_{c1}$
- 5.77±0.10±0.12    3.7k    ADAM    05A    CLEO    Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{90}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.63±0.17 OUR FIT</b>				

- 10.15±0.28 OUR AVERAGE**
- 10.17±0.07±0.27    24.9k    MENDEZ    08    CLEO     $\psi(2S) \rightarrow \gamma \chi_{c1}$
  - 12.6 ±0.3 ±3.8    3k    <sup>1</sup> ABLIKIM    04B    BES     $\psi(2S) \rightarrow J/\psi X$
  - 8.5 ±2.1    <sup>2</sup> HIMEL    80    MRK2     $\psi(2S) \rightarrow \gamma \chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.24±0.17±0.23      3.7k      <sup>3</sup> ADAM      05A CLEO Repl. by MENDEZ 08

<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>2</sup> The value for  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$  quoted in HIMEL 80 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$  and  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$ .

<sup>3</sup> Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{17}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10 <sup>-4</sup> )	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

**6.8±0.5 OUR FIT**

**7.2±0.6 OUR AVERAGE**

7.3±0.5±0.5	<sup>1</sup> ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
-------------	--------------------	----	----------------------------------------------------

7.0±0.5±0.9	<sup>2</sup> ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma\chi_{c1}$
-------------	----------------------	-----	---------------------------------------------

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})$  reported by ATHAR 07 was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54)\%$ .

<sup>2</sup> Calculated by us. ABLIKIM 06R reports  $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-) = (4.0 \pm 0.3 \pm 0.5) \times 10^{-3}$ . We use  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.4) \times 10^{-2}$ .

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \Gamma_{17}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10 <sup>-4</sup> )	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

**19.6±1.6 OUR FIT**

**13.2±2.4±3.2**

<sup>1</sup> BAI	99B	BES	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
------------------	-----	-----	-----------------------------------------------

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{44}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

**0.53±0.11 OUR FIT**

<b>0.61±0.11±0.08</b>	54	<sup>1</sup> ABLIKIM	06T	BES2 $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$
-----------------------	----	----------------------	-----	----------------------------------------------------

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ .

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \Gamma_{44}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10 <sup>-4</sup> )	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

**1.52±0.31 OUR FIT**

<b>1.13±0.40±0.29</b>	<sup>1</sup> BAI	99B	BES $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$
-----------------------	------------------	-----	---------------------------------------------------


<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

$$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{54}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.41 ± 0.35</b>				<b>OUR FIT</b>
<b>7.8 ± 0.6</b>				<b>OUR AVERAGE</b> Error includes scale factor of 1.4. See the ideogram below.
7.9 ± 0.4 ± 0.3	453	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
8.2 ± 0.7 ± 0.4	141 ± 13	<sup>1</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
4.8 $^{+1.4}_{-1.3}$ ± 0.6	18.2 $^{+5.5}_{-4.9}$	BAI	04F BES	$\psi(2S) \rightarrow \gamma\chi_{c1}(1P) \rightarrow \gamma\bar{p}p$

WEIGHTED AVERAGE  
7.8 ± 0.6 (Error scaled by 1.4)



<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c1} \rightarrow p\bar{p}) = (9.0 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$ .

$$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}} \text{ (units } 10^{-6}\text{)}$$

$$\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{78}/\Gamma \times \Gamma_{163}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.49 ± 0.09 ± 0.07</b>	258	<sup>1</sup> ABLIKIM	19BB BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

<sup>1</sup> Calculated by us. ABLIKIM 19BB reports  $B(\chi_{c1} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (1.53 \pm 0.10 \pm 0.08) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.75 \pm 0.24)\%$  and other branching fractions from PDG 18.

$$\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}} \\ \Gamma_{79} / \Gamma \times \Gamma_{163}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.42 \pm 0.07 \pm 0.06</math></b>	493	<sup>1</sup> ABLIKIM	20AE BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+$ + c.c.


<sup>1</sup> Calculated by us. ABLIKIM 20AE reports  $B(\chi_{c1} \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (1.46 \pm 0.07 \pm 0.07) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma \chi_c^0) = (9.75 \pm 0.24)\%$  and other branching fractions from PDG 20.

### MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

**$a_2 = M2 / \sqrt{E1^2 + M2^2}$  Magnetic quadrupole fractional transition amplitude**

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-6.7 \pm 0.9</math> OUR AVERAGE</b>				Error includes scale factor of 2.6. See the ideogram below.
$-7.40 \pm 0.33 \pm 0.34$	164k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$-6.26 \pm 0.63 \pm 0.24$	39k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$0.2 \pm 3.2 \pm 0.4$	2090	AMBROGIANI	02 E835	$p \bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi \gamma$
$-0.2 \begin{smallmatrix} +0.8 \\ -2.0 \end{smallmatrix}$	921	OREGLIA	82 CBAL	$\psi(2S) \rightarrow \chi_{c1} \gamma \rightarrow J/\psi \gamma \gamma$

WEIGHTED AVERAGE  
 $-6.7 \pm 0.9$  (Error scaled by 2.6)



<sup>1</sup> Correlated with  $b_2$  with correlation coefficient  $\rho_{a_2 b_2} = 0.133$ .

$$a_2 = M2 / \sqrt{E1^2 + M2^2} \text{ (units } 10^{-2}\text{)}$$

MULTIPOLE AMPLITUDES IN  $\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$  RADIATIVE DECAY $b_2 = M_2/\sqrt{E_1^2 + M_2^2}$  Magnetic quadrupole fractional transition amplitude

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.5 ± 0.4 OUR AVERAGE</b>				
2.29 ± 0.39 ± 0.27	164k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.76 ± 0.73 ± 0.23	39k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
7.7 <sup>+5.0</sup> <sub>-4.5</sub>	921	OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>1</sup> Correlated with  $a_2$  with correlation coefficient  $\rho_{a_2 b_2} = 0.133$ .

## MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

 $\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$  and  $\chi_{c1} \rightarrow \gamma J/\psi(1S)$  $a_2/b_2$  Magnetic quadrupole transition amplitude ratio

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-2.27 <sup>+0.57</sup><sub>-0.99</sub></b>	39k	<sup>1</sup> ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>1</sup> Statistical and systematic errors combined. Not independent of  $a_2(\chi_{c1})$  and  $b_2(\chi_{c1})$  values from ARTUSO 09. $\chi_{c1}(1P)$  REFERENCES

ABLIKIM	21AV	JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AE	PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20I	PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BI	PRL 119 221801	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
Also		Private Comm.	M.J. Oreglia	(EFI)
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)
TANENBAUM	75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)