


η(1405)

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See also the η(1475).

η(1405) MASS

VALUE (MeV) DOCUMENT ID
1408.8 ± 2.0 OUR AVERAGE Includes data from the 2 datablocks that follow this one.
 Error includes scale factor of 2.2. See the ideogram below.



ηππ MODE


VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
 The data in this block is included in the average printed for a previous datablock.

1405.8 ± 2.6 OUR AVERAGE Error includes scale factor of 2.3. See the ideogram below.

1409.0 ± 1.7	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
1407.0 ± 3.5	198	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$
1399.8 ± 2.2 ^{+2.8} _{-0.1}		¹ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
1392 ± 14	900 ± 375	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$
1394 ± 8	6.6 ± 2.0k	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$
1404 ± 6	9082	MANAK	00A MPS	$18 \pi^-p \rightarrow \eta\pi^+\pi^-n$

1424 ± 6	2200	ALDE	97B GAM4	100	$\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1400 ± 6		² BOLTON	92B MRK3		$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
1388 ± 4		FUKUI	91C SPEC	8.95	$\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1398 ± 6	261	³ AUGUSTIN	90 DM2		$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
1420 ± 5		ANDO	86 SPEC	8	$\pi^- p \rightarrow \eta \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1404.0 ± 11.0	195	ABLIKIM	19BABES3		$e^+ e^- \rightarrow \psi(2S)$
1385 ± 7		BAI	99 BES		$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
1409 ± 3		⁴ AMSLER	95F CBAR		$0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta$

- ¹ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.
- ² From fit to the $a_0(980) \pi 0^-+$ partial wave.
- ³ Best fit with a single Breit Wigner.
- ⁴ Superseded by AMSLER 04B.



$K \bar{K} \pi$ MODE ($a_0(980) \pi$ or direct $K \bar{K} \pi$)

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

1413.9 ± 1.7 OUR AVERAGE Error includes scale factor of 1.1.

1413 ± 14	3651	¹ NICHITIU	02	OBLX	0	$\bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1416 ± 4 ± 2	20k	ADAMS	01B	B852	18	GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1405 ± 5		² CICALO	99	OBLX	0	$\bar{p} p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
1407 ± 5		² BERTIN	97	OBLX	0	$\bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
1416 ± 2		² BERTIN	95	OBLX	0	$\bar{p} p \rightarrow K \bar{K} \pi \pi \pi$
1416 ± 8 ⁺⁷ / ₋₅	700	³ BAI	90C	MRK3		$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

1413 ± 5 ³ RATH 89 MPS 21.4 $\pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$

••• We do not use the following data for averages, fits, limits, etc. •••

1459 ± 5 ⁴ AUGUSTIN 92 DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$

¹ Decaying dominantly directly to $K^+ K^- \pi^0$.

² Decaying into $(K \bar{K})_S \pi$, $(K \pi)_S \bar{K}$, and $a_0(980) \pi$.

³ From fit to the $a_0(980) \pi 0^-+$ partial wave. Cannot rule out a $a_0(980) \pi 1^++$ partial wave.

⁴ Excluded from averaging because averaging would be meaningless.

$\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1403 \pm 17 OUR AVERAGE Error includes scale factor of 1.8.

1390 \pm 12	235 \pm 91	AMSLEER	04B CBAR	0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \eta$
---------------	--------------	---------	----------	--

1424 \pm 10 \pm 11	547	BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
------------------------	-----	-----	----------	--

••• We do not use the following data for averages, fits, limits, etc. •••

1401 \pm 18		^{1,2} AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+ \pi^- \gamma \gamma$
---------------	--	-------------------------	--------	--

1432 \pm 8		² COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+ \pi^- 2\gamma$
--------------	--	----------------------	---------	--

¹ Best fit with a single Breit Wigner.

² This peak in the $\gamma\rho$ channel may not be related to the $\eta(1405)$.

4π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

1420 \pm 20		BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
---------------	--	------	---------	---

1489 \pm 12	3270	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
---------------	------	----------------------	---------	---------------------------------

¹ Estimated by us from various fits.

$K\bar{K}\pi$ MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

1452.7 \pm 3.3	191	^{1,2} ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$
------------------	-----	------------------------	----------	---------------------------------------

1437.6 \pm 3.2	249 \pm 35	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$
------------------	--------------	------------------------	----------	--

1445.9 \pm 5.7	62 \pm 18	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
------------------	-------------	------------------------	----------	---

1442 \pm 10	410	¹ BAI	98C BES	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
---------------	-----	------------------	---------	---

1445 \pm 8	693	¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
--------------	-----	-----------------------	--------	---

1433 \pm 8	296	¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
--------------	-----	-----------------------	--------	---

1413 \pm 8	500	¹ DUCH	89 ASTE	$\bar{p} p \rightarrow \pi^+ \pi^- K^\pm \pi^\mp K^0$
--------------	-----	-------------------	---------	---

1453 \pm 7	170	¹ RATH	89 MPS	21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
--------------	-----	-------------------	--------	--

1419 \pm 1	8800	¹ BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
--------------	------	---------------------	--------	---

1424 \pm 3	620	¹ REEVES	86 SPEC	6.6 $p\bar{p} \rightarrow K \bar{K} \pi X$
--------------	-----	---------------------	---------	--

1421 \pm 2		¹ CHUNG	85 SPEC	8 $\pi^- p \rightarrow K \bar{K} \pi n$
--------------	--	--------------------	---------	---

1440 $\begin{smallmatrix} +20 \\ -15 \end{smallmatrix}$	174	¹ EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
---	-----	----------------------	----------	---

1440 $\begin{smallmatrix} +10 \\ -15 \end{smallmatrix}$		¹ SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
---	--	----------------------	---------	---

1425 \pm 7	800	^{1,3} BAILLON	67 HBC	0 $\bar{p} p \rightarrow K \bar{K} \pi \pi \pi$
--------------	-----	------------------------	--------	---

¹ These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

² Systematic uncertainty not evaluated.

³ From best fit of 0^-+ partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.


$\eta(1405)$ WIDTH

VALUE (MeV)

DOCUMENT ID

50.1 ± 2.6 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.7. See the ideogram below.

WEIGHTED AVERAGE
50.1 ± 2.6 (Error scaled by 1.7)



$\eta\pi\pi$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.


52.6 ± 3.2 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

48.3 ± 5.2	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
55.0 ± 11.0	198	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$
52.8 ± 7.6 ^{+0.1} _{-7.6}		¹ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
55 ± 11	900	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$
55 ± 12	6.6k	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma$
80 ± 21	9.0k	MANAK	00A MPS	$18 \pi^-p \rightarrow \eta\pi^+\pi^-n$
85 ± 18	2.2k	ALDE	97B GAM4	$100 \pi^-p \rightarrow \eta\pi^0\pi^0n$
47 ± 13		² BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
59 ± 4		FUKUI	91C SPEC	$8.95 \pi^-p \rightarrow \eta\pi^+\pi^-n$
53 ± 11		³ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
31 ± 7		ANDO	86 SPEC	$8 \pi^-p \rightarrow \eta\pi^+\pi^-n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

79.0 ± 16.0 195 ABLIKIM 19BA BES3 $e^+e^- \rightarrow \psi(2S)$
 86 ± 10 4 AMSLER 95F CBAR $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$

- ¹ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.
- ² From fit to the $a_0(980)\pi 0^-+$ partial wave.
- ³ From $\eta\pi^+\pi^-$ mass distribution - mainly $a_0(980)\pi$ - no spin-parity determination available.
- ⁴ Superseded by AMSLER 04B.



$K\bar{K}\pi$ MODE ($a_0(980)\pi$ or direct $K\bar{K}\pi$)

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT


The data in this block is included in the average printed for a previous datablock.

48 ± 4 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.

51 ± 6	3651	¹ NICHITIU	02	OBLX	$0 \bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0$
42 ± 10 ± 9	20k	ADAMS	01B	B852	$18 \text{ GeV } \pi^-p \rightarrow K^+K^-\pi^0n$
50 ± 4		CICALO	99	OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
48 ± 5		² BERTIN	97	OBLX	$0.0 \bar{p}p \rightarrow K^\pm(K^0)\pi^\mp \pi^+ \pi^-$
50 ± 4		² BERTIN	95	OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
75 ± 9		AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
91 ⁺⁶⁷⁺¹⁵ -31-38		³ BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
19 ± 7		³ RATH	89	MPS	$21.4 \pi^-p \rightarrow n K_S^0 K_S^0 \pi^0$

- ¹ Decaying dominantly directly to $K^+K^-\pi^0$.
- ² Decaying into $(K\bar{K})_S\pi$, $(K\pi)_S\bar{K}$, and $a_0(980)\pi$.

³ From fit to the $a_0(980)\pi^0\pi^+\pi^-$ partial wave, but $a_0(980)\pi^+\pi^+\pi^-$ cannot be excluded.



$\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
89 ±17	OUR AVERAGE	Error includes scale factor of 1.7.		
64 ±18	235 ± 91	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
101.0 ± 8.8 ± 8.8	547	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
174 ±44		AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$
90 ±26		¹ COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+\pi^-2\gamma$

¹ This peak in the $\gamma\rho$ channel may not be related to the $\eta(1405)$.

4 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
160 ±30		BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
144 ±13	3270	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

$K\bar{K}\pi$ MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
45.9 ± 8.2	191	^{1,2} ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$
48.9 ± 9.0	249 ± 35	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$
34.2 ± 18.5	62 ± 18	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
93 ± 14	296	¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$

105 ± 10	693	¹ AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
62 ± 16	500	¹ DUCH	89	ASTE	$\bar{p}p \rightarrow K \bar{K} \pi \pi$
100 ± 11	170	¹ RATH	89	MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
66 ± 2	8800	¹ BIRMAN	88	MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
60 ± 10	620	¹ REEVES	86	SPEC	$6.6 p \bar{p} \rightarrow K K \pi X$
60 ± 10		¹ CHUNG	85	SPEC	$8 \pi^- p \rightarrow K \bar{K} \pi n$
55 ⁺²⁰ ₋₃₀	174	¹ EDWARDS	82E	CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
50 ⁺³⁰ ₋₂₀		¹ SCHARRE	80	MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
80 ± 10	800	^{1,3} BAILLON	67	HBC	$0.0 \bar{p}p \rightarrow K \bar{K} \pi \pi$

¹ These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

² Systematic uncertainty not evaluated.

³ From best fit to 0^-+ partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.

$\eta(1405)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K \bar{K} \pi$	seen	
Γ_2 $\eta \pi \pi$	seen	
Γ_3 $a_0(980)\pi$	seen	
Γ_4 $\eta(\pi\pi)_S\text{-wave}$	seen	
Γ_5 $f_0(980)\pi^0 \rightarrow \pi^+ \pi^- \pi^0$	not seen	
Γ_6 $f_0(980)\eta$	seen	
Γ_7 4π	seen	
Γ_8 $\rho\rho$	<58 %	99.85%
Γ_9 $\gamma\gamma$		
Γ_{10} $\rho^0\gamma$	seen	
Γ_{11} $\phi\gamma$		
Γ_{12} $K^*(892)K$	seen	

$\eta(1405)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K \bar{K} \pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_9/\Gamma$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.035	90	^{1,2} AHOHE	05	CLE2	$10.6 e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$
--------	----	----------------------	----	------	--

¹ Using $\eta(1405)$ mass and width 1410 MeV and 51 MeV, respectively.

² Assuming three-body phase-space decay to $K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_9/\Gamma$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

<0.095	95	ACCIARRI	01G	L3	$183\text{--}202 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
--------	----	----------	-----	----	--

$\Gamma(\rho^0\gamma) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{10}\Gamma_9/\Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.5	95	ALTHOFF	84E TASS	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$	

 $\eta(1405)$ BRANCHING RATIOS

$\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$					Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
1.09±0.48		¹ AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.5	90	EDWARDS	83B CBAL	$J/\psi \rightarrow \eta\pi\pi\gamma$	
<1.1	90	SCHARRE	80 MRK2	$J/\psi \rightarrow \eta\pi\pi\gamma$	
<1.5	95	FOSTER	68B HBC	0.0 $\bar{p}p$	

¹ Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$.

$\Gamma(\rho^0\gamma)/\Gamma(\eta\pi\pi)$					Γ_{10}/Γ_2
VALUE		DOCUMENT ID	TECN	COMMENT	
0.111±0.064		AMSLER	04B CBAR	0 $\bar{p}p$	

$\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}\pi)$					Γ_3/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 0.15		¹ BERTIN	95 OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$	
~ 0.8	500	¹ DUCH	89 ASTE	$\bar{p}p \rightarrow \pi^+\pi^-\pi^\pm\pi^\mp K^0$	
~ 0.75		¹ REEVES	86 SPEC	$6.6 \bar{p}p \rightarrow KK\pi X$	

¹ Assuming that the $a_0(980)$ decays only into $K\bar{K}$.

$\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$					Γ_3/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.29±0.10		ABELE	98E CBAR	$0 \bar{p}p \rightarrow \eta\pi^0\pi^0\pi^0$	
0.19±0.04	2200	¹ ALDE	97B GAM4	$100 \pi^-p \rightarrow \eta\pi^0\pi^0n$	
0.56±0.04±0.03		¹ AMSLER	95F CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$	

¹ Assuming that the $a_0(980)$ decays only into $\eta\pi$.

$\Gamma(a_0(980)\pi)/\Gamma(\eta(\pi\pi)_{\text{S-wave}})$					Γ_3/Γ_4
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.91±0.12		ANISOVICH	01 SPEC	$0.0 \bar{p}p \rightarrow \eta\pi^+\pi^-\pi^+\pi^-$	
0.15±0.04	9082	¹ MANAK	00A MPS	$18 \pi^-p \rightarrow \eta\pi^+\pi^-n$	
0.70±0.12±0.20		² BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	

¹ Statistical error only.² Assuming that the $a_0(980)$ decays only into $\eta\pi$.

$\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$ Γ_{10}/Γ_1

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.0152±0.0038	¹ COFFMAN 90	MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
¹ Using $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi) = 4.2 \times 10^{-3}$ and $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma\gamma\rho^0) = 6.4 \times 10^{-5}$.			

 $\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$ Γ_9/Γ_1

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.78 × 10⁻³	90	¹ ABLIKIM 180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma$
¹ Using results from BAI 00D.				

 $\Gamma(\eta(\pi\pi)_{S\text{-wave}})/\Gamma(\eta\pi\pi)$ Γ_4/Γ_2

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.81±0.04	2200	ALDE 97B	GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

 $\Gamma(f_0(980)\eta)/\Gamma(\eta\pi\pi)$ Γ_6/Γ_2

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.32±0.07	¹ ANISOVICH 00	SPEC	0.9–1.2 $\bar{p}p \rightarrow \eta 3\pi^0$
¹ Using preliminary Crystal Barrel data.			

 $\Gamma(f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	¹ ABLIKIM 17AJ	BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$
¹ ABLIKIM 17AJ reports $B(\psi(2S) \rightarrow \gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0) < 5.0 \times 10^{-7}$.			

 $\Gamma(\rho\rho)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.58	99.85	^{1,2} AMSLER 04B	CBAR	0 $\bar{p}p$
¹ Assuming that the $\eta(1405)$ decays are saturated by the $\pi\pi\eta$, $K\bar{K}\pi$ and $\rho\rho$ modes.				
² Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$.				

 $\Gamma(K^*(892)K)/\Gamma(a_0(980)\pi)$ Γ_{12}/Γ_3

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.084±0.024	¹ ADAMS 01B	B852	18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$
¹ Statistical error only.			

 $\Gamma(\phi\gamma)/\Gamma(\rho^0\gamma)$ Γ_{11}/Γ_{10}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.09±0.03		¹ ABLIKIM 18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
0.13±0.04		² ABLIKIM 18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

<0.77 95 ³BAI 04J BES2 $J/\psi \rightarrow \gamma\gamma K^+ K^-$

¹ Constructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18l reports the inverse as 11.10 ± 3.5 .

² Destructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18l reports the inverse as 7.53 ± 2.49 .

³ Calculated by us from $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \phi\gamma\gamma) < 0.82 \times 10^{-4}$ and $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \rho^0\gamma\gamma) = (1.07 \pm 0.17 \pm 0.11) \times 10^{-4}$.

$\eta(1405)$ REFERENCES

ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18l	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)
AMSLER	04B	EPJ C33 23	C. AMSler <i>et al.</i>	(Crystal Barrel Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH	01	NP A690 567	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00	PL B472 168	A.V. Anisovich <i>et al.</i>	
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
CICALO	99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)
ABELE	98E	NP B514 45	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BAI	98C	PL B440 217	J.Z. Bai <i>et al.</i>	(BES Collab.)
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 60 458.		
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
AMSLER	95F	PL B358 389	C. AMSler <i>et al.</i>	(Crystal Barrel Collab.)
BERTIN	95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
DUCH	89	ZPHY C45 223	K.D. Duch <i>et al.</i>	(ASTERIX Collab.) JP
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) IJP
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
	Also	PRL 50 219	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
FOSTER	68B	NP B8 174	M. Foster <i>et al.</i>	(CERN, CDEF)
BAILLON	67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)