


$\eta(1405)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See also the $\eta(1475)$.

$\eta(1405)$ MASS

VALUE (MeV) DOCUMENT ID
1408.8 ± 2.0 OUR AVERAGE Includes data from the 2 datablocks that follow this one.
 Error includes scale factor of 2.2. See the ideogram below.



$\eta\pi\pi$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
 The data in this block is included in the average printed for a previous datablock.

1405.8 ± 2.6 OUR AVERAGE Error includes scale factor of 2.3. See the ideogram below.

| | | | | |
|--|------------|----------------------|----------|---|
| 1409.0 ± 1.7 | 743 | ABLIKIM | 12E BES3 | $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$ |
| 1407.0 ± 3.5 | 198 | ABLIKIM | 12E BES3 | $J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$ |
| 1399.8 ± 2.2 ^{+2.8} _{-0.1} | | ¹ ABLIKIM | 11J BES3 | $J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$ |
| 1392 ± 14 | 900 ± 375 | AMSLER | 04B CBAR | $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$ |
| 1394 ± 8 | 6.6 ± 2.0k | AMSLER | 04B CBAR | $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$ |
| 1404 ± 6 | 9082 | MANAK | 00A MPS | $18 \pi^-p \rightarrow \eta\pi^+\pi^-n$ |


| | | | | |
|---|------|-----------------------|----------|--|
| 1424 ± 6 | 2200 | ALDE | 97B GAM4 | 100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$ |
| 1400 ± 6 | | ² BOLTON | 92B MRK3 | $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$ |
| 1388 ± 4 | | FUKUI | 91C SPEC | 8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$ |
| 1398 ± 6 | 261 | ³ AUGUSTIN | 90 DM2 | $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$ |
| 1420 ± 5 | | ANDO | 86 SPEC | 8 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 1404.0 ± 11.0 | 195 | ABLIKIM | 19BABES3 | $e^+ e^- \rightarrow \psi(2S)$ |
| 1385 ± 7 | | BAI | 99 BES | $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$ |
| 1409 ± 3 | | ⁴ AMSLER | 95F CBAR | 0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta$ |

¹ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.

² From fit to the $a_0(980) \pi 0^- +$ partial wave.

³ Best fit with a single Breit Wigner.

⁴ Superseded by AMSLER 04B.



$K \bar{K} \pi$ MODE ($a_0(980) \pi$ or direct $K \bar{K} \pi$)

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

1413.9 ± 1.7 OUR AVERAGE Error includes scale factor of 1.1.

| | | | | |
|---|------|-----------------------|----------|---|
| 1413 ± 14 | 3651 | ¹ NICHITIU | 02 OBLX | 0 $\bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| 1416 ± 4 ± 2 | 20k | ADAMS | 01B B852 | 18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 1405 ± 5 | | ² CICALO | 99 OBLX | 0 $\bar{p} p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$ |
| 1407 ± 5 | | ² BERTIN | 97 OBLX | 0 $\bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$ |
| 1416 ± 2 | | ² BERTIN | 95 OBLX | 0 $\bar{p} p \rightarrow K \bar{K} \pi \pi \pi$ |
| 1416 ± 8 $\begin{smallmatrix} +7 \\ -5 \end{smallmatrix}$ | 700 | ³ BAI | 90C MRK3 | $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |

1413 ± 5 ³ RATH 89 MPS 21.4 $\pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$

••• We do not use the following data for averages, fits, limits, etc. •••

1459 ± 5 ⁴ AUGUSTIN 92 DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$

¹ Decaying dominantly directly to $K^+ K^- \pi^0$.

² Decaying into $(K \bar{K})_S \pi$, $(K \pi)_S \bar{K}$, and $a_0(980) \pi$.

³ From fit to the $a_0(980) \pi 0^-+$ partial wave. Cannot rule out a $a_0(980) \pi 1^++$ partial wave.

⁴ Excluded from averaging because averaging would be meaningless.

$\pi\pi\gamma$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

1403 \pm 17 OUR AVERAGE Error includes scale factor of 1.8.

| | | | | |
|---------------|--------------|--------|----------|--|
| 1390 \pm 12 | 235 \pm 91 | AMSLER | 04B CBAR | $0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \eta$ |
|---------------|--------------|--------|----------|--|

| | | | | |
|------------------------|-----|-----|----------|--|
| 1424 \pm 10 \pm 11 | 547 | BAI | 04J BES2 | $J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$ |
|------------------------|-----|-----|----------|--|

••• We do not use the following data for averages, fits, limits, etc. •••

| | | | | |
|---------------|--|-------------------------|--------|--|
| 1401 \pm 18 | | ^{1,2} AUGUSTIN | 90 DM2 | $J/\psi \rightarrow \pi^+ \pi^- \gamma \gamma$ |
|---------------|--|-------------------------|--------|--|

| | | | | |
|--------------|--|----------------------|---------|--|
| 1432 \pm 8 | | ² COFFMAN | 90 MRK3 | $J/\psi \rightarrow \pi^+ \pi^- 2\gamma$ |
|--------------|--|----------------------|---------|--|

¹ Best fit with a single Breit Wigner.

² This peak in the $\gamma\rho$ channel may not be related to the $\eta(1405)$.

4π MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

••• We do not use the following data for averages, fits, limits, etc. •••

| | | | | |
|---------------|--|------|---------|---|
| 1420 \pm 20 | | BUGG | 95 MRK3 | $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$ |
|---------------|--|------|---------|---|

| | | | | |
|---------------|------|----------------------|---------|---------------------------------|
| 1489 \pm 12 | 3270 | ¹ BISELLO | 89B DM2 | $J/\psi \rightarrow 4\pi\gamma$ |
|---------------|------|----------------------|---------|---------------------------------|

¹ Estimated by us from various fits.

$K\bar{K}\pi$ MODE (unresolved)

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

••• We do not use the following data for averages, fits, limits, etc. •••

| | | | | |
|------------------|-----|------------------------|----------|---------------------------------------|
| 1452.7 \pm 3.3 | 191 | ^{1,2} ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K K \pi$ |
|------------------|-----|------------------------|----------|---------------------------------------|

| | | | | |
|------------------|--------------|------------------------|----------|--|
| 1437.6 \pm 3.2 | 249 \pm 35 | ^{1,2} ABLIKIM | 08E BES2 | $J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$ |
|------------------|--------------|------------------------|----------|--|

| | | | | |
|------------------|-------------|------------------------|----------|---|
| 1445.9 \pm 5.7 | 62 \pm 18 | ^{1,2} ABLIKIM | 08E BES2 | $J/\psi \rightarrow \omega K^+ K^- \pi^0$ |
|------------------|-------------|------------------------|----------|---|

| | | | | |
|---------------|-----|------------------|---------|---|
| 1442 \pm 10 | 410 | ¹ BAI | 98C BES | $J/\psi \rightarrow \gamma K^+ K^- \pi^0$ |
|---------------|-----|------------------|---------|---|

| | | | | |
|--------------|-----|-----------------------|--------|---|
| 1445 \pm 8 | 693 | ¹ AUGUSTIN | 90 DM2 | $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |
|--------------|-----|-----------------------|--------|---|

| | | | | |
|--------------|-----|-----------------------|--------|---|
| 1433 \pm 8 | 296 | ¹ AUGUSTIN | 90 DM2 | $J/\psi \rightarrow \gamma K^+ K^- \pi^0$ |
|--------------|-----|-----------------------|--------|---|

| | | | | |
|--------------|-----|-------------------|---------|---|
| 1413 \pm 8 | 500 | ¹ DUCH | 89 ASTE | $\bar{p} p \rightarrow \pi^+ \pi^- K^\pm \pi^\mp K^0$ |
|--------------|-----|-------------------|---------|---|

| | | | | |
|--------------|-----|-------------------|--------|--|
| 1453 \pm 7 | 170 | ¹ RATH | 89 MPS | $21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$ |
|--------------|-----|-------------------|--------|--|

| | | | | |
|--------------|------|---------------------|--------|---|
| 1419 \pm 1 | 8800 | ¹ BIRMAN | 88 MPS | $8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$ |
|--------------|------|---------------------|--------|---|

| | | | | |
|--------------|-----|---------------------|---------|---|
| 1424 \pm 3 | 620 | ¹ REEVES | 86 SPEC | $6.6 p \bar{p} \rightarrow K \bar{K} \pi X$ |
|--------------|-----|---------------------|---------|---|

| | | | | |
|--------------|--|--------------------|---------|---|
| 1421 \pm 2 | | ¹ CHUNG | 85 SPEC | $8 \pi^- p \rightarrow K \bar{K} \pi n$ |
|--------------|--|--------------------|---------|---|

| | | | | |
|---|-----|----------------------|----------|---|
| 1440 $\begin{smallmatrix} +20 \\ -15 \end{smallmatrix}$ | 174 | ¹ EDWARDS | 82E CBAL | $J/\psi \rightarrow \gamma K^+ K^- \pi^0$ |
|---|-----|----------------------|----------|---|

| | | | | |
|---|--|----------------------|---------|---|
| 1440 $\begin{smallmatrix} +10 \\ -15 \end{smallmatrix}$ | | ¹ SCHARRE | 80 MRK2 | $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |
|---|--|----------------------|---------|---|

| | | | | |
|--------------|-----|------------------------|--------|---|
| 1425 \pm 7 | 800 | ^{1,3} BAILLON | 67 HBC | $0 \bar{p} p \rightarrow K \bar{K} \pi \pi \pi$ |
|--------------|-----|------------------------|--------|---|

¹ These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

² Systematic uncertainty not evaluated.

³ From best fit of 0^-+ partial wave, 50% $K^*(892) K$, 50% $a_0(980) \pi$.


$\eta(1405)$ WIDTH

VALUE (MeV)

DOCUMENT ID

50.1 ± 2.6 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.7. See the ideogram below.

WEIGHTED AVERAGE
50.1 ± 2.6 (Error scaled by 1.7)



$\eta\pi\pi$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

52.6 ± 3.2 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

| | | | | | |
|--|------|-----------------------|-----|------|---|
| 48.3 ± 5.2 | 743 | ABLIKIM | 12E | BES3 | $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$ |
| 55.0 ± 11.0 | 198 | ABLIKIM | 12E | BES3 | $J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$ |
| 52.8 ± 7.6 ^{+0.1} _{-7.6} | | ¹ ABLIKIM | 11J | BES3 | $J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$ |
| 55 ± 11 | 900 | AMSLER | 04B | CBAR | $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$ |
| 55 ± 12 | 6.6k | AMSLER | 04B | CBAR | $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma$ |
| 80 ± 21 | 9.0k | MANAK | 00A | MPS | $18 \pi^-p \rightarrow \eta\pi^+\pi^-n$ |
| 85 ± 18 | 2.2k | ALDE | 97B | GAM4 | $100 \pi^-p \rightarrow \eta\pi^0\pi^0n$ |
| 47 ± 13 | | ² BOLTON | 92B | MRK3 | $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$ |
| 59 ± 4 | | FUKUI | 91C | SPEC | $8.95 \pi^-p \rightarrow \eta\pi^+\pi^-n$ |
| 53 ± 11 | | ³ AUGUSTIN | 90 | DM2 | $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$ |
| 31 ± 7 | | ANDO | 86 | SPEC | $8 \pi^-p \rightarrow \eta\pi^+\pi^-n$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

79.0 ± 16.0 195 ABLIKIM 19BA BES3 $e^+e^- \rightarrow \psi(2S)$
 86 ± 10 4 AMSLER 95F CBAR $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$


¹ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

² From fit to the $a_0(980)\pi 0^-+$ partial wave.

³ From $\eta\pi^+\pi^-$ mass distribution - mainly $a_0(980)\pi$ - no spin-parity determination available.

⁴ Superseded by AMSLER 04B.

WEIGHTED AVERAGE
 52.6 ± 3.2 (Error scaled by 1.3)



$\eta(1405)$ width $\eta\pi\pi$ mode (MeV)

$K\bar{K}\pi$ MODE ($a_0(980)\pi$ or direct $K\bar{K}\pi$)

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.


48 ± 4 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.

| | | | | | |
|--|------|-----------------------|-----|------|--|
| 51 ± 6 | 3651 | ¹ NICHITIU | 02 | OBLX | $0 \bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0$ |
| 42 ± 10 ± 9 | 20k | ADAMS | 01B | B852 | $18 \text{ GeV } \pi^-p \rightarrow K^+K^-\pi^0n$ |
| 50 ± 4 | | CICALO | 99 | OBLX | $0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$ |
| 48 ± 5 | | ² BERTIN | 97 | OBLX | $0.0 \bar{p}p \rightarrow K^\pm(K^0)\pi^\mp \pi^+ \pi^-$ |
| 50 ± 4 | | ² BERTIN | 95 | OBLX | $0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$ |
| 75 ± 9 | | AUGUSTIN | 92 | DM2 | $J/\psi \rightarrow \gamma K\bar{K}\pi$ |
| 91 ⁺ 67 ⁺ 15 -31-38 | | ³ BAI | 90C | MRK3 | $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |
| 19 ± 7 | | ³ RATH | 89 | MPS | $21.4 \pi^-p \rightarrow n K_S^0 K_S^0 \pi^0$ |

¹ Decaying dominantly directly to $K^+K^-\pi^0$.

² Decaying into $(K\bar{K})_S\pi$, $(K\pi)_S\bar{K}$, and $a_0(980)\pi$.

³ From fit to the $a_0(980)\pi^0\pi^+\pi^-$ partial wave, but $a_0(980)\pi^+\pi^+\pi^-$ cannot be excluded.



$\pi\pi\gamma$ MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|--------------------|-------------------------------------|----------|---|
| 89 ±17 | OUR AVERAGE | Error includes scale factor of 1.7. | | |
| 64 ±18 | 235 ± 91 | AMSLER | 04B CBAR | $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$ |
| 101.0 ± 8.8 ± 8.8 | 547 | BAI | 04J BES2 | $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 174 ±44 | | AUGUSTIN | 90 DM2 | $J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$ |
| 90 ±26 | | ¹ COFFMAN | 90 MRK3 | $J/\psi \rightarrow \pi^+\pi^-2\gamma$ |

¹ This peak in the $\gamma\rho$ channel may not be related to the $\eta(1405)$.

4 π MODE

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|----------------------|---------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 160 ±30 | | BUGG | 95 MRK3 | $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$ |
| 144 ±13 | 3270 | ¹ BISELLO | 89B DM2 | $J/\psi \rightarrow 4\pi\gamma$ |

¹ Estimated by us from various fits.

$K\bar{K}\pi$ MODE (unresolved)

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|------------------------|----------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 45.9 ± 8.2 | 191 | ^{1,2} ABLIKIM | 13M BES3 | $\psi(2S) \rightarrow \omega K K \pi$ |
| 48.9 ± 9.0 | 249 ± 35 | ^{1,2} ABLIKIM | 08E BES2 | $J/\psi \rightarrow \omega K_S^0 K^+\pi^- + c.c.$ |
| 34.2 ± 18.5 | 62 ± 18 | ^{1,2} ABLIKIM | 08E BES2 | $J/\psi \rightarrow \omega K^+ K^- \pi^0$ |
| 93 ± 14 | 296 | ¹ AUGUSTIN | 90 DM2 | $J/\psi \rightarrow \gamma K^+ K^- \pi^0$ |

| | | | | | |
|----------------------------------|------|------------------------|-----|------|---|
| 105 ± 10 | 693 | ¹ AUGUSTIN | 90 | DM2 | $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |
| 62 ± 16 | 500 | ¹ DUCH | 89 | ASTE | $\bar{p}p \rightarrow K \bar{K} \pi \pi$ |
| 100 ± 11 | 170 | ¹ RATH | 89 | MPS | $21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$ |
| 66 ± 2 | 8800 | ¹ BIRMAN | 88 | MPS | $8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$ |
| 60 ± 10 | 620 | ¹ REEVES | 86 | SPEC | $6.6 p \bar{p} \rightarrow K K \pi X$ |
| 60 ± 10 | | ¹ CHUNG | 85 | SPEC | $8 \pi^- p \rightarrow K \bar{K} \pi n$ |
| 55 ⁺²⁰ ₋₃₀ | 174 | ¹ EDWARDS | 82E | CBAL | $J/\psi \rightarrow \gamma K^+ K^- \pi^0$ |
| 50 ⁺³⁰ ₋₂₀ | | ¹ SCHARRE | 80 | MRK2 | $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ |
| 80 ± 10 | 800 | ^{1,3} BAILLON | 67 | HBC | $0.0 \bar{p}p \rightarrow K \bar{K} \pi \pi$ |

¹ These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

² Systematic uncertainty not evaluated.

³ From best fit to 0^-+ partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.

$\eta(1405)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|--|--------------------------------|------------------|
| Γ_1 $K \bar{K} \pi$ | seen | |
| Γ_2 $\eta \pi \pi$ | seen | |
| Γ_3 $a_0(980)\pi$ | seen | |
| Γ_4 $\eta(\pi\pi)_S\text{-wave}$ | seen | |
| Γ_5 $f_0(980)\pi^0 \rightarrow \pi^+ \pi^- \pi^0$ | not seen | |
| Γ_6 $f_0(980)\eta$ | seen | |
| Γ_7 4π | seen | |
| Γ_8 $\rho\rho$ | <58 % | 99.85% |
| Γ_9 $\gamma\gamma$ | | |
| Γ_{10} $\rho^0\gamma$ | seen | |
| Γ_{11} $\phi\gamma$ | | |
| Γ_{12} $K^*(892)K$ | seen | |

$\eta(1405) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K \bar{K} \pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1 \Gamma_9 / \Gamma$

| VALUE (keV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------|-----|-------------|------|---------|
|-------------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------|----|----------------------|----|------|--|
| <0.035 | 90 | ^{1,2} AHOHE | 05 | CLE2 | $10.6 e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$ |
|--------|----|----------------------|----|------|--|

¹ Using $\eta(1405)$ mass and width 1410 MeV and 51 MeV, respectively.

² Assuming three-body phase-space decay to $K_S^0 K^\pm \pi^\mp$.

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2 \Gamma_9 / \Gamma$

| VALUE (keV) | CL% | DOCUMENT ID | TECN | COMMENT |
|-------------|-----|-------------|------|---------|
|-------------|-----|-------------|------|---------|

| | | | | | |
|--------|----|----------|-----|----|--|
| <0.095 | 95 | ACCIARRI | 01G | L3 | $183\text{--}202 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$ |
|--------|----|----------|-----|----|--|

| $\Gamma(\rho^0\gamma) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{10}\Gamma_9/\Gamma$ |
|---|-----|-------------|------|---------|---|
| VALUE (keV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <1.5 | 95 | ALTHOFF | 84E | TASS | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$ |

 $\eta(1405)$ BRANCHING RATIOS

| $\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$ | | | | | Γ_2/Γ_1 |
|---|-----|---------------------|------|---------|---|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT | |
| 1.09±0.48 | | ¹ AMSLER | 04B | CBAR | $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <0.5 | 90 | EDWARDS | 83B | CBAL | $J/\psi \rightarrow \eta\pi\pi\gamma$ |
| <1.1 | 90 | SCHARRE | 80 | MRK2 | $J/\psi \rightarrow \eta\pi\pi\gamma$ |
| <1.5 | 95 | FOSTER | 68B | HBC | 0.0 $\bar{p}p$ |

¹ Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$.

| $\Gamma(\rho^0\gamma)/\Gamma(\eta\pi\pi)$ | | | | | Γ_{10}/Γ_2 |
|---|--|-------------|------|---------|------------------------|
| VALUE | | DOCUMENT ID | TECN | COMMENT | |
| 0.111±0.064 | | AMSLER | 04B | CBAR | 0 $\bar{p}p$ |

| $\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}\pi)$ | | | | | Γ_3/Γ_1 |
|---|------|---------------------|------|---------|---|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| ~ 0.15 | | ¹ BERTIN | 95 | OBLX | $0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$ |
| ~ 0.8 | 500 | ¹ DUCH | 89 | ASTE | $\bar{p}p \rightarrow \pi^+\pi^-\pi^\pm\pi^\mp K^0$ |
| ~ 0.75 | | ¹ REEVES | 86 | SPEC | $6.6 p\bar{p} \rightarrow KK\pi X$ |

¹ Assuming that the $a_0(980)$ decays only into $K\bar{K}$.

| $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$ | | | | | Γ_3/Γ_2 |
|---|------|---------------------|------|---------|---|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 0.29±0.10 | | ABELE | 98E | CBAR | $0 p\bar{p} \rightarrow \eta\pi^0\pi^0\pi^0$ |
| 0.19±0.04 | 2200 | ¹ ALDE | 97B | GAM4 | $100 \pi^-p \rightarrow \eta\pi^0\pi^0n$ |
| 0.56±0.04±0.03 | | ¹ AMSLER | 95F | CBAR | $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$ |

¹ Assuming that the $a_0(980)$ decays only into $\eta\pi$.

| $\Gamma(a_0(980)\pi)/\Gamma(\eta(\pi\pi)_{\text{S-wave}})$ | | | | | Γ_3/Γ_4 |
|---|------|--------------------|------|---------|---|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| 0.91±0.12 | | ANISOVICH | 01 | SPEC | $0.0 \bar{p}p \rightarrow \eta\pi^+\pi^-\pi^+\pi^-$ |
| 0.15±0.04 | 9082 | ¹ MANAK | 00A | MPS | $18 \pi^-p \rightarrow \eta\pi^+\pi^-n$ |
| 0.70±0.12±0.20 | | ² BAI | 99 | BES | $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$ |

¹ Statistical error only.² Assuming that the $a_0(980)$ decays only into $\eta\pi$.

$\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$ Γ_{10}/Γ_1

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|----------------------|------|--|
| 0.0152±0.0038 | | ¹ COFFMAN | 90 | MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |
| ¹ Using $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi)=4.2 \times 10^{-3}$ and $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma\gamma\rho^0)=6.4 \times 10^{-5}$. | | | | |

 $\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$ Γ_9/Γ_1

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|----------------------|------|--|
| <1.78 × 10⁻³ | 90 | ¹ ABLIKIM | 180 | BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma$ |
| ¹ Using results from BAI 00D. | | | | |

 $\Gamma(\eta(\pi\pi)_{S\text{-wave}})/\Gamma(\eta\pi\pi)$ Γ_4/Γ_2

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.81±0.04 | 2200 | ALDE | 97B | GAM4 100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$ |

 $\Gamma(f_0(980)\eta)/\Gamma(\eta\pi\pi)$ Γ_6/Γ_2

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.32±0.07 | ¹ ANISOVICH | 00 | SPEC 0.9–1.2 $\bar{p}p \rightarrow \eta 3\pi^0$ |
| ¹ Using preliminary Crystal Barrel data. | | | |

 $\Gamma(f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--|----------------------|------|---|
| not seen | ¹ ABLIKIM | 17AJ | BES3 $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$ |
| ¹ ABLIKIM 17AJ reports $B(\psi(2S) \rightarrow \gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0) < 5.0 \times 10^{-7}$. | | | |

 $\Gamma(\rho\rho)/\Gamma_{\text{total}}$ Γ_8/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-------|-----------------------|------|-------------------|
| <0.58 | 99.85 | ^{1,2} AMSLER | 04B | CBAR 0 $\bar{p}p$ |
| ¹ Assuming that the $\eta(1405)$ decays are saturated by the $\pi\pi\eta$, $K\bar{K}\pi$ and $\rho\rho$ modes. | | | | |
| ² Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$. | | | | |

 $\Gamma(K^*(892)K)/\Gamma(a_0(980)\pi)$ Γ_{12}/Γ_3

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|--------------------|------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 0.084±0.024 | ¹ ADAMS | 01B | B852 18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$ |
| ¹ Statistical error only. | | | |

 $\Gamma(\phi\gamma)/\Gamma(\rho^0\gamma)$ Γ_{11}/Γ_{10}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------|------|--|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 0.09±0.03 | | ¹ ABLIKIM | 181 | BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$ |
| 0.13±0.04 | | ² ABLIKIM | 181 | BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$ |

<0.77 95 ^3BAI 04J BES2 $J/\psi \rightarrow \gamma\gamma K^+ K^-$ ¹ Constructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18l reports the inverse as 11.10 ± 3.5 .² Destructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18l reports the inverse as 7.53 ± 2.49 .³ Calculated by us from $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \phi\gamma\gamma) < 0.82 \times 10^{-4}$ and $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \rho^0\gamma\gamma) = (1.07 \pm 0.17 \pm 0.11) \times 10^{-4}$.

$\eta(1405)$ REFERENCES

| | | | | |
|-----------|------|-----------------------------|------------------------------|------------------------------|
| ABLIKIM | 19BA | PR D100 092003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18l | PR D97 051101 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 18O | PR D97 072014 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 17AJ | PR D96 112008 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13M | PR D87 092006 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12E | PRL 108 182001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 11J | PRL 107 182001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 08E | PR D77 032005 | M. Ablikim <i>et al.</i> | (BES Collab.) |
| AHOHE | 05 | PR D71 072001 | R. Ahohe <i>et al.</i> | (CLEO Collab.) |
| AMSLER | 04B | EPJ C33 23 | C. AMSler <i>et al.</i> | (Crystal Barrel Collab.) |
| BAI | 04J | PL B594 47 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| NICHITIU | 02 | PL B545 261 | F. Nichitiu <i>et al.</i> | (OBELIX Collab.) |
| ACCIARRI | 01G | PL B501 1 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ADAMS | 01B | PL B516 264 | G.S. Adams <i>et al.</i> | (BNL E852 Collab.) |
| ANISOVICH | 01 | NP A690 567 | A.V. Anisovich <i>et al.</i> | |
| ANISOVICH | 00 | PL B472 168 | A.V. Anisovich <i>et al.</i> | |
| BAI | 00D | PL B476 25 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| MANAK | 00A | PR D62 012003 | J.J. Manak <i>et al.</i> | (BNL E852 Collab.) |
| BAI | 99 | PL B446 356 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| CICALO | 99 | PL B462 453 | C. Cicalo <i>et al.</i> | (OBELIX Collab.) |
| ABELE | 98E | NP B514 45 | A. Abele <i>et al.</i> | (Crystal Barrel Collab.) |
| BAI | 98C | PL B440 217 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ALDE | 97B | PAN 60 386 | D. Alde <i>et al.</i> | (GAMS Collab.) |
| | | Translated from YAF 60 458. | | |
| BERTIN | 97 | PL B400 226 | A. Bertin <i>et al.</i> | (OBELIX Collab.) |
| AMSLER | 95F | PL B358 389 | C. AMSler <i>et al.</i> | (Crystal Barrel Collab.) |
| BERTIN | 95 | PL B361 187 | A. Bertin <i>et al.</i> | (OBELIX Collab.) |
| BUGG | 95 | PL B353 378 | D.V. Bugg <i>et al.</i> | (LOQM, PNPI, WASH) |
| AUGUSTIN | 92 | PR D46 1951 | J.E. Augustin, G. Cosme | (DM2 Collab.) |
| BOLTON | 92B | PRL 69 1328 | T. Bolton <i>et al.</i> | (Mark III Collab.) |
| FUKUI | 91C | PL B267 293 | S. Fukui <i>et al.</i> | (SUGI, NAGO, KEK, KYOT+) |
| AUGUSTIN | 90 | PR D42 10 | J.E. Augustin <i>et al.</i> | (DM2 Collab.) |
| BAI | 90C | PRL 65 2507 | Z. Bai <i>et al.</i> | (Mark III Collab.) |
| COFFMAN | 90 | PR D41 1410 | D.M. Coffman <i>et al.</i> | (Mark III Collab.) |
| BISELLO | 89B | PR D39 701 | G. Busetto <i>et al.</i> | (DM2 Collab.) |
| DUCH | 89 | ZPHY C45 223 | K.D. Duch <i>et al.</i> | (ASTERIX Collab.) JP |
| RATH | 89 | PR D40 693 | M.G. Rath <i>et al.</i> | (NDAM, BRAN, BNL, CUNY+) |
| BIRMAN | 88 | PRL 61 1557 | A. Birman <i>et al.</i> | (BNL, FSU, IND, MASD) JP |
| ANDO | 86 | PRL 57 1296 | A. Ando <i>et al.</i> | (KEK, KYOT, NIRS, SAGA+) IJP |
| REEVES | 86 | PR D34 1960 | D.F. Reeves <i>et al.</i> | (FLOR, BNL, IND+) JP |
| CHUNG | 85 | PRL 55 779 | S.U. Chung <i>et al.</i> | (BNL, FLOR, IND+) JP |
| ALTHOFF | 84E | PL 147B 487 | M. Althoff <i>et al.</i> | (TASSO Collab.) |
| EDWARDS | 83B | PRL 51 859 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |
| EDWARDS | 82E | PRL 49 259 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |
| | Also | PRL 50 219 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |
| SCHARRE | 80 | PL 97B 329 | D.L. Scharre <i>et al.</i> | (SLAC, LBL) |
| FOSTER | 68B | NP B8 174 | M. Foster <i>et al.</i> | (CERN, CDEF) |
| BAILLON | 67 | NC 50A 393 | P.H. Baillon <i>et al.</i> | (CERN, CDEF, IRAD) |