

$\eta_c(2S)$ $I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers are quark model predictions.

 $\eta_c(2S)$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|----------|-------------------------------------|----------|---|
| 3637.5±1.1 OUR AVERAGE | | Error includes scale factor of 1.2. | | |
| 3635.1±3.7±2.9 | 106 | XU | 18 BELL | $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$ |
| 3633.6±1.7±0.6 | 106 | ¹ AAIJ | 17ADLHCb | $p p \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$ |
| 3636.4±4.1±0.7 | 365 | ² AAIJ | 17BBLHCb | $p p \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-)X$ |
| 3637.0±5.7±3.4 | 178 | ^{3,4} LEES | 14E BABR | $\gamma\gamma \rightarrow K^+ K^- \pi^0$ |
| 3635.1±5.8±2.1 | 47 | ^{3,5} LEES | 14E BABR | $\gamma\gamma \rightarrow K^+ K^- \eta$ |
| 3646.9±1.6±3.6 | 57 ± 17 | ABLIKIM | 13K BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ |
| 3637.6±2.9±1.6 | 127 ± 18 | ⁶ ABLIKIM | 12G BES3 | $\psi(2S) \rightarrow \gamma K^0 K\pi, K K\pi^0$ |
| 3638.5±1.5±0.8 | 624 | ³ DEL-AMO-SA..11M | BABR | $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$ |
| 3640.5±3.2±2.5 | 1201 | ³ DEL-AMO-SA..11M | BABR | $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |
| 3636.1 ^{+3.9 +0.7} _{-4.2 -2.0} | 128 | ⁷ VINOKUROVA 11 | BELL | $B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$ |
| 3626 ± 5 ± 6 | 311 | ⁸ ABE | 07 BELL | $e^+ e^- \rightarrow J/\psi(c\bar{c})$ |
| 3645.0±5.5 ^{+4.9} _{-7.8} | 121 ± 27 | AUBERT | 05C BABR | $e^+ e^- \rightarrow J/\psi c\bar{c}$ |
| 3642.9±3.1±1.5 | 61 | ASNER | 04 CLEO | $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 3639 ± 7 | 98 ± 52 | ⁹ AUBERT | 06E BABR | $B^\pm \rightarrow K^\pm X c\bar{c}$ |
| 3630.8±3.4±1.0 | 112 ± 24 | ¹⁰ AUBERT | 04D BABR | $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$ |
| 3654 ± 6 ± 8 | 39 ± 11 | ¹¹ CHOI | 02 BELL | $B \rightarrow K K_S K^- \pi^+$ |
| 3594 ± 5 | | ¹² EDWARDS | 82C CBAL | $e^+ e^- \rightarrow \gamma X$ |

¹AAIJ 17AD report $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$ MeV. We use the current value $m_{\psi(2S)} = 3686.097 \pm 0.025$ MeV to obtain the quoted mass.

²From a fit of the $\phi\phi$ invariant mass with the width of $\eta_c(2S)$ fixed to the PDG 16 value.

³Ignoring possible interference with continuum.

⁴With a width fixed to 11.3 MeV.

⁵With a width fixed to 11.3 MeV. Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

⁷Accounts for interference with non-resonant continuum.

⁸From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

⁹From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁰Superseded by DEL-AMO-SANCHEZ 11M.

¹¹Superseded by VINOKUROVA 11.

¹²Assuming mass of $\psi(2S) = 3686$ MeV.

$\eta_c(2S)$ WIDTH

| VALUE (MeV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|--------------|------------------------------------|--|---|
| 11.3^{+ 3.2}_{- 2.9} OUR AVERAGE | | | | | |
| 9.9 \pm 4.8 \pm 2.9 | | 57 \pm 17 | ABLIKIM | 13K BES3 | $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ |
| 16.9 \pm 6.4 \pm 4.8 | | 127 \pm 18 | ¹³ ABLIKIM | 12G BES3 | $\psi(2S) \rightarrow \gamma K^0 K\pi,$ $K K\pi^0$ |
| 13.4 \pm 4.6 \pm 3.2 | | 624 | ¹⁴ DEL-AMO-SA..11M BABR | $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$ | |
| 6.6 ^{+ 8.4 + 2.6} _{- 5.1 - 0.9} | | 128 | ¹⁵ VINOKUROVA 11 | BELL | $B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$ |
| 6.3 \pm 12.4 \pm 4.0 | | 61 | ASNER | 04 CLEO | $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| < 23 | 90 | 98 \pm 52 | ¹⁶ AUBERT | 06E BABR | $B^\pm \rightarrow K^\pm X_{c\bar{c}}$ |
| 22 \pm 14 | | 121 \pm 27 | AUBERT | 05C BABR | $e^+ e^- \rightarrow J/\psi c\bar{c}$ |
| 17.0 \pm 8.3 \pm 2.5 | | 112 \pm 24 | ¹⁷ AUBERT | 04D BABR | $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$ |
| < 55 | 90 | 39 \pm 11 | ¹⁸ CHOI | 02 BELL | $B \rightarrow K K_S K^- \pi^+$ |
| < 8.0 | 95 | | ¹⁹ EDWARDS | 82C CBAL | $e^+ e^- \rightarrow \gamma X$ |
| ¹³ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes. | | | | | |
| ¹⁴ Ignoring possible interference with continuum. | | | | | |
| ¹⁵ Accounts for interference with non-resonant continuum. | | | | | |
| ¹⁶ From the fit of the kaon momentum spectrum. Systematic errors not evaluated. | | | | | |
| ¹⁷ Superseded by DEL-AMO-SANCHEZ 11M. | | | | | |
| ¹⁸ For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11. | | | | | |
| ¹⁹ For a mass value of 3594 ± 5 MeV | | | | | |

$\eta_c(2S)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Confidence level |
|--|--------------------------------|------------------|
| Γ_1 hadrons | | not seen |
| Γ_2 $K\bar{K}\pi$ | (1.9 \pm 1.2) % | |
| Γ_3 $K\bar{K}\eta$ | (5 \pm 4) $\times 10^{-3}$ | |
| Γ_4 $2\pi^+ 2\pi^-$ | not seen | |
| Γ_5 $\rho^0 \rho^0$ | not seen | |
| Γ_6 $3\pi^+ 3\pi^-$ | not seen | |
| Γ_7 $K^+ K^- \pi^+ \pi^-$ | not seen | |
| Γ_8 $K^{*0} \bar{K}^{*0}$ | not seen | |
| Γ_9 $K^+ K^- \pi^+ \pi^- \pi^0$ | (1.4 \pm 1.0) % | |
| Γ_{10} $K^+ K^- 2\pi^+ 2\pi^-$ | not seen | |
| Γ_{11} $K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$ | seen | |
| Γ_{12} $2K^+ 2K^-$ | not seen | |
| Γ_{13} $\phi\phi$ | not seen | |

| | | | | |
|---------------|------------------------|--------------------------------|-----|--|
| Γ_{14} | $p\bar{p}$ | seen | | |
| Γ_{15} | $p\bar{p}\pi^+\pi^-$ | seen | | |
| Γ_{16} | $\gamma\gamma$ | $(1.9 \pm 1.3) \times 10^{-4}$ | | |
| Γ_{17} | $\gamma J/\psi(1S)$ | $< 1.4 \quad \%$ | 90% | |
| Γ_{18} | $\pi^+\pi^-\eta$ | not seen | | |
| Γ_{19} | $\pi^+\pi^-\eta'$ | not seen | | |
| Γ_{20} | $\pi^+\pi^-\eta_c(1S)$ | $< 25 \quad \%$ | 90% | |

 $\eta_c(2S)$ PARTIAL WIDTHS **$\Gamma(\gamma\gamma)$** **Γ_{16}**

| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|---------|---|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.44 ± 0.14 | 106 | 20 XU | 18 BELL | $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$ |
| 1.3 ± 0.6 | | 21 ASNER | 04 CLEO | $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ |

20 Assuming that the branching fraction into $\eta'\pi^+\pi^-$ is the same as for $\eta_c(1S)$.21 They measure $\Gamma(\eta_c(2S)\gamma\gamma)$ $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma)$ $B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV. **$\Gamma(\gamma\gamma) \times \Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$** **$\Gamma_{16}\Gamma_{19}/\Gamma$**

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|------|-------------|---------|--|
| $5.6^{+1.2}_{-1.1} \pm 1.1$ | 106 | XU | 18 BELL | $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$ |

 $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$** **$\Gamma_4\Gamma_{16}/\Gamma$**

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------|-----|-------------|---------|---|
| < 6.5 | 90 | UEHARA | 08 BELL | $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$ |

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_2\Gamma_{16}/\Gamma$**

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------|------|--------------------|------|--|
| $41^{+4}_{-6} \pm 6$ | 624 | 22 DEL-AMO-SA..11M | BABR | $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$ |

22 Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_7\Gamma_{16}/\Gamma$**

| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT |
|------------|-----|-------------|---------|--|
| < 5.0 | 90 | UEHARA | 08 BELL | $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$ |

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_9\Gamma_{16}/\Gamma$**

| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------|------|--------------------|------|--|
| $30^{+6}_{-5} \pm 5$ | 1201 | 23 DEL-AMO-SA..11M | BABR | $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$ |

23 Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

| $\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{12}\Gamma_{16}/\Gamma$ |
|--|-----|-------------|-----------|--|--|
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <2.9 | 90 | UEHARA | 08 | BELL | $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+ K^-)$ |
| $\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{20}\Gamma_{16}/\Gamma$ |
| VALUE (eV) | CL% | DOCUMENT ID | TECN | COMMENT | |
| <133 | 90 | LEES | 12AE BABR | $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$ | |

 $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

| $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$ |
|---|-----|---------------|------|-------------------------------------|--|
| VALUE (units 10^{-8}) | CL% | DOCUMENT ID | TECN | COMMENT | |
| < 5.6 | 90 | AMBROGIANI 01 | E835 | $\bar{p}p \rightarrow \gamma\gamma$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| < 8.0 | 90 | AMBROGIANI 01 | E835 | $\bar{p}p \rightarrow \gamma\gamma$ | |
| <12.0 | 90 | AMBROGIANI 01 | E835 | $\bar{p}p \rightarrow \gamma\gamma$ | |
| ²⁴ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis. | | | | | |
| ²⁵ For a total width $\Gamma=5$ MeV. | | | | | |
| ²⁶ For the resonance mass region 3589–3599 MeV/ c^2 . | | | | | |
| ²⁷ For the resonance mass region 3575–3660 MeV/ c^2 . | | | | | |

 $\eta_c(2S)$ BRANCHING RATIOS

| $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ | | | | | Γ_1/Γ |
|---|-------------|----------|--|--|-------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | | |
| not seen | ABREU | 980 DLPH | $e^+ e^- \rightarrow e^+ e^- + \text{hadrons}$ | | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| seen | 28 EDWARDS | 82C CBAL | $e^+ e^- \rightarrow \gamma X$ | | |
| ²⁸ For a mass value of 3594 \pm 5 MeV | | | | | |

| $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ | | | | | Γ_2/Γ |
|--|--------------|-------------|-----------|--|-------------------|
| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 1.9\pm0.4\pm1.1 | 59 ± 12 | 29 AUBERT | 08AB BABR | $B \rightarrow \eta_c(2S) K \rightarrow K\bar{K}\pi K$ | |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | | |
| seen | 127 \pm 18 | ABLIKIM | 13K BES3 | $\psi(2S) \rightarrow \gamma K\bar{K}\pi$ | |
| seen | 39 \pm 11 | 30 CHOI | 02 BELL | $B \rightarrow K K_S K^- \pi^+$ | |
| ²⁹ Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S) K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S) K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$. | | | | | |
| ³⁰ For a mass value of 3654 \pm 6 MeV | | | | | |

| $\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$ | | | | | Γ_3/Γ_2 |
|--|------|-------------|----------|---|---------------------|
| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 27.3\pm7.0\pm9.0 | 225 | 31 LEES | 14E BABR | $\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$ | |
| ³¹ LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$, which we divide by 3 to account for isospin symmetry. | | | | | |

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|--|
| not seen | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \eta_c(2S)$ |

 Γ_4/Γ $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|--|
| not seen | ABLIKIM | 11H | BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$ |

 Γ_5/Γ $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|--|
| not seen | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \eta_c(2S)$ |

 Γ_7/Γ $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---------------------------|-------------|--------------------|-------------|--|
| 0.73 ± 0.17 ± 0.17 | 1201 | 32 DEL-AMO-SA..11M | BABR | $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$ |

³² We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

 Γ_9/Γ_2 $\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|--|
| not seen | ABLIKIM | 11H | BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$ |

 Γ_8/Γ $\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|--|
| seen | 57 ± 17 | ABLIKIM | 13K | BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$ |

 Γ_{11}/Γ $\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|--|
| not seen | UEHARA | 08 | BELL $\gamma\gamma \rightarrow \eta_c(2S)$ |

 Γ_{12}/Γ $\Gamma(\phi\phi)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|--------------------|-------------|--|
| not seen | ABLIKIM | 11H | BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$ |

 Γ_{13}/Γ $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|---|
| seen | 106 | 33 AAIJ | 17AD LHCb | $p\bar{p} \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$ |

 Γ_{14}/Γ

³³ AAIJ 17AD report a 6.4 standard deviation signal, with $B(B^+ \rightarrow \eta_c(2S) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$.

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|-------------|--------------------|-------------|---------------------------------------|
| seen | 110 | 34 CHILIKIN | 19 | BELL $e^+ e^- \rightarrow \gamma(4S)$ |

 Γ_{15}/Γ

³⁴ CHILIKIN 19 reports signals in $B^+ \rightarrow \eta_c(2S) K^+$ and $B^0 \rightarrow \eta_c(2S) K_S^0$ with 12.3 and 5.9 standard deviations, respectively.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{16}/Γ

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--|------------|--------------------|-------------|---|
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ | | | | |
| $<4 \times 10^{-4}$ | 90 | 35 WICHT | 08 | BELL $B^\pm \rightarrow K^\pm \gamma\gamma$ |
| not seen | | AMBROGIANI | 01 | E835 $\bar{p}p \rightarrow \gamma\gamma$ |
| <0.01 | 90 | LEE | 85 | CBAL $\psi' \rightarrow \text{photons}$ |
| 35 WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c(2S)K^+) = 4.4 \times 10^{-4}$. | | | | |

 $\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma(K\bar{K}\pi)$ Γ_{20}/Γ_2

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|-----------------|------------|--------------------|-------------|---|
| <3.33 | 90 | 36 LEES | 12AE BABR | $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$ |

36 We divided the reported limit by 3 to take into account isospin relations.

 $\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS $\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ $<11.8 \times 10^{-6}$ 90 37 CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma K^+ K^- \eta$ 37 CRONIN-HENNESSY 10 reports a limit of $< 5.9 \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 account for isospin symmetry. It assumes $\Gamma(\eta_c(2S)) = 14$ MeV. It also gives the analytic dependence of limits on width. $\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

<14.6 × 10⁻⁶ 90 38 CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$ 38 Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width. $\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

<12.7 × 10⁻⁷ 90 ABLIKIM 11H BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$ $\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$

| <u>VALUE</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|------------|--------------------|-------------|----------------|
|--------------|------------|--------------------|-------------|----------------|

<13.2 × 10⁻⁶ 90 39 CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$ 39 Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_7/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------------|------|---|
| $<9.6 \times 10^{-6}$ | 90 | 40 CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$ |

⁴⁰ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_8/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------|----------|---|
| $<19.6 \times 10^{-7}$ | 90 | ABLIKIM | 11H BES3 | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$ |

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_9/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------------|------|---|
| $<43.0 \times 10^{-6}$ | 90 | 41 CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$ |

⁴¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_{10}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------------|------|---|
| $<9.7 \times 10^{-6}$ | 90 | 42 CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$ |

⁴² Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_{11}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| VALUE (units 10^{-6}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|---------|-------------|------|--|
| $7.03 \pm 2.10 \pm 0.7$ | 60 | ABLIKIM | 13K BES3 | CLEO | $\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------|----|-------------------|------|--|
| < 15.2 | 90 | 43 CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$ |
|----------|----|-------------------|------|--|

⁴³ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\frac{\Gamma(\eta_c(2S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}}{\Gamma_{13}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|----------|---|
| $<7.8 \times 10^{-7}$ | 90 | ABLIKIM | 11H BES3 | $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$ |

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{14}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------|------|---|
| $<1.4 \times 10^{-6}$ | 90 | ABLIKIM | 13V | BES3 $\psi(2S) \rightarrow \gamma p\bar{p}$ |

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{17}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|------|-------------|------|---|
| $<9.7 \times 10^{-6}$ | 90 | 33 | 44 ABLIKIM | 17N | BES3 $\psi(2S) \rightarrow \gamma\gamma J/\psi$ |

⁴⁴ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{18}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------------|------|---|
| $<4.3 \times 10^{-6}$ | 90 | 45 CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$ |

⁴⁵ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{19}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|------------------------|-----|-------------------|------|--|
| $<14.2 \times 10^{-6}$ | 90 | 46 CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta'$ |

⁴⁶ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{20}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|-------------------|------|---|
| $<1.7 \times 10^{-4}$ | 90 | 47 CRONIN-HEN..10 | CLEO | $\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta_c(1S)$ |

⁴⁷ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$\eta_c(2S)$ REFERENCES

| | | | | |
|---------------|------|----------------|-----------------------------------|------------------|
| CHILIKIN | 19 | PR D100 012001 | K. Chilikin <i>et al.</i> | (BELLE Collab.) |
| XU | 18 | PR D98 072001 | Q.N. Xu <i>et al.</i> | (BELLE Collab.) |
| AAIJ | 17AD | PL B769 305 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 17BB | EPJ C77 609 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| ABLIKIM | 17N | PR D95 072004 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| PDG | 16 | CP C40 100001 | C. Patrignani <i>et al.</i> | (PDG Collab.) |
| LEES | 14E | PR D89 112004 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 13K | PR D87 052005 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 13V | PR D88 112001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| ABLIKIM | 12G | PRL 109 042003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| LEES | 12AE | PR D86 092005 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 11H | PR D84 091102 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| DEL-AMO-SA... | 11M | PR D84 012004 | P. del Amo Sanchez <i>et al.</i> | (BABAR Collab.) |
| VINOKUROVA | 11 | PL B706 139 | A. Vinokurova <i>et al.</i> | (BELLE Collab.) |
| CRONIN-HEN... | 10 | PR D81 052002 | D. Cronin-Hennessey <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 08AB | PR D78 012006 | B. Aubert <i>et al.</i> | (BABAR Collab.) |

| | | | | |
|------------|-----|---------------|------------------------------|---------------------|
| UEHARA | 08 | EPJ C53 1 | S. Uehara <i>et al.</i> | (BELLE Collab.) |
| WICHT | 08 | PL B662 323 | J. Wicht <i>et al.</i> | (BELLE Collab.) |
| ABE | 07 | PRL 98 082001 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| AUBERT | 06E | PRL 96 052002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 05C | PR D72 031101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| ABE | 04G | PR D70 071102 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| ASNER | 04 | PRL 92 142001 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 04D | PRL 92 142002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| ABE,K | 02 | PRL 89 142001 | K. Abe <i>et al.</i> | (BELLE Collab.) |
| CHOI | 02 | PRL 89 102001 | S.-K. Choi <i>et al.</i> | (BELLE Collab.) |
| AMBROGIANI | 01 | PR D64 052003 | M. Ambrogiani <i>et al.</i> | (FNAL E835 Collab.) |
| ABREU | 98O | PL B441 479 | P. Abreu <i>et al.</i> | (DELPHI Collab.) |
| ARMSTRONG | 95F | PR D52 4839 | T.A. Armstrong <i>et al.</i> | (FNAL, FERR, GENO+) |
| LEE | 85 | SLAC 282 | R.A. Lee | (SLAC) |
| EDWARDS | 82C | PRL 48 70 | C. Edwards <i>et al.</i> | (CIT, HARV, PRIN+) |