

$f_1(1285)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

 $f_1(1285)$ MASS

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|--------------------|---|-----------|--|
| 1281.9 ± 0.5 | OUR AVERAGE | Error includes scale factor of 1.8. See the ideogram below. | | |
| 1281.0 ± 0.8 | | DICKSON | 16 CLAS | 2.55 $\gamma p \rightarrow \eta \pi^+ \pi^- p$ |
| 1287.4 ± 3.0 | 87 | ABLIKIM | 15P BES3 | $J/\psi \rightarrow K^+ K^- 3\pi$ |
| 1281.16 ± 0.39 ± 0.45 | | ¹ LEES | 12X BABR | $\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$ |
| 1285.1 ± 1.0 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.6 \\ 0.3 \end{smallmatrix}$ | | ² ABLIKIM | 11J BES3 | $J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$ |
| 1281 ± 2 ± 1 | | AUBERT | 07AU BABR | 10.6 $e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$ |
| 1276.1 ± 8.1 ± 8.0 | 203 | BAI | 04J BES2 | $J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$ |
| 1274 ± 6 | 237 | ABDALLAH | 03H DLPH | 91.2 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$ |
| 1280 ± 4 | | ACCIARRI | 01G L3 | |
| 1288 ± 4 ± 5 | 20k | ADAMS | 01B B852 | 18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 1284 ± 6 | 1400 | ALDE | 97B GAM4 | 100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$ |
| 1281 ± 1 | | BARBERIS | 97B OMEG | 450 $pp \rightarrow pp 2(\pi^+ \pi^-)$ |
| 1281 ± 1 | | BARBERIS | 97C OMEG | 450 $pp \rightarrow pp K_S^0 K^\pm \pi^\mp$ |
| 1280 ± 2 | | ³ ANTINORI | 95 OMEG | 300,450 $pp \rightarrow pp 2(\pi^+ \pi^-)$ |
| 1282.2 ± 1.5 | | LEE | 94 MPS2 | 18 $\pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$ |
| 1279 ± 5 | | FUKUI | 91C SPEC | 8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$ |
| 1278 ± 2 | 140 | ARMSTRONG | 89 OMEG | 300 $pp \rightarrow K \bar{K} \pi pp$ |
| 1278 ± 2 | | ARMSTRONG | 89G OMEG | 85 $\pi^+ p \rightarrow 4\pi \pi p, pp \rightarrow 4\pi pp$ |
| 1280.1 ± 2.1 | 60 | RATH | 89 MPS | 21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$ |
| 1285 ± 1 | 4750 | ⁴ BIRMAN | 88 MPS | 8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$ |
| 1280 ± 1 | 504 | BITYUKOV | 88 SPEC | 32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 1280 ± 4 | | ANDO | 86 SPEC | 8 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$ |
| 1277 ± 2 | 420 | REEVES | 86 SPEC | 6.6 $p\bar{p} \rightarrow KK\pi X$ |
| 1285 ± 2 | | CHUNG | 85 SPEC | 8 $\pi^- p \rightarrow NK\bar{K}\pi$ |
| 1279 ± 2 | 604 | ARMSTRONG | 84 OMEG | 85 $\pi^+ p \rightarrow K\bar{K}\pi\pi p, pp \rightarrow K\bar{K}\pi pp$ |
| 1286 ± 1 | | CHAUVAT | 84 SPEC | ISR 31.5 pp |
| 1278 ± 4 | | EVANGELIS... | 81 OMEG | 12 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$ |
| 1283 ± 3 | 103 | DIONISI | 80 HBC | 4 $\pi^- p \rightarrow K\bar{K}\pi n$ |
| 1282 ± 2 | 320 | NACASCH | 78 HBC | 0.7,0.76 $\bar{p}p \rightarrow K\bar{K}3\pi$ |
| 1279 ± 5 | 210 | GRASSLER | 77 HBC | 16 $\pi^\mp p$ |
| 1286 ± 3 | 180 | DUBOC | 72 HBC | 1.2 $\bar{p}p \rightarrow 2K4\pi$ |
| 1283 ± 5 | | DAHL | 67 HBC | 1.6–4.2 $\pi^- p$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|--------------------|-----|------------------------------------|------------|------|--|
| 1289.3 ± 2.8 | 234 | ABLIKIM | 19BA | BES3 | $e^+e^- \rightarrow \psi(2S)$ |
| 1284.2 ± 2.2 | | ⁵ AAIJ | 14Y | LHCB | $\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+\pi^-)$ |
| 1281.9 ± 0.5 | | ⁵ SOSA | 99 | SPEC | $pp \rightarrow p_{\text{slow}}(K_S^0 K^+\pi^-) p_{\text{fast}}$ |
| 1282.8 ± 0.6 | | ⁵ SOSA | 99 | SPEC | $pp \rightarrow p_{\text{slow}}(K_S^0 K^-\pi^+) p_{\text{fast}}$ |
| 1270 ± 10 | | AMELIN | 95 | VES | $37 \pi^- N \rightarrow \pi^-\pi^+\pi^-\gamma N$ |
| 1280 ± 2 | | ABATZIS | 94 | OMEG | $450 pp \rightarrow pp 2(\pi^+\pi^-)$ |
| 1282 ± 4 | | ARMSTRONG | 93C | E760 | $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$ |
| 1270 ± 6 ± 10 | | ARMSTRONG | 92C | OMEG | $300 pp \rightarrow pp \pi^+\pi^-\gamma$ |
| 1281 ± 1 | | ARMSTRONG | 89E | OMEG | $300 pp \rightarrow pp 2(\pi^+\pi^-)$ |
| 1279 ± 6 ± 10 | 16 | BECKER | 87 | MRK3 | $e^+e^- \rightarrow \phi K \bar{K} \pi$ |
| 1286 ± 9 | | GIDAL | 87 | MRK2 | $e^+e^- \rightarrow e^+e^-\eta \pi^+\pi^-$ |
| 1287 ± 5 ~ 1279 | 353 | BITYUKOV ⁶ TORNQVIST | 84B 82B | SPEC | $32 \pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 1275 ± 6 | 31 | BROMBERG | 80 | SPEC | $100 \pi^- p \rightarrow K \bar{K} \pi X$ |
| 1288 ± 9 | 200 | GURTU | 79 | HBC | $4.2 K^- p \rightarrow n \eta 2\pi$ |
| ~ 1275.0 | 46 | ⁷ STANTON | 79 | CNTR | $8.5 \pi^- p \rightarrow n 2\gamma 2\pi$ |
| 1271 ± 10 | 34 | CORDEN | 78 | OMEG | $12-15 \pi^- p \rightarrow K^+ K^- \pi n$ |
| 1295 ± 12 | 85 | CORDEN | 78 | OMEG | $12-15 \pi^- p \rightarrow n 5\pi$ |
| 1292 ± 10 | 150 | DEFOIX | 72 | HBC | $0.7 \bar{p}p \rightarrow 7\pi$ |
| 1280 ± 3 | 500 | ⁸ THUN | 72 | MMS | $13.4 \pi^- p$ |
| 1303 ± 8 | | BARDADIN-... | 71 | HBC | $8 \pi^+ p \rightarrow p 6\pi$ |
| 1283 ± 6 | | BOESEBECK | 71 | HBC | $16.0 \pi p \rightarrow p 5\pi$ |
| 1270 ± 10 | | CAMPBELL | 69 | DBC | $2.7 \pi^+ d$ |
| 1285 ± 7 | | LORSTAD | 69 | HBC | $0.7 \bar{p}p, 4,5\text{-body}$ |
| 1290 ± 7 | | D'ANDLAU | 68 | HBC | $1.2 \bar{p}p, 5-6 \text{ body}$ |

¹ Using the $2\pi^+2\pi^-$ and $\pi^+\pi^-\eta$ modes of $f_1(1285)$ decay.

² The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

³ Supersedes ABATZIS 94, ARMSTRONG 89E.

⁴ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.


⁵ No systematic error given.

⁶ From a unitarized quark-model calculation.

⁷ From phase shift analysis of $\eta \pi^+ \pi^-$ system.

⁸ Seen in the missing mass spectrum.

WEIGHTED AVERAGE
 1281.9 ± 0.5 (Error scaled by 1.8)



$f_1(1285)$ WIDTH

Only experiments giving width error less than 20 MeV are kept for averaging.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|---|------|---|
| 22.7 ± 1.1 OUR AVERAGE | | Error includes scale factor of 1.5. See the ideogram below. | | |
| 18.4 ± 1.4 | | DICKSON 16 | CLAS | $2.55 \gamma p \rightarrow \eta \pi^+ \pi^- p$ |
| 18.3 ± 6.3 | 87 | ABLIKIM 15P | BES3 | $J/\psi \rightarrow K^+ K^- 3\pi$ |
| $22.0 \pm 3.1^{+2.0}_{-1.5}$ | | ¹ ABLIKIM 11J | BES3 | $J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$ |
| $35 \pm 6 \pm 4$ | | AUBERT 07AU | BABR | $10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$ |
| $40.0 \pm 8.6 \pm 9.3$ | 203 | BAI 04J | BES2 | $J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$ |

| | | | | | | |
|---|------|-----------------------|------|------|-----------|--|
| 29 ± 12 | 237 | ABDALLAH | 03H | DLPH | 91.2 | $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$ |
| 45 ± 9 ± 7 | 20k | ADAMS | 01B | B852 | 18 GeV | $\pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 55 ± 18 | 1400 | ALDE | 97B | GAM4 | 100 | $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$ |
| 24 ± 3 | | BARBERIS | 97B | OMEG | 450 | $pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 20 ± 2 | | BARBERIS | 97C | OMEG | 450 | $pp \rightarrow ppK_S^0 K^\pm \pi^\mp$ |
| 36 ± 5 | | ² ANTINORI | 95 | OMEG | 300,450 | $pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 29.0 ± 4.1 | | LEE | 94 | MPS2 | 18 | $\pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$ |
| 25 ± 4 | 140 | ARMSTRONG | 89 | OMEG | 300 | $pp \rightarrow K \bar{K} \pi pp$ |
| 22 ± 2 | 4750 | ³ BIRMAN | 88 | MPS | 8 | $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$ |
| 25 ± 4 | 504 | BITYUKOV | 88 | SPEC | 32.5 | $\pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| 19 ± 5 | | ANDO | 86 | SPEC | 8 | $\pi^- p \rightarrow \eta \pi^+ \pi^- n$ |
| 32 ± 8 | 420 | REEVES | 86 | SPEC | 6.6 | $p\bar{p} \rightarrow KK\pi X$ |
| 22 ± 2 | | CHUNG | 85 | SPEC | 8 | $\pi^- p \rightarrow NK\bar{K}\pi$ |
| 32 ± 3 | 604 | ARMSTRONG | 84 | OMEG | 85 | $\pi^+ p \rightarrow K\bar{K}\pi\pi p,$ $pp \rightarrow K\bar{K}\pi pp$ |
| 24 ± 3 | | CHAUVAT | 84 | SPEC | ISR 31.5 | pp |
| 29 ± 10 | 103 | DIONISI | 80 | HBC | 4 | $\pi^- p \rightarrow K\bar{K}\pi n$ |
| 28.3 ± 6.7 | 320 | NACASCH | 78 | HBC | 0.7, 0.76 | $\bar{p}p \rightarrow K\bar{K}3\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | | |
| 17.1 ± 3.4 | 234 | ABLIKIM | 19BA | BES3 | | $e^+e^- \rightarrow \psi(2S)$ |
| 32.4 ± 5.8 | | ⁴ AAIJ | 14Y | LHCB | | $\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$ |
| 18.2 ± 1.2 | | ⁴ SOSA | 99 | SPEC | | $pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$ p_{fast} |
| 19.4 ± 1.5 | | ⁴ SOSA | 99 | SPEC | | $pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+)$ p_{fast} |
| 40 ± 5 | | ABATZIS | 94 | OMEG | 450 | $pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 31 ± 5 | | ARMSTRONG | 89E | OMEG | 300 | $pp \rightarrow pp2(\pi^+ \pi^-)$ |
| 41 ± 12 | | ARMSTRONG | 89G | OMEG | 85 | $\pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$ |
| 17.9 ± 10.9 | 60 | RATH | 89 | MPS | 21.4 | $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$ |
| 14 ⁺²⁰ / ₋₁₄ ± 10 | 16 | BECKER | 87 | MRK3 | | $e^+e^- \rightarrow \phi K\bar{K}\pi$ |
| 26 ± 12 | | EVANGELIS... | 81 | OMEG | 12 | $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$ |
| 25 ± 15 | 200 | GURTU | 79 | HBC | 4.2 | $K^- p \rightarrow n\eta 2\pi$ |
| ~ 10 | | ⁵ STANTON | 79 | CNTR | 8.5 | $\pi^- p \rightarrow n2\gamma 2\pi$ |
| 24 ± 18 | 210 | GRASSLER | 77 | HBC | 16 | $\pi^\mp p$ |
| 28 ± 5 | 150 | ⁶ DEFOIX | 72 | HBC | 0.7 | $\bar{p}p \rightarrow 7\pi$ |
| 46 ± 9 | 180 | ⁶ DUBOC | 72 | HBC | 1.2 | $\bar{p}p \rightarrow 2K4\pi$ |
| 37 ± 5 | 500 | ⁷ THUN | 72 | MMS | 13.4 | $\pi^- p$ |
| 10 ± 10 | | BOESEBECK | 71 | HBC | 16.0 | $\pi p \rightarrow p5\pi$ |
| 30 ± 15 | | CAMPBELL | 69 | DBC | 2.7 | $\pi^+ d$ |
| 60 ± 15 | | ⁶ LORSTAD | 69 | HBC | 0.7 | $\bar{p}p, 4,5\text{-body}$ |
| 35 ± 10 | | ⁶ DAHL | 67 | HBC | 1.6–4.2 | $\pi^- p$ |

¹ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

² Supersedes ABATZIS 94, ARMSTRONG 89E.

³ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.


⁴ No systematic error given.

⁵ From phase shift analysis of $\eta\pi^+\pi^-$ system.

⁶ Resolution is not unfolded.

⁷ Seen in the missing mass spectrum.

WEIGHTED AVERAGE
 22.7 ± 1.1 (Error scaled by 1.5)



$f_1(1285)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|--|--------------------------------|-----------------------------------|
| Γ_1 4π | $(32.7 \pm 1.9) \%$ | S=1.2 |
| Γ_2 $\pi^0\pi^0\pi^+\pi^-$ | $(21.8 \pm 1.3) \%$ | S=1.2 |
| Γ_3 $2\pi^+2\pi^-$ | $(10.9 \pm 0.6) \%$ | S=1.2 |
| Γ_4 $\rho^0\pi^+\pi^-$ | $(10.9 \pm 0.6) \%$ | S=1.2 |
| Γ_5 $\rho^0\rho^0$ | seen | |
| Γ_6 $4\pi^0$ | $< 7 \times 10^{-4}$ | CL=90% |
| Γ_7 $\eta\pi^+\pi^-$ | $(35 \pm 15) \%$ | |
| Γ_8 $\eta\pi\pi$ | $(52.2 \pm 2.0) \%$ | S=1.2 |
| Γ_9 $a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$] | $(38 \pm 4) \%$ | |

| | | | |
|---------------|---|--------------------------------|--------|
| Γ_{10} | $\eta\pi\pi$ [excluding $a_0(980)\pi$] | $(14 \pm 4) \%$ | |
| Γ_{11} | $K\bar{K}\pi$ | $(9.0 \pm 0.4) \%$ | S=1.1 |
| Γ_{12} | $K\bar{K}^*(892)$ | not seen | |
| Γ_{13} | $\pi^+\pi^-\pi^0$ | $(3.0 \pm 0.9) \times 10^{-3}$ | |
| Γ_{14} | $\rho^\pm\pi^\mp$ | $< 3.1 \times 10^{-3}$ | CL=95% |
| Γ_{15} | $\gamma\rho^0$ | $(6.1 \pm 1.0) \%$ | S=1.7 |
| Γ_{16} | $\phi\gamma$ | $(7.4 \pm 2.6) \times 10^{-4}$ | |
| Γ_{17} | e^+e^- | $< 9.4 \times 10^{-9}$ | CL=90% |
| Γ_{18} | $\gamma\gamma^*$ | | |
| Γ_{19} | $\gamma\gamma$ | | |

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 18 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 24.0$ for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | | |
|----------|-------|-------|----------|----------|
| x_9 | -30 | | | |
| x_{10} | -12 | -88 | | |
| x_{11} | 22 | -10 | -4 | |
| x_{15} | -25 | -7 | -3 | -27 |
| | x_1 | x_9 | x_{10} | x_{11} |

$f_1(1285) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

| $\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ | | $\Gamma_8\Gamma_{19}/\Gamma = (\Gamma_9+\Gamma_{10})\Gamma_{19}/\Gamma$ | | |
|--|-----|---|------|--|
| VALUE (keV) | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.62 | 95 | GIDAL | 87 | MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$ |

| $\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}}$ | | $\Gamma_8\Gamma_{18}/\Gamma = (\Gamma_9+\Gamma_{10})\Gamma_{18}/\Gamma$ | | |
|---|------|---|------|--|
| VALUE (keV) | EVTS | DOCUMENT ID | TECN | COMMENT |
| 1.4 \pm 0.4 OUR AVERAGE | | Error includes scale factor of 1.4. | | |
| 1.18 \pm 0.25 \pm 0.20 | 26 | ^{1,2} AIHARA | 88B | TPC $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$ |
| 2.30 \pm 0.61 \pm 0.42 | | ^{1,3} GIDAL | 87 | MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 1.8 \pm 0.3 \pm 0.3 | 420 | ⁴ ACHARD | 02B | L3 183–209 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$ |

¹ Assuming a ρ -pole form factor.

² Published value multiplied by $\eta\pi\pi$ branching ratio 0.49.

³ Published value divided by 2 and multiplied by the $\eta\pi\pi$ branching ratio 0.49.

⁴ Published value multiplied by the $\eta\pi\pi$ branching ratio 0.52.

$f_1(1285)$ BRANCHING RATIOS $\Gamma(K\bar{K}\pi)/\Gamma(4\pi)$ Γ_{11}/Γ_1

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------------|------------------------|------|---|
| 0.274±0.017 OUR FIT | | | Error includes scale factor of 1.4. |
| 0.271±0.016 OUR AVERAGE | | | Error includes scale factor of 1.2. |
| 0.265±0.014 | ¹ BARBERIS | 97C | OMEG 450 $pp \rightarrow p\rho K_S^0 K^\pm \pi^\mp$ |
| 0.28 ±0.05 | ² ARMSTRONG | 89E | OMEG 300 $pp \rightarrow p\rho f_1(1285)$ |
| 0.37 ±0.03 ±0.05 | ³ ARMSTRONG | 89G | OMEG 85 $\pi p \rightarrow 4\pi X$ |

¹ Using $2(\pi^+ \pi^-)$ data from BARBERIS 97B.² Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.³ 4π consistent with being entirely $\rho\pi\pi$. $\Gamma(\pi^0 \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma = \frac{2}{3}\Gamma_1/\Gamma$

| VALUE | DOCUMENT ID |
|----------------------------|-------------|
| 0.218±0.013 OUR FIT | |

Error includes scale factor of 1.2.

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$

| VALUE | DOCUMENT ID |
|----------------------------|-------------|
| 0.109±0.006 OUR FIT | |

Error includes scale factor of 1.2.

 $\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma = \frac{1}{3}\Gamma_1/\Gamma$

| VALUE | DOCUMENT ID |
|----------------------------|-------------|
| 0.109±0.006 OUR FIT | |

Error includes scale factor of 1.2.

 $\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2\pi^+ 2\pi^-)$ Γ_4/Γ_3

| VALUE | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|------------------------|
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1.0±0.4 | GRASSLER | 77 | HBC 16 GeV $\pi^\pm p$ |

 $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_5/Γ

| VALUE | DOCUMENT ID | COMMENT |
|-------------|-------------|---------------------------------------|
| seen | BARBERIS | 00C 450 $pp \rightarrow p_f 4\pi p_s$ |

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---|
| <7 | 90 | ALDE | 87 | GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$ |

 $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma(\eta \pi^+ \pi^-)$ Γ_{13}/Γ_7

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------|------|-----------------------|------|---|
| 0.86±0.16±0.20 | 2.3k | ¹ DOROFEEV | 11 | VES $\pi^- N \rightarrow \pi^- f_1(1285) N$ |

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+ \pi^-$ mass spectrum. $\Gamma(\eta \pi \pi)/\Gamma_{\text{total}}$ $\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$

| VALUE | DOCUMENT ID |
|----------------------------|-------------|
| 0.522±0.020 OUR FIT | |

Error includes scale factor of 1.2.

$\Gamma(4\pi)/\Gamma(\eta\pi\pi)$

$\Gamma_1/\Gamma_8 = \Gamma_1/(\Gamma_9+\Gamma_{10})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.63±0.06 OUR FIT Error includes scale factor of 1.3.**0.41±0.14 OUR AVERAGE**0.37±0.11±0.11 BOLTON 92 MRK3 $J/\psi \rightarrow \gamma f_1(1285)$ 0.64±0.40 GURTU 79 HBC 4.2 $K^- p$

••• We do not use the following data for averages, fits, limits, etc. •••

0.93±0.30 ¹ GRASSLER 77 HBC 16 $\pi^\mp p$ ¹ Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states. $\Gamma(2\pi^+2\pi^-)/\Gamma(\eta\pi\pi)$

Γ_3/Γ_8

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.28±0.02±0.02 ¹ LEES 12X BABR $\tau^- \rightarrow \pi^- f_1(1285)\nu_\tau$ ¹ Assuming $B(f_1(1285) \rightarrow \pi\pi\eta) = 3/2 B(f_1(1285) \rightarrow \pi^+\pi^-\eta)$. $\Gamma(a_0(980)\pi [\text{ignoring } a_0(980) \rightarrow K\bar{K}])/ \Gamma(\eta\pi\pi)$

$\Gamma_9/\Gamma_8 = \Gamma_9/(\Gamma_9+\Gamma_{10})$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

0.72±0.08 OUR FIT**0.72±0.07 OUR AVERAGE**0.74±0.02±0.09 DICKSON 16 CLAS $\gamma p \rightarrow f_1(1285)p$ 0.72±0.15 GURTU 79 HBC 4.2 $K^- p$ 0.6 $\begin{matrix} +0.3 \\ -0.2 \end{matrix}$ CORDEN 78 OMEG 12–15 $\pi^- p$

••• We do not use the following data for averages, fits, limits, etc. •••

>0.69 95 ACHARD 02B L3 183–209 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$ 0.28±0.07 ALDE 97B GAM4 100 $\pi^- p \rightarrow \eta\pi^0\pi^0n$ 1.0 ±0.3 GRASSLER 77 HBC 16 $\pi^\mp p$ $\Gamma(K\bar{K}\pi)/\Gamma(\eta\pi\pi)$

$\Gamma_{11}/\Gamma_8 = \Gamma_{11}/(\Gamma_9+\Gamma_{10})$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

0.172±0.012 OUR FIT Error includes scale factor of 1.1.**0.176±0.012 OUR AVERAGE**0.216±0.010±0.031 DICKSON 16 CLAS $\gamma p \rightarrow f_1(1285)p$ 0.166±0.01 ±0.008 BARBERIS 98c OMEG 450 $pp \rightarrow p_f f_1(1285)p_S$ 0.42 ±0.15 GURTU 79 HBC 4.2 $K^- p$ 0.5 ±0.2 ¹ CORDEN 78 OMEG 12–15 $\pi^- p$ 0.20 ±0.08 ² DEFOIX 72 HBC 0.7 $\bar{p}p \rightarrow 7\pi$ 0.16 ±0.08 CAMPBELL 69 DBC 2.7 $\pi^+ d$ ¹ CORDEN 78 assumes low-mass $\eta\pi\pi$ region is dominantly 1^{++} . See BARBERIS 98c and MANAK 00A for discussion.² $K\bar{K}$ system characterized by the $l = 1$ threshold enhancement. (See under $a_0(980)$). $\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$

Γ_{12}/Γ

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

not seen NACASCH 78 HBC 0.7,0.76 $\bar{p}p \rightarrow K\bar{K}3\pi$

••• We do not use the following data for averages, fits, limits, etc. •••

seen ¹ ACHARD 07 L3 183–209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$ ¹ A clear signal of 19.8 ± 4.4 events observed at high Q^2 .

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

| VALUE (%) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------|------|--------------------------|------|---|
| 0.30±0.055±0.074 | 2.3k | ¹ DOROFEEV 11 | VES | $\pi^- N \rightarrow \pi^- f_1(1285) N$ |

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum. The systematic error includes the uncertainty on the partial width $f_1 \rightarrow \eta\pi\pi$ obtained from PDG 10 data.

$\Gamma(\rho^\pm\pi^\mp)/\Gamma_{\text{total}}$ Γ_{14}/Γ

| VALUE (%) | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------|-----|-------------|------|---|
| <0.31 | 95 | DOROFEEV 11 | VES | $\pi^- N \rightarrow \pi^- f_1(1285) N$ |

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

| VALUE (units 10^{-2}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|-------------------------------------|
| 6.1±1.0 OUR FIT | | | | Error includes scale factor of 1.7. |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------------|----|------------------------|------|---|
| $2.8\pm 0.7\pm 0.6$ | | ¹ AMELIN 95 | VES | $37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$ |
| <5 | 95 | BITYUKOV 91B | SPEC | $32 \pi^- p \rightarrow \pi^+ \pi^- \gamma n$ |

¹ Not an independent measurement.

$\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------------------|------|---|
| 0.55±0.10 OUR FIT | | | Error includes scale factor of 1.5. |
| 0.45±0.18 | ¹ COFFMAN 90 | MRK3 | $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |

¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.

$\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$ $\Gamma_8/\Gamma_{15} = (\Gamma_9 + \Gamma_{10})/\Gamma_{15}$


| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------------|-------------|------|-------------------------------------|
| 8.6±1.6 OUR FIT | | | Error includes scale factor of 1.9. |

8.5±2.0 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

| | | | |
|----------------------|----------------------------|----------|---|
| 21.3 ± 4.4 | DICKSON 16 | CLAS | $\gamma p \rightarrow f_1(1285) p$ |
| $10.0\pm 1.0\pm 2.0$ | BARBERIS 98C | OMEG 450 | $pp \rightarrow p_f f_1(1285) p_s$ |
| 7.5 ± 1.0 | ¹ ARMSTRONG 92C | OMEG 300 | $pp \rightarrow pp\pi^+\pi^-\gamma, pp\eta\pi^+\pi^-$ |

¹ Published value multiplied by 1.5.

WEIGHTED AVERAGE
8.5±2.0 (Error scaled by 2.2)



$\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$

Γ_{15}/Γ_{11}

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|--|-----|-------------------------|------|---|
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| >0.035 | 90 | ¹ COFFMAN 90 | MRK3 | $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ |
| ¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) < 0.72 \times 10^{-3}$. | | | | |

$\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$

Γ_{16}/Γ_{11}

| VALUE (units 10^{-2}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|--------------|------|---|
| 0.82±0.21±0.20 | | 19 | BITYUKOV 88 | SPEC | 32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <0.50 | 95 | | BARBERIS 98C | OMEG | 450 $pp \rightarrow p_f f_1(1285) p_s$ |
| <0.93 | 95 | | AMELIN 95 | VES | 37 $\pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$ |

$\Gamma(e^+e^-)/\Gamma_{total}$

Γ_{17}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------------------|------|-------------------------------------|
| <9.4 × 10⁻⁹ | 90 | ¹ ACHASOV 20 | SND | $e^+e^- \rightarrow \eta\pi^0\pi^0$ |
| ¹ ACHASOV 20 reports two candidate events corresponding to a significance of 2.5 σ and the branching fraction of $(5.1^{+3.7}_{-2.7}) \times 10^{-9}$. | | | | |

$f_1(1285)$ REFERENCES

| | | | | |
|--------------|------|------------------------|--------------------------------------|---------------------------------|
| ACHASOV | 20 | PL B800 135074 | M.N. Achasov <i>et al.</i> | (SND Collab.) |
| ABLIKIM | 19BA | PR D100 092003 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| DICKSON | 16 | PR C93 065202 | R. Dickson <i>et al.</i> | (JLab CLAS Collab.) |
| ABLIKIM | 15P | PR D92 012007 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| AAIJ | 14Y | PRL 112 091802 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| LEES | 12X | PR D86 092010 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| ABLIKIM | 11J | PRL 107 182001 | M. Ablikim <i>et al.</i> | (BESIII Collab.) |
| DOROFEEV | 11 | EPJ A47 68 | V. Dorofeev <i>et al.</i> | (SERP, MIPT) |
| PDG | 10 | JP G37 075021 | K. Nakamura <i>et al.</i> | (PDG Collab.) |
| ACHARD | 07 | JHEP 0703 018 | P. Achard <i>et al.</i> | (L3 Collab.) |
| AUBERT | 07AU | PR D76 092005 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BAI | 04J | PL B594 47 | J.Z. Bai <i>et al.</i> | (BES Collab.) |
| ABDALLAH | 03H | PL B569 129 | J. Abdallah <i>et al.</i> | (DELPHI Collab.) |
| ACHARD | 02B | PL B526 269 | P. Achard <i>et al.</i> | (L3 Collab.) |
| ACCIARRI | 01G | PL B501 1 | M. Acciarri <i>et al.</i> | (L3 Collab.) |
| ADAMS | 01B | PL B516 264 | G.S. Adams <i>et al.</i> | (BNL E852 Collab.) |
| BARBERIS | 00C | PL B471 440 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| MANAK | 00A | PR D62 012003 | J.J. Manak <i>et al.</i> | (BNL E852 Collab.) |
| SOSA | 99 | PRL 83 913 | M. Sosa <i>et al.</i> | |
| BARBERIS | 98C | PL B440 225 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| ALDE | 97B | PAN 60 386 | D. Alde <i>et al.</i> | (GAMS Collab.) |
| | | Translated from YAF 60 | 458. | |
| BARBERIS | 97B | PL B413 217 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| BARBERIS | 97C | PL B413 225 | D. Barberis <i>et al.</i> | (WA 102 Collab.) |
| AMELIN | 95 | ZPHY C66 71 | D.V. Amelin <i>et al.</i> | (VES Collab.) |
| ANTINORI | 95 | PL B353 589 | F. Antinori <i>et al.</i> | (ATHU, BARI, BIRM+) |
| ABATZIS | 94 | PL B324 509 | S. Abatzis <i>et al.</i> | (ATHU, BARI, BIRM+) |
| LEE | 94 | PL B323 227 | J.H. Lee <i>et al.</i> | (BNL, IND, KYUN, MASD+) |
| ARMSTRONG | 93C | PL B307 394 | T.A. Armstrong <i>et al.</i> | (FNAL, FERR, GENO+) |
| ARMSTRONG | 92C | ZPHY C54 371 | T.A. Armstrong <i>et al.</i> | (ATHU, BARI, BIRM+) |
| BOLTON | 92 | PL B278 495 | T. Bolton <i>et al.</i> | (Mark III Collab.) |
| BITYUKOV | 91B | SJNP 54 318 | S.I. Bityukov <i>et al.</i> | (SERP) |
| | | Translated from YAF 54 | 529. | |
| FUKUI | 91C | PL B267 293 | S. Fukui <i>et al.</i> | (SUGI, NAGO, KEK, KYOT+) |
| COFFMAN | 90 | PR D41 1410 | D.M. Coffman <i>et al.</i> | (Mark III Collab.) |
| ARMSTRONG | 89 | PL B221 216 | T.A. Armstrong <i>et al.</i> | (CERN, CDEF, BIRM+) JPC |
| ARMSTRONG | 89E | PL B228 536 | T.A. Armstrong, M. Benayoun | (ATHU, BARI, BIRM+) |
| ARMSTRONG | 89G | ZPHY C43 55 | T.A. Armstrong <i>et al.</i> | (CERN, BIRM, BARI+) |
| RATH | 89 | PR D40 693 | M.G. Rath <i>et al.</i> | (NDAM, BRAN, BNL, CUNY+) |
| AIHARA | 88B | PL B209 107 | H. Aihara <i>et al.</i> | (TPC-2 γ Collab.) |
| BIRMAN | 88 | PRL 61 1557 | A. Birman <i>et al.</i> | (BNL, FSU, IND, MASD) JP |
| BITYUKOV | 88 | PL B203 327 | S.I. Bityukov <i>et al.</i> | (SERP) |
| MIR | 88 | Photon-Photon 88, 126 | R. Mir | (Mark III Collab.) |
| Conference | | | | |
| ALDE | 87 | PL B198 286 | D.M. Alde <i>et al.</i> | (LANL, BRUX, SERP, LAPP) |
| BECKER | 87 | PRL 59 186 | J.J. Becker <i>et al.</i> | (Mark III Collab.) |
| GIDAL | 87 | PRL 59 2012 | G. Gidal <i>et al.</i> | (LBL, SLAC, HARV) |
| ANDO | 86 | PRL 57 1296 | A. Ando <i>et al.</i> | (KEK, KYOT, NIRS, SAGA+) IJP |
| REEVES | 86 | PR D34 1960 | D.F. Reeves <i>et al.</i> | (FLOR, BNL, IND+) JP |
| CHUNG | 85 | PRL 55 779 | S.U. Chung <i>et al.</i> | (BNL, FLOR, IND+) JP |
| ARMSTRONG | 84 | PL 146B 273 | T.A. Armstrong <i>et al.</i> | (ATHU, BARI, BIRM+) JP |
| BITYUKOV | 84B | PL 144B 133 | S.I. Bityukov <i>et al.</i> | (SERP) |
| CHAUVAT | 84 | PL 148B 382 | P. Chauvat <i>et al.</i> | (CERN, CLER, UCLA+) |
| TORNQVIST | 82B | NP B203 268 | N.A. Tornqvist | (HELS) |
| EVANGELIS... | 81 | NP B178 197 | C. Evangelista <i>et al.</i> | (BARI, BONN, CERN+) |
| BROMBERG | 80 | PR D22 1513 | C.M. Bromberg <i>et al.</i> | (CIT, FNAL, ILLC+) |
| DIONISI | 80 | NP B169 1 | C. Dionisi <i>et al.</i> | (CERN, MADR, CDEF+) |
| GURTU | 79 | NP B151 181 | A. Gurtu <i>et al.</i> | (CERN, ZEEM, NIJM, OXF) |
| STANTON | 79 | PRL 42 346 | N.R. Stanton <i>et al.</i> | (OSU, CARL, MCGI+) JP |
| CORDEN | 78 | NP B144 253 | M.J. Corden <i>et al.</i> | (BIRM, RHEL, TELA+) JP |
| NACASCH | 78 | NP B135 203 | R. Nacasch <i>et al.</i> | (PARIS, MADR, CERN) |
| GRASSLER | 77 | NP B121 189 | H. Grassler <i>et al.</i> | (AACH3, BERL, BONN+) |
| DEFOIX | 72 | NP B44 125 | C. Defoix <i>et al.</i> | (CDEF, CERN) |
| DUBOC | 72 | NP B46 429 | J. Duboc <i>et al.</i> | (PARIS, LIVP) |
| THUN | 72 | PRL 28 1733 | R. Thun <i>et al.</i> | (STON, NEAS) |
| BARDADIN-... | 71 | PR D4 2711 | M. Bardadin-Otwinowska <i>et al.</i> | (WARS) |
| BOESEBECK | 71 | PL 34B 659 | K. Boesebeck | (AACH, BERL, BONN, CERN, CRAC+) |
| CAMPBELL | 69 | PRL 22 1204 | J.H. Campbell <i>et al.</i> | (PURD) |

| | | | | |
|----------|----|-------------|---------------------------|-------------------------|
| LORSTAD | 69 | NP B14 63 | B. Lorstad <i>et al.</i> | (CDEF, CERN) JP |
| D'ANDLAU | 68 | NP B5 693 | C. d'Andlau <i>et al.</i> | (CDEF, CERN, IRAD+) IJP |
| DAHL | 67 | PR 163 1377 | O.I. Dahl <i>et al.</i> | (LRL) IJP |
