

$f_2(2150)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called T_0 . **$f_2(2150)$ MASS** **$f_2(2150)$ MASS, COMBINED MODES (MeV)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2157±12 OUR AVERAGE Includes data from the datablock that follows this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2170±6 80k ¹UMAN 06 E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$ ¹Statistical error only. **$\eta\eta$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

2157±12 OUR AVERAGE

2151±16	BARBERIS	00E	450 $pp \rightarrow p_f \eta \eta p_s$
2175±20	PROKOSHKIN	95D	GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $pp \rightarrow pp 2\eta$
2130±35	SINGOVSKI	94	GAM4 450 $pp \rightarrow pp 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2140±30 ²ABELE 99B CBAR 1.94 $\bar{p}p \rightarrow \pi^0 \eta \eta$ 2104±20 ³ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$ ²Spin not determined.³No J^{PC} determination. **$\eta\pi\pi$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2135±20±45 ⁴ADOMEIT 96 CBAR 0 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$ ⁴ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance. **$\bar{p}p \rightarrow \pi\pi$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2090 ⁵OAKDEN 94 RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$ ~ 2120 ⁶OAKDEN 94 RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$ ~ 2170 ⁷MARTIN 80B RVUE~ 2150 ⁷MARTIN 80C RVUE~ 2150 ⁸DULUDE 78B OSPK 1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$ ⁵OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.⁶From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2139^{+8}_{-9}	⁹ EVANGELIS...	97	SPEC	0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
~ 2190	⁹ CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{9,10} COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{9,11} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel

⁹ Isospins 0 and 1 not separated.
¹⁰ From a fit to the total elastic cross section.
¹¹ Referred to as T or \bar{T} region by ALSPECTOR 73.

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2200 ± 13	VLADIMIRSK...06	SPEC 40	$\pi^- p \rightarrow K_S^0 K_S^0 n$
2150 ± 20	ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
2130 ± 35	BARBERIS 99	OMEG 450	$pp \rightarrow p_S p_f K^+ K^-$


$f_2(2150)$ WIDTH

$f_2(2150)$ WIDTH, COMBINED MODES (MeV)

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
152±30 OUR AVERAGE Includes data from the datablock that follows this one. Error includes scale factor of 1.4. See the ideogram below.

• • • We do not use the following data for averages, fits, limits, etc. • • •
 182 ± 11 80k ¹² UMAN 06 E835 $5.2 \bar{p}p \rightarrow \eta\eta\pi^0$

WEIGHTED AVERAGE
 152 ± 30 (Error scaled by 1.4)



¹² Statistical error only.
 $f_2(2150)$ width, combined modes (MeV)

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

152±30 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

280±70	BARBERIS	00E	450 $p p \rightarrow p_f \eta \eta p_s$
150±35	PROKOSHKIN	95D	GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p p \rightarrow p p 2\eta$
130±30	SINGOVSKI	94	GAM4 450 $p p \rightarrow p p 2\eta$


• • • We do not use the following data for averages, fits, limits, etc. • • •

310±50	¹³ ABELE	99B	CBAR 1.94 $\bar{p} p \rightarrow \pi^0 \eta \eta$
203±10	¹⁴ ARMSTRONG	93C	E760 $\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$

¹³ Spin not determined.

¹⁴ No J^{PC} determination.

WEIGHTED AVERAGE
152±30 (Error scaled by 1.4)



$\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

250±25±45	¹⁵ ADOMEIT	96	CBAR 0	1.94 $\bar{p} p \rightarrow \eta 3\pi^0$
-----------	-----------------------	----	--------	--

¹⁵ ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

$\bar{p} p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

250 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 70	16 OAKDEN	94	RVUE	0.36–1.55	$\bar{p}p \rightarrow \pi\pi$
~ 250	17 MARTIN	80B	RVUE		
~ 250	17 MARTIN	80C	RVUE		
~ 250	18 DULUDE	78B	OSPK	1–2	$\bar{p}p \rightarrow \pi^0\pi^0$

¹⁶ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

¹⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

¹⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

56^{+31}_{-16}	¹⁹ EVANGELIS...	97	SPEC	0.6–2.4	$\bar{p}p \rightarrow K_S^0 K_S^0$
135 ± 75	^{20,21} COUPLAND	77	CNTR	0	$0.7\text{--}2.4 \bar{p}p \rightarrow \bar{p}p$
98 ± 8	²¹ ALSPECTOR	73	CNTR		$\bar{p}p$ S channel

¹⁹ Isospin 0 and 2 not separated.

²⁰ From a fit to the total elastic cross section.

²¹ Isospins 0 and 1 not separated.

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

91 ± 62	VLADIMIRSK..06	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$
150 ± 30	ABLIKIM	04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
270 ± 50	BARBERIS	99	OMEG	450 $pp \rightarrow p_S p_f K^+ K^-$

$f_2(2150)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	
Γ_2 $\eta\eta$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $f_2(1270)\eta$	seen
Γ_5 $a_2(1320)\pi$	seen
Γ_6 $p\bar{p}$	seen

$f_2(2150)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\eta\eta)$ Γ_3/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

1.28 ± 0.23 BARBERIS 00E 450 $pp \rightarrow p_f \eta \eta p_S$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	95	²² PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $pp \rightarrow p p 2\eta$
------	----	------------------------------	------	---

²² Using data from ARMSTRONG 89D.

$\Gamma(\pi\pi)/\Gamma(\eta\eta)$ Γ_1/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.33	95	²³ PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p p \rightarrow p p 2\eta$
-------	----	------------------------------	------	--

²³ Derived from a $\pi^0\pi^0/\eta\eta$ limit. $\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$ Γ_4/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.79±0.11	²⁴ ADOMEIT 96	CBAR	1.94 $\bar{p} p \rightarrow \eta 3\pi^0$
------------------	--------------------------	------	--

²⁴ Using $B(a_2(1320) \rightarrow \eta\pi) = 0.145$ $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	73	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
-------------	----	--------------	------	---

 $f_2(2150)$ REFERENCES

ALEXANDER 10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSK... 06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)
	Translated from YAF 69 515.		
ABLIKIM 04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
ANISOVICH 00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS 00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ABELE 99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BARBERIS 99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
EVANGELIS... 97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
MARTIN 97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)
ADOMEIT 96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KLOET 96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
PROKOSHKIN 95D	PD 40 495	Y.D. Prokoshkin	(SERP) IGJPC
	Translated from DANS 344 469.		
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN 94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
SINGOVSKI 94	NC A107 1911	A.V. Singovsky	(SERP)
ARMSTRONG 93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG 89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE 78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)