

**$h_c(1P)$** 

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Quantum numbers are quark model prediction,  $C = -$  established by  $\eta_c \gamma$  decay.

 **$h_c(1P)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3525.38±0.11 OUR AVERAGE</b>				
3525.31±0.11±0.14	832	<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
3525.40±0.13±0.18	3679	ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
3525.20±0.18±0.12	1282	<sup>2</sup> DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3525.8 ±0.2 ±0.2	13	ANDREOTTI	05B E835	$\bar{p} p \rightarrow \eta_c \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3525.6 ±0.5	92 <sup>+23</sup> <sub>-22</sub>	ADAMS	09 CLEO	$\psi(2S) \rightarrow 2(\pi^+ \pi^- \pi^0)$
3524.4 ±0.6 ±0.4	168 ± 40	<sup>3</sup> ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3527 ±8	42	ANTONIAZZI	94 E705	300 $\pi^\pm$ , $p$ Li → $J/\psi \pi^0 X$
3526.28±0.18±0.19	59	<sup>4</sup> ARMSTRONG	92D E760	$\bar{p} p \rightarrow J/\psi \pi^0$
3525.4 ±0.8 ±0.4	5	BAGLIN	86 SPEC	$\bar{p} p \rightarrow J/\psi X$

<sup>1</sup>With floating width.<sup>2</sup>Combination of exclusive and inclusive analyses for the reaction  $\psi(2S) \rightarrow \pi^0 h_c \rightarrow \pi^0 \eta_c \gamma$ . This result is the average of DOBBS 08A and ROSNER 05.<sup>3</sup>Superseded by DOBBS 08A.<sup>4</sup>Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the  $\psi(2S)$  mass from AULCHENKO 03. **$h_c(1P)$  WIDTH**

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.70±0.28±0.22</b>					
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 1.44	90	3679	<sup>2</sup> ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
< 1		13	ANDREOTTI	05B E835	$\bar{p} p \rightarrow \eta_c \gamma$
< 1.1	90	59	ARMSTRONG	92D E760	$\bar{p} p \rightarrow J/\psi \pi^0$

<sup>1</sup>With floating mass.<sup>2</sup>The central value is  $\Gamma = 0.73 \pm 0.45 \pm 0.28$  MeV. **$h_c(1P)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $J/\psi(1S) \pi^0$		
$\Gamma_2$ $J/\psi(1S) \pi \pi$	not seen	
$\Gamma_3$ $J/\psi(1S) \pi^+ \pi^-$	< 2.3 × 10 <sup>-3</sup>	90%
$\Gamma_4$ $p \bar{p}$	< 1.5 × 10 <sup>-4</sup>	90%
$\Gamma_5$ $p \bar{p} \pi^+ \pi^-$	(2.9±0.6) × 10 <sup>-3</sup>	

$\Gamma_6$	$p\bar{p}\pi^0\pi^0$	$< 5 \times 10^{-4}$	90%
$\Gamma_7$	$\pi^+\pi^-\pi^0$	$(1.6\pm 0.5) \times 10^{-3}$	
$\Gamma_8$	$\pi^+\pi^-\pi^0\eta$	$(7.2\pm 2.3) \times 10^{-3}$	
$\Gamma_9$	$2\pi^+2\pi^-\pi^0$	$(8.1\pm 1.8) \times 10^{-3}$	
$\Gamma_{10}$	$3\pi^+3\pi^-\pi^0$	$< 9 \times 10^{-3}$	90%
$\Gamma_{11}$	$K^+K^-\pi^+\pi^-$	$< 6 \times 10^{-4}$	90%
$\Gamma_{12}$	$K^+K^-\pi^+\pi^-\pi^0$	$(3.2\pm 0.8) \times 10^{-3}$	
$\Gamma_{13}$	$K^+K^-\pi^+\pi^-\eta$	$< 2.3 \times 10^{-3}$	90%
$\Gamma_{14}$	$K^+K^-\pi^0$	$< 6 \times 10^{-4}$	90%
$\Gamma_{15}$	$K^+K^-\pi^0\eta$	$< 2.1 \times 10^{-3}$	90%
$\Gamma_{16}$	$K^+K^-\eta$	$< 9 \times 10^{-4}$	90%
$\Gamma_{17}$	$2K^+2K^-\pi^0$	$< 2.4 \times 10^{-4}$	90%
$\Gamma_{18}$	$K_S^0 K^\pm \pi^\mp$	$< 6 \times 10^{-4}$	90%
$\Gamma_{19}$	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(2.8\pm 1.0) \times 10^{-3}$	

### Radiative decays

$\Gamma_{20}$	$\gamma\eta$	$(4.7\pm 2.1) \times 10^{-4}$
$\Gamma_{21}$	$\gamma\eta'(958)$	$(1.5\pm 0.4) \times 10^{-3}$
$\Gamma_{22}$	$\gamma\eta_c(1S)$	$(50 \pm 9) \%$

## $h_c(1P)$ PARTIAL WIDTHS

### $h_c(1P) \Gamma(i)\Gamma(\bar{p}p)/\Gamma(\text{total})$

$\Gamma(\gamma\eta_c(1S)) \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{22}\Gamma_4/\Gamma$

VALUE (eV)      EVTS      DOCUMENT ID      TECN      COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.0±4.5      13      <sup>1</sup> ANDREOTTI 05B E835       $\bar{p}p \rightarrow \eta_c\gamma$

<sup>1</sup> Assuming  $\Gamma = 1$  MeV.

### $h_c(1P)$ BRANCHING RATIOS

$\Gamma(J/\psi(1S)\pi\pi)/\Gamma(J/\psi(1S)\pi^0)$   $\Gamma_2/\Gamma_1$

VALUE      CL%      DOCUMENT ID      TECN      COMMENT

**<0.18**      90      ARMSTRONG 92D E760       $\bar{p}p \rightarrow J/\psi\pi^0$

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$

VALUE      CL%      DOCUMENT ID      TECN      COMMENT

**<2.3 × 10<sup>-3</sup>**      90      <sup>1</sup> ABLIKIM 18M BES3       $\psi(2S) \rightarrow \pi^0\pi^+\pi^- J/\psi$

<sup>1</sup> ABLIKIM 18M reports  $[\Gamma(h_c(1P) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.0 \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-4}$	90	<sup>1</sup> ABLIKIM 13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

<sup>1</sup> ABLIKIM 13V reports  $[\Gamma(h_c(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 1.3 \times 10^{-7}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.6 \pm 0.5 \pm 0.2$		101	<sup>1</sup> ABLIKIM 19AG	BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.2$	90		<sup>2</sup> ADAMS 09	CLEO	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
--------	----	--	-----------------------	------	--

<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (1.38 \pm 0.35 \pm 0.17) \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ADAMS 09 reports  $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 0.19 \times 10^{-5}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .

 $\Gamma(\pi^+\pi^-\pi^0\eta)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$7.2 \pm 2.0 \pm 1.0$	35	<sup>1</sup> ABLIKIM 20AH	BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0\eta)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (6.2 \pm 1.6 \pm 0.7) \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.81 \pm 0.18</math> OUR AVERAGE</b>				
$0.74 \pm 0.14 \pm 0.11$	254	<sup>1</sup> ABLIKIM 19AG	BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
$2.2^{+0.8}_{-0.6} \pm 0.3$	92	<sup>2</sup> ADAMS 09	CLEO	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow 2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (6.40 \pm 0.81 \pm 0.87) \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ADAMS 09 reports  $[\Gamma(h_c(1P) \rightarrow 2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (1.88^{+0.48+0.47}_{-0.45-0.30}) \times 10^{-5}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(3\pi^+3\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9 \times 10^{-3}$	90	<sup>1</sup> ABLIKIM 19AG	BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.029                      90                      <sup>2</sup> ADAMS                      09                      CLEO                       $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$   
<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow 3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)]$   
 $< 7.5 \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .  
<sup>2</sup> ADAMS 09 reports  $[\Gamma(h_c(1P) \rightarrow 3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)]$   
 $< 2.5 \times 10^{-5}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .

### $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.9 \pm 0.5 \pm 0.4</math></b>	230	<sup>1</sup> ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)]$   
 $= (2.49 \pm 0.27 \pm 0.28) \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)$   
 $= (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 5 \times 10^{-4}</math></b>	90	12	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)]$   
 $< 4.4 \times 10^{-7}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .

### $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{11}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 6 \times 10^{-4}</math></b>	90	<sup>1</sup> ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)]$   
 $< 0.5 \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .

### $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.2 \pm 0.7 \pm 0.5</math></b>	80	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)]$   
 $= (2.8 \pm 0.5 \pm 0.3) \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

### $\Gamma(K^+K^-\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ $\Gamma_{13}/\Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>&lt; 2.3 \times 10^{-3}</math></b>	90	24	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow K^+K^-\pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0)]$   
 $< 2.0 \times 10^{-6}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-4}$	90	20	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 4.8 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .					

 $\Gamma(2K^+ 2K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-4}$	90	11	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow 2K^+ 2K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.1 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .					

 $\Gamma(K^+ K^- \pi^0 \eta)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-3}$	90	20	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^0 \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 1.8 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .					

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<9 \times 10^{-4}$	90	18	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 7.5 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .					

 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-4}$	90	17	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 4.8 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 8.6 \times 10^{-4}$ .					

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 0.9 \pm 0.4$	41	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (2.4 \pm 0.7 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

## RADIATIVE DECAYS

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7 \pm 1.5 \pm 1.4$	18	ABLIKIM	16i BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$					$\Gamma_{21}/\Gamma$
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.52±0.27±0.29</b>	44	ABLIKIM 16i	BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta'(958)$	

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$					$\Gamma_{22}/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>50± 9 OUR AVERAGE</b>					
53± 7±8	3679	<sup>1</sup> ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$	
48± 6±7		<sup>2</sup> DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
48± 6±7	1282	<sup>3</sup> DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$	
46±12±7	168	<sup>4</sup> ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$	

<sup>1</sup> ABLIKIM 10B reports  $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.58 \pm 0.40 \pm 0.50) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Average of DOBBS 08A and ROSNER 05. DOBBS 08A reports  $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.16 \pm 0.30 \pm 0.37) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> DOBBS 08A reports  $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.19 \pm 0.32 \pm 0.45) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> ROSNER 05 reports  $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.0 \pm 0.8 \pm 0.7) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (8.6 \pm 1.3) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

## $h_c(1P)$ REFERENCES

ABLIKIM 20AH	PR D102 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 19AG	PR D99 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 18M	PR D97 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 16i	PRL 116 251802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADAMS 09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)
DOBBS 08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ANDREOTTI 05B	PR D72 032001	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ROSNER 05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
AULCHENKO 03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
ANTONIAZZI 94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)
ARMSTRONG 93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ARMSTRONG 92D	PRL 69 2337	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN 86	PL B171 135	C. Baglin <i>et al.</i>	(LAPP, CERN, TORI, STRB+)