

$\phi(2170)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the review on "Spectroscopy of Light Meson Resonances."

 **$\phi(2170)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2162 ± 7</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.1.		
2176 ± 24 ± 3		<sup>1</sup> ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\eta$
2163.5 ± 6.2 ± 3.0		<sup>2</sup> ABLIKIM	21T BES3	$e^+e^- \rightarrow \phi\eta$
2177.5 ± 4.8 ± 19.5		<sup>3</sup> ABLIKIM	20M BES3	$e^+e^- \rightarrow \eta'\phi$
2126.5 ± 16.8 ± 12.4		<sup>4</sup> ABLIKIM	20S BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2273.7 ± 5.7 ± 19.3		<sup>5</sup> ABLIKIM	21AP BES3	$e^+e^- \rightarrow K_S^0 K_L^0$
2135 ± 8 ± 9	95	ABLIKIM	19I BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$
2239.2 ± 7.1 ± 11.3		<sup>6</sup> ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
2200 ± 6 ± 5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
2180 ± 8 ± 8		<sup>7,8</sup> LEES	12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
2079 ± 13 $\begin{smallmatrix} +79 \\ -28 \end{smallmatrix}$	4.8k	<sup>9</sup> SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2186 ± 10 ± 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
2125 ± 22 ± 10	483	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$
2192 ± 14	116	<sup>10</sup> AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2169 ± 20	149	<sup>10</sup> AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
2175 ± 10 ± 15	201	<sup>8,11</sup> AUBERT, BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

<sup>1</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from  $\omega(1420)$  and  $\omega(1650)/\phi(1680)$ .

<sup>2</sup> From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ( $\phi(1680)$  and  $\phi(2170)$ ) and a nonresonant term.

<sup>3</sup> From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

<sup>4</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

<sup>5</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to  $\rho(2150)$ .

<sup>6</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .

<sup>7</sup> Fit includes interference with the  $\phi(1680)$ .

<sup>8</sup> From the  $\phi f_0(980)$  component.


<sup>9</sup> From a fit with two incoherent Breit-Wigners.

<sup>10</sup> From the  $K^+K^- f_0(980)$  component.

<sup>11</sup> Superseded by LEES 12F.

## $\phi(2170)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>100 <math>\begin{smallmatrix} +31 \\ -23 \end{smallmatrix}</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 2.5. See the ideogram below.		
89 $\pm 50 \pm 5$		<sup>1</sup> ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\eta$
31.1 $\begin{smallmatrix} +21.1 \\ -11.6 \end{smallmatrix} \pm 1.1$		<sup>2</sup> ABLIKIM	21T BES3	$e^+e^- \rightarrow \phi\eta$
149.0 $\pm 15.6 \pm 8.9$		<sup>3</sup> ABLIKIM	20M BES3	$e^+e^- \rightarrow \eta'\phi$
106.9 $\pm 32.1 \pm 28.1$		<sup>4</sup> ABLIKIM	20S BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
86 $\pm 44 \pm 51$		<sup>5</sup> ABLIKIM	21AP BES3	$e^+e^- \rightarrow K_S^0 K_L^0$
104 $\pm 24 \pm 12$	95	ABLIKIM	19I BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$
139.8 $\pm 12.3 \pm 20.6$		<sup>6</sup> ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
104 $\pm 15 \pm 15$	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
77 $\pm 15 \pm 10$		<sup>7,8</sup> LEES	12F BABR	10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
192 $\pm 23 \begin{smallmatrix} +25 \\ -61 \end{smallmatrix}$	4.8k	<sup>9</sup> SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
65 $\pm 23 \pm 17$	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
61 $\pm 50 \pm 13$	483	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
71 $\pm 21$	116	<sup>10</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
102 $\pm 27$	149	<sup>10</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
58 $\pm 16 \pm 20$	201	<sup>8,11</sup> AUBERT,BE	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi\pi\gamma$



<sup>1</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from  $\omega(1420)$  and  $\omega(1650)/\phi(1680)$ .

- <sup>2</sup> From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ( $\phi(1680)$  and  $\phi(2170)$ ) and a nonresonant term.
- <sup>3</sup> From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.
- <sup>4</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.
- <sup>5</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to  $\rho(2150)$ .
- <sup>6</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .
- <sup>7</sup> Fit includes interference with the  $\phi(1680)$ .
- <sup>8</sup> From the  $\phi f_0(980)$  component.
- <sup>9</sup> From a fit with two incoherent Breit-Wigners.
- <sup>10</sup> From the  $K^+ K^- f_0(980)$  component.
- <sup>11</sup> Superseded by LEES 12F.
- $\phi(2170)$  WIDTH (MeV)

### $\phi(2170)$ DECAY MODES

	Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$	$e^+ e^-$	seen
$\Gamma_2$	$\phi\eta$	
$\Gamma_3$	$\omega\eta$	
$\Gamma_4$	$\phi\eta'$	
$\Gamma_5$	$\phi\pi\pi$	
$\Gamma_6$	$\phi f_0(980)$	seen
$\Gamma_7$	$K_S^0 K_L^0$	
$\Gamma_8$	$K^+ K^- \pi^+ \pi^-$	
$\Gamma_9$	$K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
$\Gamma_{10}$	$K^+ K^- \pi^0 \pi^0$	
$\Gamma_{11}$	$K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
$\Gamma_{12}$	$K^{*0} K^\pm \pi^\mp$	not seen
$\Gamma_{13}$	$K^*(892)^0 \bar{K}^*(892)^0$	not seen
$\Gamma_{14}$	$K^*(892)^+ K^*(892)^-$	
$\Gamma_{15}$	$K(1460)^+ K^- + \text{c.c.}$	
$\Gamma_{16}$	$K_1(1270)^+ K^- + \text{c.c.}$	
$\Gamma_{17}$	$K_1(1400)^+ K^- + \text{c.c.}$	

### $\phi(2170) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\phi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV) <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

••• We do not use the following data for averages, fits, limits, etc. •••

0.24<sup>+0.12</sup><sub>-0.07</sub> <sup>1</sup> ABLIKIM 21T BES3  $e^+ e^- \rightarrow \phi\eta$

1.7 ± 0.7 ± 1.3      483      AUBERT      08S      BABR      10.6 e<sup>+</sup>e<sup>-</sup> → φηγ

<sup>1</sup> From a solution of the fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes (φ(1680) and φ(2170)) and a nonresonant term. The other solution gives 10.11<sup>+3.87</sup><sub>-3.13</sub> eV.

**Γ(ωη) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>3</sub>Γ<sub>1</sub>/Γ**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>0.43 ± 0.15 ± 0.04</b>	<sup>1</sup> ABLIKIM	21A	BES3 e <sup>+</sup> e <sup>-</sup> → ωη

<sup>1</sup> For constructive interference with ω(1420) and ω(1650)/φ(1680). For destructive interference: 1.25 ± 0.48 ± 0.18 eV.

**Γ(φη') × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>4</sub>Γ<sub>1</sub>/Γ**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>7.1 ± 0.7 ± 0.7</b>	<sup>1</sup> ABLIKIM	20M	BES3 e <sup>+</sup> e <sup>-</sup> → η'φ

<sup>1</sup> From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

**Γ(φf<sub>0</sub>(980)) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>6</sub>Γ<sub>1</sub>/Γ**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ± 0.3 ± 0.3</b>		<sup>1,2</sup> LEES	12F	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → φπ <sup>+</sup> π <sup>-</sup> γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.5 ± 0.8 ± 0.4	201	<sup>2,3</sup> AUBERT, BE	06D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> ππγ
-----------------	-----	---------------------------	-----	---

<sup>1</sup> From a fit with constructive interference with the φ(1680). In a fit with destructive interference, the value is larger by a factor of 12.

<sup>2</sup> From the φf<sub>0</sub>(980) component.

<sup>3</sup> Superseded by LEES 12F.

**Γ(K<sub>S</sub><sup>0</sup>K<sub>L</sub><sup>0</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>7</sub>Γ<sub>1</sub>/Γ**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>0.9 ± 0.6 ± 0.7</b>	<sup>1</sup> ABLIKIM	21AP	BES3 e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>L</sub> <sup>0</sup>

<sup>1</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to ρ(2150).

**Γ(K\*(892)<sup>+</sup>K\*(892)<sup>-</sup>) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>14</sub>Γ<sub>1</sub>/Γ**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.9</b>	90	<sup>1</sup> ABLIKIM	20S	BES3 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>0</sup> π <sup>0</sup>

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

**Γ(K(1460)<sup>+</sup>K<sup>-</sup> + c.c.) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>      Γ<sub>15</sub>Γ<sub>1</sub>/Γ**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>3.0 ± 3.8</b>	<sup>1</sup> ABLIKIM	20S	BES3 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>0</sup> π <sup>0</sup>

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

$$\Gamma(K_1(1270)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{16} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<12.5	90	<sup>1</sup> ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives an upper limit value of 297.6 eV.

$$\Gamma(K_1(1400)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{17} \Gamma_1 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$4.7 \pm 3.3$	<sup>1</sup> ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives a value of  $98.8 \pm 7.8$  eV.

$$\phi(2170) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$$

$$\Gamma(\phi \pi \pi) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_5 / \Gamma \times \Gamma_1 / \Gamma$$

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.65 \pm 0.15 \pm 0.18$	4.8k	<sup>1</sup> SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
--------------------------	------	-------------------	---------	---

<sup>1</sup> Multiplied by 3/2 to take into account the  $\phi \pi^0 \pi^0$  mode. Using  $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$ .

$$\phi(2170) \text{ BRANCHING RATIOS}$$

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_9 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{11} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

$$\Gamma(K^{*0} K^\pm \pi^\mp) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	AUBERT	07AK BABR	$10.6 \text{ GeV } e^+ e^-$

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{13} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

## $\phi(2170)$ REFERENCES

ABLIKIM	21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AP	PR D104 092014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21T	PR D104 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20M	PR D102 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20S	PRL 124 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)

---