

$\psi(4660)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as $Y(4660)$; was $X(4660)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

 $\psi(4660)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4630 ± 6	OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
4651.0 ± 37.8 ± 2.1	1	ABLIKIM 21AJ	BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4619.8 ⁺ ₋ 8.9 ± 8.0 ± 2.3	66	2 JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$
4625.9 ⁺ ₋ 6.2 ± 6.0 ± 0.4	89	3 JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
4652 ± 10 ± 11	279	4 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4669 ± 21 ± 3	37	5 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4634 ⁺ ₋ 8 ± 7 ⁺ ₋ 5 ± 8	142	6 PAKHLOVA 08B	BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4652.5 ± 3.4 ± 1.1	7	DAI 17	RVUE	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
4645.2 ± 9.5 ± 6.0	8	ZHANG 17B	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4646.4 ± 9.7 ± 4.8	9	ZHANG 17C	RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
4661 ⁺ ₋ 9 ± 8 ± 6	44	10 LIU 08H	RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4664 ± 11 ± 5	44	WANG 07D	BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

¹ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.

² Using $D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-$ decays.

³ From a fit of a Breit-Wigner convolved with a Gaussian.

⁴ From a two-resonance fit. Supersedes WANG 07D.

⁵ From a two-resonance fit.

⁶ The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.


⁷ The pole parameters are extracted from the speed plot.

⁸ From a three-resonance fit.

⁹ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

¹⁰ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.


WEIGHTED AVERAGE
 4630 ± 6 (Error scaled by 1.4)



$\psi(4660)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
72^{+14}_{-12}		OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.
$155.4 \pm 24.8 \pm 0.8$		¹ ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$47.0^{+31.3}_{-14.8} \pm 4.6$	66	² JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^-(2573)^-$
$49.8^{+13.9}_{-11.5} \pm 4.0$	89	³ JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}^-(2536)^-$
$68 \pm 11 \pm 5$	279	⁴ WANG	15A BELL	$10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
$104 \pm 48 \pm 10$	37	⁵ LEES	14F BABR	$10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
$92^{+40}_{-24} \pm 10_{-21}$	142	⁶ PAKHLOVA	08B BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$62.6 \pm 5.6 \pm 4.3$		⁷ DAI	17 RVUE	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
$113.8 \pm 18.1 \pm 3.4$		⁸ ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$103.5 \pm 15.6 \pm 4.0$		⁹ ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
$42^{+17}_{-12} \pm 6$	44	¹⁰ LIU	08H RVUE	$10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
$48 \pm 15 \pm 3$	44	WANG	07D BELL	$10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
¹ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.				
² Using $D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-$ decays.				
³ From a fit of a Breit-Wigner convolved with a Gaussian.				
⁴ From a two-resonance fit. Supersedes WANG 07D.				

- ⁵ From a two-resonance fit.
- ⁶ The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+\Lambda_c^-$ states are not necessarily the same.
- ⁷ The pole parameters are extracted from the speed plot.
- ⁸ From a three-resonance fit.
- ⁹ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.
- ¹⁰ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.



$\psi(4660)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	not seen
Γ_2 $\psi(2S)\pi^+\pi^-$	seen
Γ_3 $J/\psi\eta$	not seen
Γ_4 $D^0D^{*-}\pi^+$	not seen
Γ_5 $\chi_{c1}\gamma$	not seen
Γ_6 $\chi_{c2}\gamma$	not seen
Γ_7 $\Lambda_c^+\Lambda_c^-$	seen
Γ_8 $D_s^+D_{s1}(2536)^-$	seen
Γ_9 $D_s^+D_{s2}^*(2573)^-$	

$\psi(4660) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$ $\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_2 \Gamma_1 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.7±3.8		¹ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
11.2±3.2		² ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4.7±4.2		³ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
11.3±3.3		⁴ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
2.0±0.3±0.2	279	⁵ WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
8.1±1.1±1.0	279	⁶ WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.7±1.3±0.5	37	⁷ LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.5±1.7±0.7	37	⁸ LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.2 ^{+0.7} _{-0.6}	44	⁹ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
5.9±1.6	44	¹⁰ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
3.0±0.9±0.3	44	⁷ WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.6±1.8±0.8	44	⁸ WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

¹ Solution I of four equivalent solutions in a fit using three interfering resonances.² Solution II of four equivalent solutions in a fit using three interfering resonances.³ Solution III of four equivalent solutions in a fit using three interfering resonances.⁴ Solution IV of four equivalent solutions in a fit using three interfering resonances.⁵ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.⁶ Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.⁷ Solution I of two equivalent solutions in a fit using two interfering resonances.⁸ Solution II of two equivalent solutions in a fit using two interfering resonances.⁹ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.¹⁰ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances. $\Gamma(J/\psi\eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_3 \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.94	90	WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

 $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_5 \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.45	90	¹ HAN	15 BELL	10.58 $e^+ e^- \rightarrow \chi_{c1} \gamma$

¹ Using $B(\eta \rightarrow \gamma \gamma) = (39.41 \pm 0.21)\%$. $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_6 \Gamma_1 / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	¹ HAN	15 BELL	10.58 $e^+ e^- \rightarrow \chi_{c2} \gamma$

¹ Using $B(\eta \rightarrow \gamma \gamma) = (39.41 \pm 0.21)\%$. $\Gamma(D_s^+ D_{s1}(2536)^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_8 \Gamma_1 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
14.3^{+2.8}_{-2.6} ± 1.5	89	¹ JIA	19A BELL	$e^+ e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$

¹ Assuming $B(D_{s1}(2536)^- \rightarrow \bar{D}^{*0} K^-) = 1$.

$\Gamma(D_s^+ D_{s2}^*(2573)^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_9 \Gamma_1 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$14.7^{+5.9}_{-4.5} \pm 3.6$	66	¹ JIA	20	BELL	$e^+ e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$
¹ Assuming $B(D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-) = 1$.					

 $\psi(4660)$ BRANCHING RATIOS

$\Gamma(D^0 D^{*-} \pi^+) / \Gamma(\psi(2S) \pi^+ \pi^-)$					Γ_4 / Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
< 10	90	PAKHLOVA	09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$
$\Gamma(D^0 D^{*-} \pi^+) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_4 / \Gamma \times \Gamma_1 / \Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 0.37 \times 10^{-6}$	90	¹ PAKHLOVA	09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$
¹ Using $4664 \pm 11 \pm 5$ MeV for the mass of $\psi(4660)$.					

$\Gamma(\Lambda_c^+ \Lambda_c^-) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_7 / \Gamma \times \Gamma_1 / \Gamma$
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.68^{+0.16+0.29}_{-0.15-0.30}$	142	¹ PAKHLOVA	08B	BELL	$e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
¹ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.					

 $\psi(4660)$ REFERENCES

ABLIKIM	21AJ	PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
JIA	20	PR D101 091101	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	19A	PR D100 111103	S. Jia <i>et al.</i>	(BELLE Collab.)
DAI	17	PR D96 116001	L.-Y. Dai, J. Haidenbauer, U.-G. Meissner	(JULI+)
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang	
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan	
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
PAKHLOVA	08B	PRL 101 172001	C. Pakhlova <i>et al.</i>	(BELLE Collab.)
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)