Quark and Lepton Compositeness, Searches for

The latest unpublished results are described in the "Quark and Lepton Compositeness" review.

See the related review(s):

Searches for Quark and Lepton Compositeness

CONTENTS:

- Scale Limits for Contact Interactions: $\Lambda(eeee)$
- Scale Limits for Contact Interactions: $\Lambda(ee\mu\mu)$
- Scale Limits for Contact Interactions: $\Lambda(ee\tau\tau)$
- Scale Limits for Contact Interactions: $\Lambda(\ell\ell\ell\ell)$
- Scale Limits for Contact Interactions: $\Lambda(eeqq)$
- Scale Limits for Contact Interactions: $\Lambda(\mu \mu q q)$
- Scale Limits for Contact Interactions: $\Lambda(\ell\nu\ell\nu)$
- Scale Limits for Contact Interactions: $\Lambda(e \nu q q)$
- Scale Limits for Contact Interactions: $\Lambda(q q q q)$
- Scale Limits for Contact Interactions: $\Lambda(\nu \nu q q)$
- Mass Limits for Excited $e(e^*)$
 - Limits for Excited $e(e^*)$ from Pair Production
 - Limits for Excited e (e*) from Single Production
 - Limits for Excited e (e^*) from $e^+e^- \rightarrow \gamma\gamma$
 - Indirect Limits for Excited e (e*)

Mass Limits for Excited μ (μ *)

- Limits for Excited μ (μ^*) from Pair Production
- Limits for Excited μ (μ^*) from Single Production
- Indirect Limits for Excited μ (μ^*)

Mass Limits for Excited τ (τ^*)

- Limits for Excited τ (τ^*) from Pair Production
- Limits for Excited τ (τ^*) from Single Production

Mass Limits for Excited Neutrino (ν^*)

- Limits for Excited ν (ν^*) from Pair Production
- Limits for Excited ν (ν^*) from Single Production

Mass Limits for Excited $q(q^*)$

- Limits for Excited $q(q^*)$ from Pair Production
- Limits for Excited $q(q^*)$ from Single Production

Mass Limits for Color Sextet Quarks (q_6)

Mass Limits for Color Octet Charged Leptons (ℓ_8)

Mass Limits for Color Octet Neutrinos (ν_8)

Mass Limits for W_8 (Color Octet W Boson)

SCALE LIMITS for Contact Interactions: $\Lambda(eeee)$

Limits are for Λ_{II}^{\pm} only. For other cases, see each reference.

$$\frac{\Lambda_{LL}^{+}(\text{TeV})}{\textbf{>8.3}} \quad \frac{\Lambda_{LL}^{-}(\text{TeV})}{\textbf{>10.3}} \quad \frac{CL\%}{95} \qquad \frac{DOCUMENT~ID}{1} \quad \frac{TECN}{\text{RVUE}} \quad \frac{COMMENT}{E_{cm}} = 192-208~\text{GeV}$$

https://pdg.lbl.gov

Page 1

• • • We do not use the following data for averages, fits, limits, etc. • •

>4.5	>7.0	95	² SCHAEL	07A	ALEP	$E_{\rm cm} = 189 - 209 \; {\rm GeV}$
>5.3	>6.8	95	ABDALLAH	06 C	DLPH	$E_{\rm cm} = 130-207 \; {\rm GeV}$
>4.7	>6.1	95	³ ABBIENDI	04G	OPAL	$E_{\rm cm} = 130-207 \; {\rm GeV}$
>4.3	>4.9	95	ACCIARRI	00 P	L3	$E_{\rm cm} = 130 - 189 \; {\rm GeV}$

 $^{^{1}\,\}mathrm{A}$ combined analysis of the data from ALEPH, DELPHI, L3, and OPAL.

SCALE LIMITS for Contact Interactions: $\Lambda(ee\mu\mu)$

Limits are for Λ^{\pm}_{LL} only. For other cases, see each reference.

$\Lambda_{\it LL}^+({ m TeV})$	$\Lambda_{LL}^-({\sf TeV})$	CL%	DOCUMENT ID		TECN	COMMENT
>6.6	>9.5	95	¹ SCHAEL	07A	ALEP	$E_{\rm cm} = 189 - 209 \; {\rm GeV}$
> 8.5	>3.8	95				$E_{\rm cm} = 130 - 189 {\rm GeV}$
• • • We	e do not us	e the fo	ollowing data for aver	ages,	fits, lim	nits, etc. • • •
>7.3	>7.6	95	ABDALLAH	06 C	DLPH	$E_{\rm cm} = 130 - 207 \; {\rm GeV}$
>8.1	>7.3	95				$E_{\rm cm} = 130-207 {\rm GeV}$
			, ,			

 $^{^1}$ SCHAEL 07A limits are from $R_c,~Q_{FB}^{depl}$, and hadronic cross section measurements. 2 ABBIENDI 04G limits are from $e^+\,e^-\to~\mu\mu$ cross section at $\sqrt{s}=$ 130–207 GeV.

SCALE LIMITS for Contact Interactions: $\Lambda(ee\tau\tau)$

Limits are for Λ_{II}^{\pm} only. For other cases, see each reference.

Λ_{LL}^+ (TeV)	$\Lambda_{LL}^{-}(\text{TeV})$	CL%	DOCUMENT ID		TECN	COMMENT
>7.9	>5.8	95	¹ SCHAEL	07A	ALEP	E _{cm} = 189–209 GeV
>7.9	>4.6	95				$E_{\rm cm} = 130-207 {\rm GeV}$
>4.9	>7.2	95	² ABBIENDI	04 G	OPAL	$E_{\rm cm} = 130-207 \; {\rm GeV}$
• • • W	e do not us	e the fo	ollowing data for ave	rages,	fits, lim	nits, etc. • • •
>5.4	>4.7	95	ACCIARRI	00 P	L3	$E_{\rm cm} = 130 - 189 \; {\rm GeV}$

 $^{^1}$ SCHAEL 07A limits are from R_c , Q_{FB}^{depl} , and hadronic cross section measurements.

SCALE LIMITS for Contact Interactions: $\Lambda(\ell\ell\ell\ell)$

Lepton universality assumed. Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

Λ_{LL}^+ (TeV)	$\Lambda_{LL}^{-}(\text{TeV})$	CL%	DOCUMENT ID	TECI	COMMENT
>7.9	> 10.3	95	¹ SCHAEL 0	7A ALE	P E _{cm} = 189–209 GeV
>9.1	>8.2	95	ABDALLAH 0	6c DLP	H E_{cm}^{em} = 130–207 GeV

 $^{^2}$ SCHAEL 07A limits are from $R_c,~Q_{FB}^{depl},$ and hadronic cross section measurements. 3 ABBIENDI 04G limits are from $e^+\,e^-\to~e^+\,e^-$ cross section at $\sqrt{s}=$ 130–207 GeV.

 $^{^2}$ ABBIENDI 04G limits are from $e^+\,e^-\to~\tau\tau$ cross section at $\sqrt{s}=$ 130–207 GeV.

• • • We do not use the following data for averages, fits, limits, etc. • • •

>7.7	>9.5	95	² ABBIENDI	04 G	OPAL	$E_{\rm cm} = 130-207 \; {\rm GeV}$
			³ BABICH	03	RVUE	•
>9.0	>5.2	95	ACCIARRI	00 P	L3	$E_{cm} = 130 - 189 \text{ GeV}$

 $^{^1}$ SCHAEL 07A limits are from R_c , \mathcal{Q}_{FB}^{depl} , and hadronic cross section measurements.

SCALE LIMITS for Contact Interactions: $\Lambda(eeqq)$

Limits are for Λ_{II}^{\pm} only. For other cases, see each reference.

$\Lambda_{\it LL}^+({ m TeV})$	$\Lambda_{LL}^-({\sf TeV})$	CL%	DOCUMENT ID		TECN	COMMENT
>23.5	>26.1	95	¹ AAD	21Q	ATLS	(e e q q)
>19.5	>24.0	95	² SIRUNYAN	21N	CMS	(eeqq)
>23.5	>26.1	95	³ AAD	20 AP	ATLS	(eeqq)
> 4.5	>12.8	95	⁴ ABRAMOWIC	Z19	ZEUS	(eeqq)
>16.8	>23.9	95	⁵ SIRUNYAN	19 AC	CMS	(eeqq)
>24	>37	95	⁶ AABOUD	17AT	ATLS	(eeqq)
> 8.4	>10.2	95	⁷ ABDALLAH	09	DLPH	(eebb)
> 9.4	>5.6	95	⁸ SCHAEL	07A	ALEP	(eecc)
> 9.4	>4.9	95	⁷ SCHAEL	07A	ALEP	(eebb)
>23.3	>12.5	95	⁹ CHEUNG	01 B	RVUE	(eeuu)
>11.1	>26.4	95	⁹ CHEUNG	01 B	RVUE	(eedd)
>23.3 >11.1	>12.5	95	⁹ CHEUNG	01 B	RVUE	(e e u u)

• • We do not use the following data for averages, fits, limits, etc.

> 7.1	>7.1	95	¹⁰ AAD	21 AU	ATLS	(eebs)
>15.5	>19.5	95	¹¹ AABOUD		ATLS	(eeqq)
>13.5	>18.3	95	¹² KHACHATRY.	15AE	CMS	(eeqq)
>16.4	>20.7	95	¹³ AAD	14 BE	ATLS	(eeqq)
> 9.5	>12.1	95	¹⁴ AAD	13E	ATLS	(eeqq)
>10.1	>9.4	95	¹⁵ AAD	12 AB	ATLS	(eeqq)
> 4.2	>4.0	95	¹⁶ AARON	11 C	H1	(eeqq)
> 3.8	>3.8	95	¹⁷ ABDALLAH	11	DLPH	(eetc)
>12.9	>7.2	95	¹⁸ SCHAEL	07A	ALEP	(eeqq)
> 3.7	>5.9	95	¹⁹ ABULENCIA	06L	CDF	(eeqq)

 $^{^1}$ AAD 21Q limits are from $p\,p$ collisions at $\sqrt{s}=13$ TeV. A frequentist statistical framework is used to remove the prior dependence. 2 SIRUNYAN 21N limits are from $e^+\,e^-$ mass distribution in $p\,p$ collisions at $\sqrt{s}=13$

 $^{^2}$ ABBIENDI 04G limits are from $e^+e^-\to \ell^+\ell^-$ cross section at $\sqrt{s}=130$ –207 GeV. 3 BABICH 03 obtain a bound $-0.175~{\rm TeV}^{-2}<1/\Lambda_{LL}^2<0.095~{\rm TeV}^{-2}$ (95%CL) in a model independent analysis allowing all of Λ_{LL} , Λ_{LR} , Λ_{RL} , Λ_{RR} to coexist.

³ AAD 20AP limits are from e^+e^- mass distribution in pp collisions at $\sqrt{s}=13$ TeV. ⁴ ABRAMOWICZ 19 limits are from Q² spectrum measurements of $e^\pm p \to e^\pm X$. ⁵ SIRUNYAN 19AC limits are from e^+e^- mass distribution in pp collisions at $\sqrt{s}=13$

 $^{^6}$ AABOUD 17AT limits are from pp collisions at $\sqrt{s}=13$ TeV. The quoted limit uses a uniform positive prior in $1/\Lambda^2$.

 $^{^7}$ ABDALLAH 09 and SCHAEL 07A limits are from R_b , A_{FB}^b .

 $^{^8}$ SCHAEL 07A limits are from R_c , Q_{FB}^{depl} , and hadronic cross section measurements.

 $^{^9\,\}mathrm{CHEUNG}$ 01B is an update of BARGER 98E.

- 10 AAD 21AU search for new phenomena in final states with e^+e^- and one or no b-tagged jets in pp collisions at $\sqrt{s}=13$ TeV. The quoted limits assume $g_*^2=4$ π .
- 11 AABOUD 160 limits are from pp collisions at $\sqrt{s}=13$ TeV. The quoted limit uses a uniform positive prior in $1/\Lambda^2$.
- 12 KHACHATRYAN 15AE limit is from e^+e^- mass distribution in pp collisions at $E_{\rm cm}=$
- 13 AAD 14BE limits are from pp collisions at $\sqrt{s}=8$ TeV. The quoted limit uses a uniform positive prior in $1/\Lambda^2$.
- 14 AAD 13E limis are from e^+e^- mass distribution in pp collisions at $E_{\rm cm}=7$ TeV.
- ¹⁵ AAD 12AB limis are from e^+e^- mass distribution in pp collisions at $E_{\rm cm}=7$ TeV.
- 16 AARON 11C limits are from Q^2 spectrum measurements of $e^{\pm} p \rightarrow e^{\pm} X$.
- 17 ABDALLAH 11 limit is from $e^+\,e^-\to~t\,\overline{c}$ cross section. $\varLambda_{LL}=\varLambda_{LR}=\varLambda_{RL}=\varLambda_{RR}$
- $^{18}\,\mathsf{SCHAEL}$ 07A limit assumes quark flavor universality of the contact interactions.
- 19 ABULENCIA 06L limits are from $p\overline{p}$ collisions at $\sqrt{s}=1.96$ TeV.

SCALE LIMITS for Contact Interactions: $\Lambda(\mu \mu q q)$

$\Lambda_{\it LL}^+({ m TeV})$	$\Lambda_{LL}^-({\sf TeV})$	CL%	DOCUMENT ID		TECN	COMMENT
>22.3	>32.7	95	¹ AAD	21Q	ATLS	$(\mu \mu q q)$
>23.3	> 40.0	95	² SIRUNYAN	21N	CMS	$(\mu \mu q q)$
>22.3	>32.7	95	³ AAD	20 AP	ATLS	$(\mu \mu q q)$
>20.4	>30.4	95	⁴ SIRUNYAN	19 AC	CMS	$(\mu \mu q q)$
>20	>30	95	⁵ AABOUD	1 7 AT	ATLS	$(\mu \mu q q)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 8.5	>8.5	95	⁶ AAD	21AU ATLS	$(\mu \mu b s)$
>15.8	>21.8	95	⁷ AABOUD	16∪ ATLS	
>12.0	>15.2	95		Y15AE CMS	$(\mu \mu q q)$
>12.5	>16.7	95		14BE ATLS	$(\mu \mu q q)$
> 9.6	>12.9	95	¹⁰ AAD		$(\mu \mu q q)$ (isosinglet)
> 9.5	>13.1	95	¹¹ CHATRCHY	AN 13K CMS	$(\mu \mu q q)$ (isosinglet)
> 8.0	>7.0	95	¹² AAD	12AB ATLS	$(\mu \mu q q)$ (isosinglet)

- 1 AAD 21Q limits are from pp collisions at $\sqrt{s}=13$ TeV. A frequentist statistical framework is used to remove the prior dependence.
- 2 SIRUNYAN 21N limits are from $\mu^+\mu^-$ mass distribution in pp collisions at $\sqrt{s}=13$ TeV. ³ AAD 20AP limits are from $\mu^+\mu^-$ mass distribution in pp collisions at $\sqrt{s}=13$ TeV.
- ⁴ SIRUNYAN 19AC limits are from $\mu^+\mu^-$ mass distribution in pp collisions at $\sqrt{s}=13$
- 5 AABOUD 17AT limits are from pp collisions at $\sqrt{s}=13$ TeV. The quoted limit uses a uniform positive prior in $1/\Lambda^2$.
- 6 AAD 21AU search for new phenomena in final states with $\mu^+\mu^-$ and one or no b-tagged jets in pp collisions at $\sqrt{s}=13$ TeV. The quoted limits assume $g_s^2=4$ π .
- 7 AABOUD 160 limits are from pp collisions at $\sqrt{s}=13$ TeV. The quoted limit uses a uniform positive prior in $1/\Lambda^2$.
- ⁸ KHACHATRYAN 15AE limit is from $\mu^+\mu^-$ mass distribution in pp collisions at $E_{\rm cm}=$
- 9 AAD 14BE limits are from $p\,p$ collisions at $\sqrt{s}=8$ TeV. The quoted limit uses a uniform positive prior in $1/\Lambda^2$.

Created: 8/11/2022 09:39

 10 AAD 13E limis are from $\mu^+\mu^-$ mass distribution in pp collisions at $E_{\rm cm}=$ 7 TeV.

¹¹ CHATRCHYAN 13K limis are from $\mu^+\mu^-$ mass distribution in pp collisions at $E_{\rm cm}=$ 7 TeV. $^{12}\,\mathrm{AAD}$ 12AB limis are from $\mu^+\,\mu^-$ mass distribution in pp collisions at $E_\mathrm{cm}=$ 7 TeV.

SCALE LIMITS for Contact Interactions: $\Lambda(\ell\nu\ell\nu)$

VALUE (TeV)	CL%	DOCUMENT ID		TECN	COMMENT
>3.10	90	¹ JODIDIO	86	SPEC	$\Lambda_{LR}^{\pm}(u_{\mu} u_{e}\mue)$
• • • We do not use the	following	data for averages	s, fits,	limits, e	etc. • • •
>3.8		² DIAZCRUZ	94	RVUE	$\Lambda_{LL}^+(au u_ aue u_e)$
>8.1		² DIAZCRUZ	94	RVUE	$\Lambda_{LL}^-(au u_ au\mathrm{e} u_\mathrm{e})$
>4.1		³ DIAZCRUZ	94	RVUE	$\Lambda_{LL}^+(au u_ au\mu u_\mu)$
>6.5		³ DIAZCRUZ	94	RVUE	$\Lambda_{LL}^-(au u_{ au}\mu u_{\mu})$

¹ JODIDIO 86 limit is from $\mu^+ \to \overline{\nu}_{\mu} \, e^+ \nu_e$. Chirality invariant interactions $L = (g^2/\Lambda^2)$ $\left[\eta_{LL} \; (\overline{\nu}_{\mu L} \gamma^{\alpha} \mu_{L}) \; (\overline{e}_{L} \gamma_{\alpha} \nu_{e\, L}) \; + \; \eta_{LR} \; (\overline{\nu}_{\mu L} \gamma^{\alpha} \nu_{e\, L} \; (\overline{e}_{R} \gamma_{\alpha} \mu_{R}) \right] \; \text{with} \; g^{2}/4\pi = 1 \; \text{and} \; (\overline{e}_{L} \gamma_{\alpha} \mu_{R}) \;$ $(\eta_{LL},\eta_{LR})=(0,\pm 1)$ are taken. No limits are given for Λ_{LL}^{\pm} with $(\eta_{LL},\eta_{LR})=(\pm 1,0)$. For more general constraints with right-handed neutrinos and chirality nonconserving contact interactions, see their text.

SCALE LIMITS for Contact Interactions: $\Lambda(e\nu qq)$

VALUE (TeV)	CL%	DOCUMENT ID		TECN
>2.81	95	¹ AFFOLDER	011	CDF

¹ AFFOLDER 001 bound is for a scalar interaction $\overline{q}_R q_I \overline{\nu} e_I$.

SCALE LIMITS for Contact Interactions: $\Lambda(qqqq)$

			•		
Λ_{LL}^+ (TeV)	$\Lambda_{LL}^{-}(\text{TeV})$	CL%	DOCUMENT ID	TECN	COMMENT
>13.1 none 17.4-	-29.5 >21.8	95	¹ AABOUD	17AK ATLS	pp dijet angl.
\bullet \bullet We do not	use the following	ng data f	or averages, fits, lim	its, etc. ● ●	•
			² AABOUD	18AV ATLS	$pp ightarrow t \overline{t} t \overline{t}$
>12.8	>17.5	95	³ SIRUNYAN	18DD CMS	<i>pp</i> dijet angl.
>11.5	>14.7	95	⁴ SIRUNYAN	17F CMS	<i>pp</i> dijet angl.
>12.0	>17.5	95	⁵ AAD	16s ATLS	<i>pp</i> dijet angl.
			⁶ AAD	15AR ATLS	$pp ightarrow t \overline{t} t \overline{t}$
			⁷ AAD	15BY ATLS	$pp ightarrow t \overline{t} t \overline{t}$
> 8.1	>12.0	95	⁸ AAD	15L ATLS	<i>pp</i> dijet angl.
> 9.0	>11.7	95	⁹ KHACHATRY		pp dijet angl.
> 5		95	¹⁰ FABBRICHES	I 14 RVUE	$q \overline{q} t \overline{t}$

¹ AABOUD 17AK limit is from dijet angular distribution in pp collisions at $\sqrt{s} = 13$ TeV. u, d, and s quarks are assumed to be composite.

Created: 8/11/2022 09:39

 $^{^2}$ DIAZCRUZ 94 limits are from $\Gamma(au o e
u
u)$ and assume flavor-dependent contact interactions with $\Lambda(\tau \nu_{\tau} e \nu_{e}) \ll \Lambda(\mu \nu_{\mu} e \nu_{e})$.

 $^{^3}$ DIAZCRUZ 94 limits are from $\Gamma(au o \mu
u
u)$ and assume flavor-dependent contact interactions with $\Lambda(\tau \nu_{\tau} \mu \nu_{\mu}) \ll \Lambda(\mu \nu_{\mu} e \nu_{e})$.

- 2 AABOUD 18AV obtain limit on t_R compositeness $2\pi/\Lambda_{RR}^2<1.6~{\rm TeV}^{-2}$ at 95% CL from $t\overline{t}\,t\overline{t}$ production in the pp collisions at $E_{\rm cm}=13~{\rm TeV}.$
- ³ SIRUNYAN 18DD limit is from dijet angular distribution in pp collisions at $\sqrt{s}=13$ TeV.
- ⁴ SIRUNYAN 17F limit is from dijet angular cross sections in pp collisions at $E_{\rm cm}=13$ TeV. All quarks are assumed to be composite.
- ⁵ AAD 16S limit is from dijet angular selections in pp collisions at $E_{\rm cm}=13$ TeV. u,d, and s quarks are assumed to be composite.
- 6 AAD 15AR obtain limit on the t_R compositeness $2\pi/\varLambda_{RR}^2<6.6~{\rm TeV}^{-2}$ at 95% CL from the $t\overline{t}\,t\overline{t}$ production in the $p\,p$ collisions at $E_{\rm cm}=8~{\rm TeV}.$
- 7 AAD 15BY obtain limit on the t_R compositeness $2\pi/\Lambda_{RR}^2 < 15.1~{\rm TeV}^{-2}$ at 95% CL from the $t\overline{t}\,t\overline{t}$ production in the $p\,p$ collisions at $E_{\rm cm}=8~{\rm TeV}.$
- ⁸ AAD 15L limit is from dijet angular distribution in pp collisions at $E_{\rm cm}=8$ TeV. u,d, and s quarks are assumed to be composite.
- ⁹ KHACHATRYAN 15J limit is from dijet angular distribution in pp collisions at $E_{\rm cm}=$ 8 TeV. $u,\,d,\,s,\,c$, and b quarks are assumed to be composite.
- ¹⁰ FABBRICHESI 14 obtain bounds on chromoelectric and chromomagnetic form factors of the top-quark using $pp \to t\bar{t}$ and $p\bar{p} \to t\bar{t}$ cross sections. The quoted limit on the $q\bar{q}t\bar{t}$ contact interaction is derived from their bound on the chromoelectric form factor.

SCALE LIMITS for Contact Interactions: $\Lambda(\nu\nu qq)$

Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

MASS LIMITS for Excited e (e*)

Most e^+e^- experiments assume one-photon or Z exchange. The limits from some e^+e^- experiments which depend on λ have assumed transition couplings which are chirality violating $(\eta_L = \eta_R)$. However they can be interpreted as limits for chirality-conserving interactions after multiplying the coupling value λ by $\sqrt{2}$; see Note.

Excited leptons have the same quantum numbers as other ortholeptons. See also the searches for ortholeptons in the "Searches for Heavy Leptons" section.

Limits for Excited e (e*) from Pair Production

These limits are obtained from $e^+e^- \to e^{*+}e^{*-}$ and thus rely only on the (electroweak) charge of e^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the e^* coupling is assumed to be of sequential type. Possible t channel contribution from transition magnetic coupling is neglected. All limits assume a dominant $e^* \to e\gamma$ decay except the limits from $\Gamma(Z)$.

Created: 8/11/2022 09:39

For limits prior to 1987, see our 1992 edition (Physical Review D45 S1 (1992)).

VALUE (GeV) CL% DOCUMENT ID TECN COMMENT
$$^{\prime}$$
 >103.2 95 1 ABBIENDI 02G OPAL $^{\prime}$ $^{\prime}$

¹ MCFARLAND 98 assumed a flavor universal interaction. Neutrinos were mostly of muon type.

• • • We do not use the following data for averages, fits, limits, etc. • • •

>102.8 95 2 ACHARD 03B L3 $e^+e^-
ightarrow e^*e^*$ Homodoublet type

Limits for Excited e (e*) from Single Production

These limits are from $e^+e^- \to e^*e$, $W \to e^*\nu$, or $ep \to e^*X$ and depend on transition magnetic coupling between e and e^* . All limits assume $e^* \to e\gamma$ decay except as noted. Limits from LEP, UA2, and H1 are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda - m_{e^*}$ plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review D45 S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>5600	95	¹ SIRUNYAN	20AJ CMS	$pp \rightarrow ee^*X$
• • • We do not	use the followin	g data for average	es, fits, limits,	etc. • • •
>4800	95	² AABOUD	19AZ ATLS	$pp \rightarrow ee^*X$
>3900	95	³ SIRUNYAN	19z CMS	$pp \rightarrow ee^*X$
>2450	95	⁴ KHACHATRY	'16AQ CMS	$pp \rightarrow ee^*X$
>3000	95	⁵ AAD	15AP ATLS	$p p ightarrow e^{(*)} e^* X$
>2200	95	⁶ AAD	13BB ATLS	$pp \rightarrow ee^*X$
>1900	95	⁷ CHATRCHYA		$pp \rightarrow ee^*X$
>1870	95	⁸ AAD	12AZ ATLS	$p p ightarrow e^{(*)} e^* X$

- ¹ SIRUNYAN 20AJ search for e^* production in 2e2j final states in pp collisions at $\sqrt{s}=13$ TeV. The quoted limit assumes $\Lambda=m_{e^*},\ f=f'=1$. The contact interaction is included. See their Fig.11 for exclusion limits in $m_{e^*}-\Lambda$ plane.
- ² AABOUD 19AZ search for single e^* production in pp collisions at $\sqrt{s}=13$ TeV. The limit quoted above is from $e^* \to e \, q \, \overline{q}$ and $e^* \to \nu \, W$ decays assuming f=f'=1 and $m_{e^*}=\Lambda$. The contact interaction is included in e^* production and decay amplitudes. See their Fig.6 for exclusion limits in $m_{e^*}-\Lambda$ plane.
- ³ SIRUNYAN 19Z search for e^* production in $\ell\ell\gamma$ final states in pp collisions at $\sqrt{s}=13$ TeV. The quoted limit assumes $\Lambda=m_{e^*}$, f=f'=1. The contact interaction is included in the e^* production and decay amplitudes.
- ⁴ KHACHATRYAN 16AQ search for single e^* production in pp collisions at $\sqrt{s}=8$ TeV. The limit above is from the $e^* \to e\gamma$ search channel assuming f=f'=1, $m_{e^*}=\Lambda$. See their Table 7 for limits in other search channels or with different assumptions.
- ⁵ AAD 15AP search for e^* production in evens with three or more charged leptons in pp collisions at $\sqrt{s}=8$ TeV. The quoted limit assumes $\Lambda=m_{e^*}$, f=f'=1. The contact interaction is included in the e^* production and decay amplitudes.
- ⁶ AAD 13BB search for single e^* production in pp collisions with $e^* \to e\gamma$ decay. f = f' = 1, and e^* production via contact interaction with $\Lambda = m_{a^*}$ are assumed.
- ⁷ CHATRCHYAN 13AE search for single e^* production in pp collisions with $e^* \to e\gamma$ decay. f = f' = 1, and e^* production via contact interaction with $\Lambda = m_{e^*}$ are assumed.
- ⁸ AAD 12AZ search for e^* production via four-fermion contact interaction in pp collisions with $e^* \to e\gamma$ decay. The quoted limit assumes $\Lambda = m_{e^*}$. See their Fig. 8 for the exclusion plot in the mass-coupling plane.

¹ From e^+e^- collisions at $\sqrt{s}=183-209$ GeV. f=f' is assumed.

² From e^+e^- collisions at $\sqrt{s}=189$ –209 GeV. f=f' is assumed. ACHARD 03B also obtain limit for f=-f': $m_{e^*}>96.6$ GeV.

Limits for Excited $e(e^*)$ from $e^+e^- \rightarrow \gamma\gamma$

These limits are derived from indirect effects due to e^* exchange in the t channel and depend on transition magnetic coupling between e and e^* . All limits are for $\lambda_{\gamma}=1$. All limits except ABE 89J and ACHARD 02D are for nonchiral coupling with $\eta_{L}=\eta_{R}=1$. We choose the chiral coupling limit as the best limit and list it in the Summary Table.

For limits prior to 1987, see our 1992 edition (Physical Review D45 S1 (1992)).

•	•		`		` //	
VALUE (GeV)	CL%	DOCUMENT ID		TECN	COMMENT	
>356	95	$^{ m 1}$ ABDALLAH	04N	DLPH	\sqrt{s} = 161–208 GeV	
• • • We do not use the	ne following	data for averages	s, fits,	limits, e	etc. • • •	
>310	95	ACHARD	02 D	L3	\sqrt{s} = 192–209 GeV	

¹ ABDALLAH 04N also obtain a limit on the excited electron mass with ee^* chiral coupling, $m_{a^*} > 295$ GeV at 95% CL.

Indirect Limits for Excited e (e*)

These limits make use of loop effects involving e^* and are therefore subject to theoretical uncertainty.

MASS LIMITS for Excited μ (μ^*)

Limits for Excited μ (μ *) from Pair Production

These limits are obtained from $e^+e^- \to \mu^{*+}\mu^{*-}$ and thus rely only on the (electroweak) charge of μ^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the μ^* coupling is assumed to be of sequential type. All limits assume a dominant $\mu^* \to \mu \gamma$ decay except the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review D45 S1 (1992)).

$$>$$
102.8 95 2 ACHARD 03B L3 $e^+e^-
ightarrow \, \mu^*\mu^*$ Homodoublet type

 $^{^1}$ DORENBOSCH 89 obtain the limit $\lambda_{\gamma}^2\Lambda_{\rm cut}^2/m_{e^*}^2<2.6$ (95% CL), where $\Lambda_{\rm cut}$ is the cutoff scale, based on the one-loop calculation by GRIFOLS 86. If one assumes that $\Lambda_{\rm cut}=1$ TeV and $\lambda_{\gamma}=1$, one obtains $m_{e^*}>620$ GeV. However, one generally expects

 $[\]lambda_{\gamma} \approx m_{e^*}/\Lambda_{\rm cut}$ in composite models. 2 GRIFOLS 86 uses $\nu_{\mu}\,e \to \nu_{\mu}\,e$ and $\overline{\nu}_{\mu}\,e \to \overline{\nu}_{\mu}\,e$ data from CHARM Collaboration to derive mass limits which depend on the scale of compositeness.

³ RENARD 82 derived from g-2 data limits on mass and couplings of e^* and μ^* . See figures 2 and 3 of the paper.

¹ From e^+e^- collisions at $\sqrt{s}=183$ –209 GeV. f=f' is assumed.

 $^{^2}$ From e^+e^- collisions at $\sqrt{s}=$ 189–209 GeV. f=f' is assumed. ACHARD 03B also obtain limit for $f=-f'\colon m_{\mu^*}>$ 96.6 GeV.

Limits for Excited μ (μ *) from Single Production

These limits are from $e^+e^- \to \mu^*\mu$ and depend on transition magnetic coupling between μ and μ^* . All limits assume $\mu^* \to \mu\gamma$ decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda - m_{\mu^*}$ plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review D45 S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>5700	95	¹ SIRUNYAN 20A	AJ CMS	$pp \rightarrow \mu \mu^* X$
• • • We do not use th	e followin	g data for averages, fit	s, limits,	etc. • • •
>3800	95	² SIRUNYAN 197		
>2800	95			$pp ightarrow \mu \mu^* X$
>2470	95	⁴ KHACHATRY16		
>3000	95		AP ATLS	$pp o \mu^{ig(st)} \mu^st X$
>2200	95			$pp \rightarrow \mu \mu^* X$
>1900	95	⁷ CHATRCHYAN 13A		
>1750	95	⁸ AAD 12A	AZ ATLS	$p p o \mu^{ig(st)} \mu^st X$

- 1 SIRUNYAN 20AJ search for μ^* production in $2\mu 2j$ final states in $p\,p$ collisions at $\sqrt{s}=13$ TeV. The quoted limit assumes $\Lambda=m_{\mu^*}$, f=f'=1. The contact interaction is included. See their Fig.11 for exclusion limits in $m_{\mu^*}-\Lambda$ plane.
- ² SIRUNYAN 19Z search for μ^* production in $\ell\ell\gamma$ final states in pp collisions at $\sqrt{s}=13$ TeV. The quoted limit assumes $\Lambda=m_{\mu^*}$, f=f'=1. The contact interaction is included in the μ^* production and decay amplitudes.
- ³ AAD 16BM search for μ^* production in $\mu\mu jj$ events in pp collisions at $\sqrt{s}=8$ TeV. Both the production and decay are assumed to occur via a contact interaction with $\Lambda=m_{\mu^*}$.
- ⁴ KHACHATRYAN 16AQ search for single μ^* production in pp collisions at $\sqrt{s}=8$ TeV. The limit above is from the $\mu^*\to\mu\gamma$ search channel assuming f=f'=1, $m_{\mu^*}=\Lambda$. See their Table 7 for limits in other search channels or with different assumptions.
- ⁵ AAD 15AP search for μ^* production in evens with three or more charged leptons in pp collisions at $\sqrt{s}=8$ TeV. The quoted limit assumes $\Lambda=m_{\mu^*}$, f=f'=1. The contact interaction is included in the μ^* production and decay amplitudes.
- ⁶ AAD 13BB search for single μ^* production in pp collisions with $\mu^* \to \mu \gamma$ decay. f = f' = 1, and μ^* production via contact interaction with $\Lambda = m_{\mu^*}$ are assumed.
- ⁷ CHATRCHYAN 13AE search for single μ^* production in pp collisions with $\mu^* \to \mu \gamma$ decay. f = f' = 1, and μ^* production via contact interaction with $\Lambda = m_{\mu^*}$ are assumed.
- ⁸ AAD 12AZ search for μ^* production via four-fermion contact interaction in pp collisions with $\mu^* \to \mu \gamma$ decay. The quoted limit assumes $\Lambda = m_{\mu^*}$. See their Fig. 8 for the exclusion plot in the mass-coupling plane.

Indirect Limits for Excited μ (μ *)

These limits make use of loop effects involving μ^* and are therefore subject to theoretical uncertainty.

Created: 8/11/2022 09:39

 VALUE (GeV)
 DOCUMENT ID
 TECN
 COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • •

¹ RENARD 82 THEO
$$g-2$$
 of muon

MASS LIMITS for Excited au (au^*)

Limits for Excited τ (τ^*) from Pair Production

These limits are obtained from $e^+e^- \to \tau^{*+}\tau^{*-}$ and thus rely only on the (electroweak) charge of τ^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the τ^* coupling is assumed to be of sequential type. All limits assume a dominant $\tau^* \to \tau \gamma$ decay except the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review D45 S1 (1992)).

VALUE (GeV)	CL%	DOCUMENT ID		TECN	COMMENT
>103.2	95	¹ ABBIENDI	02G	OPAL	$e^+e^- ightarrow au^* au^*$ Homodoublet type
• • • \//o c	la not use	the following data	for a	/oragoc	fits limits etc.

• • We do not use the following data for averages, fits, limits, etc.

$$>$$
102.8 95 2 ACHARD 03B L3 $e^+e^-
ightarrow~ au^* au^*$ Homodoublet type

Limits for Excited τ (τ^*) from Single Production

These limits are from $e^+e^- \to \tau^*\tau$ and depend on transition magnetic coupling between τ and τ^* . All limits assume $\tau^* \to \tau\gamma$ decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda - m_{\tau^*}$ plane. See the original papers.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>2500	95	¹ AAD	15AP ATLS	$pp \rightarrow \tau^{(*)}\tau^*X$

ullet ullet We do not use the following data for averages, fits, limits, etc. ullet ullet

> 180	95	² ACHARD	03 B L3	$e^+e^- ightarrow au au^*$
> 185	95	³ ABBIENDI	02G OPAL	$e^+e^- ightarrow au au^*$

¹ AAD 15AP search for τ^* production in events with three or more charged leptons in pp collisions at $\sqrt{s}=8$ TeV. The quoted limit assumes $\Lambda=m_{\tau^*}$, f=f'=1. The contact interaction is included in the τ^* production and decay amplitudes.

¹ RENARD 82 derived from g-2 data limits on mass and couplings of e^* and μ^* . See figures 2 and 3 of the paper.

 $^{^{1}}$ From $e^{+}e^{-}$ collisions at $\sqrt{s}=$ 183–209 GeV. f=f' is assumed.

 $^{^2}$ From $e^+\,e^-$ collisions at $\sqrt{s}=189$ –209 GeV. f=f' is assumed. ACHARD 03B also obtain limit for $f=-f'\colon m_{\tau^*}>96.6$ GeV.

² ACHARD 03B result is from e^+e^- collisions at $\sqrt{s}=189$ –209 GeV. $f=f'=\Lambda/m_{\tau^*}$ is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane.

³ ABBIENDI 02G result is from e^+e^- collisions at $\sqrt{s}=183$ –209 GeV. $f=f'=\Lambda/m_{\tau^*}$ is assumed for τ^* coupling. See their Fig. 4c for the exclusion limit in the mass-coupling plane.

MASS LIMITS for Excited Neutrino (ν^*)

Limits for Excited ν (ν^*) from Pair Production

> 102.6

95

These limits are obtained from $e^+e^- \to \nu^* \nu^*$ and thus rely only on the (electroweak) charge of ν^* . Form factor effects are ignored unless noted. The ν^* coupling is assumed to be of sequential type unless otherwise noted. All limits assume a dominant $\nu^* \to \nu \gamma$ decay except the limits from $\Gamma(Z)$.

VALUE (GeV)CL%DOCUMENT IDTECNCOMMENT>160095 1 AAD15AP ATLS $pp \rightarrow \nu^* \nu^* X$

• • • We do not use the following data for averages, fits, limits, etc. • •

 2 ABBIENDI 04N OPAL 3 ACHARD 03B L3 ${
m e^+e^-}
ightarrow
u^*
u^*$ Homodoublet type

 1 AAD 15AP search for ν^* pair production in evens with three or more charged leptons in pp collisions at $\sqrt{s}=8$ TeV. The quoted limit assumes $\Lambda=m_{\nu^*}$, f=f'=1. The contact interaction is included in the ν^* production and decay amplitudes.

 2 From e^+e^- collisions at $\sqrt{s}=192$ –209 GeV, ABBIENDI 04N obtain limit on $\sigma(e^+e^-\to\nu^*\nu^*)$ B $^2(\nu^*\to\nu\gamma)$. See their Fig.2. The limit ranges from 20 to 45 fb for $m_{\nu^*}>$ 45 GeV.

 3 From $e^+\,e^-$ collisions at $\sqrt{s}=189$ –209 GeV. f=-f' is assumed. ACHARD 03B also obtain limit for $f=f'\colon$ $m_{\nu_e^*}>101.7$ GeV, $m_{\nu_\mu^*}>101.8$ GeV, and $m_{\nu_\tau^*}>92.9$ GeV.

See their Fig. 4 for the exclusion plot in the mass-coupling plane.

Limits for Excited ν (ν^*) from Single Production

These limits are from $e^+e^- \to \nu\nu^*$, $Z \to \nu\nu^*$, or $ep \to \nu^*X$ and depend on transition magnetic coupling between ν/e and ν^* . Assumptions about ν^* decay mode are given in footnotes.

VALUE (GeV)CL%DOCUMENT IDTECNCOMMENT>21395 1 AARON08H1 $ep \rightarrow \nu^* X$ • • • We do not use the following data for averages, fits, limits, etc. • • •>19095 2 ACHARD03BL3 $e^+e^- \rightarrow \nu\nu^*$

none 50–150 95 3 ADLOFF 02 H1 $e\,p
ightarrow \,
upper 2$ X > 158 95 4 CHEKANOV 02D ZEUS $e\,p
ightarrow \,
upper 2$ X

¹ AARON 08 search for single ν^* production in ep collisions with the decays $\nu^* \to \nu \gamma$, νZ , eW. The quoted limit assumes $f = -f' = \Lambda/m_{\nu^*}$. See their Fig. 3 and Fig. 4 for the exclusion plots in the mass-coupling plane.

² ACHARD 03B result is from e^+e^- collisions at $\sqrt{s}=189$ –209 GeV. The quoted limit is for ν_e^* . $f=-f'=\Lambda/m_{\nu^*}$ is assumed. See their Fig. 4 for the exclusion plot in the mass-coupling plane.

³ ADLOFF 02 search for single ν^* production in ep collisions with the decays $\nu^* \to \nu \gamma$, νZ , eW. The quoted limit assumes $f = -f' = \Lambda/m_{\nu^*}$. See their Fig. 1 for the exclusion plots in the mass-coupling plane.

⁴ CHEKANOV 02D search for single ν^* production in $e\,p$ collisions with the decays $\nu^* \to \nu\gamma, \ \nu Z, \ e\,W. \ f = -f' = \Lambda/m_{\nu^*}$ is assumed for the e^* coupling. CHEKANOV 02D also obtain limit for $f = f' = \Lambda/m_{\nu^*}$: $m_{\nu^*} > 135$ GeV. See their Fig. 5c and Fig. 5d for the exclusion plot in the mass-coupling plane.

MASS LIMITS for Excited $q(q^*)$

Limits for Excited $q(q^*)$ from Pair Production

These limits are mostly obtained from $e^+e^- \to q^* \overline{q}^*$ and thus rely only on the (electroweak) charge of the q^* . Form factor effects are ignored unless noted. Assumptions about the q^* decay are given in the comments and footnotes.

<i>VALUE</i> (GeV)	<u>CL%</u>	DOCUMENT ID		TECN	COMMENT
>338	95	$^{ m 1}$ AALTONEN	10H	CDF	$q^* o tW^-$
• • • We do not us	se the following	ng data for average	es, fits	s, limits,	etc. • • •
none 700-1200	95	² SIRUNYAN	18V	CMS	$pp ightarrow \ t_{3/2}^* \overline{t}_{3/2}^* ightarrow$
					ttgg
		³ BARATE	98 U	ALEP	$Z \rightarrow q^* q^*$
> 45.6	95				u or d type, $Z \rightarrow q^*q^*$
> 41.7	95	⁵ BARDADIN	92	RVUE	u -type, $\Gamma(Z)$
> 44.7	95	⁵ BARDADIN	92	RVUE	d -type, $\Gamma(Z)$
> 40.6	95	⁶ DECAMP	92	ALEP	u -type, $\Gamma(Z)$
> 44.2	95	⁶ DECAMP	92	ALEP	d -type, $\Gamma(Z)$
> 45	95	⁷ DECAMP	92	ALEP	u or d type, $Z ightarrow q^* q^*$
> 45	95	⁶ ABREU	91F	DLPH	u -type, $\Gamma(Z)$
> 45	95	⁶ ABREU	91F	DLPH	d -type, $\Gamma(Z)$

¹ AALTONEN 10H obtain limits on the q^*q^* production cross section in $p\overline{p}$ collisions. See their Fig. 3.

Limits for Excited $q(q^*)$ from Single Production

These limits are from $e^+e^- \to q^*\overline{q}$, $p\overline{p} \to q^*X$, or $pp \to q^*X$ and depend on transition magnetic couplings between q and q^* . Assumptions about q^* decay mode are given in the footnotes and comments.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>6700 (CL = 95°	%) OUR	LIMIT		
none 2000-6700	95	¹ AAD	20T ATLS	$pp \rightarrow q^*X, q^* \rightarrow qg$
none 1250-3200	95	¹ AAD	20T ATLS	$pp \rightarrow b^* X, b^* \rightarrow bg, b\gamma, bZ, tW$
none 1800-6300	95	² SIRUNYAN	20AI CMS	$pp \rightarrow q^*X, q^* \rightarrow qg$
none 1500-2600	95	³ AABOUD	18AB ATLS	$pp ightarrow \ b^* X, \ b^* ightarrow \ bg$
none 1500-5300	95	⁴ AABOUD	18BA ATLS	$pp ightarrow \ q^* X$, $q^* ightarrow \ q \gamma$
none 1000-5500	95	⁵ SIRUNYAN	18AG CMS	$pp ightarrow \ q^* X$, $q^* ightarrow \ q \gamma$
none 1000-1800	95	⁶ SIRUNYAN	18AG CMS	$pp ightarrow \ b^* X$, $b^* ightarrow \ b \gamma$
none 600-6000	95	⁷ SIRUNYAN	18BO CMS	$pp ightarrow \ q^* X, \ q^* ightarrow \ qg$
none 1200-5000	95	⁸ SIRUNYAN	18P CMS	$pp ightarrow q^* X$, $q^* ightarrow q W$
none 1200-4700	95	⁸ SIRUNYAN	18P CMS	$pp ightarrow \ q^* X, \ q^* ightarrow \ q Z$
>6000	95	⁹ AABOUD	17AK ATLS	$pp \rightarrow q^*X, q^* \rightarrow qg$

² SIRUNYAN 18V search for pair production of spin 3/2 excited top quarks. B($t_{3/2}^* \rightarrow t_g$) = 1 is assumed.

³ BARATE 98U obtain limits on the form factor. See their Fig. 16 for limits in mass-form factor plane.

⁴ ADRIANI 93M limit is valid for B($q^* \rightarrow qg$)> 0.25 (0.17) for up (down) type.

 $^{^{5}}$ BARDADIN-OTWINOWSKA 92 limit based on $\Delta\Gamma(Z)$ <36 MeV.

⁶ These limits are independent of decay modes.

⁷ Limit is for B($q^* \rightarrow qg$)+B($q^* \rightarrow q\gamma$)=1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

```
<sup>10</sup> SIRUNYAN
                                                       21AG CMS
                                                                        pp \rightarrow b^* X, b^* \rightarrow tW
>2600
                      95
                                 ^{11} KHACHATRY...17W CMS pp 
ightarrow q^* X, q^* 
ightarrow qg
                      95
none 600-5400
                                 <sup>12</sup> AABOUD
                                                       16 ATLS pp \rightarrow b^*X, b^* \rightarrow bg
none 1100-2100
                      95
                                 <sup>13</sup> AAD
                      95
                                                       16AH ATLS pp \rightarrow b^*X, b^* \rightarrow tW
>1500
                                 <sup>14</sup> AAD
>4400
                      95
                                                       16AI ATLS pp \rightarrow q^*X, q^* \rightarrow q\gamma
                                 15 AAD
                                                     16AV ATLS pp \rightarrow q^*X, q^* \rightarrow Wb
                                 <sup>16</sup> AAD
                                                       16S ATLS pp \rightarrow q^*X, q^* \rightarrow qg
>5200
                      95
                                 <sup>17</sup> KHACHATRY...161 CMS
                      95
                                                                       pp \rightarrow b^* X, b^* \rightarrow tW
>1390
                                 <sup>18</sup> KHACHATRY...16K CMS
>5000
                      95
                                                                       pp \rightarrow q^* X, q^* \rightarrow qg
                                 <sup>19</sup> KHACHATRY...16L CMS
                      95
                                                                       pp \rightarrow q^*X, q^* \rightarrow qg
none 500-1600
>4060
                      95
                                                       15V ATLS pp \rightarrow q^*X, q^* \rightarrow qg
                                 <sup>21</sup> KHACHATRY...15V CMS
>3500
                      95
                                                                       pp \rightarrow q^* X, q^* \rightarrow qg
                                                14A ATLS pp \rightarrow q^*X, q^* \rightarrow q\gamma
>3500
                      95
                                 <sup>23</sup> KHACHATRY...14 CMS
                                                                       pp \rightarrow q^*X, q^* \rightarrow qW
                      95
>3200
                                 <sup>24</sup> KHACHATRY...14 CMS pp \rightarrow q^*X, q^* \rightarrow qZ
                      95
>2900
                                 <sup>25</sup> KHACHATRY...14」 CMS
                                                                       pp \rightarrow q^* X, q^* \rightarrow q\gamma
                      95
none 700-3500
                                 ^{26} CHATRCHYAN 13AJ CMS pp 
ightarrow q^* X, \; q^* 
ightarrow q W
>2380
                      95
                      95
                                 <sup>27</sup> CHATRCHYAN 13AJ CMS pp \rightarrow q^*X, q^* \rightarrow qZ
>2150
```

- 1 AAD 20T search for resonances decaying into dijets in $p\,p$ collisions at $\sqrt{s}=13$ TeV. Assume $\Lambda=m_{g^*}$, $f_{\rm S}=f=f'=1.$
- ² SIRUNYAN 20Al search for resonances decaying into dijets in pp collisions at $\sqrt{s}=13$ TeV. Assume $\Lambda=m_{a^*}$, $f_s=f=f'=1$.
- ³ AABOUD 18AB assume $\Lambda = m_{b^*}$, $f_s = f = f' = 1$. The contact interactions are not included in b^* production and decay amplitudes.
- ⁴ AABOUD 18BA search for first-generation excited quarks (u^* and d^*) with degenerate mass, assuming $\Lambda = m_{q^*}$, $f_S = f = f' = 1$. The contact interactions are not included in q^* production and decay amplitudes.
- ⁵ SIRUNYAN 18AG search for first-generation excited quarks (u^* and d^*) with degenerate mass, assuming $\Lambda=m_{\sigma^*}$, $f_{\rm S}=f=f'=1$.
- ⁶ SIRUNYAN 18AG search for excited b quark assuming $\Lambda = m_{q^*}$, $f_s = f = f' = 1$.
- 7 SIRUNYAN 18BO assume $\Lambda=m_{q^*}$, $f_s=f=f'=1$. The contact interactions are not included in q^* production and decay amplitudes.
- ⁸ SIRUNYAN 18P use the hadronic decay of W or Z, assuming $\Lambda=m_{q^*}$, $f_{S}=f=f'=1$.
- ⁹ AABOUD 17AK assume $\Lambda=m_{q^*}$, $f_S=f=f'=1$. The contact interactions are not included in q^* production and decay amplitudes. Only the decay of $q^*\to g\,u$ and $q^*\to g\,d$ is simulated as the benchmark signals in the analysis.
- ¹⁰ SIRUNYAN 21AG search for b^* decaying to tW in pp collisions at $\sqrt{s}=13$ TeV. The limit quoted above assumes $\kappa_L^b=g_L=1, \ \kappa_R^b=g_R=0$. The limit becomes $m_{b^*}>2.8$ TeV (> 3.1 TeV) if we assume $\kappa_L^b=g_L=0, \ \kappa_R^b=g_R=1$ ($\kappa_L^b=g_L=\kappa_R^b=g_R=1$). See their Fig. 5 for limits on $\sigma \cdot B$.
- ¹¹ KHACHATRYAN 17W assume $\Lambda = m_{q^*}$, $f_s = f = f' = 1$. The contact interactions are not included in q^* production and decay amplitudes.
- ¹² AABOUD 16 assume $\Lambda = m_{b^*}$, $f_s = f = f' = 1$. The contact interactions are not included in the b^* production and decay amplitudes.

- ¹³ AAD 16AH search for b^* decaying to tW in pp collisions at $\sqrt{s}=8$ TeV. $f_g=f_L=f_R=1$ are assumed. See their Fig. 12b for limits on $\sigma \cdot B$.
- 14 AAD 16AI assume $\Lambda=m_{a^*}$, $f_{s}=f=f'=1$.
- 15 AAD 16AV search for single production of vector-like quarks decaying to $W\,b$ in $p\,p$ collisions. See their Fig. 8 for the limits on couplings and mixings.
- 16 AAD 16S assume $\Lambda=m_{q^*}$, $f_s=f=f'=1$. The contact interactions are not included in q^* production and decay amplitudes.
- ¹⁷ KHACHATRYAN 16I search for b^* decaying to tW in pp collisions at $\sqrt{s}=8$ TeV. $\kappa_L^b=g_L=1,\ \kappa_R^b=g_R=0$ are assumed. See their Fig. 8 for limits on $\sigma\cdot B$.
- 18 KHACHATRYAN 16K assume $\Lambda=m_{q^*}$, $f_s=f=f'=1$. The contact interactions are not included in q^* production and decay amplitudes.
- ¹⁹ KHACHATRYAN 16L search for resonances decaying to dijets in pp collisions at $\sqrt{s}=8$ TeV using the data scouting technique which increases the sensitivity to the low mass resonances.
- ²⁰ AAD 15V assume $\Lambda=m_{q^*}$, $f_s=f=f'=1$. The contact interactions are not included in q^* production and decay amplitudes.
- ²¹ KHACHATRYAN 15V assume $\Lambda=m_{q^*}$, $f_s=f=f'=1$. The contact interactions are not included in q^* production and decay amplitudes.
- ²² AAD 14A assume $\Lambda = m_{a^*}$, $f_s = f = f' = 1$.
- ²³ KHACHATRYAN 14 use the hadronic decay of W, assuming $\Lambda = m_{\sigma^*}$, $f_s = f = f' = 1$.
- ²⁴ KHACHATRYAN 14 use the hadronic decay of Z, assuming $\Lambda = m_{a^*}$, $f_s = f = f' = 1$.
- 25 KHACHATRYAN 14J assume $f_{\rm S}=f=f'=\Lambda\ /\ m_{g^*}$.
- 26 CHATRCHYAN 13AJ use the hadronic decay of W.
- 27 CHATRCHYAN 13AJ use the hadronic decay of Z.

MASS LIMITS for Color Sextet Quarks (q_6)

			<i>y ,</i>			
VALUE (GeV)	CL%	DOCUMENT II	כ	TECN	COMMENT	
>84	95	¹ ABE	89D	CDF	$p\overline{p} \rightarrow q_6\overline{q}_6$	<u>-</u>

¹ ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color sextet quark is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. A limit of 121 GeV is obtained for a color decuplet.

MASS LIMITS for Color Octet Charged Leptons (ℓ_8)

$$\lambda \equiv m_{\ell_{\rm N}}/\Lambda$$

VALUE (GeV)CL%DOCUMENT IDTECNCOMMENT>8695
1
 ABE89DCDFStable ℓ_8 : $p\overline{p} \rightarrow \ell_8 \overline{\ell}_8$

ullet ullet We do not use the following data for averages, fits, limits, etc. ullet ullet

² ABT 93 H1
$$e_8$$
: $e_p \rightarrow e_8$ X

¹ ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color octet lepton is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. The limit improves to 99 GeV if it always fragments into a unit-charged hadron.

 2 ABT 93 search for e_8 production via e-gluon fusion in $e\,p$ collisions with $e_8\to\,e\,g$. See their Fig. 3 for exclusion plot in the m_{e_8} – Λ plane for $m_{e_8}=$ 35–220 GeV.

MASS LIMITS for Color Octet Neutrinos (ν_8)

 $\lambda \equiv m_{\ell_8}/\Lambda$

<i>VALUE</i> (GeV)	<u>CL%</u>	<u>DOCUMENT ID</u>)	TECN	COMMENT
>110	90	¹ BARGER	89	RVUE	$\nu_8: p\overline{p} \rightarrow \nu_8\overline{\nu}_8$
\bullet \bullet We do not	use the fo	llowing data for a	verages	, fits, lin	nits, etc. • • •
none 3.8-29.8	95	² KIM	90	AMY	ν_8 : $e^+e^- o$ acoplanar jets

none 3.8–29.8 95 2 KIM 90 AMY ν_8 : $e^+e^- \to$ acoplanar jets none 9–21.9 95 3 BARTEL 87B JADE ν_8 : $e^+e^- \to$ acoplanar jets

MASS LIMITS for W_8 (Color Octet W Boson)

VALUE (GeV) DOCUMENT ID TECN COMMENT

 1 ALBAJAR 89 UA1 $p\overline{p}
ightarrow W_{8}$ X, $W_{8}
ightarrow W_{g}$

REFERENCES FOR Searches for Quark and Lepton Compositeness

AAD	21AU	PRL 127 141801	G. Aad et al.	(ATLAS	Collab.)
AAD	21Q	JHEP 2104 142	G. Aad et al.	(ATLAS	Collab.)
SIRUNYAN	21AG	JHEP 2112 106	A.M. Sirunyan et al.	` (CMS	Collab.)
SIRUNYAN	21N	JHEP 2107 208	A.M. Sirunyan et al.	(CMS	Collab.)
AAD	20AP	JHEP 2011 005	G. Aad et al.	(ATLAS	Collab.)
AAD	20T	JHEP 2003 145	G. Aad et al.	(ATLAS	Collab.)
SIRUNYAN	20AI	JHEP 2005 033	A.M. Sirunyan et al.	(CMS	Collab.)
SIRUNYAN	20AJ	JHEP 2005 052	A.M. Sirunyan et al.	(CMS	Collab.)
AABOUD	19AZ	EPJ C79 803	M. Aaboud et al.	(ATLAS	Collab.)
ABRAMOWICZ	19	PR D99 092006	H. Abramowicz et al.	(ZEUS	Collab.)
SIRUNYAN	19AC	JHEP 1904 114	A.M. Sirunyan et al.	(CMS	Collab.)
SIRUNYAN	19Z	JHEP 1904 015	A.M. Sirunyan et al.	(CMS	Collab.)
AABOUD	18AB	PR D98 032016	M. Aaboud et al.	(ATLAS	Collab.)
AABOUD	18AV	JHEP 1807 089	M. Aaboud et al.		Collab.)
AABOUD	18BA	EPJ C78 102	M. Aaboud et al.	(ATLAS	Collab.)
SIRUNYAN	18AG	PL B781 390	A.M. Sirunyan et al.	(CMS	Collab.)
SIRUNYAN	18BO	JHEP 1808 130	A.M. Sirunyan et al.	(CMS	Collab.)
SIRUNYAN	18DD	EPJ C78 789	A.M. Sirunyan <i>et al.</i>	(CMS	Collab.)
SIRUNYAN	18P	PR D97 072006	A.M. Sirunyan et al.	(CMS	Collab.)
SIRUNYAN	18V	PL B778 349	A.M. Sirunyan et al.	(CMS	Collab.)
AABOUD	17AK	PR D96 052004	M. Aaboud et al.	(ATLAS	Collab.)
AABOUD	17AT	JHEP 1710 182	M. Aaboud et al.	(ATLAS	Collab.)
KHACHATRY	17W	PL B769 520	V. Khachatryan et al.	(CMS	Collab.)
SIRUNYAN	17F	JHEP 1707 013	A.M. Sirunyan et al.	(CMS	Collab.)
AABOUD	16	PL B759 229	M. Aaboud <i>et al.</i>	(ATLAS	Collab.)
AABOUD	16U	PL B761 372	M. Aaboud <i>et al.</i>	(ATLAS	Collab.)
AAD	16AH	JHEP 1602 110	G. Aad et al.		Collab.)
AAD	16AI	JHEP 1603 041	G. Aad <i>et al</i> .	(ATLAS	Collab.)

https://pdg.lbl.gov Page 15 Created: 8/11/2022 09:39

¹BARGER 89 used ABE 89B limit for events with large missing transverse momentum. Two-body decay $\nu_8 \to \nu_g$ is assumed.

 $^{^2}$ KIM 90 is at $E_{
m cm} =$ 50–60.8 GeV. The same assumptions as in BARTEL 87B are used.

³ BARTEL 87B is at $E_{\rm cm}=46.3$ –46.78 GeV. The limit assumes the ν_8 pair production cross section to be eight times larger than that of the corresponding heavy neutrino pair production. This assumption is not valid in general for the weak couplings, and the limit can be sensitive to its SU(2)_I ×U(1)_Y quantum numbers.

^{• • •} We do not use the following data for averages, fits, limits, etc. • • •

 $^{^1}$ ALBAJAR 89 give $\sigma(W_8 \to~W+{\rm jet})/\sigma(W) < 0.019$ (90% CL) for $m_{W_8}~>$ 220 GeV.

	16BM 16S 16AQ	EPJ C76 442 NJP 18 073021 PL B754 302 JHEP 1603 125	G. Aad et al. G. Aad et al. G. Aad et al. V. Khachatryan et al.	(ATLAS Collab.) (ATLAS Collab.) (ATLAS Collab.) (CMS Collab.)
KHACHATRY KHACHATRY KHACHATRY AAD AAD	16K 16L 15AP	JHEP 1601 166 PRL 116 071801 PRL 117 031802 JHEP 1508 138 JHEP 1508 105	 V. Khachatryan et al. V. Khachatryan et al. V. Khachatryan et al. G. Aad et al. G. Aad et al. 	(CMS Collab.) (CMS Collab.) (CMS Collab.) (ATLAS Collab.) (ATLAS Collab.)
AAD AAD AAD	15BY 15L 15V	JHEP 1510 150 PRL 114 221802 PR D91 052007	G. Aad <i>et al.</i> G. Aad <i>et al.</i> G. Aad <i>et al.</i>	(ATLAS Collab.) (ATLAS Collab.) (ATLAS Collab.)
KHACHATRY KHACHATRY	15J 15V	JHEP 1504 025 PL B746 79 PR D91 052009	V. Khachatryan <i>et al.</i> V. Khachatryan <i>et al.</i> V. Khachatryan <i>et al.</i>	(CMS Collab.) (CMS Collab.) (CMS Collab.)
AAD AAD FABBRICHESI	14	PL B728 562 EPJ C74 3134 PR D89 074028	G. Aad et al. G. Aad et al. M. Fabbrichesi, M. Pinamonti, A	
KHACHATRY KHACHATRY AAD AAD	14J	JHEP 1408 173 PL B738 274 NJP 15 093011 PR D87 015010	V. Khachatryan <i>et al.</i> V. Khachatryan <i>et al.</i> G. Aad <i>et al.</i> G. Aad <i>et al.</i>	(CMS Collab.) (CMS Collab.) (ATLAS Collab.)
CHATRCHYAN CHATRCHYAN CHATRCHYAN	13AE 13AJ	PL B720 309	S. Chatrchyan et al. S. Chatrchyan et al. S. Chatrchyan et al.	(ATLAS Collab.) (CMS Collab.) (CMS Collab.) (CMS Collab.)
AAD AAD AARON	12AB	PL B712 40 PR D85 072003 PL B705 52	G. Aad et al. G. Aad et al. F. D. Aaron et al.	(ATLAS Collab.) (ATLAS Collab.) (H1 Collab.)
ABDALLAH AALTONEN ABDALLAH	11 10H 09	EPJ C71 1555 PRL 104 091801 EPJ C60 1	J. Abdallah <i>et al.</i> T. Aaltonen <i>et al.</i> J. Abdallah <i>et al.</i>	(DELPHI Collab.) (CDF Collab.) (DELPHI Collab.)
AARON SCHAEL ABDALLAH	08 07A 06C	PL B663 382 EPJ C49 411 EPJ C45 589	F.D. Aaron <i>et al.</i> S. Schael <i>et al.</i> J. Abdallah <i>et al.</i>	(H1 Collab.) (ALEPH Collab.) (DELPHI Collab.)
ABBIENDI ABBIENDI	06L 04G 04N	PRL 96 211801 EPJ C33 173 PL B602 167	A. Abulencia et al. G. Abbiendi et al. G. Abbiendi et al.	(CDF Collab.) (OPAL Collab.) (OPAL Collab.)
ABDALLAH ACHARD BABICH ABBIENDI	04N 03B 03 02G	EPJ C37 405 PL B568 23 EPJ C29 103 PL B544 57	J. Abdallah <i>et al.</i> P. Achard <i>et al.</i> A.A. Babich <i>et al.</i> G. Abbiendi <i>et al.</i>	(DELPHI Collab.) (L3 Collab.) (OPAL Collab.)
ACHARD ADLOFF CHEKANOV	02D 02D 02 02D	PL B531 28 PL B525 9 PL B549 32	P. Achard et al. C. Adloff et al. S. Chekanov et al.	(L3 Collab.) (H1 Collab.) (ZEUS Collab.)
AFFOLDER BOURILKOV CHEUNG	01I 01 01B	PRL 87 231803 PR D64 071701 PL B517 167	T. Affolder <i>et al.</i> D. Bourilkov K. Cheung	(CDF Collab.)
ACCIARRI AFFOLDER BARATE BARGER	00P 00I 98U 98E	PL B489 81 PR D62 012004 EPJ C4 571 PR D57 391	M. Acciarri et al. T. Affolder et al. R. Barate et al. V. Barger et al.	(L3 Collab.) (CDF Collab.) (ALEPH Collab.)
MCFARLAND DIAZCRUZ ABT	98 94 93	EPJ C1 509 PR D49 2149 NP B396 3	K.S. McFarland <i>et al.</i> J.L. Diaz Cruz, O.A. Sampayo I. Abt <i>et al.</i>	(CCFR/NuTeV Collab.) (CINV) (H1 Collab.)
ADRIANI BARDADIN DECAMP	93M 92 92	PRPL 236 1 ZPHY C55 163 PRPL 216 253	O. Adriani <i>et al.</i> M. Bardadin-Otwinowska D. Decamp <i>et al.</i>	(L3 Collab.) (CLER) (ALEPH Collab.)
PDG ABREU KIM	92 91F 90	PR D45 S1 NP B367 511 PL B240 243	K. Hikasa <i>et al.</i> P. Abreu <i>et al.</i> G.N. Kim <i>et al.</i>	(KEK, LBL, BOST+) (DELPHI Collab.) (AMY Collab.)
ABE ABE ABE ALBAJAR	89B 89D 89J 89	PRL 62 1825 PRL 63 1447 ZPHY C45 175 ZPHY C44 15	F. Abe <i>et al.</i> F. Abe <i>et al.</i> K. Abe <i>et al.</i> C. Albajar <i>et al.</i>	(CDF Collab.) (CDF Collab.) (VENUS Collab.) (UA1 Collab.)
BARGER DORENBOS BARTEL	89 89 87B	PL B220 464 ZPHY C41 567 ZPHY C36 15	V. Barger et al. J. Dorenbosch et al. W. Bartel et al.	(WISC, KEK) (CHARM Collab.) (JADE Collab.)
GRIFOLS	86	PL 168B 264	J.A. Grifols, S. Peris	(BARC)

JODIDIO 86	PR D34 1967	A. Jodidio et al.	(LBL, NWES, TRIU)
Also	PR D37 237 (erratum)	A. Jodidio et al.	(LBL, NWES, TRIU)
RENARD 82	PL 116B 264	F.M. Renard	(CERN)