

$\Upsilon(1S)$

$$J^{PC} = 0^{-}(1^{- -})$$

$\Upsilon(1S)$ MASS

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9460.30 ± 0.26 OUR AVERAGE	Error includes scale factor of 3.3.		
9460.51 ± 0.09 ± 0.05	¹ ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
9459.97 ± 0.11 ± 0.07	MACKAY 84	REDE	$e^+ e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
9460.60 ± 0.09 ± 0.05	^{2,3} BARU	92B	REDE $e^+ e^- \rightarrow$ hadrons
9460.59 ± 0.12	BARU	86	REDE $e^+ e^- \rightarrow$ hadrons
9460.6 ± 0.4	^{3,4} ARTAMONOV 84	REDE	$e^+ e^- \rightarrow$ hadrons
¹ Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).			
² Superseding BARU 86.			
³ Superseded by ARTAMONOV 00.			
⁴ Value includes data of ARTAMONOV 82.			

$\Upsilon(1S)$ WIDTH

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
54.02 ± 1.25 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

$\Upsilon(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\tau^+ \tau^-$	(2.60 ± 0.10) %	
Γ_2 $e^+ e^-$	(2.38 ± 0.11) %	
Γ_3 $\mu^+ \mu^-$	(2.48 ± 0.05) %	
Hadronic decays		
Γ_4 $g g g$	(81.7 ± 0.7) %	
Γ_5 $\gamma g g$	(2.2 ± 0.6) %	
Γ_6 $\eta'(958)$ anything	(2.94 ± 0.24) %	
Γ_7 $J/\psi(1S)$ anything	(5.4 ± 0.4) × 10 ⁻⁴	S=1.4
Γ_8 $J/\psi(1S)\eta_c$	< 2.2	× 10 ⁻⁶ CL=90%
Γ_9 $J/\psi(1S)\chi_{c0}$	< 3.4	× 10 ⁻⁶ CL=90%
Γ_{10} $J/\psi(1S)\chi_{c1}$	(3.9 ± 1.2) × 10 ⁻⁶	
Γ_{11} $J/\psi(1S)\chi_{c2}$	< 1.4	× 10 ⁻⁶ CL=90%
Γ_{12} $J/\psi(1S)\eta_c(2S)$	< 2.2	× 10 ⁻⁶ CL=90%
Γ_{13} $J/\psi(1S)X(3940)$	< 5.4	× 10 ⁻⁶ CL=90%
Γ_{14} $J/\psi(1S)X(4160)$	< 5.4	× 10 ⁻⁶ CL=90%
Γ_{15} $X(4350)$ anything, $X \rightarrow J/\psi(1S)\phi$	< 8.1	× 10 ⁻⁶ CL=90%

Γ_{16}	$Z_c(3900)^\pm$ anything, $Z_c \rightarrow J/\psi(1S)\pi^\pm$	< 1.3	$\times 10^{-5}$	CL=90%
Γ_{17}	$Z_c(4200)^\pm$ anything, $Z_c \rightarrow J/\psi(1S)\pi^\pm$	< 6.0	$\times 10^{-5}$	CL=90%
Γ_{18}	$Z_c(4430)^\pm$ anything, $Z_c \rightarrow J/\psi(1S)\pi^\pm$	< 4.9	$\times 10^{-5}$	CL=90%
Γ_{19}	X_{cs}^\pm anything, $X \rightarrow J/\psi K^\pm$	< 5.7	$\times 10^{-6}$	CL=90%
Γ_{20}	$\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)\pi^+\pi^-$	< 3.8	$\times 10^{-5}$	CL=90%
Γ_{21}	$\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)K^+K^-$	< 7.5	$\times 10^{-6}$	CL=90%
Γ_{22}	$\chi_{c1}(4140)$ anything, $\chi_{c1} \rightarrow J/\psi(1S)\phi$	< 5.2	$\times 10^{-6}$	CL=90%
Γ_{23}	χ_{c0} anything	< 4	$\times 10^{-3}$	CL=90%
Γ_{24}	χ_{c1} anything	$(1.90 \pm 0.35) \times 10^{-4}$		
Γ_{25}	$\chi_{c1}(1P)X_{tetra}$	< 3.78	$\times 10^{-5}$	CL=90%
Γ_{26}	χ_{c2} anything	$(2.8 \pm 0.8) \times 10^{-4}$		
Γ_{27}	$\psi(2S)$ anything	$(1.23 \pm 0.20) \times 10^{-4}$		
Γ_{28}	$\psi(2S)\eta_c$	< 3.6	$\times 10^{-6}$	CL=90%
Γ_{29}	$\psi(2S)\chi_{c0}$	< 6.5	$\times 10^{-6}$	CL=90%
Γ_{30}	$\psi(2S)\chi_{c1}$	< 4.5	$\times 10^{-6}$	CL=90%
Γ_{31}	$\psi(2S)\chi_{c2}$	< 2.1	$\times 10^{-6}$	CL=90%
Γ_{32}	$\psi(2S)\eta_c(2S)$	< 3.2	$\times 10^{-6}$	CL=90%
Γ_{33}	$\psi(2S)X(3940)$	< 2.9	$\times 10^{-6}$	CL=90%
Γ_{34}	$\psi(2S)X(4160)$	< 2.9	$\times 10^{-6}$	CL=90%
Γ_{35}	$\psi(4230)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 7.9	$\times 10^{-5}$	CL=90%
Γ_{36}	$\psi(4360)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 5.2	$\times 10^{-5}$	CL=90%
Γ_{37}	$\psi(4660)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 2.2	$\times 10^{-5}$	CL=90%
Γ_{38}	$X(4050)^\pm$ anything, $X \rightarrow \psi(2S)\pi^\pm$	< 8.8	$\times 10^{-5}$	CL=90%
Γ_{39}	$Z_c(4430)^\pm$ anything, $Z_c \rightarrow \psi(2S)\pi^\pm$	< 6.7	$\times 10^{-5}$	CL=90%
Γ_{40}	$\chi_{c1}(3872)$ anything	< 2.5	$\times 10^{-4}$	CL=90%
Γ_{41}	$Z_c(4200)^+Z_c(4200)^-$	< 2.23	$\times 10^{-5}$	CL=90%
Γ_{42}	$Z_c(3900)^\pm Z_c(4200)^\mp$	< 8.1	$\times 10^{-6}$	CL=90%
Γ_{43}	$Z_c(3900)^+Z_c(3900)^-$	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{44}	$X(4050)^+X(4050)^-$	< 1.58	$\times 10^{-5}$	CL=90%
Γ_{45}	$X(4250)^+X(4250)^-$	< 2.66	$\times 10^{-5}$	CL=90%
Γ_{46}	$X(4050)^\pm X(4250)^\mp$	< 4.42	$\times 10^{-5}$	CL=90%
Γ_{47}	$Z_c(4430)^+Z_c(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%
Γ_{48}	$X(4055)^\pm X(4055)^\mp$	< 2.33	$\times 10^{-5}$	CL=90%

Γ_{49}	$X(4055)^\pm Z_c(4430)^\mp$	< 4.55	$\times 10^{-5}$	CL=90%
Γ_{50}	$\rho\pi$	< 3.68	$\times 10^{-6}$	CL=90%
Γ_{51}	$\omega\pi^0$	< 3.90	$\times 10^{-6}$	CL=90%
Γ_{52}	$\pi^+\pi^-$	< 5	$\times 10^{-4}$	CL=90%
Γ_{53}	K^+K^-	< 5	$\times 10^{-4}$	CL=90%
Γ_{54}	$p\bar{p}$	< 5	$\times 10^{-4}$	CL=90%
Γ_{55}	$\pi^+\pi^-\pi^0$	(2.1 ± 0.8)	$\times 10^{-6}$	
Γ_{56}	ϕK^+K^-	(2.4 ± 0.5)	$\times 10^{-6}$	
Γ_{57}	$\omega\pi^+\pi^-$	(4.5 ± 1.0)	$\times 10^{-6}$	
Γ_{58}	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(4.4 ± 0.8)	$\times 10^{-6}$	
Γ_{59}	$\phi f_2'(1525)$	< 1.63	$\times 10^{-6}$	CL=90%
Γ_{60}	$\omega f_2(1270)$	< 1.79	$\times 10^{-6}$	CL=90%
Γ_{61}	$\rho(770) a_2(1320)$	< 2.24	$\times 10^{-6}$	CL=90%
Γ_{62}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	(3.0 ± 0.8)	$\times 10^{-6}$	
Γ_{63}	$K_1(1270)^\pm K^\mp$	< 2.41	$\times 10^{-6}$	CL=90%
Γ_{64}	$K_1(1400)^\pm K^\mp$	(1.0 ± 0.4)	$\times 10^{-6}$	
Γ_{65}	$b_1(1235)^\pm \pi^\mp$	< 1.25	$\times 10^{-6}$	CL=90%
Γ_{66}	$\pi^+\pi^-\pi^0\pi^0$	(1.28 ± 0.30)	$\times 10^{-5}$	
Γ_{67}	$K_S^0 K^+ \pi^- + \text{c.c.}$	(1.6 ± 0.4)	$\times 10^{-6}$	
Γ_{68}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	(2.9 ± 0.9)	$\times 10^{-6}$	
Γ_{69}	$K^*(892)^- K^+ + \text{c.c.}$	< 1.11	$\times 10^{-6}$	CL=90%
Γ_{70}	$f_1(1285)$ anything	(4.6 ± 3.1)	$\times 10^{-3}$	
Γ_{71}	$D^*(2010)^\pm$ anything	(2.52 ± 0.20)	%	
Γ_{72}	$\frac{f_1(1285)}{X_{tetra}}$	< 6.24	$\times 10^{-5}$	CL=90%
Γ_{73}	2H anything	(2.85 ± 0.25)	$\times 10^{-5}$	
Γ_{74}	Sum of 100 exclusive modes	(1.200 ± 0.017)	%	

Radiative decays

Γ_{75}	$\gamma\pi^+\pi^-$	(6.3 ± 1.8)	$\times 10^{-5}$	
Γ_{76}	$\gamma\pi^0\pi^0$	(1.7 ± 0.7)	$\times 10^{-5}$	
Γ_{77}	$\gamma\pi\pi$ (S-wave)	(4.6 ± 0.7)	$\times 10^{-5}$	
Γ_{78}	$\gamma\pi^0\eta$	< 2.4	$\times 10^{-6}$	CL=90%
Γ_{79}	γK^+K^-	[a] (1.14 ± 0.13)	$\times 10^{-5}$	
Γ_{80}	$\gamma p\bar{p}$	[b] < 6	$\times 10^{-6}$	CL=90%
Γ_{81}	$\gamma 2h^+ 2h^-$	(7.0 ± 1.5)	$\times 10^{-4}$	
Γ_{82}	$\gamma 3h^+ 3h^-$	(5.4 ± 2.0)	$\times 10^{-4}$	
Γ_{83}	$\gamma 4h^+ 4h^-$	(7.4 ± 3.5)	$\times 10^{-4}$	
Γ_{84}	$\gamma\pi^+\pi^- K^+ K^-$	(2.9 ± 0.9)	$\times 10^{-4}$	
Γ_{85}	$\gamma 2\pi^+ 2\pi^-$	(2.5 ± 0.9)	$\times 10^{-4}$	
Γ_{86}	$\gamma 3\pi^+ 3\pi^-$	(2.5 ± 1.2)	$\times 10^{-4}$	
Γ_{87}	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	(2.4 ± 1.2)	$\times 10^{-4}$	
Γ_{88}	$\gamma\pi^+\pi^- p\bar{p}$	(1.5 ± 0.6)	$\times 10^{-4}$	
Γ_{89}	$\gamma 2\pi^+ 2\pi^- p\bar{p}$	(4 ± 6)	$\times 10^{-5}$	
Γ_{90}	$\gamma 2K^+ 2K^-$	(2.0 ± 2.0)	$\times 10^{-5}$	

Γ_{91}	$\gamma\eta'(958)$	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{92}	$\gamma\eta$	< 1.0	$\times 10^{-6}$	CL=90%
Γ_{93}	$\gamma f_0(980)$	< 3	$\times 10^{-5}$	CL=90%
Γ_{94}	$\gamma f_2'(1525)$	(2.9 ± 0.6)	$\times 10^{-5}$	
Γ_{95}	$\gamma f_2(1270)$	(1.01 ± 0.06)	$\times 10^{-4}$	
Γ_{96}	$\gamma\eta(1405)$	< 8.2	$\times 10^{-5}$	CL=90%
Γ_{97}	$\gamma f_0(1500)$	< 1.5	$\times 10^{-5}$	CL=90%
Γ_{98}	$\gamma f_0(1500) \rightarrow \gamma K^+ K^-$	(1.0 ± 0.4)	$\times 10^{-5}$	
Γ_{99}	$\gamma f_0(1710)$	< 2.6	$\times 10^{-4}$	CL=90%
Γ_{100}	$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$	(1.01 ± 0.32)	$\times 10^{-5}$	
Γ_{101}	$\gamma f_0(1710) \rightarrow \gamma\pi^+\pi^-$	(5.3 ± 2.0)	$\times 10^{-6}$	
Γ_{102}	$\gamma f_0(1710) \rightarrow \gamma\pi^0\pi^0$	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{103}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{104}	$\gamma f_4(2050)$	< 5.3	$\times 10^{-5}$	CL=90%
Γ_{105}	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	< 2	$\times 10^{-4}$	CL=90%
Γ_{106}	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	< 8	$\times 10^{-7}$	CL=90%
Γ_{107}	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$	< 6	$\times 10^{-7}$	CL=90%
Γ_{108}	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	< 1.1	$\times 10^{-6}$	CL=90%
Γ_{109}	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	< 3	$\times 10^{-3}$	CL=90%
Γ_{110}	$\gamma\eta_c(1S)$	< 2.9	$\times 10^{-5}$	CL=90%
Γ_{111}	$\gamma\eta_c(2S)$	< 4	$\times 10^{-4}$	CL=90%
Γ_{112}	$\gamma\chi_{c0}$	< 6.6	$\times 10^{-5}$	CL=90%
Γ_{113}	$\gamma\chi_{c1}$	$(4.7 \begin{smallmatrix} +2.4 \\ -1.9 \end{smallmatrix})$	$\times 10^{-5}$	
Γ_{114}	$\gamma\chi_{c2}$	< 7.6	$\times 10^{-6}$	CL=90%
Γ_{115}	$\gamma\chi_{c1}(3872)$	< 4	$\times 10^{-5}$	CL=90%
Γ_{116}	$\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%
Γ_{117}	$\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi$	< 3.0	$\times 10^{-6}$	CL=90%
Γ_{118}	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	< 2.2	$\times 10^{-6}$	CL=90%
Γ_{119}	γX	[c] < 4.5	$\times 10^{-6}$	CL=90%
Γ_{120}	$\gamma X\bar{X} (m_X < 3.1 \text{ GeV})$	[d] < 1	$\times 10^{-3}$	CL=90%
Γ_{121}	$\gamma X\bar{X} (m_X < 4.5 \text{ GeV})$	[e] < 2.4	$\times 10^{-4}$	CL=90%
Γ_{122}	$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	[f] < 1.78	$\times 10^{-4}$	CL=95%
Γ_{123}	$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	[g] < 9	$\times 10^{-6}$	CL=90%
Γ_{124}	$\gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$	[a] < 1.30	$\times 10^{-4}$	CL=90%
Γ_{125}	$\gamma a_1^0 \rightarrow \gamma g g$	[h] < 1	%	CL=90%
Γ_{126}	$\gamma a_1^0 \rightarrow \gamma s\bar{s}$	[h] < 1	$\times 10^{-3}$	CL=90%

Lepton Family number (LF) violating modes

Γ_{127}	$\mu^\pm\tau^\mp$	LF	< 6.0	$\times 10^{-6}$	CL=95%
----------------	-------------------	----	---------	------------------	--------

Other decays

Γ_{128}	invisible	< 3.0	$\times 10^{-4}$	CL=90%
Γ_{129}	hadrons	(97 ± 5)	%	

- [a] $2m_\tau < M(\tau^+\tau^-) < 9.2$ GeV
 [b] 2 GeV $< m_{K^+K^-} < 3$ GeV
 [c] $X =$ scalar with $m < 8.0$ GeV
 [d] $X\bar{X} =$ vectors with $m < 3.1$ GeV
 [e] X and $\bar{X} =$ zero spin with $m < 4.5$ GeV
 [f] 1.5 GeV $< m_\chi < 5.0$ GeV
 [g] 201 MeV $< M(\mu^+\mu^-) < 3565$ MeV
 [h] 0.5 GeV $< m_\chi < 9.0$ GeV, where m_χ is the invariant mass of the hadronic final state.

$\Upsilon(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_3\Gamma_2/\Gamma$
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
31.2±1.6±1.7	KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{129}\Gamma_2/\Gamma$
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.240±0.016 OUR AVERAGE				
1.252±0.004±0.019	¹ ROSNER	06	CLEO	$9.5 e^+e^- \rightarrow \text{hadrons}$
1.187±0.023±0.031	¹ BARU	92B	MD1	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.02 ±0.05	¹ JAKUBOWSKI	88	CBAL	$e^+e^- \rightarrow \text{hadrons}$
1.37 ±0.06 ±0.09	² GILES	84B	CLEO	$e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.08 ±0.04	² ALBRECHT	82	DASP	$e^+e^- \rightarrow \text{hadrons}$
1.13 ±0.07 ±0.11	² NICZYPORUK	82	LENA	$e^+e^- \rightarrow \text{hadrons}$
1.09 ±0.25	² BOCK	80	CNTR	$e^+e^- \rightarrow \text{hadrons}$
1.35 ±0.14	³ BERGER	79	PLUT	$e^+e^- \rightarrow \text{hadrons}$

¹ Radiative corrections evaluated following KURAEV 85.

² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

³ Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$.

$\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$				Γ_2
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>			
1.340±0.018 OUR EVALUATION				

$\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$				Γ_1/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.60±0.10 OUR AVERAGE				
2.53±0.13±0.05	60k	¹ BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S) \rightarrow \tau^+\tau^-$
2.61±0.12 ^{+0.09} _{-0.13}	25k	CINABRO	94B	CLE2 $e^+e^- \rightarrow \tau^+\tau^-$
2.7 ±0.4 ±0.2		² ALBRECHT	85C	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-\tau^+\tau^-$

3.4 ± 0.4 ± 0.4 GILES 83 CLEO $e^+e^- \rightarrow \tau^+\tau^-$

¹ BESSON 07 reports $[\Gamma(\Upsilon(1S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = 1.02 \pm 0.02 \pm 0.05$ which we multiply by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.38 ± 0.11 OUR AVERAGE				
2.29 ± 0.08 ± 0.11		ALEXANDER 98	CLE2	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.42 ± 0.14 ± 0.14	307	ALBRECHT 87	ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.8 ± 0.3 ± 0.2	826	BESSON 84	CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
5.1 ± 3.0		BERGER 80c	PLUT	$e^+e^- \rightarrow e^+e^-$

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0248 ± 0.0005 OUR AVERAGE				
0.0249 ± 0.0002 ± 0.0007	345k	ADAMS 05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0249 ± 0.0008 ± 0.0013		ALEXANDER 98	CLE2	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.0212 ± 0.0020 ± 0.0010		¹ BARU 92	MD1	$e^+e^- \rightarrow \mu^+\mu^-$
0.0231 ± 0.0012 ± 0.0010		¹ KOBEL 92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0252 ± 0.0007 ± 0.0007		CHEN 89B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0261 ± 0.0009 ± 0.0011		KAARSBERG 89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.0230 ± 0.0025 ± 0.0013	86	ALBRECHT 87	ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.029 ± 0.003 ± 0.002	864	BESSON 84	CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.027 ± 0.003 ± 0.003		ANDREWS 83	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.032 ± 0.013 ± 0.003		ALBRECHT 82	DASP	$e^+e^- \rightarrow \mu^+\mu^-$
0.038 ± 0.015 ± 0.002		NICZYPORUK 82	LENA	$e^+e^- \rightarrow \mu^+\mu^-$
0.014 +0.034 -0.014		BOCK 80	CNTR	$e^+e^- \rightarrow \mu^+\mu^-$
0.022 ± 0.020		BERGER 79	PLUT	$e^+e^- \rightarrow \mu^+\mu^-$

¹ Taking into account interference between the resonance and continuum.

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_1/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.008 ± 0.023 OUR AVERAGE				
1.005 ± 0.013 ± 0.022	0.7M	¹ DEL-AMO-SA..10c	BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\Upsilon(1S)$
1.02 ± 0.02 ± 0.05	60k	BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(1S)$

¹ Allows any number of extra photons with total energy < 500 MeV.

$\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
81.7 ± 0.7	20M	¹ BESSON 06A	CLEO	$\Upsilon(1S) \rightarrow \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ from BESSON 06A and PDG 08 values of $B(\mu^+\mu^-) = (2.48 \pm 0.05)\%$ and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20 ± 0.60	400k	¹ BESSON	06A CLEO	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$

¹ Calculated using BESSON 06A values of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ and $\Gamma(g g g)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

 $\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_5/Γ_4

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.70 ± 0.01 ± 0.27	20M	BESSON	06A CLEO	$\Upsilon(1S) \rightarrow (\gamma +) \text{hadrons}$

 $\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0294 ± 0.0024 OUR AVERAGE			
0.030 ± 0.002 ± 0.002	AQUINES	06A CLE3	$\Upsilon(1S) \rightarrow \eta' \text{ anything}$
0.028 ± 0.004 ± 0.002	ARTUSO	03 CLE2	$\Upsilon(1S) \rightarrow \eta' \text{ anything}$

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.4 ± 0.4 OUR FIT					Error includes scale factor of 1.4.
5.4 ± 0.4 OUR AVERAGE					Error includes scale factor of 1.5.
5.25 ± 0.13 ± 0.25		3k	SHEN	16 BELL	$e^+ e^- \rightarrow J/\psi X$
6.4 ± 0.4 ± 0.6		730	BRIERE	04 CLEO	$e^+ e^- \rightarrow J/\psi X$
11 ± 4 ± 2			¹ FULTON	89 CLEO	$e^+ e^- \rightarrow \mu^+ \mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	ALBRECHT	92J ARG	$e^+ e^- \rightarrow e^+ e^- X, \mu^+ \mu^- X$
<17	90	MASCHMANN	90 CBAL	$e^+ e^- \rightarrow \text{hadrons}$
<200	90	NICZYPORUK	83 LENA	

¹ Using $B((J/\psi) \rightarrow \mu^+ \mu^-) = (6.9 \pm 0.9)\%$.

 $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10⁻⁶	90	YANG	14 BELL	$e^+ e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.4 × 10⁻⁶	90	YANG	14 BELL	$e^+ e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.90 ± 1.21 ± 0.23	20	YANG	14 BELL	$e^+ e^- \rightarrow J/\psi X$

 $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10⁻⁶	90	YANG	14 BELL	$e^+ e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$					Γ_{12}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<2.2 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$					Γ_{13}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<5.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$					Γ_{14}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<5.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(X(4350) \text{ anything}, X \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$					Γ_{15}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<8.1 \times 10^{-6}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$
$\Gamma(Z_c(3900)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$					Γ_{16}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.3 \times 10^{-5}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$
$\Gamma(Z_c(4200)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$					Γ_{17}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<6.0 \times 10^{-5}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$
$\Gamma(Z_c(4430)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$					Γ_{18}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<4.9 \times 10^{-5}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$
$\Gamma(X_{cs}^\pm \text{ anything}, X \rightarrow J/\psi K^\pm)/\Gamma_{\text{total}}$					Γ_{19}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<5.7 \times 10^{-6}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi K^- X$
$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{20}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<3.8 \times 10^{-5}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^+ \pi^- X$
$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$					Γ_{21}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<7.5 \times 10^{-6}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$
$\Gamma(\chi_{c1}(4140) \text{ anything}, \chi_{c1} \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$					Γ_{22}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<5.2 \times 10^{-6}$	90	SHEN	16	BELL	$\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$
$\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{23}/Γ_7
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<7.4	90	BRIERE	04	CLEO	$e^+e^- \rightarrow J/\psi X$

$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.35 OUR FIT				
1.90±0.43±0.14	215	JIA	17	BELL $\Upsilon(1S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{24}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.35±0.07 OUR FIT				
0.35±0.08±0.06	52 ± 12	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$

 $\Gamma(\chi_{c1}(1P)X_{tetra})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<37.8 × 10⁻⁶	90	¹ JIA	17A	BELL $e^+e^- \rightarrow \text{hadrons}$

¹For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 37.8×10^{-6} .

 $\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{26}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.52±0.12±0.09	47 ± 11	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$

 $\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.23±0.17±0.11	215	SHEN	16	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{27}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.41±0.11±0.08	42 ± 11	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi \pi^+ \pi^- X$

 $\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.6 × 10⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.5 × 10⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.5 × 10⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.1 × 10⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-6}$	90	YANG 14	BELL	$e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-6}$	90	YANG 14	BELL	$e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(4230) \text{ anything, } \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.9 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$

 $\Gamma(\psi(4360) \text{ anything, } \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$

 $\Gamma(\psi(4660) \text{ anything, } \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$

 $\Gamma(X(4050)^\pm \text{ anything, } X \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.8 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

 $\Gamma(Z_c(4430)^\pm \text{ anything, } Z_c \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.7 \times 10^{-5}$	90	SHEN 16	BELL	$\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

 $\Gamma(\chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-4}$	90	¹ SHEN 16	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^+\pi^-X$

¹ SHEN 16 reports $[\Gamma(\Upsilon(1S) \rightarrow \chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))] < 9.5 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S)) = 3.8 \times 10^{-2}$.

 $\Gamma(Z_c(4200)^+Z_c(4200)^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<22.3 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$

¹ Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi\pi^\pm) = 1$.

 $\Gamma(Z_c(3900)^\pm Z_c(4200)^\mp)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.1 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$

¹ Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi\pi^\pm) = 1 = B(Z_c(3900)^\pm \rightarrow J/\psi\pi^\pm)$.

$\Gamma(Z_c(3900)^+ Z_c(3900)^-)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$

 $\Gamma(X(4050)^+ X(4050)^-)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<15.8 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 $\Gamma(X(4250)^+ X(4250)^-)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<26.6 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(X(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

 $\Gamma(X(4050)^\pm X(4250)^\mp)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<44.2 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

¹ Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(X(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

 $\Gamma(Z_c(4430)^+ Z_c(4430)^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(Z_c(4430)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

 $\Gamma(X(4055)^\pm X(4055)^\mp)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<23.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(X(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

 $\Gamma(X(4055)^\pm Z_c(4430)^\mp)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<45.5 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S) \pi^\pm X$

¹ Assuming $B(X(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(Z_c(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.68	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1 \times 10^3$	90	BLINOV	90	MD1 $\Upsilon(1S) \rightarrow \rho^0 \pi^0$
$<2 \times 10^2$	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
$<2.1 \times 10^3$	90	NICZYPORUK	83	LENA $\Upsilon(1S) \rightarrow \rho^0 \pi^0$

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.90	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{52}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow \pi^+\pi^-$

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{53}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow K^+K^-$

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{54}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	¹ BARU	96	MD1 $\Upsilon(1S) \rightarrow \rho\bar{\rho}$

¹Supersedes BARU 92 in this node. $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.14 ± 0.72 ± 0.34		26 ± 9	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18.4	90		ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$
-------	----	--	---------------	------	-------------------------------------

 $\Gamma(\phi K^+K^-)/\Gamma_{\text{total}}$ Γ_{56}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.36 ± 0.37 ± 0.29	56	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(K^+K^-)$

 $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.46 ± 0.67 ± 0.72	64	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(K^*(892)^0 K^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{58}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.42 ± 0.50 ± 0.58	173	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{59}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.63	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(K^+K^-)$

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{60}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.79	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$ Γ_{61}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.24	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{62}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.02 ± 0.68 ± 0.34	42	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$

$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.41	90	SHEN	12A	BELL $\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.02 ± 0.35 ± 0.22	24	SHEN	12A	BELL $\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.25	90	SHEN	12A	BELL $\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
12.8 ± 2.0 ± 2.3	143 ± 22	SHEN	13	BELL $\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.59 ± 0.33 ± 0.18	37 ± 8	SHEN	13	BELL $\gamma(1S) \rightarrow K_S^0 K^- \pi^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	¹ DOBBS	12A	$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$
------	----	--------------------	-----	--

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
2.92 ± 0.85 ± 0.37	16 ± 5	SHEN	13	BELL $\gamma(1S) \rightarrow K_S^0 K^- \pi^+$

 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.11	90	SHEN	13	BELL $\gamma(1S) \rightarrow K_S^0 K^- \pi^+$

 $\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.6 ± 2.8 ± 1.3	3.1k	JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
25.2 ± 1.3 ± 1.5	≈ 2k	¹ AUBERT	10c	BABR $\gamma(2S) \rightarrow \pi^+ \pi^- \gamma(1S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	² ALBRECHT	92J	ARG $e^+ e^- \rightarrow D^0 \pi^\pm X$
-----	----	-----------------------	-----	---

¹ For $x_p > 0.1$.

² For $x_p > 0.2$.

$\Gamma(f_1(1285)X_{tetra})/\Gamma_{total}$					Γ_{72}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<62.4 \times 10^{-6}$	90	¹ JIA	17A	BELL	$e^+e^- \rightarrow \text{hadrons}$

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.6×10^{-6} to 62.4×10^{-6} .

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{total}$					Γ_{73}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.85 ± 0.25	OUR AVERAGE				
$2.81 \pm 0.49^{+0.20}_{-0.24}$		LEES	14G	BABR	$e^+e^- \rightarrow \overline{2H} X$
$2.86 \pm 0.19 \pm 0.21$	455	ASNER	07	CLEO	$e^+e^- \rightarrow \overline{2H} X$

$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{total}$					Γ_{74}/Γ
VALUE (units 10^{-2})		DOCUMENT ID	TECN	COMMENT	
1.200 ± 0.017		^{1,2} DOBBS	12A	$\Upsilon(1S) \rightarrow \text{hadrons}$	

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(ggg, \gamma gg \rightarrow \overline{d} \text{ anything})/\Gamma(ggg, \gamma gg \rightarrow \text{anything})$				
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.36 \pm 0.23 \pm 0.25$	455	ASNER	07	CLEO $e^+e^- \rightarrow \overline{d} X$

$\Gamma(\gamma\pi^+\pi^-)/\Gamma_{total}$					Γ_{75}/Γ
VALUE (units 10^{-5})		DOCUMENT ID	TECN	COMMENT	
$6.3 \pm 1.2 \pm 1.3$		¹ ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$	

¹ For $m_{\pi\pi} > 1$ GeV.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{total}$					Γ_{76}/Γ
VALUE (units 10^{-5})		DOCUMENT ID	TECN	COMMENT	
$1.7 \pm 0.6 \pm 0.3$		¹ ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$	

¹ For $m_{\pi\pi} > 1$ GeV.

$\Gamma(\gamma\pi\pi(\text{S-wave}))/\Gamma_{total}$					Γ_{77}/Γ
VALUE (units 10^{-5})		DOCUMENT ID	TECN	COMMENT	
$4.63 \pm 0.56 \pm 0.48$		LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$	

$\Gamma(\gamma\pi^0\eta)/\Gamma_{total}$					Γ_{78}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<2.4	90	¹ BESSON	07A	CLEO $e^+e^- \rightarrow \Upsilon(1S)$	

¹ BESSON 07A obtained this limit for $0.7 < m_{\pi^0\eta} < 3$ GeV.

$$\Gamma(\gamma K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{79}/\Gamma$$

$(2 < m_{K^+ K^-} < 3 \text{ GeV})$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.14±0.08±0.10	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

$$\Gamma(\gamma p \bar{p})/\Gamma_{\text{total}} \quad \Gamma_{80}/\Gamma$$

$(2 < m_{p \bar{p}} < 3 \text{ GeV})$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.6	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma p \bar{p}$

$$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}} \quad \Gamma_{81}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.0±1.1±1.0	80 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}} \quad \Gamma_{82}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.4±1.5±1.3	39 ± 11	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}} \quad \Gamma_{83}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.4±2.5±2.5	36 ± 12	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{84}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.9±0.7±0.6	29 ± 8	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{85}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.5±0.7±0.5	26 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{86}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.5±0.9±0.8	17 ± 5	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{87}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.4±0.9±0.8	18 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma \pi^+ \pi^- p \bar{p})/\Gamma_{\text{total}} \quad \Gamma_{88}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.5±0.5±0.3	22 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(\gamma 2\pi^+ 2\pi^- p \bar{p})/\Gamma_{\text{total}} \quad \Gamma_{89}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.4±0.4±0.4	7 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.2 ± 0.2	2 ± 2	FULTON	90B	CLEO $e^+ e^- \rightarrow$ hadrons

 $\Gamma(\gamma \eta'(958))/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.9	90	ATHAR	07A	CLEO $\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \pi^+ \pi^- \eta, \gamma \rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<16	90	RICHICHI	01B	CLE2 $\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \eta \pi^+ \pi^-$

 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.0	90	ATHAR	07A	CLEO $\Upsilon(1S) \rightarrow \gamma \eta \rightarrow \gamma \gamma \gamma, \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<21	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma \eta$

 $\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3	90	¹ ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 1$.				

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.6 OUR AVERAGE					
2.13 ± 0.28 ± 0.72			¹ LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
4.1 ± 1.4 ± 0.1		17	² BESSION	11	CLEO $\Upsilon(1S) \rightarrow K_S^0 K_S^0$
3.7 $^{+0.9}_{-0.7}$ ± 0.8			ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<14	90		³ FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19.4	90		³ ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

¹ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.887 \pm 0.022$ and $B(K^0\bar{K}^0) = 1/2 B(K\bar{K})$.

² BESSION 11 reports $(4.0 \pm 1.3 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(\Upsilon(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})]$ assuming $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 3.1) \times 10^{-2}$, which we rescale to our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The result also assumes $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ and $B(f'_2(1525) \rightarrow K\bar{K}) = 4 B(f'_2(1525) \rightarrow K_S^0 K_S^0)$.

³ Assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
10.1 ± 0.6 OUR AVERAGE				
10.15 ± 0.59 $^{+0.54}_{-0.43}$		¹ LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

10.5 ± 1.6	$\begin{matrix} +1.9 \\ -1.8 \end{matrix}$	² BESSON	07A	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$
10.2 ± 0.8	± 0.7	ATHAR	06	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
8.1 ± 2.3	$\begin{matrix} +2.9 \\ -2.7 \end{matrix}$	³ ANASTASSOV	99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<21	90	³ FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<13	90	³ ALBRECHT	89	ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<81	90	SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$
¹ Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = 1/3 B(f_2(1270) \rightarrow \pi \pi)$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9})\%$.					
² Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = B(f_2(1270) \rightarrow \pi \pi)/3$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.7^{+2.5}_{-1.2})\%$.					
³ Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$.					

$\Gamma(\gamma \eta(1405))/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	¹ FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$

¹ Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$.

$\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	¹ BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.1	90	² BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$
¹ Using $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi \pi)/3$ and $B(f_0(1500) \rightarrow \pi \pi) = (0.349 \pm 0.023)\%$.				
² Calculated by us using $B(f_0(1500) \rightarrow \eta \eta) = (5.1 \pm 0.9)\%$.				

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$1.04 \pm 0.14 \pm 0.33$		¹ LEES	18A	BABR $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$

¹ LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1500) \rightarrow \gamma K \bar{K}) = (2.08 \pm 0.27 \pm 0.65) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$.

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.6	90	¹ ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.3	90	¹ FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<19	90	¹ FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90	² ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<24	90	³ SCHMITT	88	CBAL $\Upsilon(1S) \rightarrow \gamma X$

¹ Assuming $B(f_0(1710) \rightarrow K \bar{K}) = 0.38$.

² Assuming $B(f_0(1710) \rightarrow \pi \pi) = 0.04$.

³ Assuming $B(f_0(1710) \rightarrow \eta \eta) = 0.18$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{100} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

1.01 ± 0.26 ± 0.18		¹ LEES	18A BABR	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$
---------------------------	--	-------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.7	90	ATHAR	06 CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$
-------	----	-------	---------	---

¹ LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma K \bar{K}) = (2.02 \pm 0.51 \pm 0.35) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{101} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.53 ± 0.17 ± 0.11	¹ LEES	18A BABR	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
---------------------------	-------------------	----------	---

¹ LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi \pi) = (0.79 \pm 0.26 \pm 0.17) \times 10^{-5}$ assuming $B(\pi^0 \pi^0) = 1/3 B(\pi \pi)$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{102} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 1.4	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$
-------	----	--------	----------	---

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{103} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 1.8	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$
-------	----	--------	----------	---

 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{104} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 5.3	90	¹ ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
-------	----	--------------------	---------	---

¹ Assuming $B(f_4(2050) \rightarrow \pi \pi) = 0.17$.

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{105} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 0.0002	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
----------	----	------	--------	---

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{106} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 8	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
-----	----	-------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 160	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
-------	----	-------	---------	---

< 150	90	FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
-------	----	--------	----------	---

< 290	90	ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
-------	----	----------	--------	---

< 2000	90	BARU	89 MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
--------	----	------	--------	---

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{107} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 6	90	ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
-----	----	-------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 120	90	MASEK	02 CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
-------	----	-------	---------	---

$\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma p \bar{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<160	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi \phi)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

 $\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.9 $\times 10^{-5}$	90	¹ KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5.7 $\times 10^{-5}$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

 $\Gamma(\gamma \eta_c(2S))/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4 $\times 10^{-4}$	90	¹ KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

 $\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.6 $\times 10^{-5}$	90	¹ KATRENKO	20	BELL $\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.5 $\times 10^{-4}$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$ Γ_{113}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7^{+2.4+0.4}_{-1.8-0.5}$		5	¹ KATRENKO	20	BELL $\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.3	90		SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.					

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.6 $\times 10^{-6}$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3.3 $\times 10^{-5}$	90	¹ KATRENKO	20	BELL $\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

$\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-5}$	90	¹ SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

¹ SHEN 10A reports $[\Gamma(\Upsilon(1S) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 1.6 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 3.8 \times 10^{-2}$.

 $\Gamma(\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-6}$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 $\Gamma(\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 $\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 $\Gamma(\gamma X)/\Gamma_{\text{total}}$ Γ_{119}/Γ

(X = scalar with $m < 8.0$ GeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	¹ DEL-AMO-SA..11J	BABR	$e^+e^- \rightarrow \gamma + X$

••• We do not use the following data for averages, fits, limits, etc. •••

<30	90	² BALEST 95	CLEO	$e^+e^- \rightarrow \gamma + X$
-------	----	------------------------	------	---------------------------------

¹ For a noninteracting scalar X with mass $m < 8.0$ GeV.

² For a noninteracting pseudoscalar X with mass < 7.2 GeV.

 $\Gamma(\gamma X \bar{X} (m_X < 3.1 \text{ GeV}))/\Gamma_{\text{total}}$ Γ_{120}/Γ

($X \bar{X}$ = vectors with $m < 3.1$ GeV)

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	¹ BALEST 95	CLEO	$e^+e^- \rightarrow \gamma + X \bar{X}$

¹ For a noninteracting vector X with mass < 3.1 GeV.

 $\Gamma(\gamma X \bar{X} (m_X < 4.5 \text{ GeV}))/\Gamma_{\text{total}}$ Γ_{121}/Γ

X and \bar{X} = zero spin with $m < 4.5$ GeV

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	¹ DEL-AMO-SA..11J	BABR	$e^+e^- \rightarrow \gamma + X \bar{X}$

¹ For a noninteracting scalar X with mass $m < 4.5$ GeV.

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{122}/Γ

($1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.78	95	ROSNER 07A	CLEO	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ **Γ_{123}/Γ**
 (201 < M($\mu^+ \mu^-$) < 3565 MeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	¹ LOVE 08	CLEO	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.7	90	² LEES 13C	BABR	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$
------	----	-----------------------	------	---

¹ For a narrow scalar or pseudoscalar a_1^0 with 201 < M($\mu^+ \mu^-$) < 3565 MeV, excluding J/ψ . Measured 90% CL limits as a function of M($\mu^+ \mu^-$) range from 1–9 × 10⁻⁶.

² For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9200 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from 0.28–9.7 × 10⁻⁶.

$\Gamma(\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ **Γ_{124}/Γ**
 (2 m_τ < M($\tau^+ \tau^-$) < 9.2 GeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<130	90	¹ LEES 13R	BABR	$\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 50	90	² LOVE 08	CLEO	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$
------	----	----------------------	------	---

¹ For a narrow scalar a_1^0 with 2 m_τ < M(a_1^0) < 9.2 GeV, which result in a 90% CL upper limits of 0.9 × 10⁻⁵ at M(a_1^0) = 2 m_τ , ≈ 1.5 × 10⁻⁵ at M(a_1^0) = 7.5 GeV, and 13 × 10⁻⁵ at M(a_1^0) = 9.2 GeV.

² For a narrow scalar or pseudoscalar a_1^0 with 2 m_τ < M(a_1^0) < 7.5 GeV, which result in a 90% CL limits ranging from 1 × 10⁻⁵ at M(a_1^0)=2 m_τ to 5 × 10⁻⁵ at M(a_1^0)=7.5 GeV.

$\Gamma(\gamma a_1^0 \rightarrow \gamma g g)/\Gamma_{\text{total}}$ **Γ_{125}/Γ**
 (0.5 GeV < m < 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1 × 10 ⁻²	90	¹ LEES 13L	BABR	$\Upsilon(1S) \rightarrow \gamma X$

¹ For a narrow, CP -odd pseudoscalar a_1^0 searched for in 26 hadronic decay modes with invariant mass 0.5 GeV < m_X < 9.0 GeV. Measured 90% CL limit as a function of m_X range from 10⁻⁶ to 10⁻².

$\Gamma(\gamma a_1^0 \rightarrow \gamma s \bar{s})/\Gamma_{\text{total}}$ **Γ_{126}/Γ**
 (0.5 GeV < m < 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1 × 10 ⁻³	90	¹ LEES 13L	BABR	$\Upsilon(1S) \rightarrow \gamma X$

¹ For a narrow, CP -odd pseudoscalar a_1^0 searched for in 14 hadronic decay modes with invariant mass 1.5 GeV < m_X < 9.0 GeV. Measured 90% CL limit as a function of m_X range from 10⁻⁵ to 10⁻³.

———— **LEPTON FAMILY NUMBER (LF) VIOLATING MODES** ————

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ **Γ_{127}/Γ**

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	95	LOVE 08A	CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$					Γ_{128}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 3.0	90	AUBERT	09AX BABR	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<39	90	RUBIN	07 CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$	
<25	90	TAJIMA	07 BELL	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$	

 $\Upsilon(1S)$ REFERENCES

KATRENKO	20	PRL 124 122001	P. Katrenko <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
SHEN	16	PR D93 112013	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
LEES	13C	PR D87 031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
BESSON	11	PR D83 037101	D. Besson <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SA...	10C	PRL 104 191801	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AUBERT	09AX	PRL 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
BESSON	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)
TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)

BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)
