

$\Upsilon(3S)$

$$J^{PC} = 0^{-}(1^{-}-)$$

$\Upsilon(3S)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|---------------------------|----------|------------------------------|
| 10355.2±0.5 | ¹ ARTAMONOV 00 | MD1 | $e^+e^- \rightarrow$ hadrons |
| ••• We do not use the following data for averages, fits, limits, etc. ••• | | | |
| 10355.3±0.5 | ^{2,3} BARU | 86B REDE | $e^+e^- \rightarrow$ hadrons |
| ¹ Reanalysis of BARU 86B using new electron mass (COHEN 87). | | | |
| ² Reanalysis of ARTAMONOV 84. | | | |
| ³ Superseded by ARTAMONOV 00. | | | |

$m\Upsilon(3S) - m\Upsilon(2S)$

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------------------|-------------|----------|----------------------------------|
| 331.50±0.02±0.13 | LEES | 11C BABR | $e^+e^- \rightarrow \pi^+\pi^-X$ |

$\Upsilon(3S)$ WIDTH

| VALUE (keV) | DOCUMENT ID | COMMENT |
|----------------------------------|---|---------|
| 20.32±1.85 OUR EVALUATION | See the Note on "Width Determinations of the Υ States" | |

$\Upsilon(3S)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|--|----------------------------------|-----------------------------------|
| Γ_1 $\Upsilon(2S)$ anything | (10.6 ± 0.8) % | |
| Γ_2 $\Upsilon(2S)\pi^+\pi^-$ | (2.82 ± 0.18) % | S=1.6 |
| Γ_3 $\Upsilon(2S)\pi^0\pi^0$ | (1.85 ± 0.14) % | |
| Γ_4 $\Upsilon(2S)\gamma\gamma$ | (5.0 ± 0.7) % | |
| Γ_5 $\Upsilon(2S)\pi^0$ | < 5.1 × 10 ⁻⁴ | CL=90% |
| Γ_6 $\Upsilon(1S)\pi^+\pi^-$ | (4.37 ± 0.08) % | |
| Γ_7 $\Upsilon(1S)\pi^0\pi^0$ | (2.20 ± 0.13) % | |
| Γ_8 $\Upsilon(1S)\eta$ | < 1 × 10 ⁻⁴ | CL=90% |
| Γ_9 $\Upsilon(1S)\pi^0$ | < 7 × 10 ⁻⁵ | CL=90% |
| Γ_{10} $h_b(1P)\pi^0$ | < 1.2 × 10 ⁻³ | CL=90% |
| Γ_{11} $h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$ | (4.3 ± 1.4) × 10 ⁻⁴ | |
| Γ_{12} $h_b(1P)\pi^+\pi^-$ | < 1.2 × 10 ⁻⁴ | CL=90% |
| Γ_{13} $\tau^+\tau^-$ | (2.29 ± 0.30) % | |
| Γ_{14} $\mu^+\mu^-$ | (2.18 ± 0.21) % | S=2.1 |
| Γ_{15} e^+e^- | (2.18 ± 0.20) % | |
| Γ_{16} hadrons | (93 ± 12) % | |
| Γ_{17} ggg | (35.7 ± 2.6) % | |
| Γ_{18} $\underline{\gamma}gg$ | (9.7 ± 1.8) × 10 ⁻³ | |
| Γ_{19} 2H anything | (2.33 ± 0.33) × 10 ⁻⁵ | |

Radiative decays

| | | | |
|---------------|---|---------------------------------|-----------|
| Γ_{20} | $\gamma\chi_{b2}(2P)$ | $(13.1 \pm 1.6) \%$ | $S=3.4$ |
| Γ_{21} | $\gamma\chi_{b1}(2P)$ | $(12.6 \pm 1.2) \%$ | $S=2.4$ |
| Γ_{22} | $\gamma\chi_{b0}(2P)$ | $(5.9 \pm 0.6) \%$ | $S=1.4$ |
| Γ_{23} | $\gamma\chi_{b2}(1P)$ | $(10.0 \pm 1.0) \times 10^{-3}$ | $S=1.7$ |
| Γ_{24} | $\gamma\chi_{b1}(1P)$ | $(9 \pm 5) \times 10^{-4}$ | $S=1.8$ |
| Γ_{25} | $\gamma\chi_{b0}(1P)$ | $(2.7 \pm 0.4) \times 10^{-3}$ | |
| Γ_{26} | $\gamma\eta_b(2S)$ | $< 6.2 \times 10^{-4}$ | $CL=90\%$ |
| Γ_{27} | $\gamma\eta_b(1S)$ | $(5.1 \pm 0.7) \times 10^{-4}$ | |
| Γ_{28} | $\gamma A^0 \rightarrow \gamma \text{hadrons}$ | $< 8 \times 10^{-5}$ | $CL=90\%$ |
| Γ_{29} | $\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$ | [a] $< 2.2 \times 10^{-4}$ | $CL=95\%$ |
| Γ_{30} | $\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$ | $< 5.5 \times 10^{-6}$ | $CL=90\%$ |
| Γ_{31} | $\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$ | [b] $< 1.6 \times 10^{-4}$ | $CL=90\%$ |

Lepton Family number (LF) violating modes

| | | | | |
|---------------|--------------------|----|------------------------|-----------|
| Γ_{32} | $e^\pm \tau^\mp$ | LF | $< 4.2 \times 10^{-6}$ | $CL=90\%$ |
| Γ_{33} | $\mu^\pm \tau^\mp$ | LF | $< 3.1 \times 10^{-6}$ | $CL=90\%$ |

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ [b] For $m_{\tau^+ \tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. **$\Upsilon(3S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$**

| $\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_{16}\Gamma_{15}/\Gamma$ |
|---|--------------------|------|---------|---|---------------------------------|
| VALUE (keV) | DOCUMENT ID | TECN | COMMENT | | |
| 0.414 ± 0.007 OUR AVERAGE | | | | | |
| $0.413 \pm 0.004 \pm 0.006$ | ROSNER | 06 | CLEO | $10.4 e^+ e^- \rightarrow \text{hadrons}$ | |
| $0.45 \pm 0.03 \pm 0.03$ | ⁴ GILES | 84B | CLEO | $e^+ e^- \rightarrow \text{hadrons}$ | |

⁴ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

| $\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ | | | | | $\Gamma_6\Gamma_{15}/\Gamma$ |
|---|------|---------------------|------|---------|--|
| VALUE (eV) | EVTS | DOCUMENT ID | TECN | COMMENT | |
| $18.46 \pm 0.27 \pm 0.77$ | 6.4k | ⁵ AUBERT | 08BP | BABR | $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$ |

⁵ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

 $\Upsilon(3S)$ PARTIAL WIDTHS

| $\Gamma(e^+ e^-)$ | | | | | Γ_{15} |
|--|-------------|--|--|--|---------------|
| VALUE (keV) | DOCUMENT ID | | | | |
| 0.443 ± 0.008 OUR EVALUATION | | | | | |

 $\Upsilon(3S)$ BRANCHING RATIOS

| $\Gamma(\Upsilon(2S)\text{anything})/\Gamma_{\text{total}}$ | | | | | Γ_1/Γ |
|---|------|-------------------------|------|---------|---------------------------------------|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |
| 0.106 ± 0.008 OUR AVERAGE | | | | | |
| 0.1023 ± 0.0105 | 4625 | ^{6,7,8} BUTLER | 94B | CLE2 | $e^+ e^- \rightarrow \ell^+ \ell^- X$ |

0.111 ± 0.012 4891 7,8,9 BROCK 91 CLEO $e^+e^- \rightarrow \pi^+\pi^-X,$
 $\pi^+\pi^-\ell^+\ell^-$

⁶ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

⁸ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (18.5 \pm 0.8)\%$.


⁹ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_2/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------|-----------|---|
| 2.82±0.18 OUR AVERAGE | | | | Error includes scale factor of 1.6. See the ideogram below. |
| 3.00±0.02±0.14 | 543k | LEES | 11C BABR | $e^+e^- \rightarrow \pi^+\pi^-X$ |
| 2.40±0.10±0.26 | 800 | ¹⁰ AUBERT | 08BP BABR | $e^+e^- \rightarrow \gamma\pi^+\pi^-e^+e^-$ |
| 3.12±0.49 | 980 | ^{11,12} BUTLER | 94B CLE2 | $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$ |
| 2.13±0.38 | 974 | ¹³ BROCK | 91 CLEO | $e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 4.82±0.65±0.53 | 138 | ¹³ WU | 93 CUSB | $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ |
| 3.1 ± 2.0 | 5 | MAGERAS | 82 CUSB | $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ |

WEIGHTED AVERAGE
 2.82±0.18 (Error scaled by 1.6)



¹⁰ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

¹¹ From the exclusive mode.

¹² Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

¹³ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.
 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ (units 10^{-2})

$\Gamma(\Upsilon(2S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|-------------------------|-------------|--|
| 1.85±0.14 OUR AVERAGE | | | | |
| 1.82±0.09±0.12 | 4391 | ¹⁴ BHARI | 09 | CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
| 2.16±0.39 | | ^{15,16} BUTLER | 94B | CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
| 1.7 ±0.5 ±0.2 | 10 | ¹⁷ HEINTZ | 92 | CSB2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |

¹⁴ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

¹⁵ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.

¹⁶ From the exclusive mode.

¹⁷ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\Upsilon(2S)\gamma\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------------|--------------------|-------------|--|
| 0.0502±0.0069 | | | |
| ¹⁸ BUTLER | 94B | CLE2 | $e^+e^- \rightarrow \ell^+\ell^-2\gamma$ |

¹⁸ From the exclusive mode.

$\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

| <u>VALUE (units 10^{-3})</u> | <u>CL%</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------|--------------------|-------------|--|
| <0.51 | 90 | ¹⁹ HE | 08A | CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ |

¹⁹ Authors assume $B(\Upsilon(2S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

Abbreviation MM in the *COMMENT* field below stands for missing mass.

| <u>VALUE (units 10^{-2})</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|-------------|----------------------|-------------|--|
| 4.37±0.08 OUR AVERAGE | | | | |
| 4.32±0.07±0.13 | 90k | ²⁰ LEES | 11L | BABR $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ |
| 4.46±0.01±0.13 | 190k | ²¹ BHARI | 09 | CLEO $e^+e^- \rightarrow \pi^+\pi^-$ MM |
| 4.17±0.06±0.19 | 6.4k | ²² AUBERT | 08BP | BABR $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$ |
| 4.52±0.35 | 11830 | ²³ BUTLER | 94B | CLE2 $e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$ |
| 4.46±0.34±0.50 | 451 | ²³ WU | 93 | CUSB $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ |
| 4.46±0.30 | 11221 | ²³ BROCK | 91 | CLEO $e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|----------|----|---------|----|--|
| 4.9 ±1.0 | 22 | GREEN | 82 | CLEO $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ |
| 3.9 ±1.3 | 26 | MAGERAS | 82 | CUSB $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ |

²⁰ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

²¹ A weighted average of the inclusive and exclusive results.

²² Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

²³ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_6

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------|------|-------------|------|---------|
|-------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|-----------------------------|-----|----------------------|-----------|---|
| $0.577 \pm 0.026 \pm 0.060$ | 800 | ²⁴ AUBERT | 08BP BABR | $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$ |
|-----------------------------|-----|----------------------|-----------|---|

²⁴ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|-------------|------|---------|
|--------------------------|------|-------------|------|---------|

2.20 ± 0.13 OUR AVERAGE

| | | | | |
|--------------------------|------|---------------------|---------|---|
| $2.24 \pm 0.09 \pm 0.11$ | 6584 | ²⁵ BHARI | 09 CLEO | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
|--------------------------|------|---------------------|---------|---|

| | | | | |
|-----------------|----|----------------------|----------|---|
| 1.99 ± 0.34 | 56 | ²⁶ BUTLER | 94B CLE2 | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
|-----------------|----|----------------------|----------|---|

| | | | | |
|-----------------------|----|----------------------|---------|---|
| $2.2 \pm 0.4 \pm 0.3$ | 33 | ²⁷ HEINTZ | 92 CSB2 | $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$ |
|-----------------------|----|----------------------|---------|---|

²⁵ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

²⁶ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality.

²⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_6

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------------------|---------------------|---------|-----------------------------------|
| 0.501 ± 0.043 | ²⁸ BHARI | 09 CLEO | $e^+e^- \rightarrow \Upsilon(3S)$ |
|-------------------|---------------------|---------|-----------------------------------|

²⁸ Not independent of other values reported by BHARI 09.

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

| | | | | |
|-----------------------------|----|--------------------|----------|---|
| <0.1 | 90 | ²⁹ LEES | 11L BABR | $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$ |
|-----------------------------|----|--------------------|----------|---|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|-------------------------|-----------|--|
| <0.8 | 90 | ^{29,30} AUBERT | 08BP BABR | $e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$ |
|--------|----|-------------------------|-----------|--|

| | | | | |
|---------|----|------------------|----------|---------------------------------------|
| <0.18 | 90 | ³¹ HE | 08A CLEO | $e^+e^- \rightarrow \ell^+\ell^-\eta$ |
|---------|----|------------------|----------|---------------------------------------|

| | | | | |
|--------|----|-------|---------|---------------------------------------|
| <2.2 | 90 | BROCK | 91 CLEO | $e^+e^- \rightarrow \ell^+\ell^-\eta$ |
|--------|----|-------|---------|---------------------------------------|

²⁹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³⁰ Using $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

³¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_8/Γ_6

| VALUE (units 10^{-2}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

| | | | | |
|------------------------------|----|--------------------|----------|---|
| <0.23 | 90 | ³² LEES | 11L BABR | $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$ |
|------------------------------|----|--------------------|----------|---|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|--------|----|----------------------|-----------|--|
| <1.9 | 90 | ³³ AUBERT | 08BP BABR | $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$ |
|--------|----|----------------------|-----------|--|

³² Not independent of other values reported by LEES 11L.

³³ Not independent of other values reported by AUBERT 08BP.

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

| VALUE (units 10^{-3}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|------|---------|
|--------------------------|-----|-------------|------|---------|

| | | | | |
|------------------------------|----|------------------|----------|---|
| <0.07 | 90 | ³⁴ HE | 08A CLEO | $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$ |
|------------------------------|----|------------------|----------|---|

³⁴ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

$\Gamma(h_b(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|------------------|------|--|
| $<1.2 \times 10^{-3}$ | 90 | ³⁵ GE | 11 | CLEO $\Upsilon(3S) \rightarrow \pi^0$ anything |

³⁵ Assuming $M(h_b(1P)) = 9900$ MeV and $\Gamma(h_b(1P)) = 0$ MeV, and allowing $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$ to vary from 0–100%.

 $\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

| VALUE (units 10^{-4}) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---|
| $4.3 \pm 1.1 \pm 0.9$ | LEES | 11K | BABR $\Upsilon(3S) \rightarrow \eta_b \gamma \pi^0$ |

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|----------------------|------|--|
| < 1.2 | 90 | ³⁶ LEES | 11C | BABR $e^+e^- \rightarrow \pi^+\pi^- X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| < 18 | | ³⁶ BUTLER | 94B | CLE2 $e^+e^- \rightarrow \pi^+\pi^- X$ |
| < 15 | | ³⁶ BROCK | 91 | CLEO $e^+e^- \rightarrow \pi^+\pi^- X$ |

³⁶ For $M(h_b(1P)) = 9900$ MeV.

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--------------------------|------|----------------------|------|---|
| $2.29 \pm 0.21 \pm 0.22$ | 15k | ³⁷ BESSON | 07 | CLEO $e^+e^- \rightarrow \Upsilon(3S) \rightarrow \tau^+\tau^-$ |

³⁷ BESSON 07 reports $[\Gamma(\Upsilon(3S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$ which we multiply by our best value $B(\Upsilon(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.


 $\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_{13}/Γ_{14}

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|--|
| 0.968 ± 0.016 OUR AVERAGE | | | | |
| $0.966 \pm 0.008 \pm 0.014$ | 2.2M | LEES | 20E | BABR $e^+e^- \rightarrow \Upsilon(3S)$ |
| $1.05 \pm 0.08 \pm 0.05$ | 15k | BESSON | 07 | CLEO $e^+e^- \rightarrow \Upsilon(3S)$ |

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|---|
| 0.0218 ± 0.0021 OUR AVERAGE | | | | Error includes scale factor of 2.1. See the ideogram below. |
| $0.0239 \pm 0.0007 \pm 0.0010$ | 81k | ADAMS | 05 | CLEO $e^+e^- \rightarrow \mu^+\mu^-$ |
| $0.0202 \pm 0.0019 \pm 0.0033$ | | CHEN | 89B | CLEO $e^+e^- \rightarrow \mu^+\mu^-$ |
| $0.0173 \pm 0.0015 \pm 0.0011$ | | KAARSBERG | 89 | CSB2 $e^+e^- \rightarrow \mu^+\mu^-$ |
| $0.033 \pm 0.013 \pm 0.007$ | 1096 | ANDREWS | 83 | CLEO $e^+e^- \rightarrow \mu^+\mu^-$ |

WEIGHTED AVERAGE
 0.0218 ± 0.0021 (Error scaled by 2.1)



$\Gamma(g g g)/\Gamma_{total}$

Γ_{17}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------------------|------|--------------------------|------|------------------------------------|
| 35.7 ± 2.6 | 3M | ³⁸ BESSON 06A | CLEO | $\Upsilon(3S) \rightarrow$ hadrons |

³⁸ Calculated using BESSON 06A value of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and the PDG 08 values of $B(\Upsilon(2S) + \text{anything}) = (10.6 \pm 0.8)\%$, $B(\pi^+ \pi^- \Upsilon(1S)) = (4.40 \pm 0.10)\%$, $B(\pi^0 \pi^0 \Upsilon(1S)) = (2.20 \pm 0.13)\%$, $B(\gamma \chi_{b2}(2P)) = (13.1 \pm 1.6)\%$, $B(\gamma \chi_{b1}(2P)) = (12.6 \pm 1.2)\%$, $B(\gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$, $B(\gamma \chi_{b0}(1P)) = (0.30 \pm 0.11)\%$, $B(\mu^+ \mu^-) = (2.18 \pm 0.21)\%$, and $R_{hadrons} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(\gamma g g)/\Gamma_{total}$ BESSON 06A value.

$\Gamma(\gamma g g)/\Gamma_{total}$

Γ_{18}/Γ

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-----------------------------------|------|--------------------------|------|---|
| 0.97 ± 0.18 | 60k | ³⁹ BESSON 06A | CLEO | $\Upsilon(3S) \rightarrow \gamma +$ hadrons |

³⁹ Calculated using BESSON 06A values of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and $\Gamma(g g g)/\Gamma_{total}$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(g g g)/\Gamma_{total}$ BESSON 06A value.

$\Gamma(\gamma g g)/\Gamma(g g g)$

Γ_{18}/Γ_{17}

| VALUE (units 10^{-2}) | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|------|-------------|------|---|
| $2.72 \pm 0.06 \pm 0.49$ | 3M | BESSON 06A | CLEO | $\Upsilon(3S) \rightarrow (\gamma +)$ hadrons |

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{total}$

Γ_{19}/Γ


| VALUE (units 10^{-5}) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|--|
| $2.33 \pm 0.15^{+0.31}_{-0.28}$ | LEES 14G | BABR | $e^+ e^- \rightarrow \overline{2H} \chi$ |

$\Gamma(\gamma\chi_{b2}(2P))/\Gamma_{\text{total}}$

Γ_{20}/Γ

VALUE EVTS DOCUMENT ID TECN COMMENT

WEIGHTED AVERAGE
0.131±0.016 (Error scaled by 3.4)



0.131 ± 0.016 OUR AVERAGE Error includes scale factor of 3.4. See the ideogram below.

| | | | | | |
|--------------------------|-------|----------------------|----|------|-------------------------------|
| 0.1579 ± 0.0017 ± 0.0073 | 568k | ARTUSO | 05 | CLEO | $e^+e^- \rightarrow \gamma X$ |
| 0.111 ± 0.005 ± 0.004 | 10319 | ⁴⁰ HEINTZ | 92 | CSB2 | $e^+e^- \rightarrow \gamma X$ |
| 0.135 ± 0.003 ± 0.017 | 30741 | MORRISON | 91 | CLE2 | $e^+e^- \rightarrow \gamma X$ |

⁴⁰Supersedes NARAIN 91.

$\Gamma(\gamma\chi_{b2}(2P))/\Gamma_{\text{total}}$

$\Gamma(\gamma\chi_{b1}(2P))/\Gamma_{\text{total}}$


Γ_{21}/Γ

VALUE EVTS DOCUMENT ID TECN COMMENT

0.126 ± 0.012 OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.

| | | | | | |
|---|-------|----------------------|----|------|-------------------------------|
| 0.1454 ± 0.0018 ± 0.0073 | 537k | ARTUSO | 05 | CLEO | $e^+e^- \rightarrow \gamma X$ |
| 0.115 ± 0.005 ± 0.005 | 11147 | ⁴¹ HEINTZ | 92 | CSB2 | $e^+e^- \rightarrow \gamma X$ |
| 0.105 ^{+0.003} _{-0.002} ± 0.013 | 25759 | MORRISON | 91 | CLE2 | $e^+e^- \rightarrow \gamma X$ |


WEIGHTED AVERAGE
 0.126 ± 0.012 (Error scaled by 2.4)



⁴¹Supersedes NARAIN 91.
 $\Gamma(\gamma\chi_{b1}(2P))/\Gamma_{\text{total}}$

| $\Gamma(\gamma\chi_{b0}(2P))/\Gamma_{\text{total}}$ | | | | | | Γ_{22}/Γ |
|--|--------------------|---|-------------|----------------|-------------------------------|----------------------|
| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> | | |
| 0.059 ± 0.006 | OUR AVERAGE | Error includes scale factor of 1.4. See the ideogram below. | | | | |
| $0.0677 \pm 0.0020 \pm 0.0065$ | 225k | ARTUSO | 05 | CLEO | $e^+e^- \rightarrow \gamma X$ | |
| $0.060 \pm 0.004 \pm 0.006$ | 4959 | ⁴² HEINTZ | 92 | CSB2 | $e^+e^- \rightarrow \gamma X$ | |
| $0.049 \begin{smallmatrix} +0.003 \\ -0.004 \end{smallmatrix} \pm 0.006$ | 9903 | MORRISON | 91 | CLE2 | $e^+e^- \rightarrow \gamma X$ | |

⁴²Supersedes NARAIN 91.



$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$ Γ_{23}/Γ

| VALUE (units 10^{-3}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|--|-----|------|-------------------------------------|------|--|
| 10.0 ± 1.0 OUR AVERAGE | | | Error includes scale factor of 1.7. | | |
| $8.0 \pm 1.3 \pm 0.4$ | | 126 | ^{43,44} KORNICER | 11 | CLEO $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$ |
| $10.5 \pm 0.3^{+0.7}_{-0.6}$ | | 9.7k | LEES | 11J | BABR $\Upsilon(3S) \rightarrow X\gamma$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | | |
|-------------|----|----------------------|-----|------|--|
| <19 seen | 90 | ⁴⁵ ASNER | 08A | CLEO | $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$ |
| | | ⁴⁶ HEINTZ | 92 | CSB2 | $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$ |

⁴³ Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.

⁴⁴ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S))]$
 $= (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$ which we divide by our best value $B(\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S)) = (18.0 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁵ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))]$
 $< 27.1 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

⁴⁶ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\Upsilon(1S) \rightarrow \ell^+\ell^-$.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_{24}/Γ

| VALUE (units 10^{-3}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|---------------------------|------|---|
| 0.9±0.5 OUR AVERAGE Error includes scale factor of 1.8. | | | | | |
| 1.5±0.4±0.1 | | 50 | ^{47,48} KORNICER | 11 | CLEO $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$ |
| 0.5±0.3 ^{+0.2} _{-0.1} | | | LEES | 11J | BABR $\Upsilon(3S) \rightarrow X\gamma$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <1.7 | 90 | | ⁴⁹ ASNER | 08A | CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$ |
| seen | | | ⁵⁰ HEINTZ | 92 | CSB2 $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$ |
| ⁴⁷ Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$. | | | | | |
| ⁴⁸ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma\Upsilon(1S))] = (5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$ which we divide by our best value $B(\chi_{b1}(1P) \rightarrow \gamma\Upsilon(1S)) = (35.2 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. | | | | | |
| ⁴⁹ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] < 2.5 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$. | | | | | |
| ⁵⁰ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\Upsilon(1S) \rightarrow \ell^+\ell^-$. | | | | | |

 $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$ Γ_{25}/Γ

| VALUE (units 10^{-2}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|---------------------|------|---|
| 0.27±0.04 OUR AVERAGE | | | | | |
| 0.27±0.04±0.02 | | 2.3k | LEES | 11J | BABR $\Upsilon(3S) \rightarrow X\gamma$ |
| 0.30±0.04±0.10 | | 8.7k | ARTUSO | 05 | CLEO $e^+e^- \rightarrow \gamma X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <0.8 | 90 | | ⁵¹ ASNER | 08A | CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$ |
| ⁵¹ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 21.9 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$. | | | | | |

 $\Gamma(\gamma\eta_b(2S))/\Gamma_{\text{total}}$ Γ_{26}/Γ

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|------|---|
| < 6.2 | | | | |
| | 90 | ARTUSO | 05 | CLEO $e^+e^- \rightarrow \gamma X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| <19 | 90 | LEES | 11J | BABR $\Upsilon(3S) \rightarrow X\gamma$ |

 $\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ Γ_{27}/Γ

| VALUE (units 10^{-4}) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------------|-------------------------|------|--|
| 5.1±0.7 OUR AVERAGE | | | | | |
| 7.1±1.8±1.3 | | 2.3 ± 0.5k | ⁵² BONVICINI | 10 | CLEO $\Upsilon(3S) \rightarrow \gamma X$ |
| 4.8±0.5±0.6 | | 19 ± 3k | ⁵² AUBERT | 09AQ | BABR $\Upsilon(3S) \rightarrow \gamma X$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | | |
| <8.5 | 90 | | LEES | 11J | BABR $\Upsilon(3S) \rightarrow X\gamma$ |
| 4.8±0.5±1.2 | | 19 ± 3k | ^{52,53} AUBERT | 08V | BABR $\Upsilon(3S) \rightarrow \gamma X$ |
| <4.3 | 90 | | ⁵⁴ ARTUSO | 05 | CLEO $e^+e^- \rightarrow \gamma X$ |
| ⁵² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. | | | | | |
| ⁵³ Systematic error re-evaluated by AUBERT 09AQ. | | | | | |
| ⁵⁴ Superseded by BONVICINI 10. | | | | | |

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$ Γ_{28}/Γ
 (0.3 GeV < m_{A^0} < 7 GeV)

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|---------------------|-----|--------------------|----------|--|
| $<8 \times 10^{-5}$ | 90 | ⁵⁵ LEES | 11H BABR | $\Upsilon(3S) \rightarrow \gamma \text{hadrons}$ |

⁵⁵ For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{29}/Γ
 (1.5 GeV < m_X < 5.0 GeV)

| VALUE (units 10^{-4}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|-------------------------------|
| <2.2 | 95 | ROSNER | 07A CLEO | $e^+e^- \rightarrow \gamma X$ |

$\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|----------------------|----------|--|
| <5.5 | 90 | ⁵⁶ AUBERT | 09Z BABR | $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$ |

⁵⁶ For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from 0.27 – 5.5×10^{-6} .

$\Gamma(\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-----------------------|-----|----------------------|----------|--|
| $<1.6 \times 10^{-4}$ | 90 | ⁵⁷ AUBERT | 09P BABR | $e^+e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$ |

⁵⁷ For a narrow scalar or pseudoscalar a_1^0 with $M(\tau^+ \tau^-)$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. Measured 90% CL limits as a function of $M(\tau^+ \tau^-)$ range from 1.5 – 16×10^{-5} .

———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{32}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|-------------------------------------|
| <4.2 | 90 | LEES | 10B BABR | $e^+e^- \rightarrow e^\pm \tau^\mp$ |

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{33}/Γ

| VALUE (units 10^{-6}) | CL% | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-----|-------------|----------|---------------------------------------|
| <3.1 | 90 | LEES | 10B BABR | $e^+e^- \rightarrow \mu^\pm \tau^\mp$ |
| <20.3 | 95 | LOVE | 08A CLEO | $e^+e^- \rightarrow \mu^\pm \tau^\mp$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Upsilon(3S)$ REFERENCES

| | | | | |
|----------|-----|----------------|---------------------------|-----------------|
| LEES | 20E | PRL 125 241801 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 14G | PR D89 111102 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| GE | 11 | PR D84 032008 | J.Y. Ge <i>et al.</i> | (CLEO Collab.) |
| KORNICER | 11 | PR D83 054003 | M. Kornicer <i>et al.</i> | (CLEO Collab.) |
| LEES | 11C | PR D84 011104 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 11H | PRL 107 221803 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 11J | PR D84 072002 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| LEES | 11K | PR D84 091101 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |

| | | | | |
|-------------|------|---|------------------------------|--------------------|
| LEES | 11L | PR D84 092003 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| BONVICINI | 10 | PR D81 031104 | G. Bonvicini <i>et al.</i> | (CLEO Collab.) |
| LEES | 10B | PRL 104 151802 | J.P. Lees <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 09AQ | PRL 103 161801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 09P | PRL 103 181801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 09Z | PRL 103 081803 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| BHARI | 09 | PR D79 011103 | S.R. Bhari <i>et al.</i> | (CLEO Collab.) |
| ASNER | 08A | PR D78 091103 | D.M. Asner <i>et al.</i> | (CLEO Collab.) |
| AUBERT | 08BP | PR D78 112002 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 08V | PRL 101 071801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| HE | 08A | PRL 101 192001 | Q. He <i>et al.</i> | (CLEO Collab.) |
| LOVE | 08A | PRL 101 201601 | W. Love <i>et al.</i> | (CLEO Collab.) |
| PDG | 08 | PL B667 1 | C. Amsler <i>et al.</i> | (PDG Collab.) |
| BESSON | 07 | PRL 98 052002 | D. Besson <i>et al.</i> | (CLEO Collab.) |
| ROSNER | 07A | PR D76 117102 | J.L. Rosner <i>et al.</i> | (CLEO Collab.) |
| BESSON | 06A | PR D74 012003 | D. Besson <i>et al.</i> | (CLEO Collab.) |
| ROSNER | 06 | PRL 96 092003 | J.L. Rosner <i>et al.</i> | (CLEO Collab.) |
| ADAMS | 05 | PRL 94 012001 | G.S. Adams <i>et al.</i> | (CLEO Collab.) |
| ARTUSO | 05 | PRL 94 032001 | M. Artuso <i>et al.</i> | (CLEO Collab.) |
| ARTAMONOV | 00 | PL B474 427 | A.S. Artamonov <i>et al.</i> | |
| BUTLER | 94B | PR D49 40 | F. Butler <i>et al.</i> | (CLEO Collab.) |
| WU | 93 | PL B301 307 | Q.W. Wu <i>et al.</i> | (CUSB Collab.) |
| HEINTZ | 92 | PR D46 1928 | U. Heintz <i>et al.</i> | (CUSB II Collab.) |
| BROCK | 91 | PR D43 1448 | I.C. Brock <i>et al.</i> | (CLEO Collab.) |
| HEINTZ | 91 | PRL 66 1563 | U. Heintz <i>et al.</i> | (CUSB Collab.) |
| MORRISON | 91 | PRL 67 1696 | R.J. Morrison <i>et al.</i> | (CLEO Collab.) |
| NARAIN | 91 | PRL 66 3113 | M. Narain <i>et al.</i> | (CUSB Collab.) |
| CHEN | 89B | PR D39 3528 | W.Y. Chen <i>et al.</i> | (CLEO Collab.) |
| KAARSBERG | 89 | PRL 62 2077 | T.M. Kaarsberg <i>et al.</i> | (CUSB Collab.) |
| BUCHMUEL... | 88 | HE e^+e^- Physics 412 | W. Buchmueller, S. Cooper | (HANN, DESY, MIT) |
| | | Editors: A. Ali and P. Soeding, World Scientific, Singapore | | |
| COHEN | 87 | RMP 59 1121 | E.R. Cohen, B.N. Taylor | (RISC, NBS) |
| BARU | 86B | ZPHY C32 622 (erratum) | S.E. Baru <i>et al.</i> | (NOVO) |
| KURAEV | 85 | SJNP 41 466 | E.A. Kuraev, V.S. Fadin | (NOVO) |
| | | Translated from YAF 41 733. | | |
| ARTAMONOV | 84 | PL 137B 272 | A.S. Artamonov <i>et al.</i> | (NOVO) |
| GILES | 84B | PR D29 1285 | R. Giles <i>et al.</i> | (CLEO Collab.) |
| ANDREWS | 83 | PRL 50 807 | D.E. Andrews <i>et al.</i> | (CLEO Collab.) |
| GREEN | 82 | PRL 49 617 | J. Green <i>et al.</i> | (CLEO Collab.) |
| MAGERAS | 82 | PL 118B 453 | G. Mageras <i>et al.</i> | (COLU, CORN, LSU+) |