

$\Xi_c(2930)$  $I(J^P) = ?(?^?)$  Status: \*\*

OMITTED FROM SUMMARY TABLE

 $\Xi_c(2930)$  MASSES $\Xi_c(2930)^+$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2942.3 ± 4.4 ± 1.5</b>	21	LI	18D	BELL $e^+e^-$ at $\Upsilon(4S)$

 $\Xi_c(2930)^0$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2938.55 ± 0.21 ± 0.22</b>	10.4k	<sup>1</sup> AAIJ	20X	LHCB $pp$ at 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

2928.9 ± 3.0  $\begin{smallmatrix} + 0.9 \\ - 12.0 \end{smallmatrix}$  61 LI 18A BELL  $e^+e^-$  at  $\Upsilon(4S)$

2931 ± 3 ± 5 34 AUBERT 08H BABR  $\Upsilon(4S) \rightarrow B\bar{B}$

<sup>1</sup>AAIJ 20X reports  $2938.55 \pm 0.21 \pm 0.17 \pm 0.14$  MeV where the last uncertainty is due to the  $\Lambda_c^+$  mass. Observes that the broader resonance at 2930 MeV seen in  $B^- \rightarrow K^- \Lambda_c^+ \bar{\Lambda}_c^-$  by LI 18A and AUBERT 08H resolves into two narrower peaks at approximately 2939 MeV and 2923 MeV.

 $\Xi_c(2930)^+ - \Xi_c(2930)^0$  MASS DIFFERENCE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.4 ± 5.3 <math>\begin{smallmatrix} + 1.7 \\ - 12.1 \end{smallmatrix}</math></b>	21	<sup>1</sup> LI	18D	BELL $e^+e^-$ at $\Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>This LI 18D value is not independent of the mass measurements.

 $\Xi_c(2930)$  WIDTHS $\Xi_c(2930)^+$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>14.8 ± 8.8 ± 2.5</b>	21	LI	18D	BELL $e^+e^-$ at $\Upsilon(4S)$

 $\Xi_c(2930)^0$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.2 ± 0.8 ± 1.1</b>	10.4k	<sup>1</sup> AAIJ	20X	LHCB $pp$ at 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5 ± 8.4  $\begin{smallmatrix} + 5.9 \\ - 7.9 \end{smallmatrix}$  61 LI 18A BELL  $e^+e^-$  at  $\Upsilon(4S)$

36 ± 7 ± 11 34 AUBERT 08H BABR  $\Upsilon(4S) \rightarrow B\bar{B}$

<sup>1</sup>AAIJ 20X observes that the broader resonance at 2930 MeV seen in  $B^- \rightarrow K^- \Lambda_c^+ \bar{\Lambda}_c^-$  by LI 18A and AUBERT 08H resolves into two narrower peaks at approximately 2939 MeV and 2923 MeV.

$\Xi_c(2930)$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \Lambda_c^+ K^-$	seen
$\Gamma_2 \quad \Lambda_c^+ K_S^0$	seen

 $\Xi_c(2930)$  BRANCHING RATIOS

$\Gamma(\Lambda_c^+ K^-)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>seen</b>	10.4k	AAIJ	20X	LHCB	$pp$ at 13 TeV
seen	61	LI	18A	BELLE	Significance 5.1 std
seen	34	AUBERT	08H	BABR	$e^+e^-$ at $\Upsilon(4S)$

  

$\Gamma(\Lambda_c^+ K_S^0)/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<b>seen</b>	21	LI	18D	BELLE	Significance 4.1 std

 $\Xi_c(2930)$  REFERENCES

AAIJ	20X	PRL 124 222001	R. Aaij <i>et al.</i>	(LHCb Collab.)
LI	18A	EPJ C78 252	Y.B. Li <i>et al.</i>	(BELLE Collab.)
LI	18D	EPJ C78 928	Y.B. Li <i>et al.</i>	(BELLE Collab.)
AUBERT	08H	PR D77 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)