

# SEARCHES not in other sections

## Magnetic Monopole Searches

The most sensitive experiments obtain negative results.

Best cosmic-ray supermassive monopole flux limit:

$$< 1.4 \times 10^{-16} \text{ cm}^{-2} \text{sr}^{-1} \text{s}^{-1} \quad \text{for } 1.1 \times 10^{-4} < \beta < 1$$

## Supersymmetric Particle Searches

All supersymmetric mass bounds here are model dependent.

The limits assume:

1)  $\tilde{\chi}_1^0$  is the lightest supersymmetric particle; 2)  $R$ -parity is conserved, unless stated otherwise;

See the Particle Listings for a Note giving details of supersymmetry.

$\tilde{\chi}_i^0$  — neutralinos (mixtures of  $\tilde{\gamma}$ ,  $\tilde{Z}^0$ , and  $\tilde{H}_i^0$ )

Mass  $m_{\tilde{\chi}_1^0} > 0 \text{ GeV}$ , CL = 95%

[general MSSM, non-universal gaugino masses]

Mass  $m_{\tilde{\chi}_1^0} > 46 \text{ GeV}$ , CL = 95%

[all  $\tan\beta$ , all  $m_0$ , all  $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$ ]

Mass  $m_{\tilde{\chi}_2^0} > 62.4 \text{ GeV}$ , CL = 95%

[ $1 < \tan\beta < 40$ , all  $m_0$ , all  $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$ ]

Mass  $m_{\tilde{\chi}_3^0} > 99.9 \text{ GeV}$ , CL = 95%

[ $1 < \tan\beta < 40$ , all  $m_0$ , all  $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$ ]

Mass  $m_{\tilde{\chi}_4^0} > 116 \text{ GeV}$ , CL = 95%

[ $1 < \tan\beta < 40$ , all  $m_0$ , all  $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$ ]

$\tilde{\chi}_i^\pm$  — charginos (mixtures of  $\tilde{W}^\pm$  and  $\tilde{H}_i^\pm$ )

Mass  $m_{\tilde{\chi}_1^\pm} > 94 \text{ GeV}$ , CL = 95%

[ $\tan\beta < 40$ ,  $m_{\tilde{\chi}_1^\pm} - m_{\tilde{\chi}_1^0} > 3 \text{ GeV}$ , all  $m_0$ ]

Mass  $m_{\tilde{\chi}_1^\pm} > 1000 \text{ GeV}$ , CL = 95%

[ $2\ell + \cancel{E}_T$ , Tchi1chi1C,  $m_{\tilde{\chi}_1^0} = 0 \text{ GeV}$ ]

$\tilde{\chi}^\pm$  — long-lived chargino

Mass  $m_{\tilde{\chi}^\pm} > 620$  GeV, CL = 95% [stable  $\tilde{\chi}^\pm$ ]

$\tilde{\nu}$  — sneutrino

Mass  $m > 41$  GeV, CL = 95% [model independent]

Mass  $m > 94$  GeV, CL = 95%

[CMSSM,  $1 \leq \tan\beta \leq 40$ ,  $m_{\tilde{e}_R} - m_{\tilde{\chi}_1^0} > 10$  GeV]

Mass  $m > 3400$  GeV, CL = 95% [R-Parity Violating]

$[\tilde{\nu}_\tau \rightarrow e\mu, \lambda_{312} = \lambda_{321} = 0.07, \lambda'_{311} = 0.11]$

$\tilde{e}$  — scalar electron (selectron)

Mass  $m > 107$  GeV, CL = 95% [all  $m_{\tilde{e}_L} - m_{\tilde{\chi}_1^0}$ ]

Mass  $m > 700$  GeV, CL = 95%

$[2\ell + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L} \text{ and } \tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}]$

Mass  $m > 250$  GeV, CL = 95%

$[\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{e}_R, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}]$

Mass  $m > 410$  GeV, CL = 95% [R-Parity Violating]

$[\geq 4\ell^\pm, \tilde{\ell} \rightarrow l\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu]$

$\tilde{\mu}$  — scalar muon (smuon)

Mass  $m > 700$  GeV, CL = 95%

$[2\ell + \cancel{E}_T, m_{\tilde{\ell}_R} = m_{\tilde{\ell}_L} \text{ and } \tilde{\ell} = \tilde{e}, \tilde{\mu}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}]$

Mass  $m > 210$ , CL = 95%

$[\ell^\pm \ell^\mp + \cancel{E}_T, \tilde{\mu}_R, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}]$

Mass  $m > 94$  GeV, CL = 95%

[CMSSM,  $1 \leq \tan\beta \leq 40$ ,  $m_{\tilde{\mu}_R} - m_{\tilde{\chi}_1^0} > 10$  GeV]

Mass  $m > 410$  GeV, CL = 95% [R-Parity Violating]

$[\geq 4\ell^\pm, \tilde{\ell} \rightarrow l\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell^\pm \ell^\mp \nu]$

$\tilde{\tau}$  — scalar tau (stau)

Mass  $m > 81.9$  GeV, CL = 95%

$[m_{\tilde{\tau}_R} - m_{\tilde{\chi}_1^0} > 15 \text{ GeV, all } \theta_\tau, \text{B}(\tilde{\tau} \rightarrow \tau \tilde{\chi}_1^0) = 100\%]$

Mass  $m > 90$  GeV, CL = 95%

[RPV,  $\tilde{\tau}_R$ , indirect,  $\Delta m > 5$  GeV]

Mass  $m > 286$  GeV, CL = 95% [long-lived  $\tilde{\tau}$ ]

$\tilde{q}$  — squarks of the first two quark generations

Mass  $m > 1.220 \times 10^3$  GeV, CL = 95%

$[\text{jets} + \cancel{E}_T, \text{Tsqk1, 1 non-degenerate } \tilde{q}, m_{\tilde{\chi}_1^0} = 0 \text{ GeV}]$

Mass  $m > 1.600 \times 10^3$  GeV, CL = 95% [R-Parity Violating]

$[\tilde{q} \rightarrow q\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow \ell\ell\nu, \lambda_{121}, \lambda_{122} \neq 0, m_{\tilde{g}} = 2400 \text{ GeV}]$

$\tilde{q}$  — long-lived squark

Mass  $m > 1340$ , CL = 95% [ $\tilde{t}$  R-hadrons]

Mass  $m > 1250$ , CL = 95% [ $\tilde{b}$  R-hadrons]

$\tilde{b}$  — scalar bottom (sbottom)

Mass  $m > 1.270 \times 10^3$  GeV, CL = 95%

[ $b$ -jets +  $\cancel{E}_T$ , T<sub>sb0t1</sub>,  $m_{\tilde{\chi}_1^0} = 0$  GeV]

Mass  $m > 307$  GeV, CL = 95% [R-Parity Violating]

[ $\tilde{b} \rightarrow td$  or  $ts$ ,  $\lambda''_{332}$  or  $\lambda''_{331}$  coupling]

$\tilde{t}$  — scalar top (stop)

Mass  $m > 1.310 \times 10^3$  GeV, CL = 95%

[jets +  $\cancel{E}_T$ , T<sub>stop1</sub>,  $m_{\tilde{\chi}_1^0} < 300$  GeV]

Mass  $m > 1100$  GeV, CL = 95% [R-Parity Violating]

[ $\tilde{t} \rightarrow be$ , T<sub>stop2RPV</sub>, prompt]

$\tilde{g}$  — gluino

Mass  $m > 2.300 \times 10^3$  GeV, CL = 95%

[jets +  $\cancel{E}_T$ , T<sub>glu1A</sub>,  $m_{\tilde{\chi}_1^0} < 200$  GeV]

Mass  $m > 2.260 \times 10^3$  GeV, CL = 95% [R-Parity Violating]

[ $\geq 4\ell$ ,  $\lambda_{12k} \neq 0$ ,  $m_{\tilde{\chi}_1^0} > 1000$  GeV]

## Technicolor

The limits for technicolor (and top-color) particles are quite varied depending on assumptions. See the Technicolor section of the full *Review* (the data listings).

## Quark and Lepton Compositeness, Searches for

**Scale Limits  $\Lambda$  for Contact Interactions  
(the lowest dimensional interactions with four fermions)**

If the Lagrangian has the form

$$\pm \frac{g^2}{2\Lambda^2} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L$$

(with  $g^2/4\pi$  set equal to 1), then we define  $\Lambda \equiv \Lambda_{LL}^\pm$ . For the full definitions and for other forms, see the Note in the Listings on Searches for Quark and Lepton Compositeness in the full *Review* and the original literature.

$\Lambda_{LL}^+(eeee)$	> 8.3 TeV, CL = 95%
$\Lambda_{LL}^-(eeee)$	> 10.3 TeV, CL = 95%
$\Lambda_{LL}^+(ee\mu\mu)$	> 8.5 TeV, CL = 95%
$\Lambda_{LL}^-(ee\mu\mu)$	> 9.5 TeV, CL = 95%
$\Lambda_{LL}^+(ee\tau\tau)$	> 7.9 TeV, CL = 95%
$\Lambda_{LL}^-(ee\tau\tau)$	> 7.2 TeV, CL = 95%
$\Lambda_{LL}^+(\ell\ell\ell\ell)$	> 9.1 TeV, CL = 95%
$\Lambda_{LL}^-(\ell\ell\ell\ell)$	> 10.3 TeV, CL = 95%
$\Lambda_{LL}^+(eeqq)$	> 24 TeV, CL = 95%
$\Lambda_{LL}^-(eeqq)$	> 37 TeV, CL = 95%
$\Lambda_{LL}^+(eeuu)$	> 23.3 TeV, CL = 95%
$\Lambda_{LL}^-(eeuu)$	> 12.5 TeV, CL = 95%
$\Lambda_{LL}^+(eedd)$	> 11.1 TeV, CL = 95%
$\Lambda_{LL}^-(eedd)$	> 26.4 TeV, CL = 95%
$\Lambda_{LL}^+(eecc)$	> 9.4 TeV, CL = 95%
$\Lambda_{LL}^-(eecc)$	> 5.6 TeV, CL = 95%
$\Lambda_{LL}^+(eebb)$	> 9.4 TeV, CL = 95%
$\Lambda_{LL}^-(eebb)$	> 10.2 TeV, CL = 95%
$\Lambda_{LL}^+(\mu\mu qq)$	> 22.3 TeV, CL = 95%
$\Lambda_{LL}^-(\mu\mu qq)$	> 40.0 TeV, CL = 95%
$\Lambda(\ell\nu\ell\nu)$	> 3.10 TeV, CL = 90%
$\Lambda(e\nu qq)$	> 2.81 TeV, CL = 95%
$\Lambda_{LL}^+(qqqq)$	> 13.1 none 17.4–29.5 TeV, CL = 95%
$\Lambda_{LL}^-(qqqq)$	> 21.8 TeV, CL = 95%
$\Lambda_{LL}^+(\nu\nu qq)$	> 5.0 TeV, CL = 95%
$\Lambda_{LL}^-(\nu\nu qq)$	> 5.4 TeV, CL = 95%

## Excited Leptons

The limits from  $\ell^{*+} \ell^{*-}$  do not depend on  $\lambda$  (where  $\lambda$  is the  $\ell \ell^*$  transition coupling). The  $\lambda$ -dependent limits assume chiral coupling.

$e^{*\pm}$  — excited electron

Mass  $m > 103.2$  GeV, CL = 95% (from  $e^* e^*$ )

Mass  $m > 5.600 \times 10^3$  GeV, CL = 95% (from  $e e^*$ )

Mass  $m > 356$  GeV, CL = 95% (if  $\lambda_\gamma = 1$ )

$\mu^{*\pm}$  — excited muon

Mass  $m > 103.2$  GeV, CL = 95% (from  $\mu^* \mu^*$ )

Mass  $m > 5.700 \times 10^3$  GeV, CL = 95% (from  $\mu \mu^*$ )

$\tau^{*\pm}$  — excited tau

Mass  $m > 103.2$  GeV, CL = 95% (from  $\tau^* \tau^*$ )

Mass  $m > 2.500 \times 10^3$  GeV, CL = 95% (from  $\tau \tau^*$ )

$\nu^*$  — excited neutrino

Mass  $m > 1.600 \times 10^3$  GeV, CL = 95% (from  $\nu^* \nu^*$ )

Mass  $m > 213$  GeV, CL = 95% (from  $\nu^* X$ )

$q^*$  — excited quark

Mass  $m > 338$  GeV, CL = 95% (from  $q^* q^*$ )

Mass  $m > 6700$  GeV, CL = 95% (from  $q^* X$ )

## Color Sextet and Octet Particles

Color Sextet Quarks ( $q_6$ )

Mass  $m > 84$  GeV, CL = 95% (Stable  $q_6$ )

Color Octet Charged Leptons ( $\ell_8$ )

Mass  $m > 86$  GeV, CL = 95% (Stable  $\ell_8$ )

Color Octet Neutrinos ( $\nu_8$ )

Mass  $m > 110$  GeV, CL = 90% ( $\nu_8 \rightarrow \nu g$ )

## Extra Dimensions

Please refer to the Extra Dimensions section of the full *Review* for a discussion of the model-dependence of these bounds, and further constraints.

### Constraints on the radius of the extra dimensions, for the case of two-flat dimensions of equal radii

(direct tests of Newton's law)

$$R < 3.8 \mu\text{m}, \text{ CL} = 95\% \quad (pp \rightarrow jG)$$

$$R < 0.16\text{--}916 \text{ nm} \quad (\text{astrophysics; limits depend on technique and assumptions})$$

### Constraints on the fundamental gravity scale

$$M_{TT} > 9.02 \text{ TeV}, \text{ CL} = 95\% \quad (pp \rightarrow \text{dijet, angular distribution})$$

$$M_c > 4.16 \text{ TeV}, \text{ CL} = 95\% \quad (pp \rightarrow \ell\bar{\ell})$$

### Constraints on the Kaluza-Klein graviton in warped extra dimensions

$$M_G > 4.78 \text{ TeV}, \text{ CL} = 95\% \quad (pp \rightarrow e^+e^-, \mu^+\mu^-)$$

### Constraints on the Kaluza-Klein gluon in warped extra dimensions

$$M_{g_{KK}} > 3.8 \text{ TeV}, \text{ CL} = 95\% \quad (g_{KK} \rightarrow t\bar{t})$$

---

## WIMP and Dark Matter Searches

No confirmed evidence found for galactic

WIMPs from the GeV to the TeV mass scales and down to  $1 \times 10^{-10}$  pb spin independent cross section at  $M = 100 \text{ GeV}$ .

---