

Ω BARYONS ($S = -3, I = 0$)

$$\Omega^- = sss$$

Ω^-

$$I(J^P) = 0(\frac{3}{2}^+)$$

$J^P = \frac{3}{2}^+$ is the quark-model prediction; and $J = 3/2$ is fairly well established.

Mass $m = 1672.45 \pm 0.29$ MeV
 $(m_{\Omega^-} - m_{\bar{\Omega}^+}) / m_{\Omega^-} = (-1 \pm 8) \times 10^{-5}$
 Mean life $\tau = (0.821 \pm 0.011) \times 10^{-10}$ s
 $c\tau = 2.461$ cm
 $(\tau_{\Omega^-} - \tau_{\bar{\Omega}^+}) / \tau_{\Omega^-} = 0.00 \pm 0.05$
 Magnetic moment $\mu = -2.02 \pm 0.05 \mu_N$

Decay parameters

$\alpha(\Omega^-) \alpha_-(\Lambda)$ FOR $\Omega^- \rightarrow \Lambda K^- = 0.0115 \pm 0.0015$
 $\Lambda K^- \quad \alpha = 0.0157 \pm 0.0021$
 $\Lambda K^-, \bar{\Lambda} K^+ (\alpha + \bar{\alpha}) / (\alpha - \bar{\alpha}) = -0.02 \pm 0.13$
 $\Xi^0 \pi^- \quad \alpha = 0.09 \pm 0.14$
 $\Xi^- \pi^0 \quad \alpha = 0.05 \pm 0.21$

| Ω^- DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | p (MeV/c) |
|--|--------------------------------------|------------------|----------------|
| ΛK^- | $(67.8 \pm 0.7) \%$ | | 211 |
| $\Xi^0 \pi^-$ | $(23.6 \pm 0.7) \%$ | | 294 |
| $\Xi^- \pi^0$ | $(8.6 \pm 0.4) \%$ | | 289 |
| $\Xi^- \pi^+ \pi^-$ | $(3.7^{+0.7}_{-0.6}) \times 10^{-4}$ | | 189 |
| $\Xi(1530)^0 \pi^-$ | $< 7 \times 10^{-5}$ | 90% | 17 |
| $\Xi^0 e^- \bar{\nu}_e$ | $(5.6 \pm 2.8) \times 10^{-3}$ | | 319 |
| $\Xi^- \gamma$ | $< 4.6 \times 10^{-4}$ | 90% | 314 |
| $\Delta S = 2$ forbidden ($S2$) modes | | | |
| $\Lambda \pi^-$ | $S2 \quad < 2.9 \times 10^{-6}$ | 90% | 449 |

$\Omega(2012)^-$

$$I(J^P) = 0(?^-)$$

Mass $m = 2012.4 \pm 0.9$ MeV
 Full width $\Gamma = 6.4^{+3.0}_{-2.6}$ MeV

Branching fractions are given relative to the one **DEFINED AS 1**.

| $\Omega(2012)^-$ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | p (MeV/c) |
|------------------------------|--------------------------------|------------------|-------------|
| $\Xi^0 K^-$ | DEFINED AS 1 | | 403 |
| $\Xi^- \bar{K}^0$ | 0.83 ± 0.21 | | 392 |
| $\Xi^0 \pi^0 K^-$ | < 0.30 | 90% | 245 |
| $\Xi^0 \pi^- \bar{K}^0$ | < 0.21 | 90% | 230 |
| $\Xi^- \pi^0 \bar{K}^0$ | < 0.7 | 90% | 226 |
| $\Xi^- \pi^+ K^-$ | < 0.08 | 90% | 224 |

$\Omega(2250)^-$

$$I(J^P) = 0(?^?)$$

Mass $m = 2252 \pm 9$ MeV
 Full width $\Gamma = 55 \pm 18$ MeV

| $\Omega(2250)^-$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\Xi^- \pi^+ K^-$ | seen | 532 |
| $\Xi(1530)^0 K^-$ | seen | 437 |