

Z_b(10650)

$$I^G(J^{PC}) = 1^+(1^{+-})$$

I, G, C need confirmation.

was X(10650)[±]

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Observed by BONDAR 12 in $\Upsilon(5S)$ decays to $\Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses.

Z_b(10650) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10652.2 ± 1.5	¹ BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10656.7 ± 5.0 ^{+1.1} _{-3.1}	² GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10650.7 ± 1.5 ^{+0.5} _{-0.2}	² GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10651.2 ± 1.0 ^{+0.4} _{-0.3}	² GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10657 ± 6 ± 3	³ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10651 ± 2 ± 3	³ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10652 ± 1 ± 2	³ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10654 ± 3 ⁺¹ ₋₂	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10651 ⁺² ₋₃ ⁺³ ₋₂	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹ Average of the BONDAR 12 measurements in separate channels.

² Correlated with the corresponding result from BONDAR 12.

³ Superseded by the average measurement of BONDAR 12.

Z_b(10650) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
11.5 ± 2.2	⁴ BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
12.1 ^{+11.3} _{-4.8} ^{+2.7} _{-0.6}	⁵ GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
14.2 ± 3.7 ⁺ ₋ ^{0.9} _{0.4}	⁵ GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
9.3 ± 2.2 ⁺ ₋ ^{0.3} _{0.5}	⁵ GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
16.3 ± 9.8 ⁺ ₋ ^{6.0} _{2.0}	⁶ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
13.3 ± 3.3 ⁺ ₋ ^{4.0} _{3.0}	⁶ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
8.4 ± 2.0 ± 2.0	⁶ BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

$20.9^{+5.4}_{-4.7} + 2.1_{-5.7}$	⁶ BONDAR	12	BELL	$e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$
$19 \pm 7 + 11_{-7}$	⁶ BONDAR	12	BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$

⁴ Average of the BONDAR 12 measurements in separate channels.

⁵ Correlated with the corresponding result from BONDAR 12.

⁶ Superseded by the average measurement of BONDAR 12.

$Z_b(10650)^+$ DECAY MODES

$Z_b(10650)^-$ decay modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma(1S)\pi^+$	$(1.7^{+0.8}_{-0.6}) \times 10^{-3}$
$\Gamma_2 \quad \gamma(2S)\pi^+$	$(1.4^{+0.6}_{-0.4}) \%$
$\Gamma_3 \quad \gamma(3S)\pi^+$	$(1.6^{+0.7}_{-0.5}) \%$
$\Gamma_4 \quad h_b(1P)\pi^+$	$(8.4^{+2.9}_{-2.4}) \%$
$\Gamma_5 \quad h_b(2P)\pi^+$	$(15 \pm 4) \%$
$\Gamma_6 \quad B^+\bar{B}^0$	not seen
$\Gamma_7 \quad B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	not seen
$\Gamma_8 \quad B^{*+}\bar{B}^{*0}$	$(74^{+4}_{-6}) \%$

$Z_b(10650)$ BRANCHING RATIOS

$\Gamma(\gamma(1S)\pi^+)/\Gamma_{\text{total}} \qquad \qquad \qquad \Gamma_1/\Gamma$

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$1.7^{+0.7+0.3}_{-0.6-0.2}$	⁷ GARMASH	16	BELL $e^+ e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
-----------------------------	----------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$
------	---------	----	---

seen	BONDAR	12	BELL $e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$
------	--------	----	---

⁷ Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+\gamma(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

$\Gamma(\gamma(2S)\pi^+)/\Gamma_{\text{total}} \qquad \qquad \qquad \Gamma_2/\Gamma$

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

$1.39^{+0.48+0.34}_{-0.38-0.23}$	⁸ GARMASH	16	$e^+ e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
----------------------------------	----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$
------	---------	----	---

seen	BONDAR	12	BELL $e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$
------	--------	----	---

⁸ Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+\gamma(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

$\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.63^{+0.53+0.39}_{-0.42-0.28}$	⁹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

⁹ Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

$\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$ Γ_4/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.41^{+2.43+1.49}_{-2.12-1.06}$	¹⁰ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	¹¹ MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
seen	¹² BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

¹⁰ Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

¹¹ Using e^+e^- energies near the $\Upsilon(11020)$.

¹² Using e^+e^- energies near the $\Upsilon(10860)$.

$\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$14.7^{+3.2+2.8}_{-2.8-2.3}$	¹³ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	¹⁴ MIZUK	16	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$
seen	¹⁵ BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹³ Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

¹⁴ Using e^+e^- energies near the $\Upsilon(11020)$.

¹⁵ Using e^+e^- energies near the $\Upsilon(10860)$.

$\Gamma(B^+\bar{B}^0)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^0$

$[\Gamma(B^+\bar{B}^{*0}) + \Gamma(B^{*+}\bar{B}^0)]/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$

$\Gamma(B^{*+}\bar{B}^{*0})/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$73.7^{+3.4+2.7}_{-4.4-3.5}$	161	¹⁶ GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$

¹⁶ Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+} \bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16. Using the mass and width of the $Z_b(10650)$ from BONDAR 12.

$\Gamma(B^{*+}\bar{B}^{*0})/[\Gamma(\gamma(1S)\pi^+) + \Gamma(\gamma(2S)\pi^+) + \Gamma(\gamma(3S)\pi^+) + \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)]$					$\Gamma_8/(\Gamma_1+\Gamma_2+\Gamma_3+\Gamma_4+\Gamma_5)$
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.80^{+0.69+0.54}_{-0.40-0.36}$	161	¹⁷ GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$
----------------------------------	-----	-----------------------	----	------	--

¹⁷ Combined with the results of BONDAR 12 and MIZUK 16. Not independent from $Z_b(10650)$ branching fractions to $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+} \bar{B}^{*0}$.

$Z_b(10650)$ REFERENCES

GARMASH	16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)
GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)