

**$K_2^*(1430)$** 

$$I(J^P) = \frac{1}{2}(2^+)$$

We consider that phase-shift analyses provide more reliable determinations of the mass and width.

 **$K_2^*(1430)$  T-MATRIX POLE  $\sqrt{s}$** 

Note that  $\Gamma = -2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1424 ± 4) − i (66 ± 2) OUR ESTIMATE</b>			
(1424 ± 4) − i (66 ± 2)	<sup>1</sup> PELAEZ	17 RVUE	$\pi K \rightarrow \pi K$
<sup>1</sup> Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.			

 **$K_2^*(1430)$  MASS****CHARGED ONLY, WITH FINAL STATE  $K\pi$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1427.3 ± 1.5 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.			
1432.7 ± 0.7 <sup>+2.2</sup> <sub>-2.3</sub>	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
1420 ± 4	1587	BAUBILLIER	84B HBC	−	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
1436 ± 5.5	400	<sup>1,2</sup> CLELAND	82 SPEC	+	$30 K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1500	<sup>1,2</sup> CLELAND	82 SPEC	+	$50 K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1200	<sup>1,2</sup> CLELAND	82 SPEC	−	$50 K^+ p \rightarrow K_S^0 \pi^- p$
1423 ± 5	935	TOAFF	81 HBC	−	$6.5 K^- p \rightarrow \bar{K}^0 \pi^- p$
1428.0 ± 4.6		<sup>3</sup> MARTIN	78 SPEC	+	$10 K^\pm p \rightarrow K_S^0 \pi p$
1423.8 ± 4.6		<sup>3</sup> MARTIN	78 SPEC	−	$10 K^\pm p \rightarrow K_S^0 \pi p$
1420.0 ± 3.1	1400	AGUILAR-...	71B HBC	−	$3.9, 4.6 K^- p$
1425 ± 8.0	225	<sup>1,2</sup> BARNHAM	71C HBC	+	$K^+ p \rightarrow K^0 \pi^+ p$
1416 ± 10	220	CRENNELL	69D DBC	−	$3.9 K^- N \rightarrow \bar{K}^0 \pi^- N$
1414 ± 13.0	60	<sup>1</sup> LIND	69 HBC	+	$9 K^+ p \rightarrow K^0 \pi^+ p$
1427 ± 12	63	<sup>1</sup> SCHWEING...	68 HBC	−	$5.5 K^- p \rightarrow \bar{K} \pi N$
1423 ± 11.0	39	<sup>1</sup> BASSANO	67 HBC	−	$4.6-5.0 K^- p \rightarrow \bar{K}^0 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1428 ± 2	4300	<sup>4</sup> ABLIKIM	22L BES3		$2.0-3.08 e^+ e^- \rightarrow K^+ K^- \pi^0$
1423.4 ± 2 ± 3	24809 ± 820	<sup>5</sup> BIRD	89 LASS	−	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$


<sup>1</sup> Errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>2</sup> Number of events in peak re-evaluated by us.

<sup>3</sup> Systematic error added by us.

<sup>4</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

<sup>5</sup> From a partial wave amplitude analysis.



**NEUTRAL ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1432.4 ± 1.3 OUR AVERAGE</b>				
1431.2 ± 1.8 ± 0.7		<sup>1</sup> ASTON 88	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 4 ± 6		<sup>1</sup> ASTON 87	LASS	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1433 ± 6 ± 10		<sup>1</sup> ASTON 84B	LASS	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
1471 ± 12		<sup>1</sup> BAUBILLIER 82B	HBC	8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
1428 ± 3		<sup>1</sup> ASTON 81C	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 2		<sup>1</sup> ESTABROOKS 78	ASPK	13 $K^\pm p \rightarrow pK\pi$
1440 ± 10		<sup>1</sup> BOWLER 77	DBC	5.5 $K^+ d \rightarrow K\pi pp$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1428.5 ± 3.9	1786 ± 127	<sup>2</sup> AUBERT 07AK	BABR	10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
1420 ± 7	300	HENDRICK 76	DBC	8.25 $K^+ N \rightarrow K^+ \pi N$
1421.6 ± 4.2	800	MCCUBBIN 75	HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$
1420.1 ± 4.3		<sup>3</sup> LINGLIN 73	HBC	2-13 $K^+ p \rightarrow K^+ \pi^- X$
1419.1 ± 3.7	1800	AGUILAR-... 71B	HBC	3.9,4.6 $K^- p$
1416 ± 6	600	CORDS 71	DBC	9 $K^+ n \rightarrow K^+ \pi^- p$
1421.1 ± 2.6	2200	DAVIS 69	HBC	12 $K^+ p \rightarrow K^+ \pi^- X$

<sup>1</sup> From phase shift or partial-wave analysis.


<sup>2</sup> Systematic errors not estimated.<sup>3</sup> From pole extrapolation, using world  $K^+ p$  data summary tape. **$K_2^*(1430)$  WIDTH****CHARGED ONLY, WITH FINAL STATE  $K\pi$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>100.0 ± 2.2 OUR FIT</b>	Error includes scale factor of 1.1.				
<b>100.0 ± 2.2 OUR AVERAGE</b>	Error includes scale factor of 1.1.				
102.5 ± 1.6 <sup>+3.1</sup> <sub>-2.8</sub>	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
109 ± 22	400	<sup>1,2</sup> CLELAND	82 SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$
124 ± 12.8	1500	<sup>1,2</sup> CLELAND	82 SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$
113 ± 12.8	1200	<sup>1,2</sup> CLELAND	82 SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$
85 ± 16	935	TOAFF	81 HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
96.5 ± 3.8		MARTIN	78 SPEC	+	10 $K^\pm p \rightarrow K_S^0 \pi p$
97.7 ± 4.0		MARTIN	78 SPEC	-	10 $K^\pm p \rightarrow K_S^0 \pi p$
94.7 <sup>+15.1</sup> <sub>-12.5</sub>	1400	AGUILAR-...	71B HBC	-	3.9,4.6 $K^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
107 ± 4	4300	<sup>3</sup> ABLIKIM	22L BES3		2.0-3.08 $e^+ e^- \rightarrow K^+ K^- \pi^0$
98 ± 4 ± 4	25k	<sup>4</sup> BIRD	89 LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

<sup>1</sup> Errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.<sup>2</sup> Number of events in peak re-evaluated by us.<sup>3</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.<sup>4</sup> From a partial wave amplitude analysis.**NEUTRAL ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>109 ± 5 OUR AVERAGE</b>	Error includes scale factor of 1.9. See the ideogram below.			
116.5 ± 3.6 ± 1.7		<sup>1</sup> ASTON	88 LASS	11 $K^- p \rightarrow K^- \pi^+ n$
129 ± 15 ± 15		<sup>1</sup> ASTON	87 LASS	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
131 ± 24 ± 20		<sup>1</sup> ASTON	84B LASS	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
143 ± 34		<sup>1</sup> BAUBILLIER	82B HBC	8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
98 ± 8		<sup>1</sup> ASTON	81C LASS	11 $K^- p \rightarrow K^- \pi^+ n$
140 ± 30		<sup>1</sup> ETKIN	80 SPEC	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
98 ± 5		<sup>1</sup> ESTABROOKS	78 ASPK	13 $K^\pm p \rightarrow pK\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
113.7 ± 9.2	1786 ± 127	<sup>2</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
125 ± 29	300	<sup>3</sup> HENDRICK	76 DBC	8.25 $K^+ N \rightarrow K^+ \pi N$
116 ± 18	800	MCCUBBIN	75 HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$
61 ± 14		<sup>4</sup> LINGLIN	73 HBC	2-13 $K^+ p \rightarrow K^+ \pi^- X$
116.6 <sup>+10.3</sup> <sub>-15.5</sub>	1800	AGUILAR-...	71B HBC	3.9,4.6 $K^- p$
144 ± 24.0	600	<sup>3</sup> CORDS	71 DBC	9 $K^+ n \rightarrow K^+ \pi^- p$
101 ± 10	2200	DAVIS	69 HBC	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$

- <sup>1</sup> From phase shift or partial-wave analysis.
- <sup>2</sup> Systematic errors not estimated.
- <sup>3</sup> Errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.
- <sup>4</sup> From pole extrapolation, using world  $K^+ \rho$  data summary tape.



### $K_2^*(1430)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $K\pi$	$(49.9 \pm 1.2) \%$	
$\Gamma_2$ $K^*(892)\pi$	$(24.7 \pm 1.5) \%$	
$\Gamma_3$ $K^*(892)\pi\pi$	$(13.4 \pm 2.2) \%$	
$\Gamma_4$ $K\rho$	$(8.7 \pm 0.8) \%$	S=1.2
$\Gamma_5$ $K\omega$	$(2.9 \pm 0.8) \%$	
$\Gamma_6$ $K^+\gamma$	$(2.4 \pm 0.5) \times 10^{-3}$	S=1.1
$\Gamma_7$ $K\eta$	$(1.5^{+3.4}_{-1.0}) \times 10^{-3}$	S=1.3
$\Gamma_8$ $K\omega\pi$	$< 7.2 \times 10^{-4}$	CL=95%
$\Gamma_9$ $K^0\gamma$	$< 9 \times 10^{-4}$	CL=90%

## CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 10 branching ratios uses 32 measurements and one constraint to determine 8 parameters. The overall fit has a  $\chi^2 = 21.1$  for 25 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-9						
$x_3$	-40	-73					
$x_4$	-8	36	-52				
$x_5$	-11	-3	-26	-7			
$x_6$	-1	-1	-1	-1	0		
$x_7$	-4	-7	-5	-5	-2	0	
$\Gamma$	0	0	0	0	0	-11	0
	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$	$x_7$

	Mode	Rate (MeV)	Scale factor
$\Gamma_1$	$K\pi$	$49.9 \pm 1.6$	
$\Gamma_2$	$K^*(892)\pi$	$24.7 \pm 1.6$	
$\Gamma_3$	$K^*(892)\pi\pi$	$13.5 \pm 2.3$	
$\Gamma_4$	$K\rho$	$8.7 \pm 0.8$	1.2
$\Gamma_5$	$K\omega$	$2.9 \pm 0.8$	
$\Gamma_6$	$K^+\gamma$	$0.24 \pm 0.05$	1.1
$\Gamma_7$	$K\eta$	$0.15^{+0.34}_{-0.10}$	1.3

### $K_2^*(1430)$ PARTIAL WIDTHS

#### $\Gamma(K^+\gamma)$ $\Gamma_6$

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>240 \pm 50</math> OUR FIT</b>	Error includes scale factor of 1.1.			
<b><math>240 \pm 45</math></b>	CIHANGIR	82	SPEC	+
				200 $K^+ Z \rightarrow Z K^+ \pi^0$ , $Z K_S^0 \pi^+$

#### $\Gamma(K^0\gamma)$ $\Gamma_9$

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt; 5.4</b>	90	ALAVI-HARATI02B	KTEV		$K + A \rightarrow K^* + A$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<84	90	CARLSMITH	87	SPEC	0
					60-200 $K_L^0 A \rightarrow$ $K_S^0 \pi^0 A$

**$K_2^*(1430)$  BRANCHING RATIOS** **$\Gamma(K\pi)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

**0.499±0.012 OUR FIT****0.488±0.014 OUR AVERAGE**0.485±0.006±0.020 <sup>1</sup> ASTON 88 LASS 0 11  $K^- p \rightarrow K^- \pi^+ n$ 0.49 ±0.02 <sup>1</sup> ESTABROOKS 78 ASPK ± 13  $K^\pm p \rightarrow pK\pi$ <sup>1</sup> From phase shift analysis. **$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$   $\Gamma_2/\Gamma_1$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

**0.496±0.034 OUR FIT****0.47 ±0.04 OUR AVERAGE**0.44 ±0.09 ASTON 84B LASS 0 11  $K^- p \rightarrow \bar{K}^0 2\pi n$ 0.62 ±0.19 LAUSCHER 75 HBC 0 10,16  $K^- p \rightarrow K^- \pi^+ n$ 0.54 ±0.16 DEHM 74 DBC 0 4.6  $K^+ N$ 0.47 ±0.08 AGUILAR-... 71B HBC 3.9,4.6  $K^- p$ 0.47 ±0.10 BASSANO 67 HBC -0 4.6,5.0  $K^- p$ 0.45 ±0.13 BADIER 65C HBC - 3  $K^- p$  **$\Gamma(K\omega)/\Gamma(K\pi)$   $\Gamma_5/\Gamma_1$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

**0.059±0.017 OUR FIT****0.070±0.035 OUR AVERAGE**0.05 ±0.04 AGUILAR-... 71B HBC 3.9,4.6  $K^- p$ 0.13 ±0.07 BASSOMPIE... 69 HBC 0 5  $K^+ p$  **$\Gamma(K\rho)/\Gamma(K\pi)$   $\Gamma_4/\Gamma_1$** 


<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

**0.174±0.017 OUR FIT** Error includes scale factor of 1.2.**0.150<sup>+0.029</sup><sub>-0.017</sub> OUR AVERAGE**0.18 ±0.05 ASTON 84B LASS 0 11  $K^- p \rightarrow \bar{K}^0 2\pi n$ 0.02 <sup>+0.10</sup><sub>-0.02</sub> DEHM 74 DBC 0 4.6  $K^+ N$ 0.16 ±0.05 AGUILAR-... 71B HBC 3.9,4.6  $K^- p$ 0.14 ±0.10 BASSANO 67 HBC -0 4.6,5.0  $K^- p$ 0.14 ±0.07 BADIER 65C HBC - 3  $K^- p$  **$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$   $\Gamma_4/\Gamma_2$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

**0.350±0.031 OUR FIT** Error includes scale factor of 1.4.**0.354±0.033 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.0.293±0.032±0.020 ASTON 87 LASS 0 11  $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$ 0.38 ±0.09 BAUBILLIER 82B HBC 0 8.25  $K^- p \rightarrow NK_S^0 \pi \pi$ 0.39 ±0.03 DAUM 81C CNTR 63  $K^- p \rightarrow K^- 2\pi p$

WEIGHTED AVERAGE  
 $0.354 \pm 0.033$  (Error scaled by 1.4)



**$\Gamma(K\omega)/\Gamma(K^*(892)\pi)$   $\Gamma_5/\Gamma_2$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>0.118 \pm 0.034</math> OUR FIT</b>				
<b><math>0.10 \pm 0.04</math></b>	FIELD	67	HBC	— 3.8 $K^- p$

**$\Gamma(K\eta)/\Gamma(K^*(892)\pi)$   $\Gamma_7/\Gamma_2$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>0.006^{+0.014}_{-0.004}</math> OUR FIT</b>	Error includes scale factor of 1.2.			
<b><math>0.07 \pm 0.04</math></b>	FIELD	67	HBC	— 3.8 $K^- p$

**$\Gamma(K\eta)/\Gamma(K\pi)$   $\Gamma_7/\Gamma_1$**

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>0.0030^{+0.0070}_{-0.0020}</math> OUR FIT</b>		Error includes scale factor of 1.3.			
<b>0 ± 0.0056</b>		<sup>1</sup> ASTON	88B	LASS	— 11 $K^- p \rightarrow K^- \eta p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.04	95	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p$
<0.065		<sup>2</sup> BASSOMPIE...	69	HBC	5.0 $K^+ p$
<0.02		BISHOP	69	HBC	3.5 $K^+ p$

<sup>1</sup> ASTON 88B quote < 0.0092 at CL=95%. We convert this to a central value and 1 sigma error in order to be able to use it in our constrained fit.

<sup>2</sup> Restated by us.

$\Gamma(K^*(892)\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

**0.134 ± 0.022 OUR FIT****0.12 ± 0.04** <sup>1</sup> GOLDBERG 76 HBC - 3  $K^- p \rightarrow p \bar{K}^0 \pi \pi$ <sup>1</sup> Assuming  $\pi\pi$  system has isospin 1, which is supported by the data. $\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$   $\Gamma_3/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

**0.27 ± 0.05 OUR FIT****0.21 ± 0.08** <sup>1,2</sup> JONGEJANS 78 HBC - 4  $K^- p \rightarrow p \bar{K}^0 \pi \pi$ <sup>1</sup> Restated by us.<sup>2</sup> Assuming  $\pi\pi$  system has isospin 1, which is supported by the data. $\Gamma(K\omega\pi)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**<0.72** 95 0 JONGEJANS 78 HBC 4  $K^- p \rightarrow p \bar{K}^0 4\pi$  **$K_2^*(1430)$  REFERENCES**

ABLIKIM 22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PELAEZ 17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira	
AUBERT 07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
ALAVI-HARATI 02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
BIRD 89	SLAC-332	P.F. Bird	(SLAC)
ASTON 88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON 88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON 87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
CARLSMITH 87	PR D36 3502	D. Carlsmith <i>et al.</i>	(EFI, SACL)
ASTON 84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)
BAUBILLIER 84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
BAUBILLIER 82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
CIHANGIR 82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)
CLELAND 82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
ASTON 81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
DAUM 81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
TOAFF 81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)
ETKIN 80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
ESTABROOKS 78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)
Also	PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)
JONGEJANS 78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)
MARTIN 78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)
BOWLER 77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)
GOLDBERG 76	LNC 17 253	J. Goldberg	(HAIF)
HENDRICK 76	NP B112 189	K. Hendrickx <i>et al.</i>	(MONS, SACL, PARIS+)
LAUSCHER 75	NP B86 189	P. Lauscher <i>et al.</i>	(ABCLV Collab.) JP
MCCUBBIN 75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)
DEHM 74	NP B75 47	G. Dehm <i>et al.</i>	(MPIM, BRUX, MONS, CERN)
LINGLIN 73	NP B55 408	D. Linglin	(CERN)
AGUILAR-... 71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)
BARNHAM 71C	NP B28 171	K.W.J. Barnham <i>et al.</i>	(BIRM, GLAS)
CORDS 71	PR D4 1974	D. Cords <i>et al.</i>	(PURD, UCD, IUPU)
BASSOMPIE... 69	NP B13 189	G. Bassompierre <i>et al.</i>	(CERN, BRUX) JP
BISHOP 69	NP B9 403	J.M. Bishop <i>et al.</i>	(WISC)
CRENNELL 69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)
DAVIS 69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)
LIND 69	NP B14 1	V.G. Lind <i>et al.</i>	(LRL) JP
SCHWEING... 68	PR 166 1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)
Also	Thesis	F.L. Schweingruber	(NWES, NWES)
BASSANO 67	PRL 19 968	D. Bassano <i>et al.</i>	(BNL, SYRA)
FIELD 67	PL 24B 638	J.H. Field <i>et al.</i>	(UCSD)
BADIER 65C	PL 19 612	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)