

$\Sigma(1670) 3/2^-$ $I(J^P) = 1(\frac{3}{2}^-)$ Status: ****

For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** 1 (1982).

Results from production experiments are listed separately in the next entry.

$\Sigma(1670)$ POLE POSITION

REAL PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|-------------------------|
| 1655 to 1675 (≈ 1662) OUR ESTIMATE | | | |
| 1661 ± 3 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| 1669^{+7}_{-7} | ¹ KAMANO 15 | DPWA | $\bar{K}N$ multichannel |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 1674 | ZHANG 13A | DPWA | $\bar{K}N$ multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | |

–2×IMAGINARY PART

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|------------------------|------|-------------------------|
| 45 to 65 (≈ 55) OUR ESTIMATE | | | |
| 52 ± 6 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| 64^{+10}_{-14} | ¹ KAMANO 15 | DPWA | $\bar{K}N$ multichannel |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 54 | ZHANG 13A | DPWA | $\bar{K}N$ multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | |

$\Sigma(1670)$ POLE RESIDUES

The normalized residue is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow N\bar{K}$

| MODULUS | PHASE ($^\circ$) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------------------|------------------------|------|-------------------------|
| 0.10 \pm 0.02 | –31 \pm 12 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.129 | –20 | ¹ KAMANO 15 | DPWA | $\bar{K}N$ multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | | |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma\pi$

| MODULUS | PHASE ($^\circ$) | DOCUMENT ID | TECN | COMMENT |
|---|--------------------------------|------------------------|------|-------------------------|
| 0.25 \pm 0.05 | –25 \pm 10 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.249 | –21 | ¹ KAMANO 15 | DPWA | $\bar{K}N$ multichannel |
| ¹ From the preferred solution A in KAMANO 15. | | | | |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda\pi$

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|---|------------------------|-------------|-------------------------|
| 0.09 ± 0.03 | -52 ± 12 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| • • • | We do not use the following data for averages, fits, limits, etc. | • • • | | |
| 0.0818 | -7 | ¹ KAMANO 15 | DPWA | $\bar{K}N$ multichannel |

¹From the preferred solution A in KAMANO 15.**Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Xi K$**

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.02 ± 0.01 | 160 ± 20 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma\sigma$

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.08 ± 0.03 | -25 ± 15 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma(1385)\pi$, S-wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------|---|------------------------|-------------|-------------------------|
| • • • | We do not use the following data for averages, fits, limits, etc. | • • • | | |
| 0.228 | 167 | ¹ KAMANO 15 | DPWA | $\bar{K}N$ multichannel |

¹From the preferred solution A in KAMANO 15.**Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma(1385)\pi$, D-wave**

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|----------------|---|--------------------|-------------|-------------------------|
| • • • | We do not use the following data for averages, fits, limits, etc. | • • • | | |
| 0.0915 | 141 | KAMANO 15 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda(1405)\pi$

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.03 ± 0.02 | 160 ± 15 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda(1520)\pi$, P-wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.04 ± 0.02 | 120 ± 20 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda(1520)\pi$, F-wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.01 ± 0.01 | | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Delta\bar{K}$, S-wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.01 ± 0.01 | | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow N\bar{K}^*(892)$, S=3/2, S-wave

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------------------|--------------------|-------------|-------------------------|
| 0.05 ± 0.03 @ 50 ± 60 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow N\bar{K}^*(892)$, S=3/2, D-wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.01 ± 0.01 | | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

Normalized residue in $N\bar{K} \rightarrow \Sigma(1670) \rightarrow N\bar{K}^*(892)$, $S=1/2$, D -wave

| <u>MODULUS</u> | <u>PHASE ($^\circ$)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|------------------|------------------------------------|--------------------|-------------|-------------------------|
| 0.03±0.02 | | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |

 $\Sigma(1670)$ MASS

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------------|-------------|--|
| 1665 to 1685 (≈ 1675) OUR ESTIMATE | | | |
| 1665 ± 3 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| 1678 ± 2 | ZHANG 13A | DPWA | $\bar{K}N$ multichannel |
| 1673 ± 1 | GAO 12 | DPWA | $\bar{K}N \rightarrow \Lambda\pi$ |
| 1665.1± 4.1 | KOISO 85 | DPWA | $K^-p \rightarrow \Sigma\pi$ |
| 1682 ± 5 | GOPAL 80 | DPWA | $\bar{K}N \rightarrow \bar{K}N$ |
| 1679 ±10 | ALSTON-... | DPWA | $\bar{K}N \rightarrow \bar{K}N$ |
| 1670 ± 5 | GOPAL 77 | DPWA | $\bar{K}N$ multichannel |
| 1670 ± 6 | HEPP 76B | DPWA | $K^-N \rightarrow \Sigma\pi$ |
| 1685 ±20 | BAILLON 75 | IPWA | $\bar{K}N \rightarrow \Lambda\pi$ |
| 1659 $^{+12}_{-5}$ | VANHORN 75 | DPWA | $K^-p \rightarrow \Lambda\pi^0$ |
| 1670 ± 2 | KANE 74 | DPWA | $K^-p \rightarrow \Sigma\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 1667 or 1668 | ¹ MARTIN 77 | DPWA | $\bar{K}N$ multichannel |
| 1650 | DEBELLEFON 76 | IPWA | $K^-p \rightarrow \Lambda\pi^0$ |
| 1671 ± 3 | PONTE 75 | DPWA | $K^-p \rightarrow \Lambda\pi^0$ (sol. 1) |
| 1655 ± 2 | PONTE 75 | DPWA | $K^-p \rightarrow \Lambda\pi^0$ (sol. 2) |

¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

 $\Sigma(1670)$ WIDTH

| <u>VALUE (MeV)</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|---|------------------------|-------------|--|
| 40 to 100 (≈ 70) OUR ESTIMATE | | | |
| 54 ± 6 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| 55 ± 4 | ZHANG 13A | DPWA | $\bar{K}N$ multichannel |
| 52 $^{+5}_{-2}$ | GAO 12 | DPWA | $\bar{K}N \rightarrow \Lambda\pi$ |
| 65.0± 7.3 | KOISO 85 | DPWA | $K^-p \rightarrow \Sigma\pi$ |
| 79 ±10 | GOPAL 80 | DPWA | $\bar{K}N \rightarrow \bar{K}N$ |
| 56 ±20 | ALSTON-... | DPWA | $\bar{K}N \rightarrow \bar{K}N$ |
| 50 ± 5 | GOPAL 77 | DPWA | $\bar{K}N$ multichannel |
| 56 ± 3 | HEPP 76B | DPWA | $K^-N \rightarrow \Sigma\pi$ |
| 85 ±25 | BAILLON 75 | IPWA | $\bar{K}N \rightarrow \Lambda\pi$ |
| 32 ±11 | VANHORN 75 | DPWA | $K^-p \rightarrow \Lambda\pi^0$ |
| 79 ± 6 | KANE 74 | DPWA | $K^-p \rightarrow \Sigma\pi$ |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | |
| 46 or 46 | ¹ MARTIN 77 | DPWA | $\bar{K}N$ multichannel |
| 80 | DEBELLEFON 76 | IPWA | $K^-p \rightarrow \Lambda\pi^0$ |
| 44 ±11 | PONTE 75 | DPWA | $K^-p \rightarrow \Lambda\pi^0$ (sol. 1) |
| 76 ± 5 | PONTE 75 | DPWA | $K^-p \rightarrow \Lambda\pi^0$ (sol. 2) |

¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

$\Sigma(1670)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|---|--------------------------------|
| Γ_1 $N\bar{K}$ | 0.06 to 0.12 |
| Γ_2 $\Lambda\pi$ | 5–15 % |
| Γ_3 $\Sigma\pi$ | 30–60 % |
| Γ_4 $\Lambda\pi\pi$ | |
| Γ_5 $\Sigma\pi\pi$ | |
| Γ_6 $\Sigma\sigma$ | (7.0 \pm 3.0) % |
| Γ_7 $\Sigma(1385)\pi$ | |
| Γ_8 $\Sigma(1385)\pi$, <i>S</i> -wave | |
| Γ_9 $\Sigma(1385)\pi$, <i>D</i> -wave | |
| Γ_{10} $N\bar{K}^*(892)$, <i>S</i> =1/2, <i>D</i> -wave | |
| Γ_{11} $N\bar{K}^*(892)$, <i>S</i> =3/2, <i>S</i> -wave | |
| Γ_{12} $N\bar{K}^*(892)$, <i>S</i> =3/2, <i>D</i> -wave | |
| Γ_{13} $\Lambda(1405)\pi$ | |
| Γ_{14} $\Lambda(1520)\pi$ | |

 $\Sigma(1670)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on Λ and Σ Resonances.

| $\Gamma(N\bar{K})/\Gamma_{\text{total}}$ | | | | Γ_1/Γ |
|---|---------------------|------|--------------------------------------|-------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| 0.06 to 0.12 OUR ESTIMATE | | | | |
| 0.10 \pm 0.02 | SARANTSEV | 19 | DPWA $\bar{K}N$ multichannel | |
| 0.062 \pm 0.007 | ZHANG | 13A | DPWA $\bar{K}N$ multichannel | |
| 0.10 \pm 0.03 | GOPAL | 80 | DPWA $\bar{K}N \rightarrow \bar{K}N$ | |
| 0.11 \pm 0.03 | ALSTON-... | 78 | DPWA $\bar{K}N \rightarrow \bar{K}N$ | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.121 | ¹ KAMANO | 15 | DPWA $\bar{K}N$ multichannel | |
| 0.08 \pm 0.03 | GOPAL | 77 | DPWA See GOPAL 80 | |
| 0.07 or 0.07 | ² MARTIN | 77 | DPWA $\bar{K}N$ multichannel | |

¹ From the preferred solution A in KAMANO 15.

² The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

| $\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$ | | | | Γ_2/Γ |
|---|---------------------|------|------------------------------|-------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| 0.09 \pm 0.02 | SARANTSEV | 19 | DPWA $\bar{K}N$ multichannel | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 0.058 | ¹ KAMANO | 15 | DPWA $\bar{K}N$ multichannel | |

¹ From the preferred solution A in KAMANO 15.

| $\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$ | | | | Γ_3/Γ |
|---|-------------|------|------------------------------|-------------------|
| VALUE | DOCUMENT ID | TECN | COMMENT | |
| 0.70 \pm 0.15 | SARANTSEV | 19 | DPWA $\bar{K}N$ multichannel | |

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.465 ¹KAMANO 15 DPWA $\bar{K}N$ multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(\Lambda\pi\pi)/\Gamma_{\text{total}}$ **Γ_4/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11 ARMENTEROS68E HBC $K^- p$ ($\Gamma_1=0.09$)

$\Gamma(\Sigma\pi\pi)/\Gamma_{\text{total}}$ **Γ_5/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.14 ¹ARMENTEROS68E HBC $K^- p, K^- d$ ($\Gamma_1=0.09$)

¹Ratio only for $\Sigma 2\pi$ system in $l = 1$, which cannot be $\Sigma(1385)$.

$\Gamma(\Sigma\sigma)/\Gamma_{\text{total}}$ **Γ_6/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

0.07±0.03 SARANTSEV 19 DPWA $\bar{K}N$ multichannel

$\Gamma(\Sigma(1385)\pi, S\text{-wave})/\Gamma_{\text{total}}$ **Γ_8/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.309 ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$ **Γ_9/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.044 ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=1/2, D\text{-wave})/\Gamma_{\text{total}}$ **Γ_{10}/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.001 ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, S\text{-wave})/\Gamma_{\text{total}}$ **Γ_{11}/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.002 ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}}$ **Γ_{12}/Γ**

| <u>VALUE</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>COMMENT</u> |
|--------------|--------------------|-------------|----------------|
|--------------|--------------------|-------------|----------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.001 ¹KAMANO 15 DPWA Multichannel

¹From the preferred solution A in KAMANO 15.

$$\Gamma(\Lambda(1405)\pi)/\Gamma_{\text{total}} \qquad \Gamma_{13}/\Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------|---------------|------|------------------------------------|
| 0.01 ± 0.01 | SARANTSEV 19 | DPWA | $\bar{K}N$ multichannel |
| < 0.06 | ARMENTEROS68E | HBC | $K^- p, K^- d$ ($\Gamma_1=0.09$) |

$$\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma(1385)\pi) \qquad \Gamma_{13}/\Gamma_7$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-----------------|-------------|------|------------------------------------|
| 0.23 ± 0.08 | BRUCKER 70 | DBC | $K^- N \rightarrow \Sigma \pi \pi$ |

$$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda\pi \qquad (\Gamma_1 \Gamma_2)^{1/2}/\Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|----------------------------|------------------------|------|-----------------------------------|
| $+0.08 \pm 0.01$ | ZHANG 13A | DPWA | Multichannel |
| $+0.081^{+0.002}_{-0.004}$ | GAO 12 | DPWA | $\bar{K}N \rightarrow \Lambda\pi$ |
| $+0.17 \pm 0.03$ | ¹ MORRIS 78 | DPWA | $K^- n \rightarrow \Lambda\pi^-$ |
| $+0.13 \pm 0.02$ | ¹ MORRIS 78 | DPWA | $K^- n \rightarrow \Lambda\pi^-$ |
| $+0.10 \pm 0.02$ | GOPAL 77 | DPWA | $\bar{K}N$ multichannel |
| $+0.06 \pm 0.02$ | BAILLON 75 | IPWA | $\bar{K}N \rightarrow \Lambda\pi$ |
| $+0.09 \pm 0.02$ | VANHORN 75 | DPWA | $K^- p \rightarrow \Lambda\pi^0$ |
| $+0.018 \pm 0.060$ | DEVENISH 74B | | Fixed- t dispersion rel. |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------|------------------------|------|---|
| $+0.08$ or $+0.08$ | ² MARTIN 77 | DPWA | $\bar{K}N$ multichannel |
| $+0.05$ | DEBELLEFON 76 | IPWA | $K^- p \rightarrow \Lambda\pi^0$ |
| $+0.08 \pm 0.01$ | PONTE 75 | DPWA | $K^- p \rightarrow \Lambda\pi^0$ (sol. 1) |
| $+0.17 \pm 0.01$ | PONTE 75 | DPWA | $K^- p \rightarrow \Lambda\pi^0$ (sol. 2) |

¹ Results are with and without an S_{11} $\Sigma(1620)$ in the fit.

² The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

$$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma\pi \qquad (\Gamma_1 \Gamma_3)^{1/2}/\Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|-------------|------|-------------------------------|
| $+0.20 \pm 0.01$ | ZHANG 13A | DPWA | Multichannel |
| $+0.20 \pm 0.02$ | KOISO 85 | DPWA | $K^- p \rightarrow \Sigma\pi$ |
| $+0.21 \pm 0.02$ | GOPAL 77 | DPWA | $\bar{K}N$ multichannel |
| $+0.20 \pm 0.01$ | HEPP 76B | DPWA | $K^- N \rightarrow \Sigma\pi$ |
| $+0.21 \pm 0.03$ | KANE 74 | DPWA | $K^- p \rightarrow \Sigma\pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|--------------------|------------------------|------|-------------------------|
| $+0.18$ or $+0.17$ | ¹ MARTIN 77 | DPWA | $\bar{K}N$ multichannel |
|--------------------|------------------------|------|-------------------------|

¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.

$$(\Gamma_i \Gamma_f)^{1/2}/\Gamma_{\text{total}} \text{ in } N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma(1385)\pi, \text{ S-wave} \qquad (\Gamma_1 \Gamma_8)^{1/2}/\Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|------------------|----------------------|------|-------------------------------------|
| $+0.11 \pm 0.03$ | PREVOST 74 | DPWA | $K^- N \rightarrow \Sigma(1385)\pi$ |
| 0.17 ± 0.02 | ¹ SIMS 68 | DBC | $K^- N \rightarrow \Lambda\pi\pi$ |

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ SIMS 68 uses only cross-section data. Result used as upper limit only.

$$\Gamma_i \Gamma_f / \Gamma_{\text{total}}^2 \text{ in } N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda(1405)\pi \qquad \Gamma_1 \Gamma_{13} / \Gamma^2$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|-------------------|----------------------|----|--|
| 0.007 ± 0.002 | ¹ BRUCKER | 70 | DBC $K^- N \rightarrow \Sigma \pi \pi$ |
|-------------------|----------------------|----|--|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | |
|-------|--------|----|----------------------------|
| <0.03 | BERLEY | 69 | HBC $K^- p$ 0.6–0.82 GeV/c |
|-------|--------|----|----------------------------|

¹ Assuming the $\Lambda(1405)\pi$ cross-section bump is due only to $3/2^-$ resonance.

$$(\Gamma_i \Gamma_f)^{1/2} / \Gamma_{\text{total}} \text{ in } N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda(1520)\pi \qquad (\Gamma_1 \Gamma_{14})^{1/2} / \Gamma$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

| | | | |
|-------------------|----------------------|----|----------------------|
| 0.081 ± 0.016 | ¹ CAMERON | 77 | DPWA P -wave decay |
|-------------------|----------------------|----|----------------------|

¹ The CAMERON 77 upper limit on F -wave decay is 0.03.

$\Sigma(1670)$ REFERENCES

| | | | | |
|--------------------------|-----|-------------------|---|-------------------------|
| SARANTSEV | 19 | EPJ A55 180 | A.V. Sarantsev <i>et al.</i> | (BONN, PNPI) |
| KAMANO | 15 | PR C92 025205 | H. Kamano <i>et al.</i> | (ANL, OSAK) |
| ZHANG | 13A | PR C88 035205 | H. Zhang <i>et al.</i> | (KSU) |
| GAO | 12 | PR C86 025201 | P. Gao, J. Shi, B.S. Zou | (BHEP, BEIJT) |
| Also | | NP A867 41 | P. Gao, B.S. Zou, A. Sibirtsev | (BHEP, BEIJT+) |
| KOISO | 85 | NP A433 619 | H. Koiso <i>et al.</i> | (TOKY, MASA) |
| PDG | 82 | PL 111B 1 | M. Roos <i>et al.</i> | (HELS, CIT, CERN) |
| GOPAL | 80 | Toronto Conf. 159 | G.P. Gopal | (RHEL) IJP |
| ALSTON-... | 78 | PR D18 182 | M. Alston-Garnjost <i>et al.</i> | (LBL, MTHO+) IJP |
| Also | | PRL 38 1007 | M. Alston-Garnjost <i>et al.</i> | (LBL, MTHO+) IJP |
| MORRIS | 78 | PR D17 55 | W.A. Morris <i>et al.</i> | (FSU) IJP |
| CAMERON | 77 | NP B131 399 | W. Cameron <i>et al.</i> | (RHEL, LOIC) IJP |
| GOPAL | 77 | NP B119 362 | G.P. Gopal <i>et al.</i> | (LOIC, RHEL) IJP |
| MARTIN | 77 | NP B127 349 | B.R. Martin, M.K. Pidcock, R.G. Moorhouse | (LOUC+) IJP |
| Also | | NP B126 266 | B.R. Martin, M.K. Pidcock | (LOUC) |
| Also | | NP B126 285 | B.R. Martin, M.K. Pidcock | (LOUC) IJP |
| DEBELLEFON | 76 | NP B109 129 | A. de Bellefon, A. Berthon | (CDEF) IJP |
| HEPP | 76B | PL 65B 487 | V. Hepp <i>et al.</i> | (CERN, HEIDH, MPIM) IJP |
| BAILLON | 75 | NP B94 39 | P.H. Baillon, P.J. Litchfield | (CERN, RHEL) IJP |
| PONTE | 75 | PR D12 2597 | R.A. Ponte <i>et al.</i> | (MASA, TENN, UCR) IJP |
| VANHORN | 75 | NP B87 145 | A.J. van Horn | (LBL) IJP |
| Also | | NP B87 157 | A.J. van Horn | (LBL) IJP |
| DEVENISH | 74B | NP B81 330 | R.C.E. Devenish, C.D. Froggatt, B.R. Martin | (DESY+) IJP |
| KANE | 74 | LBL-2452 | D.F. Kane | (LBL) IJP |
| PREVOST | 74 | NP B69 246 | J. Prevost <i>et al.</i> | (SACL, CERN, HEID) |
| BRUCKER | 70 | Duke Conf. 155 | E.B. Brucker <i>et al.</i> | (FSU) I |
| Hyperon Resonances, 1970 | | | | |
| BERLEY | 69 | PL 30B 430 | D. Berley <i>et al.</i> | (BNL) |
| ARMENTEROS | 68E | PL 28B 521 | R. Armenteros <i>et al.</i> | (CERN, HEID, SACL) I |
| SIMS | 68 | PRL 21 1413 | W.H. Sims <i>et al.</i> | (FSU, TUFTS, BRAN) |