



$I(J^P) = \frac{1}{2}(0^-)$

Quantum numbers not measured. Values shown are quark-model predictions.

See also the $B^\pm/B^0$ ADMIXTURE and $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE sections.

See the Note “Production and Decay of b -flavored Hadrons” at the beginning of the $B^\pm$ Particle Listings and the Note on “ B^0 - $\bar{B}^0$ Mixing” near the end of the B^0 Particle Listings.

B^0 MASS

The fit uses m_{B^+} , $(m_{B^0} - m_{B^+})$, and m_{B^0} to determine m_{B^+} , m_{B^0} , and the mass difference.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5279.72±0.08 OUR FIT				
5279.63±0.20 OUR AVERAGE				
5279.74±0.30±0.10		¹ AAIJ	19U LHCb	pp at 7, 8, 13 TeV
5279.6 ±0.2 ±1.0		² AAD	13U ATLS	pp at 7 TeV
5279.58±0.15±0.28		³ AAIJ	12E LHCb	pp at 7 TeV
5279.63±0.53±0.33		⁴ ACOSTA	06 CDF	$p\bar{p}$ at 1.96 TeV
5279.1 ±0.7 ±0.3	135	⁵ CSORNA	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5281.3 ±2.2 ±1.4	51	ABE	96B CDF	$p\bar{p}$ at 1.8 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5279.2 ±0.54±2.0	340	ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5278.0 ±0.4 ±2.0		BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
5279.6 ±0.7 ±2.0	40	⁶ ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.2 ±1.0 ±3.0	40	ALBRECHT	87C ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5279.5 ±1.6 ±3.0	7	⁷ ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5280.6 ±0.8 ±2.0		BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Uses $B^0 \rightarrow J/\psi p\bar{p}$ decays.				
² Measured with $B_d^0 \rightarrow J/\psi(\mu^+\mu^-) K_S^0(\pi^+\pi^-)$ decays.				
³ Uses $B^0 \rightarrow J/\psi K^0$ fully reconstructed decays.				
⁴ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+\mu^-$ decays.				
⁵ CSORNA 00 uses fully reconstructed 135 $B^0 \rightarrow J/\psi(^{\prime}) K_S^0$ events and invariant masses without beam constraint.				
⁶ ALBRECHT 90J assumes 10580 for $\Upsilon(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.				
⁷ Found using fully reconstructed decays with J/ψ . ALBRECHT 87D assume $m_{\Upsilon(4S)} = 10577$ MeV.				

$m_{B^0} - m_{B^\pm}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.05 OUR FIT			
0.33 ± 0.05 OUR AVERAGE			
$0.57 \pm 0.49 \pm 0.10$	¹ SIRUNYAN	18DF CMS	pp at 8 TeV
$0.20 \pm 0.17 \pm 0.11$	¹ AAIJ	12E LHCb	pp at 7 TeV
$0.33 \pm 0.05 \pm 0.03$	² AUBERT	08AF BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$0.53 \pm 0.67 \pm 0.14$	¹ ACOSTA	06 CDF	$p\bar{p}$ at 1.96 TeV
$0.41 \pm 0.25 \pm 0.19$	ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$-0.4 \pm 0.6 \pm 0.5$	BORTOLETTO ⁹²	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
$-0.9 \pm 1.2 \pm 0.5$	ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$2.0 \pm 1.1 \pm 0.3$	³ BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Uses exclusively reconstructed final states containing a $J/\psi \rightarrow \mu^+\mu^-$ decay.			
² Uses the B -momentum distributions in the e^+e^- rest frame.			
³ BEBEK 87 actually measure the difference between half of E_{cm} and the $B^\pm$ or B^0 mass, so the $m_{B^0} - m_{B^\pm}$ is more accurate. Assume $m_{\Upsilon(4S)} = 10580$ MeV.			

 $m_{B_H^0} - m_{B_L^0}$

See the $B^0\text{-}\bar{B}^0$ MIXING PARAMETERS section near the end of these B^0 Listings.

 B^0 MEAN LIFE

See $B^\pm/B^0/B_s^0/b$ -baryon ADMIXTURE section for data on B -hadron mean life averaged over species of bottom particles.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.517 ± 0.004 OUR EVALUATION		(Produced by HFLAV)		
$1.499 \pm 0.013 \pm 0.008$	¹	ABUDINEN	23D BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.515 \pm 0.005 \pm 0.006$	²	SIRUNYAN	18BY CMS	pp at 8 TeV
$1.534 \pm 0.019 \pm 0.021$	³	ABAZOV	15A D0	$p\bar{p}$ at 1.96 TeV
$1.499 \pm 0.013 \pm 0.005$	⁴	AAIJ	14E LHCb	pp at 7 TeV
$1.524 \pm 0.006 \pm 0.004$	⁵	AAIJ	14E LHCb	pp at 7 TeV
$1.524 \pm 0.011 \pm 0.004$	⁶	AAIJ	14R LHCb	pp at 7 TeV
$1.509 \pm 0.012 \pm 0.018$	⁷	AAD	13U ATLS	pp at 7 TeV
$1.508 \pm 0.025 \pm 0.043$	⁴	ABAZOV	12U D0	$p\bar{p}$ at 1.96 TeV
$1.507 \pm 0.010 \pm 0.008$	⁸	AALTONEN	11 CDF	$p\bar{p}$ at 1.96 TeV
$1.414 \pm 0.018 \pm 0.034$	⁹	ABAZOV	09E D0	$p\bar{p}$ at 1.96 TeV
$1.504 \pm 0.013^{+0.018}_{-0.013}$	¹⁰	AUBERT	06G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.534 \pm 0.008 \pm 0.010$	¹¹	ABE	05B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.531 \pm 0.021 \pm 0.031$	¹²	ABDALLAH	04E DLPH	$e^+e^- \rightarrow Z$
$1.523^{+0.024}_{-0.023} \pm 0.022$	¹³	AUBERT	03C BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.533 \pm 0.034 \pm 0.038$	¹⁴	AUBERT	03H BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.497 \pm 0.073 \pm 0.032$	¹⁵	ACOSTA	02C CDF	$p\bar{p}$ at 1.8 TeV
$1.529 \pm 0.012 \pm 0.029$	¹⁶	AUBERT	02H BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$1.546 \pm 0.032 \pm 0.022$		17 AUBERT	01F BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$1.541 \pm 0.028 \pm 0.023$		16 ABBIENDI,G	00B OPAL	$e^+e^- \rightarrow Z$
$1.518 \pm 0.053 \pm 0.034$		18 BARATE	00R ALEP	$e^+e^- \rightarrow Z$
$1.523 \pm 0.057 \pm 0.053$		19 ABBIENDI	99J OPAL	$e^+e^- \rightarrow Z$
$1.474 \pm 0.039^{+0.052}_{-0.051}$		18 ABE	98Q CDF	$p\bar{p}$ at 1.8 TeV
$1.52 \pm 0.06 \pm 0.04$		19 ACCIARRI	98S L3	$e^+e^- \rightarrow Z$
$1.64 \pm 0.08 \pm 0.08$		19 ABE	97J SLD	$e^+e^- \rightarrow Z$
$1.532 \pm 0.041 \pm 0.040$		20 ABREU	97F DLPH	$e^+e^- \rightarrow Z$
$1.25^{+0.15}_{-0.13} \pm 0.05$	121	15 BUSKULIC	96J ALEP	$e^+e^- \rightarrow Z$
$1.49^{+0.17}_{-0.15} \pm 0.08$		21 BUSKULIC	96J ALEP	$e^+e^- \rightarrow Z$
$1.61^{+0.14}_{-0.13} \pm 0.08$		18,22 ABREU	95Q DLPH	$e^+e^- \rightarrow Z$
$1.63 \pm 0.14 \pm 0.13$		23 ADAM	95 DLPH	$e^+e^- \rightarrow Z$
$1.53 \pm 0.12 \pm 0.08$		18,24 AKERS	95T OPAL	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.501^{+0.078}_{-0.074} \pm 0.050$		4 ABAZOV	07S D0	Repl. by ABAZOV 12U
$1.524 \pm 0.030 \pm 0.016$		4 ABULENCIA	07A CDF	Repl. by AALTONEN 11
$1.473^{+0.052}_{-0.050} \pm 0.023$		9 ABAZOV	05B D0	Repl. by ABAZOV 05W
$1.40^{+0.11}_{-0.10} \pm 0.03$		4 ABAZOV	05C D0	Repl. by ABAZOV 07S
$1.530 \pm 0.043 \pm 0.023$		9 ABAZOV	05W D0	Repl. by ABAZOV 09E
$1.54 \pm 0.05 \pm 0.02$		25 ACOSTA	05 CDF	Repl. by AALTONEN 11
$1.554 \pm 0.030 \pm 0.019$		17 ABE	02H BELL	Repl. by ABE 05B
$1.58 \pm 0.09 \pm 0.02$		15 ABE	98B CDF	Repl. by ACOSTA 02C
$1.54 \pm 0.08 \pm 0.06$		18 ABE	96C CDF	Repl. by ABE 98Q
$1.55 \pm 0.06 \pm 0.03$		26 BUSKULIC	96J ALEP	$e^+e^- \rightarrow Z$
$1.61 \pm 0.07 \pm 0.04$		18 BUSKULIC	96J ALEP	Repl. by BARATE 00R
1.62 ± 0.12		27 ADAM	95 DLPH	$e^+e^- \rightarrow Z$
$1.57 \pm 0.18 \pm 0.08$	121	15 ABE	94D CDF	Repl. by ABE 98B
$1.17^{+0.29}_{-0.23} \pm 0.16$	96	18 ABREU	93D DLPH	Sup. by ABREU 95Q
$1.55 \pm 0.25 \pm 0.18$	76	23 ABREU	93G DLPH	Sup. by ADAM 95
$1.51^{+0.24}_{-0.23} \pm 0.12$	78	18 ACTON	93C OPAL	Sup. by AKERS 95T
$1.52^{+0.20}_{-0.18} \pm 0.07$	77	18 BUSKULIC	93D ALEP	Sup. by BUSKULIC 96J
$1.20^{+0.52}_{-0.36} \pm 0.16$	15	28 WAGNER	90 MRK2	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$
$0.82^{+0.57}_{-0.37} \pm 0.27$		29 AVERILL	89 HRS	$E_{\text{cm}}^{ee} = 29 \text{ GeV}$

¹ Measured using $B^0 \rightarrow D^{(*)-} \pi^+$ decays.

² Measured using $B^0 \rightarrow J/\psi K^{*0}(892)$ and $B^0 \rightarrow J/\psi K_S^0$ decays.

³ Measured using $B^0 \rightarrow D^- \mu^+ \nu X$ decays.

⁴ Measured mean life using $B^0 \rightarrow J/\psi K_S^0$ decays.

⁵ Measured using $B^0 \rightarrow J/\psi K^{*0}$ decays.

⁶ Measured using $B^0 \rightarrow K^+ \pi^-$ decays.

⁷ Measured with $B_d^0 \rightarrow J/\psi(\mu^+ \mu^-) K_S^0(\pi^+ \pi^-)$ decays.

⁸ Measured mean life using fully reconstructed decays ($J/\psi K^{(*)}$).

- ⁹ Measured mean life using $B^0 \rightarrow J/\psi K^{*0}$ decays.
- ¹⁰ Measured using a simultaneous fit of the B^0 lifetime and $\bar{B}^0 B^0$ oscillation frequency Δm_d in the partially reconstructed $B^0 \rightarrow D^{*-} \ell \nu$ decays.
- ¹¹ Measurement performed using a combined fit of CP -violation, mixing and lifetimes.
- ¹² Measurement performed using an inclusive reconstruction and B flavor identification technique.
- ¹³ AUBERT 03C uses a sample of approximately 14,000 exclusively reconstructed $B^0 \rightarrow D^{*}(2010)^- \ell \nu$ and simultaneously measures the lifetime and oscillation frequency.
- ¹⁴ Measurement performed with decays $B^0 \rightarrow D^{*-} \pi^+$ and $B^0 \rightarrow D^{*-} \rho^+$ using a partial reconstruction technique.
- ¹⁵ Measured mean life using fully reconstructed decays.
- ¹⁶ Data analyzed using partially reconstructed $\bar{B}^0 \rightarrow D^{*+} \ell^- \bar{\nu}$ decays.
- ¹⁷ Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.
- ¹⁸ Data analyzed using $D/D^* \ell X$ event vertices.
- ¹⁹ Data analyzed using charge of secondary vertex.
- ²⁰ Data analyzed using inclusive $D/D^* \ell X$.
- ²¹ Measured mean life using partially reconstructed $D^{*-} \pi^+ X$ vertices.
- ²² ABREU 95Q assumes $B(B^0 \rightarrow D^{*-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.
- ²³ Data analyzed using vertex-charge technique to tag B charge.
- ²⁴ AKERS 95T assumes $B(B^0 \rightarrow D_s^{(*)} D^0) = 5.0 \pm 0.9\%$ to find B^+/B^0 yield.
- ²⁵ Measured using the time-dependent angular analysis of $B_d^0 \rightarrow J/\psi K^{*0}$ decays.
- ²⁶ Combined result of $D/D^* \ell X$ analysis, fully reconstructed B analysis, and partially reconstructed $D^{*-} \pi^+ X$ analysis.
- ²⁷ Combined ABREU 95Q and ADAM 95 result.
- ²⁸ WAGNER 90 tagged B^0 mesons by their decays into $D^{*-} e^+ \nu$ and $D^{*-} \mu^+ \nu$ where the D^{*-} is tagged by its decay into $\pi^- \bar{D}^0$.
- ²⁹ AVERILL 89 is an estimate of the B^0 mean lifetime assuming that $B^0 \rightarrow D^{*+} + X$ always.

 $\tau_{B^0}/\tau_{\bar{B}^0}$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.000 \pm 0.008 \pm 0.009$	¹ AAIJ	14E LHCb	pp at 7 TeV
¹ Measured using $B^0 \rightarrow J/\psi K^{*0}$ decays.			

MEAN LIFE RATIO τ_{B^+}/τ_{B^0} **τ_{B^+}/τ_{B^0} (direct measurements)**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.076 ± 0.004 OUR EVALUATION		(Produced by HFLAV)		
$1.074 \pm 0.005 \pm 0.003$	¹	AAIJ	14E LHCb	pp at 7 TeV
$1.088 \pm 0.009 \pm 0.004$	²	AALTONEN	11 CDF	$p\bar{p}$ at 1.96 TeV
$1.080 \pm 0.016 \pm 0.014$	³	ABAZOV	05D D0	$p\bar{p}$ at 1.96 TeV
$1.066 \pm 0.008 \pm 0.008$	⁴	ABE	05B BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.060 \pm 0.021 \pm 0.024$	⁵	ABDALLAH	04E DLPH	$e^+ e^- \rightarrow Z$
$1.093 \pm 0.066 \pm 0.028$	⁶	ACOSTA	02C CDF	$p\bar{p}$ at 1.8 TeV
$1.082 \pm 0.026 \pm 0.012$	⁷	AUBERT	01F BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.085 \pm 0.059 \pm 0.018$	³	BARATE	00R ALEP	$e^+ e^- \rightarrow Z$
$1.079 \pm 0.064 \pm 0.041$	⁸	ABBIENDI	99J OPAL	$e^+ e^- \rightarrow Z$

$1.110 \pm 0.056^{+0.033}_{-0.030}$	³ ABE	98Q	CDF	$p\bar{p}$ at 1.8 TeV
$1.09 \pm 0.07 \pm 0.03$	⁸ ACCIARRI	98S	L3	$e^+e^- \rightarrow Z$
$1.01 \pm 0.07 \pm 0.06$	⁸ ABE	97J	SLD	$e^+e^- \rightarrow Z$
$1.27^{+0.23}_{-0.19}^{+0.03}_{-0.02}$	⁶ BUSKULIC	96J	ALEP	$e^+e^- \rightarrow Z$
$1.00^{+0.17}_{-0.15} \pm 0.10$	^{3,9} ABREU	95Q	DLPH	$e^+e^- \rightarrow Z$
$1.06^{+0.13}_{-0.11} \pm 0.10$	¹⁰ ADAM	95	DLPH	$e^+e^- \rightarrow Z$
$0.99 \pm 0.14^{+0.05}_{-0.04}$	^{3,11} AKERS	95T	OPAL	$e^+e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.091 \pm 0.023 \pm 0.014$	⁷ ABE	02H	BELL	Repl. by ABE 05B
$1.06 \pm 0.07 \pm 0.02$	⁶ ABE	98B	CDF	Repl. by ACOSTA 02C
$1.01 \pm 0.11 \pm 0.02$	³ ABE	96C	CDF	Repl. by ABE 98Q
$1.03 \pm 0.08 \pm 0.02$	¹² BUSKULIC	96J	ALEP	$e^+e^- \rightarrow Z$
$0.98 \pm 0.08 \pm 0.03$	³ BUSKULIC	96J	ALEP	Repl. by BARATE 00R
$1.02 \pm 0.16 \pm 0.05$	²⁶⁹ ⁶ ABE	94D	CDF	Repl. by ABE 98B
$1.11^{+0.51}_{-0.39} \pm 0.11$	¹⁸⁸ ³ ABREU	93D	DLPH	Sup. by ABREU 95Q
$1.01^{+0.29}_{-0.22} \pm 0.12$	²⁵³ ¹⁰ ABREU	93G	DLPH	Sup. by ADAM 95
$1.0^{+0.33}_{-0.25} \pm 0.08$	¹³⁰ ACTON	93C	OPAL	Sup. by AKERS 95T
$0.96^{+0.19}_{-0.15}^{+0.18}_{-0.12}$	¹⁵⁴ ³ BUSKULIC	93D	ALEP	Sup. by BUSKULIC 96J

¹ Measured using $B \rightarrow J/\psi K^{(*)}$ decays.

² Measured mean life using fully reconstructed decays ($J/\psi K^{(*)}$).

³ Data analyzed using $D/D^* \mu X$ vertices.

⁴ Measurement performed using a combined fit of CP -violation, mixing and lifetimes.

⁵ Measurement performed using an inclusive reconstruction and B flavor identification technique.

⁶ Measured using fully reconstructed decays.

⁷ Events are selected in which one B meson is fully reconstructed while the second B meson is reconstructed inclusively.

⁸ Data analyzed using charge of secondary vertex.

⁹ ABREU 95Q assumes $B(B^0 \rightarrow D^{*-} \ell^+ \nu_\ell) = 3.2 \pm 1.7\%$.

¹⁰ Data analyzed using vertex-charge technique to tag B charge.

¹¹ AKERS 95T assumes $B(B^0 \rightarrow D_s^{(*)} D^0) = 5.0 \pm 0.9\%$ to find B^+/B^0 yield.

¹² Combined result of $D/D^* \ell X$ analysis and fully reconstructed B analysis.

τ_{B^+}/τ_{B^0} (inferred from branching fractions)

These measurements are inferred from the branching fractions for semileptonic decay or other spectator-dominated decays by assuming that the rates for such decays are equal for B^0 and B^+ . We do not use measurements which assume equal production of B^0 and B^+ because of the large uncertainty in the production ratio.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.04	OUR AVERAGE				
$1.07 \pm 0.04 \pm 0.03$			URQUIJO	07	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$1.067 \pm 0.041 \pm 0.033$			AUBERT,B	06Y	BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.95^{+0.117}_{-0.080} \pm 0.091$	¹ ARTUSO	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------------------	---------------------	----	------	-----------------------------------

1.15 ± 0.17 ± 0.06		² JESSOP	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.93 ± 0.18 ± 0.12		³ ATHANAS	94	CLE2	Sup. by ARTUSO 97
0.91 ± 0.27 ± 0.21		⁴ ALBRECHT	92C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
1.0 ± 0.4	29	^{4,5} ALBRECHT	92G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.89 ± 0.19 ± 0.13		⁴ FULTON	91	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
1.00 ± 0.23 ± 0.14		⁴ ALBRECHT	89L	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.49 to 2.3	90	⁶ BEAN	87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and independent of B^0 and B^+ production fraction.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ATHANAS 94 uses events tagged by fully reconstructed B^- decays and partially or fully reconstructed B^0 decays.

⁴ Assumes equal production of B^0 and B^+ .

⁵ ALBRECHT 92G data analyzed using $B \rightarrow D_s \bar{D}, D_s \bar{D}^*, D_s^* \bar{D}, D_s^* \bar{D}^*$ events.

⁶ BEAN 87B assume the fraction of $B^0 \bar{B}^0$ events at the $\Upsilon(4S)$ is 0.41.

$\Delta\Gamma_{B_d^0} / \Gamma_{B_d^0}$

$\Gamma_{B_d^0}$ and $\Delta\Gamma_{B_d^0}$ are the decay rate average and difference between two B_d^0 CP eigenstates (light – heavy). The λ_{CP} characterizes B^0 and $\bar{B}^0$ decays to states of charmonium plus K_L^0 , see the review on “ CP Violation” in the reviews section.

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
0.1 ± 1.0 OUR EVALUATION		(Produced by HFLAV)		
0.1 ± 1.0 OUR AVERAGE				
3.4 ± 2.3 ± 2.4		¹ SIRUNYAN	18BY CMS	pp at 8 TeV
– 0.1 ± 1.1 ± 0.9		² AABOUD	16G ATLS	pp at 7, 8 TeV
– 4.4 ± 2.5 ± 1.1		³ AAIJ	14E LHCb	pp at 7 TeV
1.7 ± 1.8 ± 1.1		⁴ HIGUCHI	12 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
0.8 ± 3.7 ± 1.8		⁵ AUBERT,B	04C BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0 ± 9		⁶ ABDALLAH	03B DLPH	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.50 ± 1.38		ABAZOV	14 D0	$p\bar{p}$ at 1.96 TeV
< 80	95	⁷ BEHRENS	00B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Measured using $B^0 \rightarrow J/\psi K^*(892)^0$ and $B^0 \rightarrow J/\psi K_S^0$ decays, and assuming $\beta = 21.9 \pm 0.7$ degrees. ² Measured from the ratio of decay time distributions of $B^0 \rightarrow J/\psi K_S^0$ and $B^0 \rightarrow J/\psi K^{*0}$ decays. ³ Measured using the effective lifetimes of $B^0 \rightarrow J/\psi K_S^0$ and $B^0 \rightarrow J/\psi K^{*0}$ decays. ⁴ Reports $-\Delta\Gamma_d/\Gamma_d$ using $B^0 \rightarrow J/\psi K_S^0, J/\psi K_L^0, D^-\pi^+, D^{*-}\pi^+, D^{*-}\rho^+$, and $D^{*-}\ell^+\nu$ decays. ⁵ Corresponds to 90% confidence range $[-0.084, 0.068]$. ⁶ Used the measured $\tau_{B^0} = 1.55 \pm 0.03$ ps. Corresponds to an upper limit of < 0.18 at 95% C.L. ⁷ BEHRENS 00B uses high-momentum lepton tags and partially reconstructed $\bar{B}^0 \rightarrow D^{*+}\pi^-, \rho^-$ decays to determine the flavor of the B meson. Assumes $\Delta_{md}=0.478 \pm 0.018$ ps $^{-1}$ and $\tau_{B^0}=1.548 \pm 0.032$ ps.				

B^0 DECAY MODES

$\bar{B}^0$ modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing. Modes which do not identify the charge state of the B are listed in the $B^\pm/B^0$ ADMIXTURE section.

The branching fractions listed below assume 50% $B^0\bar{B}^0$ and 50% B^+B^- production at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D , D_S , D^* , and ψ branching ratios to current values whenever this would affect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

For inclusive branching fractions, e.g., $B \rightarrow D^\pm X$, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \quad \ell^+ \nu_\ell X$	[a] (10.33 $\pm$ 0.28) %	
$\Gamma_2 \quad e^+ \nu_e X_c$	(10.1 $\pm$ 0.4) %	
$\Gamma_3 \quad \ell^+ \nu_\ell X_u$	[a] (1.51 $\pm$ 0.19) $\times 10^{-3}$	
$\Gamma_4 \quad D \ell^+ \nu_\ell X$	[a] (9.1 $\pm$ 0.8) %	
$\Gamma_5 \quad D^- \ell^+ \nu_\ell$	[a] (2.10 $\pm$ 0.07) %	
$\Gamma_6 \quad D^- \tau^+ \nu_\tau$	(9.8 $\pm$ 2.1) $\times 10^{-3}$	
$\Gamma_7 \quad D^*(2010)^- \ell^+ \nu_\ell$	[a] (4.87 $\pm$ 0.09) %	
$\Gamma_8 \quad D^*(2010)^- e^+ \nu_e$		
$\Gamma_9 \quad D^*(2010)^- \mu^+ \nu_\mu$		
$\Gamma_{10} \quad D^*(2010)^- \tau^+ \nu_\tau$	(1.48 $\pm$ 0.09) %	S=1.2
$\Gamma_{11} \quad \bar{D}^{(*)} n \pi \ell^+ \nu_\ell (n \geq 1)$	[a] (2.2 $\pm$ 0.4) %	
$\Gamma_{12} \quad \bar{D}^0 \pi^- \ell^+ \nu_\ell$	[a] (3.64 $\pm$ 0.20) $\times 10^{-3}$	
$\Gamma_{13} \quad D_0^*(2300)^- \ell^+ \nu_\ell,$ $D_0^{*-} \rightarrow \bar{D}^0 \pi^-$	[a] < 4.4 $\times 10^{-4}$	CL=90%
$\Gamma_{14} \quad D_2^*(2460)^- \ell^+ \nu_\ell,$ $D_2^{*-} \rightarrow \bar{D}^0 \pi^-$	[a] (1.41 $\pm$ 0.20) $\times 10^{-3}$	S=1.7
$\Gamma_{15} \quad \bar{D}^{*0} \pi^- \ell^+ \nu_\ell$	[a] (5.44 $\pm$ 0.28) $\times 10^{-3}$	
$\Gamma_{16} \quad D_1(2420)^- \ell^+ \nu_\ell, D_1^- \rightarrow$ $\bar{D}^{*0} \pi^-$	[a] (2.85 $\pm$ 0.25) $\times 10^{-3}$	
$\Gamma_{17} \quad D_1(2420)^- \ell^+ \nu_\ell, D_1^- \rightarrow$ $D^- \pi^+ \pi^-$	[a] (1.02 $\pm$ 0.16) $\times 10^{-3}$	
$\Gamma_{18} \quad D_1'(2430)^- \ell^+ \nu_\ell, D_1'^- \rightarrow$ $\bar{D}^{*0} \pi^-$	[a] (2.5 $\pm$ 0.6) $\times 10^{-3}$	

Γ_{19}	$D_2^*(2460)^- \ell^+ \nu_\ell, D_2^{*-} \rightarrow \bar{D}^{*0} \pi^-$	[a]	$(6.6 \pm 1.1) \times 10^{-4}$	
Γ_{20}	$D^- \pi^+ \pi^- \ell^+ \nu_\ell$	[a]	$(1.45 \pm 0.22) \times 10^{-3}$	
Γ_{21}	$D^{*-} \pi^+ \pi^- \ell^+ \nu_\ell$	[a]	$(5.1 \pm 2.3) \times 10^{-4}$	
Γ_{22}	$\rho^- \ell^+ \nu_\ell$	[a]	$(2.94 \pm 0.21) \times 10^{-4}$	
Γ_{23}	$\pi^- \ell^+ \nu_\ell$	[a]	$(1.50 \pm 0.05) \times 10^{-4}$	
Γ_{24}	$\pi^- \mu^+ \nu_\mu$			
Γ_{25}	$\pi^- \tau^+ \nu_\tau$		$< 2.5 \times 10^{-4}$	CL=90%

Inclusive modes

Γ_{26}	$K^\pm X$		$(78 \pm 8) \%$	
Γ_{27}	$D^0 X$		$(8.1 \pm 1.5) \%$	
Γ_{28}	$\bar{D}^0 X$		$(47.4 \pm 2.8) \%$	
Γ_{29}	$D^+ X$	$<$	3.9%	CL=90%
Γ_{30}	$D^- X$		$(36.9 \pm 3.3) \%$	
Γ_{31}	$D_s^+ X$		$(10.3 \pm 2.1) \%$	
Γ_{32}	$D_s^- X$	$<$	2.6%	CL=90%
Γ_{33}	$\Lambda_c^+ X$	$<$	3.1%	CL=90%
Γ_{34}	$\bar{\Lambda}_c^- X$		$(5.0 \pm 2.1) \%$	
Γ_{35}	$\bar{c} X$		$(95 \pm 5) \%$	
Γ_{36}	$c X$		$(24.6 \pm 3.1) \%$	
Γ_{37}	$\bar{c}/c X$		$(119 \pm 6) \%$	
vglue -0.07in				

D, D*, or D_s modes

Γ_{38}	$D^- \pi^+$		$(2.51 \pm 0.08) \times 10^{-3}$	
Γ_{39}	$D^- \rho^+$		$(7.6 \pm 1.2) \times 10^{-3}$	
Γ_{40}	$D^- K^0 \pi^+$		$(4.9 \pm 0.9) \times 10^{-4}$	
Γ_{41}	$D^- K^*(892)^+$		$(4.5 \pm 0.7) \times 10^{-4}$	
Γ_{42}	$D^- \omega \pi^+$		$(2.8 \pm 0.6) \times 10^{-3}$	
Γ_{43}	$D^- K^+$		$(2.05 \pm 0.08) \times 10^{-4}$	
Γ_{44}	$D^- K^+ \pi^+ \pi^-$		$(3.5 \pm 0.8) \times 10^{-4}$	
Γ_{45}	$D^- K^+ \bar{K}^0$		$(1.64 \pm 0.26) \times 10^{-4}$	
Γ_{46}	$D^- K^+ \bar{K}^*(892)^0$		$(7.7 \pm 0.6) \times 10^{-4}$	
Γ_{47}	$\bar{D}^0 \pi^+ \pi^-$		$(8.8 \pm 0.5) \times 10^{-4}$	
Γ_{48}	$D^*(2010)^- \pi^+$		$(2.66 \pm 0.07) \times 10^{-3}$	
Γ_{49}	$\bar{D}^0 K^+ K^-$		$(6.1 \pm 0.5) \times 10^{-5}$	
Γ_{50}	$D^- \pi^+ \pi^+ \pi^-$		$(6.0 \pm 0.6) \times 10^{-3}$	
Γ_{51}	$(D^- \pi^+ \pi^+ \pi^-)$ nonresonant		$(3.9 \pm 1.9) \times 10^{-3}$	
Γ_{52}	$D^- \pi^+ \rho^0$		$(1.1 \pm 1.0) \times 10^{-3}$	
Γ_{53}	$D^- a_1(1260)^+$		$(6.0 \pm 3.3) \times 10^{-3}$	
Γ_{54}	$D^*(2010)^- \pi^+ \pi^0$		$(1.5 \pm 0.5) \%$	
Γ_{55}	$D^*(2010)^- \rho^+$		$(6.8 \pm 0.9) \times 10^{-3}$	
Γ_{56}	$D^*(2010)^- K^+$		$(2.16 \pm 0.08) \times 10^{-4}$	

Γ_{57}	$D^*(2010)^- K^0 \pi^+$	$(3.0 \pm 0.8) \times 10^{-4}$	
Γ_{58}	$D^*(2010)^- K^*(892)^+$	$(3.3 \pm 0.6) \times 10^{-4}$	
Γ_{59}	$D^*(2010)^- K^+ \bar{K}^0$	$(1.8 \pm 0.4) \times 10^{-4}$	
Γ_{60}	$D^*(2010)^- K^+ \bar{K}^*(892)^0$	$(1.31 \pm 0.13) \times 10^{-3}$	
Γ_{61}	$D^*(2010)^- \pi^+ \pi^+ \pi^-$	$(7.21 \pm 0.29) \times 10^{-3}$	
Γ_{62}	$(D^*(2010)^- \pi^+ \pi^+ \pi^-)$ non-resonant	$(0.0 \pm 2.5) \times 10^{-3}$	
Γ_{63}	$D^*(2010)^- \pi^+ \rho^0$	$(5.7 \pm 3.2) \times 10^{-3}$	
Γ_{64}	$D^*(2010)^- a_1(1260)^+$	$(1.30 \pm 0.27) \%$	
Γ_{65}	$\bar{D}_1(2420)^0 \pi^- \pi^+, \bar{D}_1^0 \rightarrow D^{*-} \pi^+$	$(1.47 \pm 0.35) \times 10^{-4}$	
Γ_{66}	$D^*(2010)^- K^+ \pi^- \pi^+$	$(4.7 \pm 0.4) \times 10^{-4}$	
Γ_{67}	$D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0$	$(1.76 \pm 0.27) \%$	
Γ_{68}	$D^{*-} 3\pi^+ 2\pi^-$	$(4.7 \pm 0.9) \times 10^{-3}$	
Γ_{69}	$D^*(2010)^- \omega \pi^+$	$(2.46 \pm 0.18) \times 10^{-3}$	S=1.2
Γ_{70}	$\bar{D}_1(2430)^0 \omega, \bar{D}_1^0 \rightarrow D^{*-} \pi^+$	$(2.7 \pm 0.8) \times 10^{-4}$	
Γ_{71}	$D^{*-} \rho(1450)^+, \rho^+ \rightarrow \omega \pi^+$	$(1.07 \pm 0.40) \times 10^{-3}$	
Γ_{72}	$\bar{D}_1(2420)^0 \omega, \bar{D}_1^0 \rightarrow D^{*-} \pi^+$	$(7.0 \pm 2.2) \times 10^{-5}$	
Γ_{73}	$\bar{D}_2^*(2460)^0 \omega, \bar{D}_2^0 \rightarrow D^{*-} \pi^+$	$(4.0 \pm 1.4) \times 10^{-5}$	
Γ_{74}	$D^{*-} b_1(1235)^+, b_1^+ \rightarrow \omega \pi^+$	$< 7 \times 10^{-5}$	CL=90%
Γ_{75}	$\bar{D}^{*-} \pi^+$	[b] $(1.9 \pm 0.9) \times 10^{-3}$	
Γ_{76}	$D_1(2420)^- \pi^+, D_1^- \rightarrow D^- \pi^+ \pi^-$	$(9.9 \pm 2.0) \times 10^{-5}$	
Γ_{77}	$D_1(2420)^- \pi^+, D_1^- \rightarrow D^{*-} \pi^+ \pi^-$	$< 3.3 \times 10^{-5}$	CL=90%
Γ_{78}	$\bar{D}_2^*(2460)^- \pi^+, D_2^{*-} \rightarrow D^0 \pi^-$	$(2.38 \pm 0.16) \times 10^{-4}$	
Γ_{79}	$\bar{D}_0^*(2400)^- \pi^+, D_0^{*-} \rightarrow D^0 \pi^-$	$(7.6 \pm 0.8) \times 10^{-5}$	
Γ_{80}	$D_2^*(2460)^- \pi^+, D_2^{*-} \rightarrow D^{*-} \pi^+ \pi^-$	$< 2.4 \times 10^{-5}$	CL=90%
Γ_{81}	$\bar{D}_2^*(2460)^- \rho^+$	$< 4.9 \times 10^{-3}$	CL=90%
Γ_{82}	$D^0 \bar{D}^0$	$(1.4 \pm 0.7) \times 10^{-5}$	
Γ_{83}	$D^{*0} \bar{D}^0$	$< 2.9 \times 10^{-4}$	CL=90%
Γ_{84}	$D^- D^+$	$(2.11 \pm 0.18) \times 10^{-4}$	
Γ_{85}	$D^\pm D^{*\mp} (CP\text{-averaged})$	$(6.1 \pm 0.6) \times 10^{-4}$	
Γ_{86}	$D^- D_s^+$	$(8.1 \pm 0.6) \times 10^{-3}$	S=1.1
Γ_{87}	$\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV}$		
Γ_{88}	$D^*(2010)^- D_s^+$	$(8.2 \pm 0.8) \times 10^{-3}$	
Γ_{89}	$D^*(2010)^- D_s^+, D^{*-} \rightarrow \bar{D}^0 \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV}$		
Γ_{90}	$D_2^*(2460)^- D_s^+, D_2^{*-} \rightarrow \bar{D}^0 \pi^-$		
Γ_{91}	$D_1^*(2600)^- D_s^+, D_1^{*-} \rightarrow \bar{D}^0 \pi^-$		

Γ_{92}	$D_3^*(2750)^- D_s^+, D_3^{*-} \rightarrow \bar{D}^0 \pi^-$			
Γ_{93}	$D_1^*(2760)^- D_s^+, D_1^{*-} \rightarrow \bar{D}^0 \pi^-$			
Γ_{94}	$D_J^*(3000)^- D_s^+, D_J^{*-} \rightarrow \bar{D}^0 \pi^-$			
Γ_{95}	$T_{cs0}^*(2870)^0 \bar{D}^0, T_{cs0}^{*0} \rightarrow D_s^+ \pi^-$			
Γ_{96}	$D^- D_s^{*+}$	$(7.4 \pm 1.6) \times 10^{-3}$		
Γ_{97}	$D^*(2010)^- D_s^{*+}$	$(1.77 \pm 0.14) \%$		
Γ_{98}	$D_{s0}(2317)^- K^+, D_{s0}^- \rightarrow D_s^- \pi^0$	$(4.2 \pm 1.4) \times 10^{-5}$		
Γ_{99}	$D_{s0}(2317)^- \pi^+, D_{s0}^- \rightarrow D_s^- \pi^0$	< 2.5	$\times 10^{-5}$	CL=90%
Γ_{100}	$D_{sJ}(2457)^- K^+, D_{sJ}^- \rightarrow D_s^- \pi^0$	< 9.4	$\times 10^{-6}$	CL=90%
Γ_{101}	$D_{sJ}(2457)^- \pi^+, D_{sJ}^- \rightarrow D_s^- \pi^0$	< 4.0	$\times 10^{-6}$	CL=90%
Γ_{102}	$D_s^- D_s^+$	< 3.6	$\times 10^{-5}$	CL=90%
Γ_{103}	$D_s^{*-} D_s^+$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{104}	$D_s^{*-} D_s^{*+}$	< 2.4	$\times 10^{-4}$	CL=90%
Γ_{105}	$D_{s0}^*(2317)^+ D^-, D_{s0}^{*+} \rightarrow D_s^+ \pi^0$	$(1.05 \pm 0.16) \times 10^{-3}$		S=1.1
Γ_{106}	$D_{s0}(2317)^+ D^-, D_{s0}^+ \rightarrow D_s^{*+} \gamma$	< 9.5	$\times 10^{-4}$	CL=90%
Γ_{107}	$D_{s0}(2317)^+ D^*(2010)^-, D_{s0}^+ \rightarrow D_s^+ \pi^0$	$(1.5 \pm 0.6) \times 10^{-3}$		
Γ_{108}	$D_{sJ}(2457)^+ D^-$	$(3.5 \pm 1.1) \times 10^{-3}$		
Γ_{109}	$D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \gamma$	$(6.5 \pm 1.7) \times 10^{-4}$		
Γ_{110}	$D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^{*+} \gamma$	< 6.0	$\times 10^{-4}$	CL=90%
Γ_{111}	$D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \pi^+ \pi^-$	< 2.0	$\times 10^{-4}$	CL=90%
Γ_{112}	$D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \pi^0$	< 3.6	$\times 10^{-4}$	CL=90%
Γ_{113}	$D^*(2010)^- D_{sJ}(2457)^+$	$(9.3 \pm 2.2) \times 10^{-3}$		
Γ_{114}	$D_{sJ}(2457)^+ D^*(2010), D_{sJ}^+ \rightarrow D_s^+ \gamma$	$(2.3 \pm 0.9) \times 10^{-3}$		
Γ_{115}	$D^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+ + D^{*+} K^0$	$(2.8 \pm 0.7) \times 10^{-4}$		
Γ_{116}	$D^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+$	$(1.7 \pm 0.6) \times 10^{-4}$		
Γ_{117}	$D^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*+} K^0$	$(2.6 \pm 1.1) \times 10^{-4}$		

Γ_{118}	$D^*(2010)^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+ + D^{*+} K^0$	$(5.0 \pm 1.4) \times 10^{-4}$	
Γ_{119}	$D^*(2010)^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+$	$(3.3 \pm 1.1) \times 10^{-4}$	
Γ_{120}	$D^{*-} D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*+} K^0$	$(5.0 \pm 1.7) \times 10^{-4}$	
Γ_{121}	$D^- D_{sJ}(2573)^+, D_{sJ}^+ \rightarrow D^0 K^+$	$(3.4 \pm 1.8) \times 10^{-5}$	
Γ_{122}	$D^*(2010)^- D_{sJ}(2573)^+, D_{sJ}^+ \rightarrow D^0 K^+$	$< 2 \times 10^{-4}$	CL=90%
Γ_{123}	$D^- D_{sJ}(2700)^+, D_{sJ}^+ \rightarrow D^0 K^+$	$(7.1 \pm 1.2) \times 10^{-4}$	
Γ_{124}	$D^+ \pi^-$	$(7.3 \pm 1.2) \times 10^{-7}$	
Γ_{125}	$D_s^+ \pi^-$	$(2.03 \pm 0.18) \times 10^{-5}$	
Γ_{126}	$D_s^{*+} \pi^-$	$(2.1 \pm 0.4) \times 10^{-5}$	S=1.4
Γ_{127}	$D_s^+ \rho^-$	$< 2.4 \times 10^{-5}$	CL=90%
Γ_{128}	$D_s^{*+} \rho^-$	$(4.1 \pm 1.3) \times 10^{-5}$	
Γ_{129}	$D_s^+ a_0^-$	$< 1.9 \times 10^{-5}$	CL=90%
Γ_{130}	$D_s^{*+} a_0^-$	$< 3.6 \times 10^{-5}$	CL=90%
Γ_{131}	$D_s^+ a_1(1260)^-$	$< 2.1 \times 10^{-3}$	CL=90%
Γ_{132}	$D_s^{*+} a_1(1260)^-$	$< 1.7 \times 10^{-3}$	CL=90%
Γ_{133}	$D_s^+ a_2^-$	$< 1.9 \times 10^{-4}$	CL=90%
Γ_{134}	$D_s^{*+} a_2^-$	$< 2.0 \times 10^{-4}$	CL=90%
Γ_{135}	$D_s^- K^+$	$(2.7 \pm 0.5) \times 10^{-5}$	S=2.7
Γ_{136}	$D_s^{*-} K^+$	$(2.19 \pm 0.30) \times 10^{-5}$	
Γ_{137}	$D_{s1}(2536)^{\mp} K^{\pm}, D_{s1}^- \rightarrow \bar{D}^*(2007)^0 K^-$	$(5.1 \pm 0.6) \times 10^{-6}$	
Γ_{138}	$D_s^- K^*(892)^+$	$(3.5 \pm 1.0) \times 10^{-5}$	
Γ_{139}	$D_s^{*-} K^*(892)^+$	$(3.2 \pm 1.5 \pm 1.3) \times 10^{-5}$	
Γ_{140}	$D_s^- \pi^+ K^0$	$(9.7 \pm 1.4) \times 10^{-5}$	
Γ_{141}	$D_s^{*-} \pi^+ K^0$	$< 1.10 \times 10^{-4}$	CL=90%
Γ_{142}	$D_s^- K^+ \pi^+ \pi^-$	$(1.7 \pm 0.5) \times 10^{-4}$	
Γ_{143}	$D_s^- \pi^+ K^*(892)^0$	$< 3.0 \times 10^{-3}$	CL=90%
Γ_{144}	$D_s^{*-} \pi^+ K^*(892)^0$	$< 1.6 \times 10^{-3}$	CL=90%
Γ_{145}	$\bar{D}^0 K^0$	$(5.5 \pm 0.4) \times 10^{-5}$	
Γ_{146}	$\bar{D}^0 K^+ \pi^-$	$(8.8 \pm 1.7) \times 10^{-5}$	
Γ_{147}	$\bar{D}^0 K^*(892)^0$	$(4.5 \pm 0.6) \times 10^{-5}$	
Γ_{148}	$\bar{D}^0 K^*(1410)^0$	$< 6.7 \times 10^{-5}$	CL=90%
Γ_{149}	$\bar{D}^0 K_0^*(1430)^0$	$(7 \pm 7) \times 10^{-6}$	
Γ_{150}	$\bar{D}^0 K_2^*(1430)^0$	$(2.1 \pm 0.9) \times 10^{-5}$	

Γ_{151}	$D_0^*(2300)^- K^+, D_0^{*-} \rightarrow \bar{D}^0 \pi^-$	$(1.9 \pm 0.9) \times 10^{-5}$	
Γ_{152}	$D_2^*(2460)^- K^+, D_2^{*-} \rightarrow \bar{D}^0 \pi^-$	$(2.03 \pm 0.35) \times 10^{-5}$	
Γ_{153}	$D_3^*(2760)^- K^+, D_3^{*-} \rightarrow \bar{D}^0 \pi^-$	$< 1.0 \times 10^{-6}$	CL=90%
Γ_{154}	$\bar{D}^0 K^+ \pi^-$ nonresonant	$< 3.7 \times 10^{-5}$	CL=90%
Γ_{155}	$[K^+ K^-]_D K^*(892)^0$	$(3.7 \pm 0.5) \times 10^{-5}$	
Γ_{156}	$[\pi^+ \pi^-]_D K^*(892)^0$	$(5.0 \pm 0.8) \times 10^{-5}$	
Γ_{157}	$[\pi^+ K^-]_D K^*(892)^0$		
Γ_{158}	$[K^+ \pi^-]_D K^*(892)^0$		
Γ_{159}	$[\pi^+ \pi^- \pi^+ \pi^-]_D K^{*0}$	$(4.0 \pm 0.6) \times 10^{-5}$	
Γ_{160}	$[\pi^+ K^- \pi^+ \pi^-]_D K^{*0}$		
Γ_{161}	$[K^+ \pi^- \pi^+ \pi^-]_D K^{*0}$		
Γ_{162}	$\bar{D}^0 \pi^0$	$(2.67 \pm 0.09) \times 10^{-4}$	
Γ_{163}	$\bar{D}^0 \rho^0$	$(3.21 \pm 0.21) \times 10^{-4}$	
Γ_{164}	$\bar{D}^0 f_2$	$(1.56 \pm 0.21) \times 10^{-4}$	
Γ_{165}	$\bar{D}^0 \eta$	$(2.56 \pm 0.12) \times 10^{-4}$	
Γ_{166}	$\bar{D}^0 \eta'$	$(1.38 \pm 0.16) \times 10^{-4}$	S=1.3
Γ_{167}	$\bar{D}^0 \omega$	$(2.54 \pm 0.16) \times 10^{-4}$	
Γ_{168}	$\bar{D}^0 \phi$	$(7.7 \pm 2.3) \times 10^{-7}$	
Γ_{169}	$D^0 K^+ \pi^-$	$(5.3 \pm 3.2) \times 10^{-6}$	
Γ_{170}	$D^0 K^*(892)^0$	$(3.0 \pm 0.6) \times 10^{-6}$	
Γ_{171}	$\bar{D}^{*0} \gamma$	$< 2.5 \times 10^{-5}$	CL=90%
Γ_{172}	$\bar{D}^*(2007)^0 \pi^0$	$(2.2 \pm 0.6) \times 10^{-4}$	S=2.6
Γ_{173}	$\bar{D}^*(2007)^0 \rho^0$	$< 5.1 \times 10^{-4}$	CL=90%
Γ_{174}	$\bar{D}^*(2007)^0 \eta$	$(2.3 \pm 0.6) \times 10^{-4}$	S=2.8
Γ_{175}	$\bar{D}^*(2007)^0 \eta'$	$(1.40 \pm 0.22) \times 10^{-4}$	
Γ_{176}	$\bar{D}^*(2007)^0 \pi^+ \pi^-$	$(6.2 \pm 2.2) \times 10^{-4}$	
Γ_{177}	$\bar{D}^*(2007)^0 K^+ \pi^-$	$(5.2 \pm 1.9) \times 10^{-5}$	
Γ_{178}	$\bar{D}^*(2007)^0 K^0$	$(3.6 \pm 1.2) \times 10^{-5}$	
Γ_{179}	$\bar{D}^*(2007)^0 K^*(892)^0$	$< 6.9 \times 10^{-5}$	CL=90%
Γ_{180}	$\bar{D}^*(2007)^0 \phi$	$(2.2 \pm 0.6) \times 10^{-6}$	
Γ_{181}	$D^*(2007)^0 K^*(892)^0$	$< 4.0 \times 10^{-5}$	CL=90%
Γ_{182}	$D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-$	$(2.7 \pm 0.5) \times 10^{-3}$	
Γ_{183}	$D^*(2010)^+ D^*(2010)^-$	$(8.0 \pm 0.6) \times 10^{-4}$	
Γ_{184}	$\bar{D}^*(2007)^0 \omega$	$(3.6 \pm 1.1) \times 10^{-4}$	S=3.1
Γ_{185}	$D^*(2010)^+ D^-$	$(6.1 \pm 1.5) \times 10^{-4}$	S=1.6
Γ_{186}	$D^*(2007)^0 \bar{D}^*(2007)^0$	$< 9 \times 10^{-5}$	CL=90%
Γ_{187}	$D^- D^0 K^+$	$(1.07 \pm 0.11) \times 10^{-3}$	
Γ_{188}	$D^- D^*(2007)^0 K^+$	$(3.5 \pm 0.4) \times 10^{-3}$	
Γ_{189}	$D^*(2010)^- D^0 K^+$	$(2.47 \pm 0.21) \times 10^{-3}$	
Γ_{190}	$D^*(2010)^- D^*(2007)^0 K^+$	$(1.06 \pm 0.09) \%$	
Γ_{191}	$D^- D^+ K^0$	$(7.5 \pm 1.7) \times 10^{-4}$	

Γ_{192}	$D^*(2010)^- D^+ K^0 + D^- D^*(2010)^+ K^0$	$(6.4 \pm 0.5) \times 10^{-3}$
Γ_{193}	$D^*(2010)^- D^*(2010)^+ K^0$	$(8.1 \pm 0.7) \times 10^{-3}$
Γ_{194}	$D^{*-} D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*+} K^0$	$(8.0 \pm 2.4) \times 10^{-4}$
Γ_{195}	$\bar{D}^0 D^0 K^0$	$(2.7 \pm 1.1) \times 10^{-4}$
Γ_{196}	$D^0 \bar{D}^0 K^+ \pi^-$	$(3.5 \pm 0.5) \times 10^{-4}$
Γ_{197}	$\bar{D}^0 D^*(2007)^0 K^0 + \bar{D}^*(2007)^0 D^0 K^0$	$(1.1 \pm 0.5) \times 10^{-3}$
Γ_{198}	$\bar{D}^*(2007)^0 D^*(2007)^0 K^0$	$(2.4 \pm 0.9) \times 10^{-3}$
Γ_{199}	$(\bar{D} + \bar{D}^*)(D + D^*) K$	$(3.68 \pm 0.26) \%$
vglue -0.07in		

Charmonium modes

Γ_{200}	$\eta_c K^0$	$(9.0 \pm 1.1) \times 10^{-4}$	
Γ_{201}	$\eta_c(1S) K^+ \pi^-$	$(6.5 \pm 0.7) \times 10^{-4}$	
Γ_{202}	$\eta_c(1S) K^+ \pi^-$ (NR)	$(6.7 \pm 1.4) \times 10^{-5}$	
Γ_{203}	$T_{c\bar{c}}(4100)^- K^+, T_{c\bar{c}}^- \rightarrow \eta_c \pi^-$	$(2.2 \pm 1.1) \times 10^{-5}$	
Γ_{204}	$\eta_c(1S) K^*(1410)^0$	$(2.1 \pm 1.6) \times 10^{-4}$	
Γ_{205}	$\eta_c(1S) K_0^*(1430)^0$	$(1.8 \pm 0.4) \times 10^{-4}$	
Γ_{206}	$\eta_c(1S) K_2^*(1430)^0$	$(5.4 \pm 2.4) \times 10^{-5}$	
Γ_{207}	$\eta_c(1S) K^*(1680)^0$	$(4 \pm 4) \times 10^{-5}$	
Γ_{208}	$\eta_c(1S) K_0^*(1950)^0$	$(4.8 \pm 3.2) \times 10^{-5}$	
Γ_{209}	$\eta_c K^*(892)^0$	$(5.3 \pm 0.8) \times 10^{-4}$	S=1.7
Γ_{210}	$\eta_c(2S) K_S^0, \eta_c \rightarrow p \bar{p} \pi^+ \pi^-$	$(4.2 \pm 1.4) \times 10^{-7}$	
Γ_{211}	$\eta_c(2S) K^{*0}$	$< 3.9 \times 10^{-4}$	CL=90%
Γ_{212}	$h_c(1P) K_S^0$	$< 1.4 \times 10^{-5}$	
Γ_{213}	$h_c(1P) K^{*0}$	$< 4 \times 10^{-4}$	CL=90%
Γ_{214}	$J/\psi(1S) K^0$	$(8.91 \pm 0.21) \times 10^{-4}$	
Γ_{215}	$J/\psi(1S) K^+ \pi^-$	$(1.15 \pm 0.05) \times 10^{-3}$	
Γ_{216}	$J/\psi(1S) K^*(892)^0$	$(1.27 \pm 0.05) \times 10^{-3}$	
Γ_{217}	$J/\psi(1S) \eta K_S^0$	$(5.4 \pm 0.9) \times 10^{-5}$	
Γ_{218}	$J/\psi(1S) \eta' K_S^0$	$< 2.5 \times 10^{-5}$	CL=90%
Γ_{219}	$J/\psi(1S) \phi K^0$	$(4.9 \pm 1.0) \times 10^{-5}$	S=1.3
Γ_{220}	$J/\psi(1S) \omega K^0$	$(2.3 \pm 0.4) \times 10^{-4}$	
Γ_{221}	$\chi_{c0}(3915), \chi_{c0} \rightarrow J/\psi \omega$	$(2.1 \pm 0.9) \times 10^{-5}$	
Γ_{222}	$J/\psi(1S) K(1270)^0$	$(1.3 \pm 0.5) \times 10^{-3}$	
Γ_{223}	$J/\psi(1S) \pi^0$	$(1.65 \pm 0.08) \times 10^{-5}$	
Γ_{224}	$J/\psi(1S) \eta$	$(1.08 \pm 0.23) \times 10^{-5}$	S=1.5
Γ_{225}	$J/\psi(1S) \pi^+ \pi^-$	$(3.99 \pm 0.15) \times 10^{-5}$	
Γ_{226}	$J/\psi(1S) \pi^+ \pi^-$ nonresonant	$< 1.2 \times 10^{-5}$	CL=90%

Γ_{227}	$J/\psi(1S) f_0(500), f_0 \rightarrow \pi\pi$	$(8.8 \pm 1.2 \pm 1.6) \times 10^{-6}$	
Γ_{228}	$J/\psi(1S) f_2$	$(3.3 \pm 0.5 \pm 0.6) \times 10^{-6}$	S=1.5
Γ_{229}	$J/\psi(1S) \rho^0$	$(2.55 \pm 0.18 \pm 0.16) \times 10^{-5}$	
Γ_{230}	$J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-$	$< 1.1 \times 10^{-6}$	CL=90%
Γ_{231}	$J/\psi(1S) \rho(1450)^0, \rho^0 \rightarrow \pi\pi$	$(2.9 \pm 1.6 \pm 0.7) \times 10^{-6}$	
Γ_{232}	$J/\psi \rho(1700)^0, \rho^0 \rightarrow \pi^+ \pi^-$	$(2.0 \pm 1.3) \times 10^{-6}$	
Γ_{233}	$J/\psi(1S) \omega$	$(1.8 \pm 0.7 \pm 0.5) \times 10^{-5}$	
Γ_{234}	$J/\psi(1S) K^+ K^-$	$(2.53 \pm 0.35) \times 10^{-6}$	
Γ_{235}	$J/\psi(1S) a_0(980), a_0 \rightarrow K^+ K^-$	$(4.7 \pm 3.4) \times 10^{-7}$	
Γ_{236}	$J/\psi(1S) \phi$	$< 1.1 \times 10^{-7}$	CL=90%
Γ_{237}	$J/\psi(1S) \eta'(958)$	$(7.6 \pm 2.4) \times 10^{-6}$	
Γ_{238}	$J/\psi(1S) K^0 \pi^+ \pi^-$	$(4.5 \pm 0.4) \times 10^{-4}$	
Γ_{239}	$J/\psi(1S) K^0 K^- \pi^+ + \text{c.c.}$	$< 2.1 \times 10^{-5}$	CL=90%
Γ_{240}	$J/\psi(1S) K^0 K^+ K^-$	$(2.5 \pm 0.7) \times 10^{-5}$	S=1.8
Γ_{241}	$J/\psi(1S) K^0 K^\pm \pi^\mp$		
Γ_{242}	$J/\psi(1S) K^0 \rho^0$	$(5.4 \pm 3.0) \times 10^{-4}$	
Γ_{243}	$J/\psi(1S) K^*(892)^+ \pi^-$	$(8 \pm 4) \times 10^{-4}$	
Γ_{244}	$J/\psi(1S) \pi^+ \pi^- \pi^+ \pi^-$	$(1.44 \pm 0.12) \times 10^{-5}$	
Γ_{245}	$J/\psi(1S) f_1(1285)$	$(8.4 \pm 2.1) \times 10^{-6}$	
Γ_{246}	$J/\psi(1S) K^*(892)^0 \pi^+ \pi^-$	$(6.6 \pm 2.2) \times 10^{-4}$	
Γ_{247}	$\eta_{c2}(1D) K_S^0, \eta_{c2} \rightarrow h_c \gamma$	$< 3.5 \times 10^{-5}$	CL=90%
Γ_{248}	$\eta_{c2}(1D) \pi^- K^+, \eta_{c2} \rightarrow h_c \gamma$	$< 1.0 \times 10^{-4}$	CL=90%
Γ_{249}	$\chi_{c1}(3872)^- K^+$	$< 5 \times 10^{-4}$	CL=90%
Γ_{250}	$\chi_{c1}(3872)^- K^+, \chi_{c1}(3872)^- \rightarrow J/\psi(1S) \pi^- \pi^0$	$[c] < 4.2 \times 10^{-6}$	CL=90%
Γ_{251}	$\chi_{c1}(3872) K^0$	$(1.1 \pm 0.4) \times 10^{-4}$	
Γ_{252}	$\chi_{c1}(3872) K^*(892)^0$	$(9 \pm 5) \times 10^{-5}$	
Γ_{253}	$\chi_{c1}(3872) K^+ \pi^-$	$(1.8 \pm 0.7) \times 10^{-4}$	
Γ_{254}	$\chi_{c1}(3872) \gamma$	$< 1.2 \times 10^{-5}$	CL=90%
Γ_{255}	$T_{c\bar{c}1}(4430)^\pm K^\mp, T_{c\bar{c}1}^\pm \rightarrow \psi(2S) \pi^\pm$	$(6.0 \pm 3.0 \pm 2.4) \times 10^{-5}$	
Γ_{256}	$T_{c\bar{c}1}(4430)^\pm K^\mp, T_{c\bar{c}1}^\pm \rightarrow J/\psi \pi^\pm$	$(5.4 \pm 4.0 \pm 1.2) \times 10^{-6}$	
Γ_{257}	$T_{c\bar{c}1}(3900)^\pm K^\mp, T_{c\bar{c}1}^\pm \rightarrow J/\psi \pi^\pm$	$< 9 \times 10^{-7}$	
Γ_{258}	$T_{c\bar{c}1}(4200)^\pm K^\mp, T_{c\bar{c}1}^\pm \rightarrow J/\psi \pi^\pm$	$(2.2 \pm 1.3 \pm 0.8) \times 10^{-5}$	

Γ_{259}	$J/\psi(1S)p\bar{p}$	$(4.5 \pm 0.6) \times 10^{-7}$	
Γ_{260}	$J/\psi(1S)\gamma$	$< 1.5 \times 10^{-6}$	CL=90%
Γ_{261}	$J/\psi\mu^+\mu^-$, $J/\psi \rightarrow \mu^+\mu^-$	$< 1.0 \times 10^{-9}$	CL=95%
Γ_{262}	$J/\psi(1S)\bar{D}^0$	$< 9.6 \times 10^{-7}$	CL=90%
Γ_{263}	$\psi(2S)\pi^0$	$(1.17 \pm 0.19) \times 10^{-5}$	
Γ_{264}	$\psi(2S)K^0$	$(5.8 \pm 0.5) \times 10^{-4}$	
Γ_{265}	$\psi(2S)K^0\pi^+\pi^-$	$(2.81 \pm 0.30) \times 10^{-4}$	
Γ_{266}	$\psi(3770)K^0$, $\psi \rightarrow \bar{D}^0 D^0$	$< 1.23 \times 10^{-4}$	CL=90%
Γ_{267}	$\psi(3770)K^0$, $\psi \rightarrow D^- D^+$	$< 1.88 \times 10^{-4}$	CL=90%
Γ_{268}	$\psi(2S)\pi^+\pi^-$	$(2.24 \pm 0.35) \times 10^{-5}$	
Γ_{269}	$\psi(2S)K^+\pi^-$	$(5.8 \pm 0.4) \times 10^{-4}$	
Γ_{270}	$\psi(2S)K^*(892)^0$	$(5.9 \pm 0.4) \times 10^{-4}$	
Γ_{271}	$\chi_{c0}K^0$	$(1.9 \pm 0.4) \times 10^{-4}$	
Γ_{272}	$\chi_{c0}K^*(892)^0$	$(1.7 \pm 0.4) \times 10^{-4}$	
Γ_{273}	$\chi_{c1}\pi^0$	$(1.12 \pm 0.28) \times 10^{-5}$	
Γ_{274}	$\chi_{c1}K^0$	$(3.95 \pm 0.27) \times 10^{-4}$	
Γ_{275}	$\chi_{c1}\pi^- K^+$	$(4.97 \pm 0.30) \times 10^{-4}$	
Γ_{276}	$\chi_{c1}K^*(892)^0$	$(2.38 \pm 0.19) \times 10^{-4}$	S=1.2
Γ_{277}	$T_{c\bar{c}}(4050)^- K^+$, $T_{c\bar{c}}^- \rightarrow \chi_{c1}\pi^-$	$(3.0 \pm_{-1.8}^{4.0}) \times 10^{-5}$	
Γ_{278}	$T_{c\bar{c}}(4250)^- K^+$, $T_{c\bar{c}}^- \rightarrow \chi_{c1}\pi^-$	$(4.0 \pm_{-1.0}^{20.0}) \times 10^{-5}$	
Γ_{279}	$\chi_{c1}\pi^+\pi^- K^0$	$(3.2 \pm 0.5) \times 10^{-4}$	
Γ_{280}	$\chi_{c1}\pi^-\pi^0 K^+$	$(3.5 \pm 0.6) \times 10^{-4}$	
Γ_{281}	$\chi_{c2}K^0$	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{282}	$\chi_{c2}K^*(892)^0$	$(4.9 \pm 1.2) \times 10^{-5}$	S=1.1
Γ_{283}	$\chi_{c2}\pi^- K^+$	$(7.2 \pm 1.0) \times 10^{-5}$	
Γ_{284}	$\chi_{c2}\pi^+\pi^- K^0$	$< 1.70 \times 10^{-4}$	CL=90%
Γ_{285}	$\chi_{c2}\pi^-\pi^0 K^+$	$< 7.4 \times 10^{-5}$	CL=90%
Γ_{286}	$\psi(4660)K^0$, $\psi \rightarrow \Lambda_c^+ \Lambda_c^-$	$< 2.3 \times 10^{-4}$	CL=90%
Γ_{287}	$\psi(4230)^0 K^0$, $\psi^0 \rightarrow J/\psi\pi^+\pi^-$	$< 1.7 \times 10^{-5}$	CL=90%

***K* or *K** modes**

Γ_{288}	$K^+\pi^-$	$(2.00 \pm 0.04) \times 10^{-5}$	
Γ_{289}	$K^0\pi^0$	$(1.01 \pm 0.04) \times 10^{-5}$	
Γ_{290}	$\eta' K^0$	$(6.6 \pm 0.4) \times 10^{-5}$	S=1.4
Γ_{291}	$\eta' K^*(892)^0$	$(2.8 \pm 0.6) \times 10^{-6}$	
Γ_{292}	$\eta' K_0^*(1430)^0$	$(6.3 \pm 1.6) \times 10^{-6}$	
Γ_{293}	$\eta' K_2^*(1430)^0$	$(1.37 \pm 0.32) \times 10^{-5}$	
Γ_{294}	ηK^0	$(1.23 \pm_{-0.24}^{0.27}) \times 10^{-6}$	
Γ_{295}	$\eta K^*(892)^0$	$(1.59 \pm 0.10) \times 10^{-5}$	
Γ_{296}	$\eta K_0^*(1430)^0$	$(1.10 \pm 0.22) \times 10^{-5}$	

Γ_{297}	$\eta K_2^*(1430)^0$	$(9.6 \pm 2.1) \times 10^{-6}$	
Γ_{298}	ωK^0	$(4.8 \pm 0.4) \times 10^{-6}$	
Γ_{299}	$a_0(980)^0 K^0, a_0^0 \rightarrow \eta \pi^0$	$< 7.8 \times 10^{-6}$	CL=90%
Γ_{300}	$b_1^0 K^0, b_1^0 \rightarrow \omega \pi^0$	$< 7.8 \times 10^{-6}$	CL=90%
Γ_{301}	$a_0(980)^\pm K^\mp, a_0^\pm \rightarrow \eta \pi^\pm$	$< 1.9 \times 10^{-6}$	CL=90%
Γ_{302}	$b_1^- K^+, b_1^- \rightarrow \omega \pi^-$	$(7.4 \pm 1.4) \times 10^{-6}$	
Γ_{303}	$b_1^0 K^{*0}, b_1^0 \rightarrow \omega \pi^0$	$< 8.0 \times 10^{-6}$	CL=90%
Γ_{304}	$b_1^- K^{*+}, b_1^- \rightarrow \omega \pi^-$	$< 5.0 \times 10^{-6}$	CL=90%
Γ_{305}	$a_0(1450)^\pm K^\mp, a_0^\pm \rightarrow \eta \pi^\pm$	$< 3.1 \times 10^{-6}$	CL=90%
Γ_{306}	$K_S^0 X^0$ (Familon)	$< 5.3 \times 10^{-5}$	CL=90%
Γ_{307}	$\omega K^*(892)^0$	$(2.0 \pm 0.5) \times 10^{-6}$	
Γ_{308}	$\omega (K\pi)_0^{*0}$	$(1.84 \pm 0.25) \times 10^{-5}$	
Γ_{309}	$\omega K_0^*(1430)^0$	$(1.60 \pm 0.34) \times 10^{-5}$	
Γ_{310}	$\omega K_2^*(1430)^0$	$(1.01 \pm 0.23) \times 10^{-5}$	
Γ_{311}	$\omega K^+ \pi^-$ nonresonant	$(5.1 \pm 1.0) \times 10^{-6}$	
Γ_{312}	$K^+ \pi^- \pi^0$	$(3.78 \pm 0.32) \times 10^{-5}$	
Γ_{313}	$K^+ \rho^-$	$(7.0 \pm 0.9) \times 10^{-6}$	
Γ_{314}	$K^+ \rho(1450)^-$	$(2.4 \pm 1.2) \times 10^{-6}$	
Γ_{315}	$K^+ \rho(1700)^-$	$(6 \pm 7) \times 10^{-7}$	
Γ_{316}	$(K^+ \pi^- \pi^0)$ nonresonant	$(2.8 \pm 0.6) \times 10^{-6}$	
Γ_{317}	$(K\pi)_0^{*+} \pi^-, (K\pi)_0^{*+} \rightarrow$	$(3.4 \pm 0.5) \times 10^{-5}$	
Γ_{318}	$K^+ \pi^0$ $(K\pi)_0^{*0} \pi^0, (K\pi)_0^{*0} \rightarrow$	$(8.6 \pm 1.7) \times 10^{-6}$	
Γ_{319}	$K_2^*(1430)^0 \pi^0$	$< 4.0 \times 10^{-6}$	CL=90%
Γ_{320}	$K^*(1680)^0 \pi^0$	$< 7.5 \times 10^{-6}$	CL=90%
Γ_{321}	$K_x^{*0} \pi^0$	[d] $(6.1 \pm 1.6) \times 10^{-6}$	
Γ_{322}	$K^0 \pi^+ \pi^-$	$(4.97 \pm 0.18) \times 10^{-5}$	
Γ_{323}	$K^0 \pi^+ \pi^-$ nonresonant	$(1.39^{+0.26}_{-0.18}) \times 10^{-5}$	S=1.6
Γ_{324}	$K^0 \rho^0$	$(3.4 \pm 1.1) \times 10^{-6}$	S=2.3
Γ_{325}	$K^*(892)^+ \pi^-$	$(7.5 \pm 0.4) \times 10^{-6}$	
Γ_{326}	$K_0^*(1430)^+ \pi^-$	$(3.3 \pm 0.7) \times 10^{-5}$	S=2.0
Γ_{327}	$K_x^{*+} \pi^-$	[d] $(5.1 \pm 1.6) \times 10^{-6}$	
Γ_{328}	$K^*(1410)^+ \pi^-, K^{*+} \rightarrow$	$< 3.8 \times 10^{-6}$	CL=90%
Γ_{329}	$K^0 \pi^+$ $(K\pi)_0^{*+} \pi^-, (K\pi)_0^{*+} \rightarrow$	$(1.62 \pm 0.13) \times 10^{-5}$	
Γ_{330}	$K^0 \pi^+$ $f_0(980) K^0, f_0 \rightarrow \pi^+ \pi^-$	$(8.1 \pm 0.8) \times 10^{-6}$	S=1.3
Γ_{331}	$K^0 f_0(500)$	$(1.6^{+2.5}_{-1.6}) \times 10^{-7}$	
Γ_{332}	$K^0 f_0(1500)$	$(1.3 \pm 0.8) \times 10^{-6}$	
Γ_{333}	$f_2(1270) K^0$	$(2.7^{+1.3}_{-1.2}) \times 10^{-6}$	

Γ_{334}	$f_x(1300)K^0, f_x \rightarrow \pi^+\pi^-$	$(1.8 \pm 0.7) \times 10^{-6}$	
Γ_{335}	$K^*(892)^0\pi^0$	$(3.3 \pm 0.6) \times 10^{-6}$	
Γ_{336}	$K_2^*(1430)^+\pi^-$	$(3.65 \pm 0.34) \times 10^{-6}$	
Γ_{337}	$K^*(1680)^+\pi^-$	$(1.41 \pm 0.10) \times 10^{-5}$	
Γ_{338}	$K^+\pi^-\pi^+\pi^-$	[e] $< 2.3 \times 10^{-4}$	CL=90%
Γ_{339}	$\rho^0 K^+\pi^-$	$(2.8 \pm 0.7) \times 10^{-6}$	
Γ_{340}	$f_0(980)K^+\pi^-, f_0 \rightarrow \pi\pi$	$(1.4 \pm_{-0.6}^{+0.5}) \times 10^{-6}$	
Γ_{341}	$K^+\pi^-\pi^+\pi^-$ nonresonant	$< 2.1 \times 10^{-6}$	CL=90%
Γ_{342}	$K^*(892)^0\pi^+\pi^-$	$(5.5 \pm 0.5) \times 10^{-5}$	
Γ_{343}	$K^*(892)^0\rho^0$	$(3.9 \pm 1.3) \times 10^{-6}$	S=1.9
Γ_{344}	$K^*(892)^0 f_0(980), f_0 \rightarrow \pi\pi$	$(3.9 \pm_{-1.8}^{+2.1}) \times 10^{-6}$	S=3.9
Γ_{345}	$K_1(1270)^+\pi^-$	$< 3.0 \times 10^{-5}$	CL=90%
Γ_{346}	$K_1(1400)^+\pi^-$	$< 2.7 \times 10^{-5}$	CL=90%
Γ_{347}	$a_1(1260)^-K^+$	[e] $(1.6 \pm 0.4) \times 10^{-5}$	
Γ_{348}	$K^*(892)^+\rho^-$	$(1.03 \pm 0.26) \times 10^{-5}$	
Γ_{349}	$K_0^*(1430)^+\rho^-$	$(2.8 \pm 1.2) \times 10^{-5}$	
Γ_{350}	$K_1(1400)^0\rho^0$	$< 3.0 \times 10^{-3}$	CL=90%
Γ_{351}	$K_0^*(1430)^0\rho^0$	$(2.7 \pm 0.6) \times 10^{-5}$	
Γ_{352}	$K_0^*(1430)^0 f_0(980), f_0 \rightarrow \pi\pi$	$(2.7 \pm 0.9) \times 10^{-6}$	
Γ_{353}	$K_2^*(1430)^0 f_0(980), f_0 \rightarrow \pi\pi$	$(8.6 \pm 2.0) \times 10^{-6}$	
Γ_{354}	K^+K^-	$(7.8 \pm 1.5) \times 10^{-8}$	
Γ_{355}	$K^0\bar{K}^0$	$(1.21 \pm 0.16) \times 10^{-6}$	
Γ_{356}	$K^0K^-\pi^+$	$(6.7 \pm 0.5) \times 10^{-6}$	
Γ_{357}	$K^*(892)^\pm K^\mp$	$< 4 \times 10^{-7}$	CL=90%
Γ_{358}	$\bar{K}^{*0}K^0 + K^{*0}\bar{K}^0$	$< 9.6 \times 10^{-7}$	CL=90%
Γ_{359}	$K^+K^-\pi^0$	$(2.2 \pm 0.6) \times 10^{-6}$	
Γ_{360}	$K_S^0 K_S^0 \pi^0$	$< 9 \times 10^{-7}$	CL=90%
Γ_{361}	$K_S^0 K_S^0 \eta$	$< 1.0 \times 10^{-6}$	CL=90%
Γ_{362}	$K_S^0 K_S^0 \eta'$	$< 2.0 \times 10^{-6}$	CL=90%
Γ_{363}	$K^0 K^+ K^-$	$(2.68 \pm 0.11) \times 10^{-5}$	
Γ_{364}	$K^0\phi$	$(7.3 \pm 0.7) \times 10^{-6}$	
Γ_{365}	$f_0(980)K^0, f_0 \rightarrow K^+K^-$	$(7.0 \pm_{-3.0}^{+3.5}) \times 10^{-6}$	
Γ_{366}	$f_0(1500)K^0$	$(1.3 \pm_{-0.5}^{+0.7}) \times 10^{-5}$	
Γ_{367}	$f_2'(1525)^0 K^0$	$(3 \pm_{-4}^{+5}) \times 10^{-7}$	
Γ_{368}	$f_0(1710)K^0, f_0 \rightarrow K^+K^-$	$(4.4 \pm 0.9) \times 10^{-6}$	
Γ_{369}	$K^0 K^+ K^-$ nonresonant	$(3.3 \pm 1.0) \times 10^{-5}$	
Γ_{370}	$K_S^0 K_S^0 K_S^0$	$(6.0 \pm 0.5) \times 10^{-6}$	S=1.1
Γ_{371}	$f_0(980)K^0, f_0 \rightarrow K_S^0 K_S^0$	$(2.7 \pm 1.8) \times 10^{-6}$	
Γ_{372}	$f_0(1710)K^0, f_0 \rightarrow K_S^0 K_S^0$	$(5.0 \pm_{-2.6}^{+5.0}) \times 10^{-7}$	
Γ_{373}	$f_2(2010)K^0, f_2 \rightarrow K_S^0 K_S^0$	$(5 \pm 6) \times 10^{-7}$	

Γ_{374}	$K_S^0 K_S^0 K_S^0$ nonresonant	(1.33 ± 0.31) $\times 10^{-5}$	
Γ_{375}	$K_S^0 K_S^0 K_L^0$	< 1.6 $\times 10^{-5}$	CL=90%
Γ_{376}	$K^*(892)^0 K^+ K^-$	(2.75 ± 0.26) $\times 10^{-5}$	
Γ_{377}	$K^*(892)^0 \phi$	(1.00 ± 0.05) $\times 10^{-5}$	
Γ_{378}	$K^+ K^- \pi^+ \pi^-$ nonresonant	< 7.17 $\times 10^{-5}$	CL=90%
Γ_{379}	$K^*(892)^0 K^- \pi^+$	(4.5 ± 1.3) $\times 10^{-6}$	
Γ_{380}	$K^*(892)^0 \bar{K}^*(892)^0$	(8.3 ± 2.4) $\times 10^{-7}$	S=1.5
Γ_{381}	$K^+ K^+ \pi^- \pi^-$ nonresonant	< 6.0 $\times 10^{-6}$	CL=90%
Γ_{382}	$K^*(892)^0 K^+ \pi^-$	< 2.2 $\times 10^{-6}$	CL=90%
Γ_{383}	$K^*(892)^0 K^*(892)^0$	< 2 $\times 10^{-7}$	CL=90%
Γ_{384}	$K^*(892)^+ K^*(892)^-$	< 2.0 $\times 10^{-6}$	CL=90%
Γ_{385}	$K_1(1400)^0 \phi$	< 5.0 $\times 10^{-3}$	CL=90%
Γ_{386}	$\phi(K\pi)_0^{*0}$	(4.3 ± 0.4) $\times 10^{-6}$	
Γ_{387}	$\phi(K\pi)_0^{*0} (1.60 < m_{K\pi} < 2.15)$ [f]	< 1.7 $\times 10^{-6}$	CL=90%
Γ_{388}	$K_0^*(1430)^0 K^- \pi^+$	< 3.18 $\times 10^{-5}$	CL=90%
Γ_{389}	$K_0^*(1430)^0 \bar{K}^*(892)^0$	< 3.3 $\times 10^{-6}$	CL=90%
Γ_{390}	$K_0^*(1430)^0 \bar{K}_0^*(1430)^0$	< 8.4 $\times 10^{-6}$	CL=90%
Γ_{391}	$K_0^*(1430)^0 \phi$	(3.9 ± 0.8) $\times 10^{-6}$	
Γ_{392}	$K_0^*(1430)^0 K^*(892)^0$	< 1.7 $\times 10^{-6}$	CL=90%
Γ_{393}	$K_0^*(1430)^0 K_0^*(1430)^0$	< 4.7 $\times 10^{-6}$	CL=90%
Γ_{394}	$K^*(1680)^0 \phi$	< 3.5 $\times 10^{-6}$	CL=90%
Γ_{395}	$K^*(1780)^0 \phi$	< 2.7 $\times 10^{-6}$	CL=90%
Γ_{396}	$K^*(2045)^0 \phi$	< 1.53 $\times 10^{-5}$	CL=90%
Γ_{397}	$K_2^*(1430)^0 \rho^0$	< 1.1 $\times 10^{-3}$	CL=90%
Γ_{398}	$K_2^*(1430)^0 \phi$	(6.8 ± 0.9) $\times 10^{-6}$	S=1.2
Γ_{399}	$K_0^0 \phi \phi$	(3.7 ± 0.7) $\times 10^{-6}$	S=1.3
Γ_{400}	$\eta' \eta' K^0$	< 3.1 $\times 10^{-5}$	CL=90%
Γ_{401}	$\eta K^0 \gamma$	(7.6 ± 1.8) $\times 10^{-6}$	
Γ_{402}	$\eta' K^0 \gamma$	< 6.4 $\times 10^{-6}$	CL=90%
Γ_{403}	$K^0 \phi \gamma$	(2.7 ± 0.7) $\times 10^{-6}$	
Γ_{404}	$K^+ \pi^- \gamma$	(4.6 ± 1.4) $\times 10^{-6}$	
Γ_{405}	$K^*(892)^0 \gamma$	(4.18 ± 0.25) $\times 10^{-5}$	S=2.1
Γ_{406}	$K^*(1410) \gamma$	< 1.3 $\times 10^{-4}$	CL=90%
Γ_{407}	$K^+ \pi^- \gamma$ nonresonant	< 2.6 $\times 10^{-6}$	CL=90%
Γ_{408}	$K^*(892)^0 X(214), X \rightarrow \mu^+ \mu^-$ [g]	< 2.26 $\times 10^{-8}$	CL=90%
Γ_{409}	$K^0 \pi^+ \pi^- \gamma$	(1.99 ± 0.18) $\times 10^{-5}$	
Γ_{410}	$K^+ \pi^- \pi^0 \gamma$	(4.1 ± 0.4) $\times 10^{-5}$	
Γ_{411}	$K_1(1270)^0 \gamma$	< 5.8 $\times 10^{-5}$	CL=90%
Γ_{412}	$K_1(1400)^0 \gamma$	< 1.2 $\times 10^{-5}$	CL=90%
Γ_{413}	$K_2^*(1430)^0 \gamma$	(1.24 ± 0.24) $\times 10^{-5}$	
Γ_{414}	$K^*(1680)^0 \gamma$	< 2.0 $\times 10^{-3}$	CL=90%
Γ_{415}	$K_3^*(1780)^0 \gamma$	< 8.3 $\times 10^{-5}$	CL=90%
Γ_{416}	$K_4^*(2045)^0 \gamma$	< 4.3 $\times 10^{-3}$	CL=90%

Light unflavored meson modes

Γ_{417}	$\rho^0 \gamma$	$(8.6 \pm 1.5) \times 10^{-7}$	
Γ_{418}	$\rho^0 X(214), X \rightarrow \mu^+ \mu^-$	$[g] < 1.73$	$\times 10^{-8}$ CL=90%
Γ_{419}	$\omega \gamma$	$(4.4 \pm_{-1.6}^{+1.8}) \times 10^{-7}$	
Γ_{420}	$\phi \gamma$	< 1.0	$\times 10^{-7}$ CL=90%
Γ_{421}	$f_2(1270) \gamma, f_2 \rightarrow (KS)^0 (KS)^0$	< 3.1	$\times 10^{-7}$
Γ_{422}	$f'_2(1525) \gamma, f'_2 \rightarrow (KS)^0 (KS)^0$	< 2.1	$\times 10^{-7}$
Γ_{423}	$\pi^+ \pi^-$	$(5.37 \pm 0.20) \times 10^{-6}$	S=1.3
Γ_{424}	$\pi^0 \pi^0$	$(1.55 \pm 0.17) \times 10^{-6}$	S=1.1
Γ_{425}	$\eta \pi^0$	$(4.1 \pm 1.7) \times 10^{-7}$	
Γ_{426}	$\eta \eta$	< 1.0	$\times 10^{-6}$ CL=90%
Γ_{427}	$\eta' \pi^0$	$(1.2 \pm 0.6) \times 10^{-6}$	S=1.7
Γ_{428}	$\eta' \eta'$	< 1.7	$\times 10^{-6}$ CL=90%
Γ_{429}	$\eta' \eta$	< 1.2	$\times 10^{-6}$ CL=90%
Γ_{430}	$\eta' \rho^0$	< 1.3	$\times 10^{-6}$ CL=90%
Γ_{431}	$\eta' f_0(980), f_0 \rightarrow \pi^+ \pi^-$	< 9	$\times 10^{-7}$ CL=90%
Γ_{432}	$\eta \rho^0$	< 1.5	$\times 10^{-6}$ CL=90%
Γ_{433}	$\eta f_0(980), f_0 \rightarrow \pi^+ \pi^-$	< 4	$\times 10^{-7}$ CL=90%
Γ_{434}	$\omega \eta$	$(9.4 \pm_{-3.1}^{+4.0}) \times 10^{-7}$	
Γ_{435}	$\omega \eta'$	$(1.0 \pm_{-0.4}^{+0.5}) \times 10^{-6}$	
Γ_{436}	$\omega \rho^0$	< 1.6	$\times 10^{-6}$ CL=90%
Γ_{437}	$\omega f_0(980), f_0 \rightarrow \pi^+ \pi^-$	< 1.5	$\times 10^{-6}$ CL=90%
Γ_{438}	$\omega \omega$	$(1.40 \pm 0.25) \times 10^{-6}$	
Γ_{439}	$\phi \pi^0$	< 1.5	$\times 10^{-7}$ CL=90%
Γ_{440}	$\phi \eta$	< 5	$\times 10^{-7}$ CL=90%
Γ_{441}	$\phi \eta'$	< 5	$\times 10^{-7}$ CL=90%
Γ_{442}	$\phi \pi^+ \pi^-$	$(1.8 \pm 0.5) \times 10^{-7}$	
Γ_{443}	$\phi \rho^0$	< 3.3	$\times 10^{-7}$ CL=90%
Γ_{444}	$\phi f_0(980), f_0 \rightarrow \pi^+ \pi^-$	< 3.8	$\times 10^{-7}$ CL=90%
Γ_{445}	$\phi \omega$	< 7	$\times 10^{-7}$ CL=90%
Γ_{446}	$\phi \phi$	< 2.7	$\times 10^{-8}$ CL=90%
Γ_{447}	$a_0(980)^\pm \pi^\mp, a_0^\pm \rightarrow \eta \pi^\pm$	< 3.1	$\times 10^{-6}$ CL=90%
Γ_{448}	$a_0(1450)^\pm \pi^\mp, a_0^\pm \rightarrow \eta \pi^\pm$	< 2.3	$\times 10^{-6}$ CL=90%
Γ_{449}	$\pi^+ \pi^- \pi^0$	< 7.2	$\times 10^{-4}$ CL=90%
Γ_{450}	$\rho^0 \pi^0$	$(2.0 \pm 0.5) \times 10^{-6}$	
Γ_{451}	$\rho^\mp \pi^\pm$	$[h] (2.30 \pm 0.23) \times 10^{-5}$	
Γ_{452}	$\pi^+ \pi^- \pi^+ \pi^-$	< 1.12	$\times 10^{-5}$ CL=90%
Γ_{453}	$\rho^0 \pi^+ \pi^-$	< 8.8	$\times 10^{-6}$ CL=90%
Γ_{454}	$\rho^0 \rho^0$	$(9.6 \pm 1.5) \times 10^{-7}$	
Γ_{455}	$f_0(980) \pi^+ \pi^-, f_0 \rightarrow \pi^+ \pi^-$	< 3.0	$\times 10^{-6}$ CL=90%

Γ_{456}	$\rho^0 f_0(980), f_0 \rightarrow \pi^+ \pi^-$	$(7.8 \pm 2.5) \times 10^{-7}$	
Γ_{457}	$f_0(980) f_0(980), f_0 \rightarrow \pi^+ \pi^-,$ $f_0 \rightarrow \pi^+ \pi^-$	$< 1.9 \times 10^{-7}$	CL=90%
Γ_{458}	$f_0(980) f_0(980), f_0 \rightarrow \pi^+ \pi^-,$ $f_0 \rightarrow K^+ K^-$	$< 2.3 \times 10^{-7}$	CL=90%
Γ_{459}	$a_1(1260)^\mp \pi^\pm$	$[h] (2.6 \pm 0.5) \times 10^{-5}$	S=1.9
Γ_{460}	$a_2(1320)^\mp \pi^\pm$	$[h] < 6.3 \times 10^{-6}$	CL=90%
Γ_{461}	$\pi^+ \pi^- \pi^0 \pi^0$	$< 3.1 \times 10^{-3}$	CL=90%
Γ_{462}	$\rho^+ \rho^-$	$(2.77 \pm 0.19) \times 10^{-5}$	
Γ_{463}	$a_1(1260)^0 \pi^0$	$< 1.1 \times 10^{-3}$	CL=90%
Γ_{464}	$\omega \pi^0$	$< 5 \times 10^{-7}$	CL=90%
Γ_{465}	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	$< 9.0 \times 10^{-3}$	CL=90%
Γ_{466}	$a_1(1260)^+ \rho^-$	$< 6.1 \times 10^{-5}$	CL=90%
Γ_{467}	$a_1(1260)^0 \rho^0$	$< 2.4 \times 10^{-3}$	CL=90%
Γ_{468}	$b_1^\mp \pi^\pm, b_1^\mp \rightarrow \omega \pi^\mp$	$(1.09 \pm 0.15) \times 10^{-5}$	
Γ_{469}	$b_1^0 \pi^0, b_1^0 \rightarrow \omega \pi^0$	$< 1.9 \times 10^{-6}$	CL=90%
Γ_{470}	$b_1^- \rho^+, b_1^- \rightarrow \omega \pi^-$	$< 1.4 \times 10^{-6}$	CL=90%
Γ_{471}	$b_1^0 \rho^0, b_1^0 \rightarrow \omega \pi^0$	$< 3.4 \times 10^{-6}$	CL=90%
Γ_{472}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$	$< 3.0 \times 10^{-3}$	CL=90%
Γ_{473}	$a_1(1260)^+ a_1(1260)^-, a_1^+ \rightarrow$ $2\pi^+ \pi^-, a_1^- \rightarrow 2\pi^- \pi^+$	$(1.18 \pm 0.31) \times 10^{-5}$	
Γ_{474}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^- \pi^0$	$< 1.1 \%$	CL=90%

Baryon modes

Γ_{475}	$p \bar{p}$	$(1.27 \pm 0.14) \times 10^{-8}$	
Γ_{476}	$p \bar{p} \pi^+ \pi^-$	$(2.87 \pm 0.19) \times 10^{-6}$	
Γ_{477}	$p \bar{p} K^+ \pi^-$	$(6.3 \pm 0.5) \times 10^{-6}$	
Γ_{478}	$p \bar{p} K^0$	$(2.66 \pm 0.32) \times 10^{-6}$	
Γ_{479}	$\Theta(1540)^+ \bar{p}, \Theta^+ \rightarrow p K_S^0$	$[i] < 5 \times 10^{-8}$	CL=90%
Γ_{480}	$f_J(2220) K^0, f_J \rightarrow p \bar{p}$	$< 4.5 \times 10^{-7}$	CL=90%
Γ_{481}	$p \bar{p} K^*(892)^0$	$(1.24^{+0.28}_{-0.25}) \times 10^{-6}$	
Γ_{482}	$f_J(2220) K_0^*, f_J \rightarrow p \bar{p}$	$< 1.5 \times 10^{-7}$	CL=90%
Γ_{483}	$p \bar{p} K^+ K^-$	$(1.21 \pm 0.32) \times 10^{-7}$	
Γ_{484}	$p \bar{p} \pi^0$	$(5.0 \pm 1.9) \times 10^{-7}$	
Γ_{485}	$p \bar{p} p \bar{p}$	$(2.2 \pm 0.4) \times 10^{-8}$	
Γ_{486}	$p \bar{\Lambda} \pi^-$	$(3.16 \pm 0.24) \times 10^{-6}$	
Γ_{487}	$p \bar{\Lambda} \pi^- \gamma$	$< 6.5 \times 10^{-7}$	CL=90%
Γ_{488}	$p \bar{\Sigma}(1385)^-$	$< 2.6 \times 10^{-7}$	CL=90%
Γ_{489}	$\Delta(1232)^+ \bar{p} + \Delta(1232)^- p$	$< 1.6 \times 10^{-6}$	
Γ_{490}	$\Delta^0 \bar{\Lambda}$	$< 9.3 \times 10^{-7}$	CL=90%
Γ_{491}	$p \bar{\Lambda} K^-$	$< 8.2 \times 10^{-7}$	CL=90%
Γ_{492}	$p \bar{\Lambda} D^-$	$(2.5 \pm 0.4) \times 10^{-5}$	
Γ_{493}	$p \bar{\Lambda} D^{*-}$	$(3.4 \pm 0.8) \times 10^{-5}$	

Γ_{494}	$p\bar{\Sigma}^0\pi^-$	$(1.2 \pm 0.4) \times 10^{-6}$	
Γ_{495}	$\bar{\Lambda}\Lambda$	$< 3.2 \times 10^{-7}$	CL=90%
Γ_{496}	$\bar{\Lambda}\Lambda K^0$	$(4.8 \pm_{-0.9}^{+1.0}) \times 10^{-6}$	
Γ_{497}	$\bar{\Lambda}\Lambda K^{*0}$	$(2.5 \pm_{-0.8}^{+0.9}) \times 10^{-6}$	
Γ_{498}	$\bar{\Lambda}\Lambda D^0$	$(1.00 \pm_{-0.26}^{+0.30}) \times 10^{-5}$	
Γ_{499}	$D^0\Sigma^0\bar{\Lambda} + \text{c.c.}$	$< 3.1 \times 10^{-5}$	CL=90%
Γ_{500}	$\Delta^0\bar{\Delta}^0$	$< 1.5 \times 10^{-3}$	CL=90%
Γ_{501}	$\Delta^{++}\bar{\Delta}^{--}$	$< 1.1 \times 10^{-4}$	CL=90%
Γ_{502}	$\bar{D}^0 p\bar{p}$	$(1.04 \pm 0.07) \times 10^{-4}$	
Γ_{503}	$D_s^- \bar{\Lambda} p$	$(2.8 \pm 0.9) \times 10^{-5}$	
Γ_{504}	$\bar{D}^{*}(2007)^0 p\bar{p}$	$(9.9 \pm 1.1) \times 10^{-5}$	
Γ_{505}	$D^{*}(2010)^- p\bar{n}$	$(1.4 \pm 0.4) \times 10^{-3}$	
Γ_{506}	$D^- p\bar{p}\pi^+$	$(3.32 \pm 0.31) \times 10^{-4}$	
Γ_{507}	$D^{*}(2010)^- p\bar{p}\pi^+$	$(4.7 \pm 0.5) \times 10^{-4}$	S=1.2
Γ_{508}	$\bar{D}^0 p\bar{p}\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$	
Γ_{509}	$\bar{D}^{*0} p\bar{p}\pi^+\pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$	
Γ_{510}	$\Theta_c \bar{p}\pi^+, \Theta_c \rightarrow D^- p$	$< 9 \times 10^{-6}$	CL=90%
Γ_{511}	$\Theta_c \bar{p}\pi^+, \Theta_c \rightarrow D^{*-} p$	$< 1.4 \times 10^{-5}$	CL=90%
Γ_{512}	$\bar{\Sigma}_c^{--}\Delta^{++}$	$< 8 \times 10^{-4}$	CL=90%
Γ_{513}	$\bar{\Lambda}_c^- p\pi^+\pi^-$	$(1.01 \pm 0.14) \times 10^{-3}$	S=1.4
Γ_{514}	$\bar{\Lambda}_c^- p$	$(1.52 \pm 0.17) \times 10^{-5}$	
Γ_{515}	$\bar{\Lambda}_c^- p\pi^0$	$(1.53 \pm 0.18) \times 10^{-4}$	
Γ_{516}	$\Sigma_c(2455)^- p$	$< 2.4 \times 10^{-5}$	
Γ_{517}	$\bar{\Lambda}_c^- p\pi^+\pi^-\pi^0$	$< 5.07 \times 10^{-3}$	CL=90%
Γ_{518}	$\bar{\Lambda}_c^- p\pi^+\pi^-\pi^+\pi^-$	$< 2.74 \times 10^{-3}$	CL=90%
Γ_{519}	$\bar{\Lambda}_c^- p\pi^+\pi^- (\text{nonresonant})$	$(5.4 \pm 1.0) \times 10^{-4}$	S=1.3
Γ_{520}	$\bar{\Sigma}_c(2520)^{--} p\pi^+$	$(1.01 \pm 0.18) \times 10^{-4}$	
Γ_{521}	$\bar{\Sigma}_c(2520)^0 p\pi^-$	$< 3.1 \times 10^{-5}$	CL=90%
Γ_{522}	$\bar{\Sigma}_c(2455)^0 p\pi^-$	$(1.08 \pm 0.09) \times 10^{-4}$	
Γ_{523}	$\bar{\Sigma}_c(2455)^0 N^0, N^0 \rightarrow p\pi^-$	$(6.3 \pm 1.6) \times 10^{-5}$	
Γ_{524}	$\bar{\Sigma}_c(2455)^{--} p\pi^+$	$(1.89 \pm 0.15) \times 10^{-4}$	
Γ_{525}	$\Lambda_c^- pK^+\pi^-$	$(3.4 \pm 0.7) \times 10^{-5}$	
Γ_{526}	$\bar{\Sigma}_c(2455)^{--} pK^+, \bar{\Sigma}_c^{--} \rightarrow \bar{\Lambda}_c^- \pi^-$	$(8.7 \pm 2.5) \times 10^{-6}$	
Γ_{527}	$\Lambda_c^- pK^*(892)^0$	$< 2.42 \times 10^{-5}$	CL=90%
Γ_{528}	$\Lambda_c^- pK^+K^-$	$(2.0 \pm 0.4) \times 10^{-5}$	
Γ_{529}	$\Lambda_c^- p\phi$	$< 9 \times 10^{-6}$	CL=90%
Γ_{530}	$\Lambda_c^- p\bar{p}p$	$< 2.8 \times 10^{-6}$	
Γ_{531}	$\bar{\Lambda}_c^- \Lambda K^+$	$(4.8 \pm 1.1) \times 10^{-5}$	

Γ ₅₃₂	$\bar{\Lambda}_c^- \Lambda_c^+$		< 1.6	$\times 10^{-5}$	CL=95%
Γ ₅₃₃	$\bar{\Lambda}_c(2593)^- / \bar{\Lambda}_c(2625)^- p$		< 1.1	$\times 10^{-4}$	CL=90%
Γ ₅₃₄	$\bar{\Xi}_c^- \Lambda_c^+$		(1.1 ± 0.8)	$\times 10^{-3}$	
Γ ₅₃₅	$\bar{\Xi}_c^- \Lambda_c^+, \bar{\Xi}_c^- \rightarrow \Xi^+ \pi^- \pi^-$		(2.3 ± 1.0)	$\times 10^{-5}$	S=1.8
Γ ₅₃₆	$\bar{\Xi}_c^- \Lambda_c^+, \bar{\Xi}_c^- \rightarrow \bar{p} K^+ \pi^-$		(5.2 ± 1.6)	$\times 10^{-6}$	
Γ ₅₃₇	$\Lambda_c^+ \Lambda_c^- K^0$		(4.0 ± 0.9)	$\times 10^{-4}$	
Γ ₅₃₈	$\bar{\Lambda}_c(2910)^- p, \bar{\Lambda}_c^- \rightarrow \bar{\Sigma}_c(2455)^- \pi^+$		(1.2 ± 0.4)	$\times 10^{-5}$	
Γ ₅₃₉	$\bar{\Lambda}_c(2910)^- p, \bar{\Lambda}_c^- \rightarrow \bar{\Sigma}_c(2455)^0 \pi^-$		(10 ± 4)	$\times 10^{-6}$	
Γ ₅₄₀	$\bar{\Xi}_c(2930)^- \Lambda_c^+, \bar{\Xi}_c^- \rightarrow \Lambda_c^- K^0$		(2.4 ± 0.6)	$\times 10^{-4}$	
Γ ₅₄₁	$\Omega_c^0 \bar{\Lambda}, \Omega_c^0 \rightarrow \Omega^- \pi^+$		< 9.7	$\times 10^{-8}$	CL=95%
Γ ₅₄₂	$\Omega_c(2770)^0 \bar{\Lambda}, \Omega_c^0 \rightarrow \Omega^- \pi^+$		< 3.12	$\times 10^{-7}$	CL=95%
Γ ₅₄₃	$\Omega_c^0 \Lambda, \Omega_c^0 \rightarrow \Omega^- \pi^+$		< 9.5	$\times 10^{-8}$	CL=95%
Γ ₅₄₄	$\Omega_c(2770)^0 \Lambda, \Omega_c^0 \rightarrow \Omega^- \pi^+$		< 1.00	$\times 10^{-7}$	CL=95%
Γ ₅₄₅	$\Lambda \psi_{DS}$	[j]	< 0.13–5.2	$\times 10^{-5}$	

Lepton Family number (LF) or Lepton number (L) or Baryon number (B) violating modes, or/and $\Delta B = 1$ weak neutral current (B1) modes

Γ ₅₄₆	$\gamma \gamma$	B1	< 6.4	$\times 10^{-8}$	CL=90%
Γ ₅₄₇	$e^+ e^-$	B1	< 2.5	$\times 10^{-9}$	CL=90%
Γ ₅₄₈	$e^+ e^- \gamma$	B1	< 1.2	$\times 10^{-7}$	CL=90%
Γ ₅₄₉	$\mu^+ \mu^-$	B1	< 1.5	$\times 10^{-10}$	CL=90%
Γ ₅₅₀	$\mu^+ \mu^- \gamma$	B1			
Γ ₅₅₁	$\mu^+ \mu^- \mu^+ \mu^-$	B1	< 1.8	$\times 10^{-10}$	CL=95%
Γ ₅₅₂	$SP, S \rightarrow \mu^+ \mu^-, P \rightarrow \mu^+ \mu^-$	B1 [k]	< 6.0	$\times 10^{-10}$	CL=95%
Γ ₅₅₃	$aa, a \rightarrow \mu^+ \mu^-$	B1	< 2.3	$\times 10^{-10}$	CL=95%
Γ ₅₅₄	$\tau^+ \tau^-$	B1	< 2.1	$\times 10^{-3}$	CL=95%
Γ ₅₅₅	$\pi^0 \ell^+ \ell^-$	B1 [a]	< 3.8	$\times 10^{-8}$	CL=90%
Γ ₅₅₆	$\pi^0 e^+ e^-$	B1	< 7.9	$\times 10^{-8}$	CL=90%
Γ ₅₅₇	$\pi^0 \mu^+ \mu^-$	B1	< 5.9	$\times 10^{-8}$	CL=90%
Γ ₅₅₈	$\eta \ell^+ \ell^-$	B1 [a]	< 4.8	$\times 10^{-8}$	CL=90%
Γ ₅₅₉	$\eta e^+ e^-$	B1	< 1.05	$\times 10^{-7}$	CL=90%
Γ ₅₆₀	$\eta \mu^+ \mu^-$	B1	< 9.4	$\times 10^{-8}$	CL=90%
Γ ₅₆₁	$\rho(770)^0 e^+ e^-$	B1	< 4.55	$\times 10^{-7}$	CL=90%
Γ ₅₆₂	$\omega e^+ e^-$	B1	< 3.07	$\times 10^{-7}$	CL=90%
Γ ₅₆₃	$\omega \mu^+ \mu^-$	B1	< 2.49	$\times 10^{-7}$	CL=90%
Γ ₅₆₄	$\omega \ell^+ \ell^-$	B1	< 2.20	$\times 10^{-7}$	CL=90%
Γ ₅₆₅	$\pi^0 \nu \bar{\nu}$	B1	< 9	$\times 10^{-6}$	CL=90%
Γ ₅₆₆	$K^0 \ell^+ \ell^-$	B1 [a]	(3.3 ± 0.6)	$\times 10^{-7}$	
Γ ₅₆₇	$K^0 e^+ e^-$	B1	(2.5 ± 1.1 / 0.9)	$\times 10^{-7}$	S=1.3
Γ ₅₆₈	$K^0 \mu^+ \mu^-$	B1	(3.39 ± 0.35)	$\times 10^{-7}$	S=1.1

Γ_{569}	$K^0 \nu \bar{\nu}$	$B1$	< 2.6	$\times 10^{-5}$	CL=90%
Γ_{570}	$\rho^0 \nu \bar{\nu}$	$B1$	< 4.0	$\times 10^{-5}$	CL=90%
Γ_{571}	$K^*(892)^0 \ell^+ \ell^-$	$B1$	[a] $(9.9 \pm 1.2 \pm 1.1)$	$\times 10^{-7}$	
Γ_{572}	$K^*(892)^0 e^+ e^-$	$B1$	$(1.03 \pm 0.19 \pm 0.17)$	$\times 10^{-6}$	
Γ_{573}	$K^*(892)^0 \mu^+ \mu^-$	$B1$	(9.4 ± 0.5)	$\times 10^{-7}$	
Γ_{574}	$K^*(892)^0 \chi, \chi \rightarrow \mu^+ \mu^-$	$B1$			
Γ_{575}	$K^*(892)^0 \tau^+ \tau^-$	$B1$	< 3.1	$\times 10^{-3}$	CL=90%
Γ_{576}	$\bar{D}^{*0} \mu^+ \mu^-$	$B1$	< 4.0	$\times 10^{-8}$	CL=90%
Γ_{577}	$\pi^+ \pi^- \mu^+ \mu^-$	$B1$	(2.1 ± 0.5)	$\times 10^{-8}$	
Γ_{578}	$K^*(892)^0 \nu \bar{\nu}$	$B1$	< 1.8	$\times 10^{-5}$	CL=90%
Γ_{579}	invisible	$B1$	< 2.4	$\times 10^{-5}$	CL=90%
Γ_{580}	$\nu \bar{\nu} \gamma$	$B1$	< 1.6	$\times 10^{-5}$	CL=90%
Γ_{581}	$\phi \mu^+ \mu^-$	$B1$	< 3.2	$\times 10^{-9}$	CL=90%
Γ_{582}	$\phi \nu \bar{\nu}$	$B1$	< 1.27	$\times 10^{-4}$	CL=90%
Γ_{583}	$e^\pm \mu^\mp$	LF	[h] < 1.0	$\times 10^{-9}$	CL=90%
Γ_{584}	$\pi^0 e^\pm \mu^\mp$	LF	< 1.4	$\times 10^{-7}$	CL=90%
Γ_{585}	$K^0 e^\pm \mu^\mp$	LF	< 3.8	$\times 10^{-8}$	CL=90%
Γ_{586}	$K^*(892)^0 e^+ \mu^-$	LF	< 6.8	$\times 10^{-9}$	CL=90%
Γ_{587}	$K^*(892)^0 e^- \mu^+$	LF	< 5.7	$\times 10^{-9}$	CL=90%
Γ_{588}	$K^*(892)^0 e^\pm \mu^\mp$	LF	< 1.01	$\times 10^{-8}$	CL=90%
Γ_{589}	$K^*(892)^0 \tau^+ \mu^-$	LF	< 1.0	$\times 10^{-5}$	CL=90%
Γ_{590}	$K^*(892)^0 \tau^- \mu^+$	LF	< 8.2	$\times 10^{-6}$	CL=90%
Γ_{591}	$e^\pm \tau^\mp$	LF	[h] < 1.6	$\times 10^{-5}$	CL=90%
Γ_{592}	$\mu^\pm \tau^\mp$	LF	[h] < 1.4	$\times 10^{-5}$	CL=95%
Γ_{593}	$\rho \mu^-$	L, B	< 2.6	$\times 10^{-9}$	CL=90%
Γ_{594}	$\Lambda_c^+ \mu^-$	L, B	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{595}	$\Lambda_c^+ e^-$	L, B	< 4	$\times 10^{-6}$	CL=90%

[a] An ℓ indicates an e or a μ mode, not a sum over these modes.

[b] $\bar{D}^{**}$ represents an excited state with mass $2.2 < M < 2.8 \text{ GeV}/c^2$.

[c] $\chi_{c1}(3872)^+$ is a hypothetical charged partner of the $\chi_{c1}(3872)$.

[d] Stands for the possible candidates of $K^*(1410)$, $K_0^*(1430)$ and $K_2^*(1430)$.

[e] B^0 and B_s^0 contributions not separated. Limit is on weighted average of the two decay rates.

[f] This decay refers to the coherent sum of resonant and nonresonant $J^P = 0^+ K \pi$ components with $1.60 < m_{K \pi} < 2.15 \text{ GeV}/c^2$.

[g] $X(214)$ is a hypothetical particle of mass $214 \text{ MeV}/c^2$ reported by the HyperCP experiment, Physical Review Letters **94** 021801 (2005)

[*h*] The value is for the sum of the charge states or particle/antiparticle states indicated.

[*i*] $\Theta(1540)^+$ denotes a possible narrow pentaquark state.

[*j*] ψ_{DS} is a GeV-scale dark sector antibaryon (mass range 1–4 GeV/ c^2).

[*k*] Here *S* and *P* are the hypothetical scalar and pseudoscalar particles with masses of 2.5 GeV/ c^2 and 214.3 MeV/ c^2 , respectively.

FIT INFORMATION

An overall fit to 36 branching ratios uses 95 measurements to determine 22 parameters. The overall fit has a $\chi^2 = 70.9$ for 73 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_{10}	31										
x_{38}	0	0									
x_{50}	0	0	28								
x_{76}	0	0	4	13							
x_{125}	0	0	19	6	1						
x_{135}	0	0	7	2	0	1					
x_{214}	0	0	0	0	0	0	0				
x_{216}	0	0	0	0	0	0	0	0			
x_{264}	0	0	0	0	0	0	0	0	0		
x_{270}	0	0	0	0	0	0	0	0	0	0	19
x_{276}	0	0	0	0	0	0	0	0	0	29	0
x_{282}	0	0	0	0	0	0	0	0	0	6	0
x_{288}	0	0	0	0	0	0	0	0	0	0	0
x_{322}	0	0	0	0	0	0	0	0	0	0	0
x_{356}	0	0	0	0	0	0	0	0	0	0	0
x_{363}	0	0	0	0	0	0	0	0	0	0	0
x_{377}	0	0	0	0	0	0	0	0	0	0	0
x_{423}	0	0	0	0	0	0	0	0	0	0	0
x_{454}	0	0	0	0	0	0	0	0	0	0	0
x_{568}	0	0	0	0	0	0	0	0	0	0	0
x_{573}	0	0	0	0	0	0	0	0	0	15	0
	x_7	x_{10}	x_{38}	x_{50}	x_{76}	x_{125}	x_{135}	x_{214}	x_{216}	x_{264}	

x276	0									
x282	0	22								
x288	0	0	0							
x322	0	0	0	0						
x356	0	0	0	0	16					
x363	0	0	0	0	27	4				
x377	0	0	0	0	0	0	0			
x423	0	0	0	24	0	0	0	0		
x454	0	0	0	0	0	0	0	20	0	
x568	0	0	0	0	0	0	0	0	0	0
x573	0	4	1	0	0	0	0	0	0	0
	x270	x276	x282	x288	x322	x356	x363	x377	x423	x454
x573	0									
	x568									

B^0 BRANCHING RATIOS

For branching ratios in which the charge of the decaying B is not determined, see the $B^\pm$ section.

$\Gamma(\ell^+ \nu_\ell X) / \Gamma_{\text{total}}$

Γ_1 / Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
10.33 ± 0.28 OUR EVALUATION	(Produced by HFLAV)		
10.14 ± 0.30 OUR AVERAGE	Error includes scale factor of 1.1.		
10.46 ± 0.30 ± 0.23	¹ URQUIJO	07	BELL $e^+ e^- \rightarrow \gamma(4S)$
9.64 ± 0.27 ± 0.33	² AUBERT,B	06Y	BABR $e^+ e^- \rightarrow \gamma(4S)$
10.78 ± 0.60 ± 0.69	³ ARTUSO	97	CLE2 $e^+ e^- \rightarrow \gamma(4S)$
9.3 ± 1.1 ± 1.5	ALBRECHT	94	ARG $e^+ e^- \rightarrow \gamma(4S)$
9.9 ± 3.0 ± 0.9	HENDERSON	92	CLEO $e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.32 ± 0.36 ± 0.35	⁴ OKABE	05	BELL Repl. by URQUIJO 07
10.9 ± 0.7 ± 1.1	ATHANAS	94	CLE2 Sup. by ARTUSO 97

¹ URQUIJO 07 report a measurement of $(9.80 \pm 0.29 \pm 0.21)\%$ for the partial branching fraction of $B \rightarrow e \nu_e X_C$ decay with electron energy above 0.6 GeV. We converted the result to $B \rightarrow e \nu_e X$ branching fraction.

² The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame. The best precision on the ratio is achieved for a momentum threshold of 1.0 GeV: $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.074 \pm 0.041 \pm 0.026$.

³ ARTUSO 97 uses partial reconstruction of $B \rightarrow D^* \ell \nu_\ell$ and inclusive semileptonic branching ratio from BARISH 96B ($0.1049 \pm 0.0017 \pm 0.0043$).

⁴ The measurements are obtained for charged and neutral B mesons partial rates of semileptonic decay to electrons with momentum above 0.6 GeV/c in the B rest frame, and their ratio of $B(B^+ \rightarrow e^+ \nu_e X) / B(B^0 \rightarrow e^+ \nu_e X) = 1.08 \pm 0.05 \pm 0.02$.

$\Gamma(e^+ \nu_e X_c)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$10.08 \pm 0.30 \pm 0.22$	¹ URQUIJO	07	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measure the independent B^+ and B^0 partial branching fractions with electron threshold energies of 0.4 GeV.

 $\Gamma(\ell^+ \nu_\ell X_u)/\Gamma_{\text{total}}$ Γ_3/Γ

Requires $E_\ell^* > 1$ GeV, where E_ℓ^* is lepton energy in B rest frame.

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.51 \pm 0.10 \pm 0.16$	¹ CAO	21A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ The correlation of 50% with $B(B^+ \rightarrow \ell^+ \nu_\ell X_u)$ (lepton energy in B rest frame $E_\ell^* > 1$ GeV) was reported.

 $\Gamma(D^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_5/Γ

ℓ denotes e or μ , not the sum.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
2.10 ± 0.07 OUR EVALUATION	(Produced by HFLAV)		
2.25 ± 0.08 OUR AVERAGE			

$2.31 \pm 0.03 \pm 0.11$	¹ GLATTAUER	16	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
$2.21 \pm 0.11 \pm 0.11$	² AUBERT	10	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$2.09 \pm 0.13 \pm 0.18$	³ BARTELT	99	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
$2.35 \pm 0.20 \pm 0.44$	⁴ BUSKULIC	97	ALEP $e^+ e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2.21 \pm 0.11 \pm 0.12$	² AUBERT	08Q	BABR Repl. by AUBERT 10
$2.13 \pm 0.12 \pm 0.39$	ABE	02E	BELL Repl. by GLATTAUER 16
$1.87 \pm 0.15 \pm 0.32$	⁵ ATHANAS	97	CLE2 Repl. by BARTELT 99
$1.8 \pm 0.6 \pm 0.3$	⁶ FULTON	91	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
$2.0 \pm 0.7 \pm 0.6$	⁷ ALBRECHT	89J	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side while the other, on the signal side, is partially reconstructed from $B \rightarrow D \ell \nu$.

² Uses a fully reconstructed B meson as a tag on the recoil side.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ BUSKULIC 97 assumes fraction (B^+) = fraction (B^0) = $(37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and branching ratio of D^* and D decays.

⁵ ATHANAS 97 uses missing energy and missing momentum to reconstruct neutrino.

⁶ FULTON 91 assumes assuming equal production of B^0 and B^+ at the $\Upsilon(4S)$ and uses Mark III D and D^* branching ratios.

⁷ ALBRECHT 89J reports $0.018 \pm 0.006 \pm 0.005$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$.

 $\Gamma(D^- \ell^+ \nu_\ell)/\Gamma(\ell^+ \nu_\ell X)$ Γ_5/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
$0.230 \pm 0.011 \pm 0.011$	¹ AUBERT	10	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson on the recoil side.

 $\Gamma(D^- \ell^+ \nu_\ell)/\Gamma(D \ell^+ \nu_\ell X)$ Γ_5/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
$0.215 \pm 0.016 \pm 0.013$	¹ AUBERT	07AN	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson on the recoil side.

$\Gamma(D^- \tau^+ \nu_\tau)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.04 \pm 0.35 \pm 0.18$	¹ AUBERT	08N BABR	Repl. by AUBERT 09S
¹ Uses a fully reconstructed B meson as a tag on the recoil side.			

 $\Gamma(D^- \tau^+ \nu_\tau)/\Gamma(D^- \ell^+ \nu_\ell)$ Γ_6/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
$0.469 \pm 0.084 \pm 0.053$	^{1,2} LEES	12D BABR	$e^+ e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.489 \pm 0.165 \pm 0.069$	¹ AUBERT	09S BABR	Repl. by LEES 12D
¹ Uses a fully reconstructed B meson as a tag on the recoil side.			
² Uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ and $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ and e^+ or μ^+ as ℓ^+ .			


 $\Gamma(D^{*(2010)-} \ell^+ \nu_\ell)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
4.87 ±0.09	OUR EVALUATION	(Produced by HFLAV) This value assumes isospin symmetry.		
5.11 ±0.14	OUR FIT	Error includes scale factor of 1.4.		
5.11 ±0.15	OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
4.922±0.023±0.220		¹ ADACHI	23J BELL	$e^+e^- \rightarrow \Upsilon(4S)$
5.02 ±0.02 ±0.16		² WAHEED	21 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
5.49 ±0.16 ±0.25		³ AUBERT	08Q BABR	$e^+e^- \rightarrow \Upsilon(4S)$
4.69 ±0.04 ±0.34		⁴ AUBERT	08R BABR	$e^+e^- \rightarrow \Upsilon(4S)$
5.90 ±0.22 ±0.50		⁵ ABDALLAH	04D DLPH	$e^+e^- \rightarrow Z^0$
6.09 ±0.19 ±0.40		⁶ ADAM	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
4.70 ±0.13 ^{+0.36} _{-0.31}		⁷ ABREU	01H DLPH	$e^+e^- \rightarrow Z$
5.26 ±0.20 ±0.46		⁸ ABBIENDI	00Q OPAL	$e^+e^- \rightarrow Z$
5.53 ±0.26 ±0.52		⁹ BUSKULIC	97 ALEP	$e^+e^- \rightarrow Z$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.917±0.032±0.216		¹⁰ ADACHI	23J BELL	$e^+e^- \rightarrow \Upsilon(4S)$
4.926±0.032±0.231		¹¹ ADACHI	23J BELL	$e^+e^- \rightarrow \Upsilon(4S)$
4.90 ±0.02 ±0.16		² WAHEED	19 BELL	Repl. by WAHEED 21
4.58 ±0.03 ±0.26		² DUNGEL	10 BELL	Repl. by WAHEED 19
4.90 ±0.07 ^{+0.36} _{-0.35}		⁵ AUBERT	05E BABR	Repl. by AUBERT 08R
5.39 ±0.11 ±0.34		¹² ABDALLAH	04D DLPH	$e^+e^- \rightarrow Z^0$
4.59 ±0.23 ±0.40		¹³ ABE	02F BELL	Repl. by DUNGEL 10
6.09 ±0.19 ±0.40		¹⁴ BRIERE	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5.08 ±0.21 ±0.66		¹⁵ ACKERSTAFF	97G OPAL	Repl. by ABBIENDI 00Q
5.52 ±0.17 ±0.68		¹⁶ ABREU	96P DLPH	Repl. by ABREU 01H
4.49 ±0.32 ±0.39	376	¹⁷ BARISH	95 CLE2	Repl. by ADAM 03
5.18 ±0.30 ±0.62	410	¹⁸ BUSKULIC	95N ALEP	Sup. by BUSKULIC 97
4.5 ±0.3 ±0.4		¹⁹ ALBRECHT	94 ARG	$e^+e^- \rightarrow \Upsilon(4S)$
4.7 ±0.5 ±0.5	235	²⁰ ALBRECHT	93 ARG	$e^+e^- \rightarrow \Upsilon(4S)$
seen	398	²¹ SANGHERA	93 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
7.0 ±1.8 ±1.4		²² ANTREASYAN	90B CBAL	$e^+e^- \rightarrow \Upsilon(4S)$
		²³ ALBRECHT	89C ARG	$e^+e^- \rightarrow \Upsilon(4S)$

6.0	± 1.0	± 1.4		24	ALBRECHT	89J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
4.0	± 0.4	± 0.6		25	BORTOLETTO	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
7.0	± 1.2	± 1.9	47	26	ALBRECHT	87J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

- ¹ Uses fully reconstructed $D^{*-}\ell^+\nu$ events with $\ell = e$ or μ .
- ² WAHEED 21 uses fully reconstructed $D^{*-}\ell^+\nu$ events ($\ell = e$ or μ).
- ³ Uses a fully reconstructed B meson as a tag on the recoil side.
- ⁴ Measured using fully reconstructed D^* sample and a simultaneous fit to the Caprini-Lellouch-Neubert form factor parameters: $\rho^2 = 1.191 \pm 0.048 \pm 0.028$, $R_1(1) = 1.429 \pm 0.061 \pm 0.044$, and $R_2(1) = 0.827 \pm 0.038 \pm 0.022$.
- ⁵ Measured using fully reconstructed D^* sample.
- ⁶ Uses the combined fit of both $B^0 \rightarrow D^{*}(2010)^-\ell\nu$ and $B^+ \rightarrow \bar{D}^{*}(2007)^0\ell\nu$ samples.
- ⁷ ABREU 01H measured using about 5000 partial reconstructed D^* sample.
- ⁸ ABBIENDI 00Q assumes the fraction $B(b \rightarrow B^0) = (39.7^{+1.8}_{-2.2})\%$. This result is an average of two methods using exclusive and partial D^* reconstruction.
- ⁹ BUSKULIC 97 assumes fraction $(B^+) = \text{fraction}(B^0) = (37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and D^* and D branching fractions.
- ¹⁰ Uses fully reconstructed $D^{*-}e^+\nu$ events.
- ¹¹ Uses fully reconstructed $D^{*-}\mu^+\nu$ events.
- ¹² Combines with previous partial reconstructed D^* measurement.
- ¹³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ¹⁴ The results are based on the same analysis and data sample reported in ADAM 03.
- ¹⁵ ACKERSTAFF 97G assumes fraction $(B^+) = \text{fraction}(B^0) = (37.8 \pm 2.2)\%$ and PDG 96 values for B lifetime and branching ratio of D^* and D decays.
- ¹⁶ ABREU 96P result is the average of two methods using exclusive and partial D^* reconstruction.
- ¹⁷ BARISH 95 use $B(D^0 \rightarrow K^-\pi^+) = (3.91 \pm 0.08 \pm 0.17)\%$ and $B(D^{*+} \rightarrow D^0\pi^+) = (68.1 \pm 1.0 \pm 1.3)\%$.
- ¹⁸ BUSKULIC 95N assumes fraction $(B^+) = \text{fraction}(B^0) = 38.2 \pm 1.3 \pm 2.2\%$ and $\tau_{B^0} = 1.58 \pm 0.06$ ps. $\Gamma(D^{*-}\ell^+\nu_\ell)/\text{total} = [5.18 - 0.13(\text{fraction}(B^0) - 38.2) - 1.5(\tau_{B^0} - 1.58)]\%$.
- ¹⁹ ALBRECHT 94 assumes $B(D^{*+} \rightarrow D^0\pi^+) = 68.1 \pm 1.0 \pm 1.3\%$. Uses partial reconstruction of D^{*+} and is independent of D^0 branching ratios.
- ²⁰ ALBRECHT 93 reports $0.052 \pm 0.005 \pm 0.006$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^-\pi^+)$. We have taken their average e and μ value. They also obtain $\alpha = 2*\Gamma^0/(\Gamma^- + \Gamma^+) - 1 = 1.1 \pm 0.4 \pm 0.2$, $A_{AF} = 3/4*(\Gamma^- - \Gamma^+)/\Gamma = 0.2 \pm 0.08 \pm 0.06$ and a value of $|V_{cb}| = 0.036\text{--}0.045$ depending on model assumptions.
- ²¹ Combining $\bar{D}^{*0}\ell^+\nu_\ell$ and $\bar{D}^{*-}\ell^+\nu_\ell$ SANGHERA 93 test $V\text{--}A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4*(\Gamma^- - \Gamma^+)/\Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb} , they measure V , A_1 , and A_2 , the three form factors for the $D^*\ell\nu_\ell$ decay, where results are slightly dependent on model assumptions.
- ²² ANTREASYAN 90B is average over B and $\bar{D}^{*}(2010)$ charge states.
- ²³ The measurement of ALBRECHT 89C suggests a D^* polarization γ_L/γ_T of 0.85 ± 0.45 . or $\alpha = 0.7 \pm 0.9$.
- ²⁴ ALBRECHT 89J is ALBRECHT 87J value rescaled using $B(D^{*}(2010)^- \rightarrow D^0\pi^-) = 0.57 \pm 0.04 \pm 0.04$. Superseded by ALBRECHT 93.
- ²⁵ We have taken average of the the BORTOLETTO 89B values for electrons and muons, $0.046 \pm 0.005 \pm 0.007$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^-\pi^+)$. The measurement suggests a D^* polarization parameter value $\alpha = 0.65 \pm 0.66 \pm 0.25$.

²⁶ ALBRECHT 87J assume μ -e universality, the $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.45$, the $B(D^0 \rightarrow K^- \pi^+) = (0.042 \pm 0.004 \pm 0.004)$, and the $B(D^*(2010)^- \rightarrow D^0 \pi^-) = 0.49 \pm 0.08$. Superseded by ALBRECHT 89J.



$\Gamma(D^*(2010)^- \ell^+ \nu_\ell) / \Gamma(D \ell^+ \nu_\ell X)$	Γ_7 / Γ_4		
VALUE	DOCUMENT ID	TECN	COMMENT
0.537±0.031±0.036	¹ AUBERT	07AN BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson on the recoil side.

$\Gamma(D^*(2010)^- e^+ \nu_e) / \Gamma(D^*(2010)^- \mu^+ \nu_\mu)$	Γ_8 / Γ_9		
VALUE	DOCUMENT ID	TECN	COMMENT
1.001±0.019 OUR AVERAGE			
0.998±0.009±0.020	ADACHI	23J BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
1.011±0.035±0.024	PRIM	23 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(D^*(2010)^- \tau^+ \nu_\tau) / \Gamma_{\text{total}}$	Γ_{10} / Γ		
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.48±0.09 OUR FIT	Error includes scale factor of 1.2.		
1.48±0.18 OUR AVERAGE	Error includes scale factor of 1.1.		
1.42±0.094±0.140	¹ AAIJ	18D LHCB	pp at 7, 8 TeV
2.02 ^{+0.40} _{-0.37} ±0.37	² MATYJA	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.11±0.51 ±0.06 ³ AUBERT 08N BABR Repl. by AUBERT 09S

¹ Normalizes to $B(B^0 \rightarrow D^*(2010)^- \pi^+ \pi^- \pi^+) = (7.214 \pm 0.28) \times 10^{-3}$.

² Observed in the recoil of the accompanying B meson.

³ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(D^*(2010)^- \tau^+ \nu_\tau) / \Gamma(D^*(2010)^- \ell^+ \nu_\ell)$ Γ_{10}/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
0.290 ± 0.018 OUR FIT	Error includes scale factor of 1.2.		
0.291 ± 0.022 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
$0.267 \pm 0.012 \pm 0.020$	^{1,2} AAIJ	23W LHCB	pp at 13 TeV
$0.302 \pm 0.030 \pm 0.011$	³ SATO	16B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$0.355 \pm 0.039 \pm 0.021$	^{4,5} LEES	12D BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.260 \pm 0.015 \pm 0.019$	¹ AAIJ	23W LHCB	pp at 13 TeV
$0.291 \pm 0.019 \pm 0.029$	¹ AAIJ	18D LHCB	Repl. by AAIJ 23W
$0.336 \pm 0.027 \pm 0.030$	⁶ AAIJ	15Q LHCB	Repl. by AAIJ 23AR
$0.207 \pm 0.095 \pm 0.008$	⁴ AUBERT	09S BABR	Repl. by LEES 12D

¹ Uses $\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau$ and $\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \pi^0 \bar{\nu}_\tau$, and μ^+ as ℓ^+ .


² Combination with measurement from AAIJ 18D.

³ Uses semileptonic B decay events for tagging and $\tau^+ \rightarrow \ell^+ \nu_\ell \bar{\nu}_\tau$ mode.

⁴ Uses a fully reconstructed B meson as a tag on the recoil side.

⁵ Uses $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ and $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ and e^+ or μ^+ as ℓ^+ .

⁶ Uses $\tau^+ \rightarrow \mu^+ \nu_\mu \bar{\nu}_\tau$ and μ^+ as ℓ^+ .



$\Gamma(D^*(2010)^- \tau^+ \nu_\tau) / \Gamma(D^*(2010)^- \pi^+ \pi^- \pi^-)$ Γ_{10}/Γ_{61}

VALUE	DOCUMENT ID	TECN	COMMENT
$1.84 \pm 0.08 \pm 0.10$	^{1,2} AAIJ	23W LHCB	pp at 7, 8, 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.79 \pm 0.11 \pm 0.11$	¹ AAIJ	23W LHCb	pp at 13 TeV
$1.97 \pm 0.13 \pm 0.18$	¹ AAIJ	18D LHCb	Repl. by AAIJ 23W

¹ Uses $\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \bar{\nu}_\tau$ and $\tau^+ \rightarrow \pi^+ \pi^- \pi^+ \pi^0 \bar{\nu}_\tau$ modes.

² Combination with result from AAIJ 18D.

$\Gamma(\bar{D}^{(*)} n \pi \ell^+ \nu_\ell (n \geq 1)) / \Gamma(D \ell^+ \nu_\ell X)$ Γ_{11}/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
$0.248 \pm 0.032 \pm 0.030$	¹ AUBERT	07AN BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson on the recoil side.

$\Gamma(\bar{D}^0 \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.64 ± 0.20 OUR AVERAGE			

$3.60 \pm 0.18 \pm 0.11$ MEIER 23 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$4.3 \pm 0.8 \pm 0.3$ ¹ AUBERT 08Q BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.05 \pm 0.36 \pm 0.41$ VOSSEN 18 BELL Repl. by MEIER 23

$4.2 \pm 0.8 \pm 0.1$ ^{1,2} LIVENTSEV 08 BELL Repl. by VOSSEN 18

$3.4 \pm 1.0 \pm 0.1$ ³ LIVENTSEV 05 BELL Repl. by LIVENTSEV 08

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² LIVENTSEV 08 reports $(4.2 \pm 0.7 \pm 0.6) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow \bar{D}^0 \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \ell^+ \nu_\ell)]$ assuming $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.12 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.10 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ LIVENTSEV 05 reports $[\Gamma(B^0 \rightarrow \bar{D}^0 \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell)] = 0.15 \pm 0.03 \pm 0.03$ which we multiply by our best value $B(B^+ \rightarrow \bar{D}^0 \ell^+ \nu_\ell) = (2.26 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_0^{*}(2300)^- \ell^+ \nu_\ell, D_0^{*-} \rightarrow \bar{D}^0 \pi^-) / \Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.4	90	MEIER	23 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$44 \pm 8 \pm 6$ ¹ AUBERT 08BL BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$20 \pm 7 \pm 5$ ¹ LIVENTSEV 08 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$\Gamma(D_2^{*}(2460)^- \ell^+ \nu_\ell, D_2^{*-} \rightarrow \bar{D}^0 \pi^-) / \Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.41 ± 0.20 OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.		


$1.57 \pm 0.15 \pm 0.05$ MEIER 23 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$1.10 \pm 0.17 \pm 0.08$ ¹ AUBERT 09Y BABR $e^+ e^- \rightarrow \Upsilon(4S)$

$2.2 \pm 0.4 \pm 0.4$ ² LIVENTSEV 08 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a simultaneous fit of all B semileptonic decays without full reconstruction of events. AUBERT 09Y reports $B(B^0 \rightarrow \bar{D}_2^{*}(2460)^- \ell^+ \nu_\ell) \cdot B(\bar{D}_2^{*}(2460)^- \rightarrow \bar{D}^{(*)0} \pi^-) = (1.77 \pm 0.26 \pm 0.11) \times 10^{-3}$ and the authors have provided us the individual measurement.

² Uses a fully reconstructed B meson as a tag on the recoil side.



$\Gamma(\bar{D}^{*0} \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}$					Γ_{15} / Γ
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT		
5.44 ± 0.28 OUR AVERAGE					
$5.51 \pm 0.24 \pm 0.17$	MEIER	23	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	
$4.8 \pm 0.8 \pm 0.4$	¹ AUBERT	08Q	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$6.46 \pm 0.53 \pm 0.52$	VOSSSEN	18	BELL	Repl. by MEIER 23	
$5.5 \pm 2.2 \pm 0.2$	^{1,2} LIVENTSEV	08	BELL	Repl. by VOSSSEN 18	
$5.3 \pm 1.2 \pm 0.1$	^{3,4} LIVENTSEV	05	BELL	Repl. by LIVENTSEV 08	

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

² LIVENTSEV 08 reports $(5.6 \pm 2.1 \pm 0.8) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow \bar{D}^{*0} \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \ell^+ \nu_\ell)]$ assuming $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.12 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(B^0 \rightarrow D^- \ell^+ \nu_\ell) = (2.10 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Excludes D^{*+} contribution to $D \pi$ modes.

⁴ LIVENTSEV 05 reports $[\Gamma(B^0 \rightarrow \bar{D}^{*0} \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}] / [B(B^+ \rightarrow \bar{D}^{*0} (2007)^0 \ell^+ \nu_\ell)] = 0.10 \pm 0.02 \pm 0.01$ which we multiply by our best value $B(B^+ \rightarrow \bar{D}^{*0} (2007)^0 \ell^+ \nu_\ell) = (5.26 \pm 0.10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D_1(2420)^- \ell^+ \nu_\ell, D_1^- \rightarrow \bar{D}^{*0} \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{16} / \Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.85$\pm$0.25 OUR AVERAGE			
3.06 $\pm$ 0.50 $\pm$ 0.28	MEIER	23	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
2.78 $\pm$ 0.24 $\pm$ 0.25	¹ AUBERT	09Y	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
2.7 $\pm$ 0.4 $\pm$ 0.3	² AUBERT	08BL	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
5.4 $\pm$ 1.9 $\pm$ 0.9	² LIVENTSEV	08	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a simultaneous measurement of all B semileptonic decays without full reconstruction of events.

² Uses a fully reconstructed B meson as a tag on the recoil side.

$$\Gamma(D_1(2420)^- \ell^+ \nu_\ell, D_1^- \rightarrow D^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{17} / \Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.02$\pm$0.13$\pm$0.09	MEIER	23	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$$\Gamma(D_1'(2430)^- \ell^+ \nu_\ell, D_1'^- \rightarrow \bar{D}^{*0} \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{18} / \Gamma$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
2.5 $\pm$0.6 OUR AVERAGE				
2.06 $\pm$ 0.68 $\pm$ 0.25		MEIER	23	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
3.1 $\pm$ 0.7 $\pm$ 0.5		¹ AUBERT	08BL	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.0 90 ¹ LIVENTSEV 08 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a fully reconstructed B meson as a tag on the recoil side.

$$\Gamma(D_2^*(2460)^- \ell^+ \nu_\ell, D_2^{*-} \rightarrow \bar{D}^{*0} \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{19} / \Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
6.6$\pm$1.1 OUR AVERAGE				
5.1 $\pm$ 4.0 $\pm$ 1.0		MEIER	23	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
6.7 $\pm$ 1.2 $\pm$ 0.5		¹ AUBERT	09Y	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
7 $\pm$ 2 $\pm$ 2		² AUBERT	08BL	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30 90 ² LIVENTSEV 08 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a simultaneous fit of all B semileptonic decays without full reconstruction of events.

AUBERT 09Y reports $B(B^0 \rightarrow \bar{D}_2^*(2460)^- \ell^+ \nu_\ell) \cdot B(\bar{D}_2^*(2460)^- \rightarrow \bar{D}^{(*)0} \pi^-) = (1.77 \pm 0.26 \pm 0.11) \times 10^{-3}$ and the authors have provided us the individual measurement.

² Uses a fully reconstructed B meson as a tag on the recoil side.

$$\Gamma(D^- \pi^+ \pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}} \quad \Gamma_{20} / \Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.45$\pm$0.18$\pm$0.13	MEIER	23	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$$\Gamma(D^- \pi^+ \pi^- \ell^+ \nu_\ell) / \Gamma(D^- \ell^+ \nu_\ell) \quad \Gamma_{20} / \Gamma_5$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
5.8$\pm$1.8$\pm$1.2	¹ LEES	16	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measurement used electrons and muons as leptons.

$$\Gamma(D^{*-}\pi^+\pi^-\ell^+\nu_\ell)/\Gamma_{\text{total}} \quad \Gamma_{21}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$5.1 \pm 2.1 \pm 0.9$	MEIER 23	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

$$\Gamma(D^{*-}\pi^+\pi^-\ell^+\nu_\ell)/\Gamma(D^{*(2010)-}\ell^+\nu_\ell) \quad \Gamma_{21}/\Gamma_7$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 0.8 \pm 0.6$	¹ LEES 16	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Measurement used electrons and muons as leptons.

$$\Gamma(\rho^-\ell^+\nu_\ell)/\Gamma_{\text{total}} \quad \Gamma_{22}/\Gamma$$

$\ell = e$ or μ , not sum over e and μ modes.

“OUR EVALUATION” includes both B^0 and B^+ decays. The average assumes equality of the semileptonic decay width for these isospin conjugate states.

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$2.94 \pm 0.11 \pm 0.18$ OUR EVALUATION (Produced by HFLAV)

2.45 ± 0.32 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

$3.22 \pm 0.27 \pm 0.24$	¹ SIBIDANOV 13	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$1.75 \pm 0.15 \pm 0.27$	² DEL-AMO-SA..11c	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$2.93 \pm 0.37 \pm 0.37$	³ ADAM 07	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$2.17 \pm 0.54 \pm 0.32$	⁴ HOKUUE 07	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$2.57 \pm 0.29^{+0.53}_{-0.62}$	⁵ BEHRENS 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.14 \pm 0.21 \pm 0.56$	² AUBERT,B 050	BABR	Repl. by DEL-AMO-SANCHEZ 11c
$2.17 \pm 0.34^{+0.62}_{-0.68}$	⁶ ATHAR 03	CLE2	Repl. by ADAM 07
$3.29 \pm 0.42 \pm 0.72$	⁷ AUBERT 03E	BABR	Repl. by AUBERT,B 050
$2.69 \pm 0.41^{+0.61}_{-0.64}$	⁸ BEHRENS 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$2.5 \pm 0.4^{+0.7}_{-0.9}$	⁹ ALEXANDER 96T	CLE2	Repl. by BEHRENS 00
<4.1	90 ¹⁰ BEAN 93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ The signal events are tagged by a second B meson reconstructed in the fully hadronic decays.

² B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

³ The B^0 and B^+ results are combined assuming the isospin, B lifetimes, and relative charged/neutral B production at the $\Upsilon(4S)$.


⁴ The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \rightarrow D^{(*)}\ell\nu_\ell$.

⁵ Averaging with ALEXANDER 96T results including experimental and theoretical correlations considered, BEHRENS 00 reports systematic errors $^{+0.33}_{-0.46} \pm 0.41$, where the second error is theoretical model dependence. We combine these in quadrature.

⁶ ATHAR 03 reports systematic errors $^{+0.47}_{-0.50} \pm 0.41 \pm 0.01$, which are experimental systematic, systematic due to residual form-factor uncertainties in the signal, and systematic due to residual form-factor uncertainties in the cross-feed modes, respectively. We combine these in quadrature.

⁷ Uses isospin constraints and extrapolation to all electron energies according to five different form-factor calculations. The second error combines the systematic and theoretical uncertainties in quadrature.

- ⁸ BEHRENS 00 reports $+0.35_{-0.40} \pm 0.50$, where the second error is the theoretical model dependence. We combine these in quadrature. B^+ and B^0 decays combined using isospin symmetry: $\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu) \approx 2\Gamma(B^+ \rightarrow \omega \ell^+ \nu)$. No evidence for $\omega \ell \nu$ is reported.
- ⁹ ALEXANDER 96T reports $+0.5_{-0.7} \pm 0.5$ where the second error is the theoretical model dependence. We combine these in quadrature. B^+ and B^0 decays combined using isospin symmetry: $\Gamma(B^0 \rightarrow \rho^- \ell^+ \nu) = 2\Gamma(B^+ \rightarrow \rho^0 \ell^+ \nu) \approx 2\Gamma(B^+ \rightarrow \omega \ell^+ \nu)$. No evidence for $\omega \ell \nu$ is reported.
- ¹⁰ BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\omega \ell^+ \nu_\ell)$ with this result, they obtain a limit $<(1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow (\omega \text{ or } \rho^0) \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.08-0.13$ at 90% CL is derived as well.



$\Gamma(\pi^- \ell^+ \nu_\ell) / \Gamma_{\text{total}}$					Γ_{23} / Γ
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT		
1.50 ± 0.05 OUR EVALUATION	(Produced by HFLAV)				
1.46 ± 0.04 OUR AVERAGE					
1.49 ± 0.09 ± 0.07	¹ SIBIDANOV	13	BELL	$e^+ e^- \rightarrow \gamma(4S)$	
1.47 ± 0.05 ± 0.06	^{2,3} LEES	12AA	BABR	$e^+ e^- \rightarrow \gamma(4S)$	
1.41 ± 0.05 ± 0.07	⁴ DEL-AMO-SA...	11C	BABR	$e^+ e^- \rightarrow \gamma(4S)$	
1.49 ± 0.04 ± 0.07	² HA	11	BELL	$e^+ e^- \rightarrow \gamma(4S)$	
1.54 ± 0.17 ± 0.09	⁴ AUBERT	08AV	BABR	$e^+ e^- \rightarrow \gamma(4S)$	
1.37 ± 0.15 ± 0.11	^{5,6} ADAM	07	CLE2	$e^+ e^- \rightarrow \gamma(4S)$	
1.38 ± 0.19 ± 0.14	⁷ HOKUUE	07	BELL	$e^+ e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.53±0.18±0.12	^{1,8} CAO	23	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
1.42±0.05±0.08	² DEL-AMO-SA...	11F	BABR	Repl. by LEES 12AA
1.46±0.07±0.08	⁹ AUBERT	07J	BABR	Repl. by DEL-AMO-SANCHEZ 11F
1.33±0.17±0.11	¹⁰ AUBERT,B	06K	BABR	Repl. by AUBERT 08AV
1.38±0.10±0.18	¹¹ AUBERT,B	05O	BABR	Repl. by DEL-AMO-SANCHEZ 11C
1.33±0.18±0.13	¹² ATHAR	03	CLE2	Repl. by ADAM 07
1.8 ±0.4 ±0.4	¹³ ALEXANDER	96T	CLE2	Repl. by ATHAR 03

¹ The signal events are tagged by a second B meson reconstructed in the fully hadronic decays.

² Uses loose neutrino reconstruction technique. Assumes $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.6 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = (48.4 \pm 0.6)\%$.

³ Reports also a branching fraction value $B(B^0 \rightarrow \pi^- \ell^+ \nu) = (1.45 \pm 0.04 \pm 0.06) \times 10^{-4}$ from the decays of B^+ and B^0 that are combined using the isospin symmetry relation.

⁴ Using the isospin symmetry relation, B^+ and B^0 branching fractions are combined.

⁵ The B^0 and B^+ results are combined assuming the isospin, B lifetimes, and relative charged/neutral B production at the $\Upsilon(4S)$.

⁶ Also report the rate for $q^2 > 16 \text{ GeV}^2$ of $(0.41 \pm 0.08 \pm 0.04) \times 10^{-4}$ from which they obtain $|V_{ub}| = 3.6 \pm 0.4 \pm 0.2^{+0.6}_{-0.4}$ (last error is from theory).

⁷ The signal events are tagged by a second B meson reconstructed in the semileptonic mode $B \rightarrow D^{(*)} \ell \nu_\ell$.

⁸ This analysis provides the inclusive and exclusive measurement of $|V_{ub}|$.

⁹ The analysis uses events in which the signal B decays are reconstructed with an innovative loose neutrino reconstruction technique.

¹⁰ The signals are tagged by a second B meson reconstructed in a semileptonic or hadronic decay. The B^0 and B^+ results are combined assuming the isospin symmetry.

¹¹ B^+ and B^0 decays combined assuming isospin symmetry. Systematic errors include both experimental and form-factor uncertainties.

¹² ATHAR 03 reports systematic errors $0.11 \pm 0.01 \pm 0.07$, which are experimental systematic, systematic due to residual form-factor uncertainties in the signal, and systematic due to residual form-factor uncertainties in the cross-feed modes, respectively. We combine these in quadrature.

¹³ ALEXANDER 96T gives systematic errors $\pm 0.3 \pm 0.2$ where the second error reflects the estimated model dependence. We combine these in quadrature. Assumes isospin symmetry: $\Gamma(B^0 \rightarrow \pi^- \ell^+ \nu) = 2 \times \Gamma(B^+ \rightarrow \pi^0 \ell^+ \nu)$.

$\Gamma(\pi^- \mu^+ \nu_\mu)/\Gamma_{\text{total}}$

Γ_{24}/Γ

VALUE	DOCUMENT ID	TECN
-------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ¹ ALBRECHT 91C ARG

¹ In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(\pi^- \tau^+ \nu_\tau)/\Gamma_{\text{total}}$

Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-4}$	90	¹ HAMER	16	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^\pm X)/\Gamma_{\text{total}}$				Γ_{26}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.78 ± 0.08	¹ ALBRECHT	96D	ARG	$e^+e^- \rightarrow \gamma(4S)$
¹ Average multiplicity.				

$\Gamma(D^0 X)/\Gamma_{\text{total}}$				Γ_{27}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.081 \pm 0.014 \pm 0.005$	¹ AUBERT	07N	BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.063 \pm 0.019 \pm 0.005$	¹ AUBERT,BE	04B	BABR	Repl. by AUBERT 07N
¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.				

$\Gamma(\bar{D}^0 X)/\Gamma_{\text{total}}$				Γ_{28}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.474 \pm 0.020^{+0.020}_{-0.019}$	¹ AUBERT	07N	BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.511 \pm 0.031 \pm 0.028$	¹ AUBERT,BE	04B	BABR	Repl. by AUBERT 07N
¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.				

$\Gamma(D^0 X)/[\Gamma(D^0 X) + \Gamma(\bar{D}^0 X)]$				$\Gamma_{27}/(\Gamma_{27} + \Gamma_{28})$
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.146 \pm 0.022 \pm 0.006$	AUBERT	07N	BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.110 \pm 0.031 \pm 0.008$	AUBERT,BE	04B	BABR	Repl. by AUBERT 07N

$\Gamma(D^+ X)/\Gamma_{\text{total}}$				Γ_{29}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.039	90	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.051	90	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N
¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.				

$\Gamma(D^- X)/\Gamma_{\text{total}}$				Γ_{30}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.369 \pm 0.016^{+0.030}_{-0.027}$	¹ AUBERT	07N	BABR	$e^+e^- \rightarrow \gamma(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.397 \pm 0.030^{+0.040}_{-0.038}$	¹ AUBERT,BE	04B	BABR	Repl. by AUBERT 07N
¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.				

$$\Gamma(D^+ X)/[\Gamma(D^+ X) + \Gamma(D^- X)] \quad \Gamma_{29}/(\Gamma_{29} + \Gamma_{30})$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.058 \pm 0.028 \pm 0.006$	AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.055 \pm 0.040 \pm 0.006$	AUBERT,BE	04B	BABR Repl. by AUBERT 07N

$$\Gamma(D_s^+ X)/\Gamma_{\text{total}} \quad \Gamma_{31}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.103 \pm 0.012^{+0.017}_{-0.014}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.109 \pm 0.021^{+0.039}_{-0.024}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$$\Gamma(D_s^- X)/\Gamma_{\text{total}} \quad \Gamma_{32}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.026	90	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.087	90	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$$\Gamma(D_s^+ X)/[\Gamma(D_s^+ X) + \Gamma(D_s^- X)] \quad \Gamma_{31}/(\Gamma_{31} + \Gamma_{32})$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.879 \pm 0.066 \pm 0.005$	AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.733 \pm 0.092 \pm 0.010$	AUBERT,BE	04B	BABR Repl. by AUBERT 07N

$$\Gamma(\Lambda_c^+ X)/\Gamma_{\text{total}} \quad \Gamma_{33}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.031	90	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.038	90	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$$\Gamma(\bar{\Lambda}_c^- X)/\Gamma_{\text{total}} \quad \Gamma_{34}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.05 \pm 0.010^{+0.019}_{-0.011}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.049 \pm 0.017^{+0.018}_{-0.011}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$$\Gamma(\Lambda_c^+ X) / [\Gamma(\Lambda_c^+ X) + \Gamma(\bar{\Lambda}_c^- X)] \quad \Gamma_{33}/(\Gamma_{33}+\Gamma_{34})$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.243^{+0.119}_{-0.121} \pm 0.003$	AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.286 \pm 0.142 \pm 0.007$	AUBERT,BE	04B	BABR Repl. by AUBERT 07N

$$\Gamma(\bar{c} X) / \Gamma_{\text{total}} \quad \Gamma_{35}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.947 \pm 0.030^{+0.045}_{-0.040}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.039 \pm 0.051^{+0.063}_{-0.058}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$$\Gamma(c X) / \Gamma_{\text{total}} \quad \Gamma_{36}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.246 \pm 0.024^{+0.021}_{-0.017}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.237 \pm 0.036^{+0.041}_{-0.027}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$$\Gamma(\bar{c}/c X) / \Gamma_{\text{total}} \quad \Gamma_{37}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.193 \pm 0.030^{+0.053}_{-0.049}$	¹ AUBERT	07N	BABR $e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.276 \pm 0.062^{+0.088}_{-0.074}$	¹ AUBERT,BE	04B	BABR Repl. by AUBERT 07N

¹ Events are selected by completely reconstructing one B and searching for a reconstructed charmed particle in the rest of the event. The last error includes systematic and charm branching ratio uncertainties.

$$\Gamma(D^- \pi^+) / \Gamma_{\text{total}} \quad \Gamma_{38}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.51 ± 0.08 OUR FIT				
2.56 ± 0.08 OUR AVERAGE				
$2.48 \pm 0.01 \pm 0.10$		WAHEED	22	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$2.55 \pm 0.05 \pm 0.16$		¹ AUBERT	07H	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$3.03 \pm 0.23 \pm 0.23$		² AUBERT,BE	06J	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$2.68 \pm 0.12 \pm 0.24$		^{1,3} AHMED	02B	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
$2.7 \pm 0.6 \pm 0.5$		⁴ BORTOLETTO	092	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
$4.8 \pm 1.1 \pm 1.1$	22	⁵ ALBRECHT	90J	ARG $e^+e^- \rightarrow \Upsilon(4S)$
$5.1^{+2.8}_{-2.5} \pm 1.3_{-1.2}$	4	⁶ BEBEK	87	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.73 \pm 0.19 \pm 0.05$	^{1,7}	AUBERT,B	040	BABR	Repl. by AUBERT 07H
$2.83 \pm 0.42 \pm 0.05$	81	⁸ ALAM	94	CLE2	Repl. by AHMED 02B
$3.1 \pm 1.3 \pm 1.0$	7	⁵ ALBRECHT	88K	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

³ AHMED 02B reports an additional uncertainty on the branching ratios to account for 4.5% uncertainty on relative production of B^0 and B^+ , which is not included here.

⁴ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁵ ALBRECHT 88K assumes $B^0\bar{B}^0:B^+B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

⁶ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

⁷ AUBERT,B 040 reports $[\Gamma(B^0 \rightarrow D^- \pi^+)/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K_S^0 \pi^+)] = (42.7 \pm 2.1 \pm 2.2) \times 10^{-6}$ which we divide by our best value $B(D^+ \rightarrow K_S^0 \pi^+) = (1.561 \pm 0.031) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸ ALAM 94 reports $[\Gamma(B^0 \rightarrow D^- \pi^+)/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = (0.265 \pm 0.032 \pm 0.023) \times 10^{-3}$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- \ell^+ \nu_\ell)/\Gamma(D^- \pi^+)$

Γ_5/Γ_{38}

VALUE	DOCUMENT ID	TECN	COMMENT
$9.9 \pm 1.0 \pm 0.9$	AALTONEN	09E	CDF $p\bar{p}$ at 1.96 TeV

$\Gamma(D^- \rho^+)/\Gamma_{\text{total}}$

Γ_{39}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0076 ± 0.0012 OUR AVERAGE				
$0.0075 \pm 0.0013 \pm 0.0001$	79	¹ ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
$0.009 \pm 0.005 \pm 0.003$	9	² ALBRECHT	90J	ARG $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.022 \pm 0.012 \pm 0.009$	6	² ALBRECHT	88K	ARG $e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	---	-----------------------	-----	---------------------------------------

¹ ALAM 94 reports $[\Gamma(B^0 \rightarrow D^- \rho^+)/\Gamma_{\text{total}}] \times [B(D^+ \rightarrow K^- 2\pi^+)] = 0.000704 \pm 0.000096 \pm 0.000070$ which we divide by our best value $B(D^+ \rightarrow K^- 2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 88K assumes $B^0\bar{B}^0:B^+B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

$\Gamma(D^- K^0 \pi^+)/\Gamma_{\text{total}}$

Γ_{40}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.9 \pm 0.7 \pm 0.5$	¹ AUBERT,BE	05B	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.5 ± 0.7 OUR AVERAGE			
$4.6 \pm 0.6 \pm 0.5$	¹ AUBERT, BE 05B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$3.7 \pm 1.5 \pm 1.0$	¹ MAHAPATRA 02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^- \omega \pi^+)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.0028 \pm 0.0005 \pm 0.0004$	¹ ALEXANDER 01B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

 $\Gamma(D^- K^+)/\Gamma(D^- \pi^+)$ Γ_{43}/Γ_{38}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
8.19 ± 0.20 OUR AVERAGE			
$8.19 \pm 0.20 \pm 0.23$	WAHEED 22	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$8.22 \pm 0.11 \pm 0.25$	AAIJ 13P	LHCB	pp at 7 TeV
$6.8 \pm 1.5 \pm 0.7$	ABE 01I	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(D^- K^+ \pi^+ \pi^-)/\Gamma(D^- \pi^+ \pi^+ \pi^-)$ Γ_{44}/Γ_{50}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$5.9 \pm 1.1 \pm 0.5$	AAIJ 12T	LHCB	pp at 7 TeV

 $\Gamma(D^- K^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$1.64 \pm 0.24 \pm 0.10$		¹ ADACHI 24L	BEL2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 3.1	90	² DRUTSKOY 02	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Used 362 fb⁻¹ Belle II data and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.484 \pm 0.012$.² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^- K^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
7.7 ± 0.6 OUR AVERAGE			
$7.56 \pm 0.45 \pm 0.38$	¹ ADACHI 24L	BEL2	$e^+e^- \rightarrow \Upsilon(4S)$
$8.8 \pm 1.1 \pm 1.5$	² DRUTSKOY 02	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Used 362 fb⁻¹ Belle II data and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.484 \pm 0.012$.² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(\bar{D}^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	CL%	EVS	DOCUMENT ID	TECN	COMMENT
8.8 ± 0.5 OUR AVERAGE					
$8.95 \pm 0.15 \pm 0.52$			¹ AAIIJ 15Y	LHCB	pp at 7, 8 TeV
$8.4 \pm 0.4 \pm 0.8$			² KUZMIN 07	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.0 ± 0.6 ± 1.5	2,3	SATPATHY	03	BELL	Repl. by KUZMIN 07
< 16	90	2 ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 70	90	4 BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 340	90	5 BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
700 ± 500	5	6 BEHREND	83	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ The second uncertainty combines in quadrature all systematic uncertainties quoted in the paper. AAIJ 15Y reports $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-) = (8.46 \pm 0.14 \pm 0.49) \times 10^{-4}$ in the kinematic region $m(\bar{D}^0 \pi^\pm) > 2.1$ GeV which we corrected to the full phase-space dividing by 0.945 from Belle.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ No assumption about the intermediate mechanism is made in the analysis.

⁴ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D . The product branching fraction into $D_0^*(2340)\pi$ followed by $D_0^*(2340) \rightarrow D^0 \pi$ is < 0.0001 at 90% CL and into $D_2^*(2460)$ followed by $D_2^*(2460) \rightarrow D^0 \pi$ is < 0.0004 at 90% CL.

⁵ BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%. $B(D^0 \rightarrow K^- \pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) = (9.1 \pm 0.8 \pm 0.8)\%$ were used.

⁶ Corrected by us using assumptions: $B(D^0 \rightarrow K^- \pi^+) = (0.042 \pm 0.006)$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 50\%$. The product branching ratio is $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-)B(\bar{D}^0 \rightarrow K^+ \pi^-) = (0.39 \pm 0.26) \times 10^{-2}$.

$\Gamma(D^*(2010)^-\pi^+)/\Gamma_{\text{total}}$			Γ_{48}/Γ		
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.66±0.07 OUR AVERAGE					
2.62±0.02±0.09		¹ KROHN	23	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
2.79±0.08±0.17		² AUBERT	07H	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
2.48±0.34±0.08		^{3,4} AUBERT,BE	06J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
2.81±0.24±0.05		⁵ BRANDENB...	98	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
2.6 ±0.3 ±0.4	82	⁶ ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
3.37±0.96±0.02		⁷ BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
2.36±0.88±0.02	12	⁸ ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
2.36 ^{+1.50} _{-1.10} ±0.02	5	⁹ BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10 ± 4 ± 1	8	10 AKERS	94J	OPAL	$e^+e^- \rightarrow Z$
2.7 ± 1.4 ± 1.0	5	11 ALBRECHT	87C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
3.5 ± 2 ± 2		12 ALBRECHT	86F	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
17 ± 5 ± 5	41	13 GILES	84	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ KROHN 23 reports $(2.623 \pm 0.016 \pm 0.086) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- \pi^+)/\Gamma_{\text{total}}] / [B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0)]$ assuming $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.486 \pm 0.006$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AUBERT, BE 06J reports $[\Gamma(B^0 \rightarrow D^*(2010)^- \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 0.99 \pm 0.11 \pm 0.08$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.51 \pm 0.08) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ⁴ Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.
- ⁵ BRANDENBURG 98 assume equal production of B^+ and B^0 at $\Upsilon(4S)$ and use the D^* reconstruction technique. The first error is their experiment's error and the second error is the systematic error from the PDG 96 value of $B(D^* \rightarrow D\pi)$.
- ⁶ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-2\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.
- ⁷ BORTOLETTO 92 reports $(4.0 \pm 1.0 \pm 0.7) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^-\pi^+)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0\pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .
- ⁸ ALBRECHT 90J reports $(2.8 \pm 0.9 \pm 0.6) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^-\pi^+)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0\pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .
- ⁹ BEBEK 87 reports $(2.8^{+1.5+1.0}_{-1.2-0.6}) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^-\pi^+)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0\pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92 and ALBRECHT 90J.
- ¹⁰ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and 38% B_d production fraction.
- ¹¹ ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.
- ¹² ALBRECHT 86F uses pseudomass that is independent of D^0 and D^+ branching ratios.
- ¹³ Assumes $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.60^{+0.08}_{-0.15}$. Assumes $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 0.40 \pm 0.02$ Does not depend on D branching ratios.

$\Gamma(D^*(2010)^-\ell^+\nu_\ell)/\Gamma(D^*(2010)^-\pi^+)$				Γ_7/Γ_{48}
VALUE	DOCUMENT ID	TECN	COMMENT	
$16.5 \pm 2.3 \pm 1.1$	AALTONEN	09E	CDF	$p\bar{p}$ at 1.96 TeV

$\Gamma(\bar{D}^0 K^+ K^-)/\Gamma(\bar{D}^0 \pi^+ \pi^-)$				Γ_{49}/Γ_{47}
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.069 \pm 0.004 \pm 0.003$	AAIJ	18AZ	LHCB	pp at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.056 \pm 0.011 \pm 0.007$	¹ AAIJ	12AMLHCB	pp at 7 TeV	

¹ Superseded by AAIJ 21Q.

$\Gamma(D^-\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{50}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0060 ± 0.0006 OUR FIT				
$0.0080 \pm 0.0021 \pm 0.0014$	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^- \pi^+ \pi^+ \pi^-)/\Gamma(D^- \pi^+)$				Γ_{50}/Γ_{38}
VALUE	DOCUMENT ID	TECN	COMMENT	
2.39 ± 0.23 OUR FIT				
2.38 ± 0.11 ± 0.21	AAIJ	11E	LHCB	pp at 7 TeV

$\Gamma((D^- \pi^+ \pi^+ \pi^-) \text{ nonresonant})/\Gamma_{\text{total}}$				Γ_{51}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0039 ± 0.0014 ± 0.0013	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .				

$\Gamma(D^- \pi^+ \rho^0)/\Gamma_{\text{total}}$				Γ_{52}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0011 ± 0.0009 ± 0.0004	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .				

$\Gamma(D^- a_1(1260)^+)/\Gamma_{\text{total}}$				Γ_{53}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0060 ± 0.0022 ± 0.0024	¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .				

$\Gamma(D^*(2010)^- \pi^+ \pi^0)/\Gamma_{\text{total}}$				Γ_{54}/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0152 ± 0.0052 ± 0.0001	51	¹ ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.015 ± 0.008 ± 0.008	8	² ALBRECHT	87C ARG	$e^+e^- \rightarrow \Upsilon(4S)$
¹ ALBRECHT 90J reports $0.018 \pm 0.004 \pm 0.005$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- \pi^+ \pi^0)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .				
² ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.				

$\Gamma(D^*(2010)^- \rho^+)/\Gamma_{\text{total}}$				Γ_{55}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.8 ± 0.9 OUR AVERAGE				
6.8 ± 0.3 ± 0.9		^{1,2} CSORNA	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
16.0 ± 11.3 ± 0.1		³ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
5.89 ± 3.52 ± 0.04	19	⁴ ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		^{2,5} MATVIENKO	15 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

7.4 ± 1.0 ± 1.4	76	^{6,7} ALAM	94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
81 ± 29 ⁺⁵⁹ ₋₂₄	19	⁸ CHEN	85	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ The second error combines the systematic and theoretical uncertainties in quadrature. CSORNA 03 includes data used in ALAM 94. A full angular fit to three complex helicity amplitudes is performed.

² Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$ resonance.

³ BORTOLETTO 92 reports $0.019 \pm 0.008 \pm 0.011$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^-\rho^+)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0\pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁴ ALBRECHT 90J reports $0.007 \pm 0.003 \pm 0.003$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^-\rho^+)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0\pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁵ MATVIENKO 15 reports $B(B^0 \rightarrow D^*(2010)^-\rho^+, \rho^+ \rightarrow \omega\pi^+) = (1.48 \pm 0.27^{+0.15+0.21}_{-0.09-0.56}) \times 10^{-3}$. The last uncertainty is a model one.

⁶ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

⁷ This decay is nearly completely longitudinally polarized, $\Gamma_L/\Gamma = (93 \pm 5 \pm 5)\%$, as expected from the factorization hypothesis (ROSNER 90). The nonresonant $\pi^+\pi^0$ contribution under the ρ^+ is less than 9% at 90% CL.

⁸ Uses $B(D^* \rightarrow D^0\pi^+) = 0.6 \pm 0.15$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 0.4$. Does not depend on D branching ratios.

$\Gamma(D^*(2010)^- K^+)/\Gamma_{\text{total}}$

Γ_{56}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.16 ± 0.08 OUR AVERAGE			
2.22 ± 0.06 ± 0.08	¹ KROHN	23	BELL $e^+e^- \rightarrow \Upsilon(4S)$
2.06 ± 0.12 ± 0.06	² AUBERT	06A	BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.4 ± 0.1	³ ABE	01I	BELL Repl. by KROHN 23
-----------------	------------------	-----	------------------------

¹ KROHN 23 reports $(2.221 \pm 0.063 \pm 0.077) \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- K^+)/\Gamma_{\text{total}}] / [B(\Upsilon(4S) \rightarrow B^0\bar{B}^0)]$ assuming $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 0.486 \pm 0.006$.

² AUBERT 06A reports $[\Gamma(B^0 \rightarrow D^*(2010)^- K^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^*(2010)^-\pi^+)] = 0.0776 \pm 0.0034 \pm 0.0029$ which we multiply by our best value $B(B^0 \rightarrow D^*(2010)^-\pi^+) = (2.66 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABE 01I reports $[\Gamma(B^0 \rightarrow D^*(2010)^- K^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^*(2010)^-\pi^+)] = 0.074 \pm 0.015 \pm 0.006$ which we multiply by our best value $B(B^0 \rightarrow D^*(2010)^-\pi^+) = (2.66 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^*(2010)^- K^+)/\Gamma(D^*(2010)^- \pi^+)$ Γ_{56}/Γ_{48}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
8.22$\pm$0.29 OUR AVERAGE	Error includes scale factor of 1.3.		
8.41 $\pm$ 0.24 $\pm$ 0.13	KROHN	23 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
7.76 $\pm$ 0.34 $\pm$ 0.26	AAIJ	13AO LHCB	pp at 7 TeV

 $\Gamma(D^*(2010)^- K^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.0$\pm$0.7$\pm$0.3	¹ AUBERT,BE	05B BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^*(2010)^- K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.3$\pm$0.6 OUR AVERAGE			
3.2 $\pm$ 0.6 $\pm$ 0.3	¹ AUBERT,BE	05B BABR	$e^+e^- \rightarrow \Upsilon(4S)$
3.8 $\pm$ 1.3 $\pm$ 0.8	² MAHAPATRA	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and an unpolarized final state. $\Gamma(D^*(2010)^- K^+ \bar{K}^0)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.82$\pm$0.38$\pm$0.10		¹ ADACHI	24L BEL2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<4.7	90	² DRUTSKOY	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Used 362 fb⁻¹ Belle II data and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.484 \pm 0.012$.² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^*(2010)^- K^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
13.1 $\pm$1.3 OUR AVERAGE			
13.12 $\pm$ 1.21 $\pm$ 0.71	¹ ADACHI	24L BEL2	$e^+e^- \rightarrow \Upsilon(4S)$
12.9 $\pm$ 2.2 $\pm$ 2.5	² DRUTSKOY	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Used 362 fb⁻¹ Belle II data and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.484 \pm 0.012$.² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
7.21$\pm$0.29 OUR AVERAGE				
7.26 $\pm$ 0.11 $\pm$ 0.31		¹ LEES	16H BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.81 $\pm$ 0.23 $\pm$ 0.72		² MAJUMDER	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
6.3 $\pm$ 1.0 $\pm$ 1.1		^{3,4} ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
13.4 $\pm$ 3.6 $\pm$ 0.1		⁵ BORTOLETTO	092 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
10.1 $\pm$ 4.1 $\pm$ 0.1		⁶ ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 $\pm$ 9 $\pm$ 16		⁷ ALBRECHT	87C ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<42	90	⁸ BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.486 \pm 0.006$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁴ The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+ , the branching ratio for $\bar{D}^{*-} a_1^+$ is twice that for $\bar{D}^{*-} \pi^+ \pi^+ \pi^-$.)

⁵ BORTOLETTO 92 reports $0.0159 \pm 0.0028 \pm 0.0037$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁶ ALBRECHT 90J reports $0.012 \pm 0.003 \pm 0.004$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁷ ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

⁸ BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

$\Gamma((D^*(2010)^- \pi^+ \pi^+ \pi^-) \text{ nonresonant})/\Gamma_{\text{total}}$				Γ_{62}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0000$\pm$0.0019$\pm$0.0016	¹ BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

¹ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

$\Gamma(D^*(2010)^- \pi^+ \rho^0)/\Gamma_{\text{total}}$				Γ_{63}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.00573$\pm$0.00317$\pm$0.00004	¹ BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

¹ BORTOLETTO 92 reports $0.0068 \pm 0.0032 \pm 0.0021$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- \pi^+ \rho^0)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^*(2010)^- a_1(1260)^+)/\Gamma_{\text{total}}$				Γ_{64}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0130$\pm$0.0027 OUR AVERAGE				
0.0126 $\pm$ 0.0020 $\pm$ 0.0022	^{1,2} ALAM 94	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	
0.0152 $\pm$ 0.0070 $\pm$ 0.0001	³ BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

¹ ALAM 94 value is twice their $\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ value based on their observation that the three pions are dominantly in the $a_1(1260)$ mass range 1.0 to 1.6 GeV.

² ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

³ BORTOLETTO 92 reports $0.018 \pm 0.006 \pm 0.006$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- a_1(1260)^+)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\overline{D}_1(2420)^0 \pi^- \pi^+, \overline{D}_1^0 \rightarrow D^{*-} \pi^+)/\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-)$		Γ_{65}/Γ_{61}		
VALUE	DOCUMENT ID	TECN	COMMENT	
$(2.04 \pm 0.42 \pm 0.22) \times 10^{-2}$	AAIJ	13AO LHCB	pp at 7 TeV	

$\Gamma(D^*(2010)^- K^+ \pi^- \pi^+)/\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-)$		Γ_{66}/Γ_{61}		
VALUE	DOCUMENT ID	TECN	COMMENT	
$(6.47 \pm 0.37 \pm 0.35) \times 10^{-2}$	AAIJ	13AO LHCB	pp at 7 TeV	

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$		Γ_{67}/Γ		
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0176 ± 0.0027 OUR AVERAGE				
0.0172 ± 0.0014 ± 0.0024		¹ ALEXANDER 01B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0345 ± 0.0181 ± 0.0003	28	² ALBRECHT 90J	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The signal is consistent with all observed $\omega \pi^+$ having proceeded through the ρ'^+ resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

² ALBRECHT 90J reports $0.041 \pm 0.015 \pm 0.016$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(D^*(2010)^+ \rightarrow D^0 \pi^+)]$ assuming $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$, which we rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (67.7 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^{*-} 3\pi^+ 2\pi^-)/\Gamma_{\text{total}}$		Γ_{68}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	
4.72 ± 0.59 ± 0.71	¹ MAJUMDER 04	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$	

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^-\omega\pi^+)/\Gamma_{\text{total}}$	Γ_{69}/Γ			
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.46±0.18 OUR AVERAGE	Error includes scale factor of 1.2.			
2.31±0.11±0.14	¹ MATVIENKO	15	BELL	$e^+e^-\rightarrow\Upsilon(4S)$
2.88±0.21±0.31	¹ AUBERT	06L	BABR	$e^+e^-\rightarrow\Upsilon(4S)$
2.9 ±0.3 ±0.4	^{1,2} ALEXANDER	01B	CLE2	$e^+e^-\rightarrow\Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The signal is consistent with all observed $\omega\pi^+$ having proceeded through the ρ'^{+} resonance at mass $1349 \pm 25^{+10}_{-5}$ MeV and width $547 \pm 86^{+46}_{-45}$ MeV.

$\Gamma(\overline{D}_1(2430)^0\omega, \overline{D}_1^0 \rightarrow D^{*-}\pi^+)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.7^{+0.8}_{-0.4}$ OUR AVERAGE			

$2.5 \pm 0.4^{+0.8}_{-0.2}$ ^{1,2} MATVIENKO 15 BELL $e^+e^- \rightarrow \Upsilon(4S)$

$4.1 \pm 1.2 \pm 1.1$ ³ AUBERT 06L BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 .

² The measurement is obtained by amplitude analysis of $B^0 \rightarrow D^{*-}\omega\pi^+$. The second uncertainty combines in quadrature experimental systematic and model uncertainties.

³ Obtained by fitting the events with $\cos\theta_{D^*} < 0.5$ and scaling up the result by a factor of 4/3. No interference effects between $B^0 \rightarrow D_1'\omega$ and $D^*\omega\pi$ are assumed.

$\Gamma(D^{*-}\rho(1450)^+, \rho^+ \rightarrow \omega\pi^+)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.07^{+0.15+0.40}_{-0.31-0.13}$	^{1,2} MATVIENKO 15 BELL		$e^+e^- \rightarrow \Upsilon(4S)$

¹ Obtained by amplitude analysis of $\overline{B}^0 \rightarrow D^{*-}\omega\pi^+$. The second uncertainty combines in quadrature experimental systematic and model uncertainties.

² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.

$\Gamma(\overline{D}_1(2420)^0\omega, \overline{D}_1^0 \rightarrow D^{*-}\pi^+)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$0.7 \pm 0.2 \pm 0.1$	^{1,2} MATVIENKO 15 BELL		$e^+e^- \rightarrow \Upsilon(4S)$

¹ Obtained by amplitude analysis of $\overline{B}^0 \rightarrow D^{*-}\omega\pi^+$. The second uncertainty combines in quadrature experimental systematic and model uncertainties.

² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.

$\Gamma(\overline{D}_2^*(2460)^0\omega, \overline{D}_2^0 \rightarrow D^{*-}\pi^+)/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$0.4 \pm 0.1 \pm 0.1$	^{1,2} MATVIENKO 15 BELL		$e^+e^- \rightarrow \Upsilon(4S)$

¹ Obtained by amplitude analysis of $\overline{B}^0 \rightarrow D^{*-}\omega\pi^+$. The second uncertainty combines in quadrature experimental systematic and model uncertainties.

² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.

$\Gamma(D^{*-}b_1(1235)^+, b_1^+ \rightarrow \omega\pi^+)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.7 \times 10^{-4}$	90	¹ MATVIENKO 15 BELL		$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.

$\Gamma(\overline{D}^{*-}\pi^+)/\Gamma_{\text{total}}$ Γ_{75}/Γ

$\overline{D}^{*-}$ represents an excited state with mass $2.2 < M < 2.8$ GeV/ c^2 .

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.9 \pm 0.1$	^{1,2} AUBERT,BE 06J BABR		$e^+e^- \rightarrow \Upsilon(4S)$

¹ AUBERT, BE 06J reports $[\Gamma(B^0 \rightarrow \bar{D}^{*-} \pi^+)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = 0.77 \pm 0.22 \pm 0.29$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.51 \pm 0.08) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Uses a missing-mass method. Does not depend on D branching fractions or B^+/B^0 production rates.

$\Gamma(D_1(2420)^- \pi^+, D_1^- \rightarrow D^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$0.99^{+0.20}_{-0.25}$ OUR FIT			

$0.89 \pm 0.15^{+0.17}_{-0.32}$	¹ ABE	05A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---	------------------	-----	------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_1(2420)^- \pi^+, D_1^- \rightarrow D^- \pi^+ \pi^-)/\Gamma(D^- \pi^+ \pi^+ \pi^-)$ Γ_{76}/Γ_{50}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR FIT			
$2.1 \pm 0.5^{+0.3}_{-0.5}$	AAIJ	11E	LHCB pp at 7 TeV

$\Gamma(D_1(2420)^- \pi^+, D_1^- \rightarrow D^{*-} \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.33	90	¹ ABE	05A	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^-)/\Gamma(D^*(2010)^- \pi^+)$ Γ_{61}/Γ_{48}

VALUE	DOCUMENT ID	TECN	COMMENT
$2.64 \pm 0.04 \pm 0.13$	AAIJ	13A0	LHCB pp at 7 TeV

$\Gamma(\bar{D}_2^*(2460)^- \pi^+, D_2^{*-} \rightarrow D^0 \pi^-)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
2.38 ± 0.16 OUR AVERAGE				
$2.44 \pm 0.07 \pm 0.16$		¹ AAIJ	15Y	LHCB pp at 7, 8 TeV
$2.15 \pm 0.17 \pm 0.31$		^{2,3} KUZMIN	07	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14.7	90	² ALAM	94	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	-------------------	----	---

¹ Result obtained using the isobar formalism. The second uncertainty combines in quadrature all systematic uncertainties quoted in the paper.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Our second uncertainty combines systematics and model errors quoted in the paper.

$\Gamma(\bar{D}_0^*(2400)^- \pi^+, D_0^{*-} \rightarrow D^0 \pi^-)/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.76 ± 0.08 OUR AVERAGE			
$0.77 \pm 0.05 \pm 0.06$	¹ AAIJ	15Y	LHCB pp at 7, 8 TeV
$0.60 \pm 0.13 \pm 0.27$	^{2,3} KUZMIN	07	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Result obtained using the isobar formalism. The second uncertainty combines in quadrature all systematic uncertainties quoted in the paper.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Our second uncertainty combines systematics and model errors quoted in the paper.

$\Gamma(D_2^{*-}(2460)^-\pi^+, D_2^{*-} \rightarrow D^{*-}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	¹ ABE	05A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(\bar{D}_2^{*-}(2460)^-\rho^+)/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0049	90	¹ ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ ALAM 94 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and $B(D_2^{*-}(2460)^+ \rightarrow D^0\pi^+) = 30\%$. $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$0.14 \pm 0.06 \pm 0.03$		¹ AAIJ	13AP	LHCB pp at 7 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.43	90	² ADACHI	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$
-------	----	---------------------	----	--

<0.6	90	² AUBERT,B	06A	BABR $e^+e^- \rightarrow \Upsilon(4S)$
------	----	-----------------------	-----	--

¹ Uses $B(B^0 \rightarrow D^-D^+) = (2.11 \pm 0.31) \times 10^{-4}$ and $B(B^+ \rightarrow \bar{D}^0D_s^+) = (10.1 \pm 1.7) \times 10^{-3}$.² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^{*0}\bar{D}^0)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	¹ AUBERT,B	06A	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^-D^+)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
2.11 ± 0.18 OUR AVERAGE				
$2.12 \pm 0.16 \pm 0.18$		¹ ROHRKEN	12	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$1.97 \pm 0.20 \pm 0.20$		¹ FRATINA	07	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$2.8 \pm 0.4 \pm 0.5$		¹ AUBERT,B	06A	BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.91 \pm 0.51 \pm 0.30$		¹ MAJUMDER	05	BELL Repl. by FRATINA 07
--------------------------	--	-----------------------	----	--------------------------

< 9.4	90	¹ LIPELES	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
-------	----	----------------------	----	--

<59	90	BARATE	98Q	ALEP $e^+e^- \rightarrow Z$
-----	----	--------	-----	-----------------------------

<12	90	ASNER	97	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
-----	----	-------	----	--

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^\pm D^{*\mp} (CP\text{-averaged}))/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$6.14 \pm 0.29 \pm 0.50$	¹ ROHRKEN	12	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^-D_s^+)/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.1.			
$8.9 \pm 0.5 \pm 0.5$		¹ ADACHI	24L	BEL2 $e^+e^- \rightarrow \Upsilon(4S)$

$7.3 \pm 0.4 \pm 0.7$	² ZUPANC	07	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$6.6 \pm 1.4 \pm 0.6$	³ AUBERT	06N	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$6.8 \pm 2.4 \pm 0.6$	⁴ GIBAUT	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$10 \pm 9 \pm 1$	⁵ ALBRECHT	92G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$5.3 \pm 3.0 \pm 0.5$	⁶ BORTOLETTO92		CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

12 ± 7	³	⁷ BORTOLETTO90	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
------------	--------------	---------------------------	------	-----------------------------------

¹ Used 362 fb^{-1} Belle II data and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.484 \pm 0.012$.

² ZUPANC 07 reports $(7.5 \pm 0.2 \pm 1.1) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AUBERT 06N reports $(0.64 \pm 0.13 \pm 0.10) \times 10^{-2}$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.0462 \pm 0.0062$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ GIBAUT 96 reports $0.0087 \pm 0.0024 \pm 0.0020$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ ALBRECHT 92G reports $0.017 \pm 0.013 \pm 0.006$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., $B(D^+ \rightarrow K^- 2\pi^+) = 7.7 \pm 1.0\%$.

⁶ BORTOLETTO 92 reports $0.0080 \pm 0.0045 \pm 0.0030$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.030 \pm 0.011$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁷ BORTOLETTO 90 assume $B(D_s \rightarrow \phi \pi^+) = 2\%$. Superseded by BORTOLETTO 92.

$\Gamma(D^*(2010)^- D_s^+)/\Gamma_{\text{total}}$	Γ_{88}/Γ
VALUE (units 10^{-3})	EVTS
8.2 ± 0.8 OUR AVERAGE	

$8.3 \pm 1.0 \pm 0.6$	¹ ADACHI	24L	BEL2	$e^+e^- \rightarrow \Upsilon(4S)$
$7.3 \pm 1.3 \pm 0.7$	² AUBERT	06N	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$8.3 \pm 1.5 \pm 0.7$	³ AUBERT	03I	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$8.8 \pm 1.7 \pm 0.8$	⁴ AHMED	00B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$8 \pm 6 \pm 1$	⁵ ALBRECHT	92G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
$11 \pm 6 \pm 1$	⁶ BORTOLETTO92		CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.2 \pm 2.2 \pm 0.6$	⁷ GIBAUT	96	CLE2	Repl. by AHMED 00B
-----------------------	---------------------	----	------	--------------------

24 ± 14 3 ⁸ BORTOLETTO90 CLEO $e^+e^- \rightarrow \Upsilon(4S)$ ¹ Used 362 fb⁻¹ Belle II data and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.484 \pm 0.012$.² AUBERT 06N reports $(0.71 \pm 0.13 \pm 0.09) \times 10^{-2}$ from a measurement of $[\Gamma(B^0 \rightarrow D^{*+}(2010)^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.0462 \pm 0.0062$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.³ AUBERT 03I reports $0.0103 \pm 0.0014 \pm 0.0013$ from a measurement of $[\Gamma(B^0 \rightarrow D^{*+}(2010)^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.⁴ AHMED 00B reports $0.0110 \pm 0.0018 \pm 0.0011$ from a measurement of $[\Gamma(B^0 \rightarrow D^{*+}(2010)^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.⁵ ALBRECHT 92G reports $0.014 \pm 0.010 \pm 0.003$ from a measurement of $[\Gamma(B^0 \rightarrow D^{*+}(2010)^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^{*+}(2010)^+$ branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \rightarrow K^- 2\pi^+) = 7.1 \pm 1.0\%$, and $B(D^{*+}(2010)^+ \rightarrow D^0 \pi^+) = 55 \pm 4\%$.⁶ BORTOLETTO 92 reports $0.016 \pm 0.009 \pm 0.006$ from a measurement of $[\Gamma(B^0 \rightarrow D^{*+}(2010)^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.030 \pm 0.011$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^{*+}(2010)$.⁷ GIBAUT 96 reports $0.0093 \pm 0.0023 \pm 0.0016$ from a measurement of $[\Gamma(B^0 \rightarrow D^{*+}(2010)^- D_s^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.⁸ BORTOLETTO 90 assume $B(D_s \rightarrow \phi \pi^+) = 2\%$. Superseded by BORTOLETTO 92.

$\Gamma(D^{*+}(2010)^- D_s^+, D^{*-} \rightarrow \bar{D}^0 \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV}) / \Gamma(\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV})$ Γ_{89}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
15.8$\pm$1.3$\pm$2.4	¹ AAIJ	23B LHCb	pp at 7, 8, 13 TeV

¹ Uses simultaneous fits of $\bar{B}^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ amplitudes assuming isospin symmetry.

$$\Gamma(D_2^{*-}(2460) D_s^+, D_2^{*-} \rightarrow \bar{D}^0 \pi^-) / \Gamma(\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV})$$

Γ_{90}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$22.38 \pm 0.88 \pm 0.60$	¹ AAIJ	23B	LHCB pp at 7, 8, 13 TeV

¹ Uses simultaneous fits of $\bar{B}^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ amplitudes assuming isospin symmetry.

$$\Gamma(D_1^{*-}(2600) D_s^+, D_1^{*-} \rightarrow \bar{D}^0 \pi^-) / \Gamma(\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV})$$

Γ_{91}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$1.35 \pm 0.40 \pm 0.59$	¹ AAIJ	23B	LHCB pp at 7, 8, 13 TeV

¹ Uses simultaneous fits of $\bar{B}^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ amplitudes assuming isospin symmetry.

$$\Gamma(D_3^{*-}(2750) D_s^+, D_3^{*-} \rightarrow \bar{D}^0 \pi^-) / \Gamma(\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV})$$

Γ_{92}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$0.31 \pm 0.14 \pm 0.17$	¹ AAIJ	23B	LHCB pp at 7, 8, 13 TeV

¹ Uses simultaneous fits of $\bar{B}^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ amplitudes assuming isospin symmetry.

$$\Gamma(D_1^{*-}(2760) D_s^+, D_1^{*-} \rightarrow \bar{D}^0 \pi^-) / \Gamma(\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV})$$

Γ_{93}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$0.28 \pm 0.25 \pm 1.48$	¹ AAIJ	23B	LHCB pp at 7, 8, 13 TeV

¹ Uses simultaneous fits of $\bar{B}^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ amplitudes assuming isospin symmetry.

$$\Gamma(D_J^{*-}(3000) D_s^+, D_J^{*-} \rightarrow \bar{D}^0 \pi^-) / \Gamma(\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV})$$

Γ_{94}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$0.45 \pm 0.16 \pm 0.38$	¹ AAIJ	23B	LHCB pp at 7, 8, 13 TeV

¹ Uses simultaneous fits of $\bar{B}^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ amplitudes assuming isospin symmetry.

$$\Gamma(T_{cs0}^{*0}(2870) \bar{D}^0, T_{cs0}^{*0} \rightarrow D_s^+ \pi^-) / \Gamma(\bar{D}^0 D_s^+ \pi^-, m(\bar{D}^0 \pi^-) > 2.05 \text{ GeV})$$

Γ_{95}/Γ_{87}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$2.48 \pm 0.67 \pm 0.77$	¹ AAIJ	23B	LHCB pp at 7, 8, 13 TeV

¹ Uses simultaneous fits of $\bar{B}^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$ amplitudes assuming isospin symmetry.

$$\Gamma(D^- D_s^{*+}) / \Gamma_{\text{total}}$$

Γ_{96}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0074 ± 0.0016 OUR AVERAGE			
$0.0071 \pm 0.0016 \pm 0.0006$	¹ AUBERT	06N	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

0.0078 ± 0.0032 ± 0.0007 ² GIBAUT 96 CLE2 $e^+e^- \rightarrow \gamma(4S)$
 0.016 ± 0.012 ± 0.001 ³ ALBRECHT 92G ARG $e^+e^- \rightarrow \gamma(4S)$

¹ AUBERT 06N reports $(0.69 \pm 0.16 \pm 0.09) \times 10^{-2}$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² GIBAUT 96 reports $0.0100 \pm 0.0035 \pm 0.0022$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALBRECHT 92G reports $0.027 \pm 0.017 \pm 0.009$ from a measurement of $[\Gamma(B^0 \rightarrow D^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., $B(D^+ \rightarrow K^- 2\pi^+) = 7.7 \pm 1.0\%$.

$\Gamma(D^*(2010)^- D_s^{*+})/\Gamma_{\text{total}}$

Γ_{97}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0177 ± 0.0014 OUR AVERAGE			
0.0173 ± 0.0018 ± 0.0015	¹ AUBERT	06N	BABR $e^+e^- \rightarrow \gamma(4S)$
0.0188 ± 0.0009 ± 0.0017	² AUBERT	05V	BABR $e^+e^- \rightarrow \gamma(4S)$
0.0158 ± 0.0027 ± 0.0014	³ AUBERT	03I	BABR $e^+e^- \rightarrow \gamma(4S)$
0.015 ± 0.004 ± 0.001	⁴ AHMED	00B	CLE2 $e^+e^- \rightarrow \gamma(4S)$
0.016 ± 0.009 ± 0.001	⁵ ALBRECHT	92G	ARG $e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.016 ± 0.005 ± 0.001 ⁶ GIBAUT 96 CLE2 Repl. by AHMED 00B

¹ AUBERT 06N reports $(1.68 \pm 0.21 \pm 0.19) \times 10^{-2}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.0462 \pm 0.0062$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² A partial reconstruction technique is used and the result is independent of the particle decay rate of D_s^+ meson. It also provides a model-independent determination of $B(D_s^+ \rightarrow \phi\pi^+) = (4.81 \pm 0.52 \pm 0.38)\%$.

³ AUBERT 03I reports $0.0197 \pm 0.0015 \pm 0.0030$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AHMED 00B reports $0.0182 \pm 0.0037 \pm 0.0025$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.036$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ ALBRECHT 92G reports $0.026 \pm 0.014 \pm 0.006$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^*(2010)^+$ branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \rightarrow K^- 2\pi^+) = 7.1 \pm 1.0\%$, and $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 55 \pm 4\%$.

⁶ GIBAUT 96 reports $0.0203 \pm 0.0050 \pm 0.0036$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^*(2010)^- D_s^{*+})/\Gamma(D^*(2010)^- D_s^+)$ Γ_{97}/Γ_{88}

VALUE	DOCUMENT ID	TECN	COMMENT
$2.19 \pm 0.08 \pm 0.01$	¹ AAIJ	21S LHCb	pp at 13 TeV

¹ AAIJ 21S reports $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})/\Gamma(B^0 \rightarrow D^*(2010)^- D_s^+)] \times [B(D_s^+ \rightarrow D_s^+ \gamma)] = 2.045 \pm 0.022 \pm 0.071$ which we divide by our best value $B(D_s^+ \rightarrow D_s^+ \gamma) = (93.6 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$[\Gamma(D^*(2010)^- D_s^+) + \Gamma(D^*(2010)^- D_s^{*+})]/\Gamma_{\text{total}}$ $(\Gamma_{88} + \Gamma_{97})/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.4 OUR AVERAGE				
$2.40 \pm 0.35 \pm 0.22$		¹ AUBERT	03I BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.3 \pm 0.9 \pm 0.3$	22	² BORTOLETTO90	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ AUBERT 03I reports $(3.00 \pm 0.19 \pm 0.39) \times 10^{-2}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^+) + \Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})]/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² BORTOLETTO 90 reports $(7.5 \pm 2.0) \times 10^{-2}$ from a measurement of $[\Gamma(B^0 \rightarrow D^*(2010)^- D_s^+) + \Gamma(B^0 \rightarrow D^*(2010)^- D_s^{*+})]/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.02$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{s0}(2317)^- K^+, D_{s0}^- \rightarrow D_s^- \pi^0)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.2^{+1.4}_{-1.3} \pm 0.4$	¹ DRUTSKOY	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ DRUTSKOY 05 reports $(5.3^{+1.5}_{-1.3} \pm 1.6) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow D_{s0}(2317)^- K^+, D_{s0}^- \rightarrow D_s^- \pi^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{s0}(2317)^- \pi^+, D_{s0}^- \rightarrow D_s^- \pi^0)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	90	¹ DRUTSKOY 05	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_{sJ}(2457)^- K^+, D_{sJ}^- \rightarrow D_s^- \pi^0)/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<0.94	90	¹ DRUTSKOY 05	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_{sJ}(2457)^- \pi^+, D_{sJ}^- \rightarrow D_s^- \pi^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<0.40	90	¹ DRUTSKOY 05	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_s^- D_s^+)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3.6×10^{-5}	90	¹ ZUPANC 07	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10 $\times 10^{-5}$	90	¹ AUBERT,BE 05F	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
---	----	----------------------------	------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_s^{*-} D_s^+)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.3×10^{-4}	90	¹ AUBERT,BE 05F	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_s^{*-} D_s^{*+})/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.4×10^{-4}	90	¹ AUBERT,BE 05F	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_{s0}^*(2317)^+ D^-, D_{s0}^{*+} \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.05 ± 0.16 OUR AVERAGE	Error includes scale factor of 1.1.		

$0.98^{+0.16}_{-0.15} \pm 0.02$	^{1,2} CHOI 15A	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
---------------------------------	-------------------------	------	------------------------------------

$1.4^{+0.5}_{-0.4} \pm 0.1$	^{2,3} AUBERT,B 04S	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------------	-----------------------------	------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.69^{+0.29}_{-0.24} \pm 0.06$	^{2,4} KROKOVNY 03B	BELL	Repl. by CHOI 15A
---------------------------------	-----------------------------	------	-------------------

¹ CHOI 15A reports $(10.2^{+1.3}_{-1.2} \pm 1.0 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow D_{s0}^*(2317)^+ D^-, D_{s0}^{*+} \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow K^+ K^- \pi^+)] \times [B(D^+ \rightarrow K^- 2\pi^+)]$ assuming $B(D_s^+ \rightarrow K^+ K^- \pi^+) = (5.39 \pm 0.21) \times 10^{-2}$, $B(D^+ \rightarrow$

$K^- 2\pi^+$) = $(9.13 \pm 0.19) \times 10^{-2}$, which we rescale to our best values $B(D_s^+ \rightarrow K^+ K^- \pi^+) = (5.45 \pm 0.08) \times 10^{-2}$, $B(D^+ \rightarrow K^- 2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AUBERT,B 04s reports $(1.8 \pm 0.4_{-0.5}^{+0.7}) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_{s0}^*(2317)^+ D^-, D_{s0}^{*+} \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ KROKOVNY 03B reports $(0.86_{-0.26}^{+0.33} \pm 0.26) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_{s0}^*(2317)^+ D^-, D_{s0}^{*+} \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{s0}(2317)^+ D^-, D_{s0}^{*+} \rightarrow D_s^{*+} \gamma)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.95	90	¹ KROKOVNY 03B	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{s0}(2317)^+ D^{*+}(2010)^-, D_{s0}^{*+} \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.4_{-0.4}^{+0.5}$	¹ AUBERT,B 04s	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_{sJ}(2457)^+ D^-)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.5 ± 1.1 OUR AVERAGE			
$2.6 \pm 1.5 \pm 0.7$	¹ AUBERT 06N	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$4.8_{-1.6}^{+2.2} \pm 1.1$	^{2,3} AUBERT,B 04s	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.9_{-1.3}^{+1.5} \pm 0.9$	^{2,4} KROKOVNY 03B	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses a missing-mass method in the events that one of the B mesons is fully reconstructed.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AUBERT,B 04s reports $[\Gamma(B^0 \rightarrow D_{sJ}(2457)^+ D^-)/\Gamma_{\text{total}}] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0)] = (2.3_{-0.7}^{+1.0} \pm 0.3) \times 10^{-3}$ which we divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ KROKOVNY 03B reports $[\Gamma(B^0 \rightarrow D_{sJ}(2457)^+ D^-)/\Gamma_{\text{total}}] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0)] = (1.9_{-0.6}^{+0.7} \pm 0.2) \times 10^{-3}$ which we divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \gamma)/\Gamma_{\text{total}}$	Γ_{109}/Γ
VALUE (units 10^{-3})	DOCUMENT ID TECN COMMENT
$0.65^{+0.17}_{-0.14}$ OUR AVERAGE	

$0.64^{+0.24}_{-0.16} \pm 0.06$ 1,2 AUBERT,B 04S BABR $e^+ e^- \rightarrow \gamma(4S)$

$0.66^{+0.21}_{-0.19} \pm 0.06$ 1,3 KROKOVNY 03B BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

² AUBERT,B 04S reports $(0.8 \pm 0.2^{+0.3}_{-0.2}) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \gamma)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ KROKOVNY 03B reports $(0.82^{+0.22}_{-0.19} \pm 0.25) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \gamma)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.036 \pm 0.009$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^{*+} \gamma)/\Gamma_{\text{total}}$	Γ_{110}/Γ
VALUE (units 10^{-3})	CL% DOCUMENT ID TECN COMMENT
<0.60	90 1 KROKOVNY 03B BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_{111}/Γ
VALUE (units 10^{-3})	CL% DOCUMENT ID TECN COMMENT
<0.20	90 1 KROKOVNY 03B BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D_{sJ}(2457)^+ D^-, D_{sJ}^+ \rightarrow D_s^+ \pi^0)/\Gamma_{\text{total}}$	Γ_{112}/Γ
VALUE (units 10^{-3})	CL% DOCUMENT ID TECN COMMENT
<0.36	90 1 KROKOVNY 03B BELL $e^+ e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$\Gamma(D^*(2010)^- D_{sJ}(2457)^+)/\Gamma_{\text{total}}$	Γ_{113}/Γ
VALUE (units 10^{-3})	DOCUMENT ID TECN COMMENT
9.3 ± 2.2 OUR AVERAGE	

$8.8 \pm 2.0 \pm 1.4$ 1 AUBERT 06N BABR $e^+ e^- \rightarrow \gamma(4S)$

11 $^{+5}_{-4} \pm 3$ 2,3 AUBERT,B 04S BABR $e^+ e^- \rightarrow \gamma(4S)$

¹ Uses a missing-mass method in the events that one of the B mesons is fully reconstructed.

² AUBERT,B 04S reports $[\Gamma(B^0 \rightarrow D^*(2010)^- D_{sJ}(2457)^+)/\Gamma_{\text{total}}] \times [B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0)] = (5.5 \pm 1.2^{+2.2}_{-1.6}) \times 10^{-3}$ which we divide by our best value $B(D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0) = (48 \pm 11) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

$$\Gamma(D_{sJ}(2457)^+ D^*(2010), D_{sJ}^+ \rightarrow D_s^+ \gamma) / \Gamma_{\text{total}} \quad \Gamma_{114} / \Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.3 \pm 0.3^{+0.9}_{-0.6}$	¹ AUBERT,B	04S	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$[\Gamma(D^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+) + \Gamma(D^{*+} K^0)] / \Gamma_{\text{total}} \quad \Gamma_{115} / \Gamma = (\Gamma_{116} + \Gamma_{117}) / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.75 \pm 0.62 \pm 0.36$	^{1,2} AUSHEV	11	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses $\Gamma(D^*(2007)^0 \rightarrow D^0 \pi^0) / \Gamma(D^*(2007)^0 \rightarrow D^0 \gamma) = 1.74 \pm 0.13$ and $\Gamma(D_{s1}(2536)^+ \rightarrow D^*(2007)^0 K^+) / \Gamma(D_{s1}(2536)^+ \rightarrow D^*(2010)^+ K^0) = 1.36 \pm 0.2$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+) / \Gamma_{\text{total}} \quad \Gamma_{116} / \Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$1.71 \pm 0.48 \pm 0.32$		¹ AUBERT	08B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5 90 AUBERT 03X BABR Repl. by AUBERT 08B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*+} K^0) / \Gamma_{\text{total}} \quad \Gamma_{117} / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.61 \pm 1.03 \pm 0.31$	¹ AUBERT	08B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$[\Gamma(D^*(2010)^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+) + \Gamma(D^{*+} K^0)] / \Gamma_{\text{total}} \quad \Gamma_{118} / \Gamma = (\Gamma_{119} + \Gamma_{120}) / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$5.01 \pm 1.21 \pm 0.70$	^{1,2} AUSHEV	11	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses $\Gamma(D^*(2007)^0 \rightarrow D^0 \pi^0) / \Gamma(D^*(2007)^0 \rightarrow D^0 \gamma) = 1.74 \pm 0.13$ and $\Gamma(D_{s1}(2536)^+ \rightarrow D^*(2007)^0 K^+) / \Gamma(D_{s1}(2536)^+ \rightarrow D^*(2010)^+ K^0) = 1.36 \pm 0.2$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^*(2010)^- D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*0} K^+) / \Gamma_{\text{total}} \quad \Gamma_{119} / \Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$3.32 \pm 0.88 \pm 0.66$		¹ AUBERT	08B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7 90 AUBERT 03X BABR Repl. by AUBERT 08B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^{*-} D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*+} K^0) / \Gamma_{\text{total}} \quad \Gamma_{120} / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$5.00 \pm 1.51 \pm 0.67$	¹ AUBERT	08B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^- D_{sJ}(2573)^+, D_{sJ}^+ \rightarrow D^0 K^+)/\Gamma_{\text{total}} \quad \Gamma_{121}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$3.4 \pm 1.7 \pm 0.5$		¹ LEES	15C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----	----	--------	----------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^{*-}(2010) D_{sJ}(2573)^+, D_{sJ}^+ \rightarrow D^0 K^+)/\Gamma_{\text{total}} \quad \Gamma_{122}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	AUBERT	03X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

$$\Gamma(D^- D_{sJ}(2700)^+, D_{sJ}^+ \rightarrow D^0 K^+)/\Gamma_{\text{total}} \quad \Gamma_{123}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$7.14 \pm 0.96 \pm 0.69$		¹ LEES	15C BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{124}/\Gamma$$

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
$7.3 \pm 1.2 \pm 0.2$		^{1,2} DAS	10 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ DAS 10 reports $[\Gamma(B^0 \rightarrow D^+ \pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow D^- \pi^+)] = (2.92 \pm 0.38 \pm 0.31) \times 10^{-4}$ which we multiply by our best value $B(B^0 \rightarrow D^- \pi^+) = (2.51 \pm 0.08) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Derived using $\tan(\theta_C) f_D/f_{D_s} \sqrt{B(B^0 \rightarrow D_s^+ \pi^-)/B(B^0 \rightarrow D^- \pi^+)}$ by assuming the flavor SU(3) symmetry, where θ_C is the Cabibbo angle, f_D (f_{D_s}) is the D (D_s) meson decay constant.

$$\Gamma(D_s^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{125}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
20.3 ± 1.8 OUR FIT				
21.6 ± 2.6 OUR AVERAGE				

$19.9 \pm 2.6 \pm 1.8$		¹ DAS	10 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
------------------------	--	------------------	---------	------------------------------------

$25 \pm 4 \pm 2$		¹ AUBERT	08AJ BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
------------------	--	---------------------	-----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$14.0 \pm 3.5 \pm 1.3$		² AUBERT	07K BABR	Repl. by AUBERT 08AJ
------------------------	--	---------------------	----------	----------------------

$25 \pm 9 \pm 2$		³ AUBERT	03D BABR	Repl. by AUBERT 07K
------------------	--	---------------------	----------	---------------------

$19 \pm 9 \pm 7$		⁴ KROKOVNY	02 BELL	Repl. by DAS 10
------------------	--	-----------------------	---------	-----------------

< 220	90	⁵ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	------------------------	----------	------------------------------------

<1300	90	⁶ BORTOLETTO	90 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	-------------------------	---------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 07K reports $[\Gamma(B^0 \rightarrow D_s^+ \pi^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = (0.63 \pm 0.15 \pm 0.05) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AUBERT 03D reports $[\Gamma(B^0 \rightarrow D_s^+ \pi^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = (1.13 \pm 0.33 \pm 0.21) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ KROKOVNY 02 reports $[\Gamma(B^0 \rightarrow D_s^+ \pi^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = (0.86_{-0.30}^{+0.37} \pm 0.11) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ ALEXANDER 93B reports $< 270 \times 10^{-6}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^+ \pi^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

⁶ BORTOLETTO 90 assume $B(D_s \rightarrow \phi \pi^+) = 2\%$.

$$[\Gamma(D_s^+ \pi^-) + \Gamma(D_s^- K^+)]/\Gamma_{\text{total}} \quad (\Gamma_{125} + \Gamma_{135})/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.0 \times 10^{-3}$	90	¹ ALBRECHT 93E	ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^+ \pi^-) + \Gamma(B^0 \rightarrow D_s^- K^+)]/\Gamma_{\text{total}} \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$$\Gamma(D_s^+ \pi^-)/\Gamma(D^- \pi^+) \quad \Gamma_{125}/\Gamma_{38}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.7 OUR FIT			
$7.7 \pm 0.7 \pm 0.6$	AAIJ	21W LHCb	pp at 7, 8, 13 TeV

$$\Gamma(D_s^{*+} \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{126}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
2.1 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.4.
$1.75 \pm 0.34 \pm 0.20$		¹ JOSHI 10	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$2.6_{-0.4}^{+0.5} \pm 0.2$		¹ AUBERT 08AJ	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.9 \pm 0.7 \pm 0.3$		² AUBERT 07K	BABR	Repl. by AUBERT 08AJ
< 4.1	90	AUBERT 03D	BABR	Repl. by AUBERT 07K
< 40	90	³ ALEXANDER 93B	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 07K reports $[\Gamma(B^0 \rightarrow D_s^{*+} \pi^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = (1.32 \pm 0.27 \pm 0.15) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALEXANDER 93B reports $< 44 \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*+} \pi^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$[\Gamma(D_s^{*+} \pi^-) + \Gamma(D_s^{*-} K^+)]/\Gamma_{\text{total}}$		$(\Gamma_{126} + \Gamma_{136})/\Gamma$		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7 \times 10^{-4}$	90	¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 1.2 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*+} \pi^-) + \Gamma(B^0 \rightarrow D_s^{*-} K^+)]/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^+ \rho^-)/\Gamma_{\text{total}}$		Γ_{127}/Γ		
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT

< 2.4	90	¹ AUBERT	08AJ BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	---------------------	-----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 130	90	² ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 50	90	³ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^+ \rho^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

³ ALEXANDER 93B reports $< 6.6 \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^+ \rho^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^{*+} \rho^-)/\Gamma_{\text{total}}$		Γ_{128}/Γ		
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT

$4.1^{+1.3}_{-1.2} \pm 0.4$		¹ AUBERT	08AJ BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------------	--	---------------------	-----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 150	90	² ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
< 60	90	³ ALEXANDER	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*+} \rho^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

³ ALEXANDER 93B reports $< 7.4 \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*+} \rho^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^+ a_0^-)/\Gamma_{\text{total}}$		Γ_{129}/Γ		
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT

< 1.9	90	¹ AUBERT	06X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	---------------------	----------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_s^{*+} a_0^-)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3.6	90	¹ AUBERT	06X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_s^+ a_1(1260)^-)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.1 $\times 10^{-3}$	90	¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 3.5 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^+ a_1(1260)^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

 $\Gamma(D_s^{*+} a_1(1260)^-)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 $\times 10^{-3}$	90	¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 2.9 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*+} a_1(1260)^-)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

 $\Gamma(D_s^+ a_2^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<19	90	¹ AUBERT	06X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_s^{*+} a_2^-)/\Gamma_{\text{total}}$ Γ_{134}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	¹ AUBERT	06X BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D_s^- K^+)/\Gamma_{\text{total}}$ Γ_{135}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
27 $\pm$ 5 OUR FIT		Error includes scale factor of 2.7.		

22 $\pm$ 5 OUR AVERAGE Error includes scale factor of 1.8.

19.1 $\pm$ 2.4 $\pm$ 1.7 ¹ DAS 10 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

29 $\pm$ 4 $\pm$ 2 ¹ AUBERT 08AJ BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

27 $\pm$ 5 $\pm$ 2 ² AUBERT 07K BABR Repl. by AUBERT 08AJ

26 $\pm$ 10 $\pm$ 2 ³ AUBERT 03D BABR Repl. by AUBERT 07K

36 $\pm$ 11 $\pm$ 3 ⁴ KROKOVNY 02 BELL Repl. by DAS 10

< 190 90 ⁵ ALEXANDER 93B CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

< 1300 90 ⁶ BORTOLETTO90 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 07K reports $[\Gamma(B^0 \rightarrow D_s^- K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (1.21 \pm 0.17 \pm 0.11) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$.

Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AUBERT 03D reports $[\Gamma(B^0 \rightarrow D_s^- K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (1.16 \pm 0.36 \pm 0.24) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$.

Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ KROKOVNY 02 reports $[\Gamma(B^0 \rightarrow D_s^- K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (1.61^{+0.45}_{-0.38} \pm 0.21) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ ALEXANDER 93B reports $< 230 \times 10^{-6}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^- K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

⁶ BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$.

$\Gamma(D_s^- K^+)/\Gamma(D^- \pi^+)$ Γ_{135}/Γ_{38}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.09±0.19 OUR FIT	Error includes scale factor of 2.7.		
1.29±0.05±0.08	AAIJ	15AC LHCb	pp at 7, 8 TeV

$\Gamma(D_s^{*-} K^+)/\Gamma_{\text{total}}$ Γ_{136}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
2.19±0.30 OUR AVERAGE				
2.02±0.33±0.22		¹ JOSHI	10 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
2.4 ±0.4 ±0.2		¹ AUBERT	08AJ BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.2 ±0.6 ±0.2		² AUBERT	07K BABR	Repl. by AUBERT 08AJ
< 2.5	90	AUBERT	03D BABR	Repl. by AUBERT 07K
<14	90	³ ALEXANDER	93B CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 07K reports $[\Gamma(B^0 \rightarrow D_s^{*-} K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (0.97 \pm 0.24 \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ALEXANDER 93B reports $< 17 \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*-} K^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_{s1}(2536)^\mp K^\pm, D_{s1}^- \rightarrow \bar{D}^*(2007)^0 K^-)/\Gamma(\bar{D}^0 K^+ K^-)$ Γ_{137}/Γ_{49}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
8.4±0.3±0.6	AAIJ	23AY LHCb	pp at 7, 8, 13 TeV

$\Gamma(D_s^- K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{138}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
3.5^{+1.0}_{-0.9}±0.4		¹ AUBERT	08AJ BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<280	90	² ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 80	90	³ ALEXANDER	93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 4.6 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^- K^*(892)^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

³ ALEXANDER 93B reports $< 9.7 \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^- K^*(892)^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^{*-} K^*(892)^+)/\Gamma_{\text{total}}$		Γ_{139}/Γ			
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
$3.2^{+1.4}_{-1.2} \pm 0.4$		¹ AUBERT	08AJ	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<350	90	² ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 90	90	³ ALEXANDER	93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 5.8 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*-} K^*(892)^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

³ ALEXANDER 93B reports $< 11.0 \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*-} K^*(892)^+)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^- \pi^+ K^0)/\Gamma_{\text{total}}$		Γ_{140}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
0.97 ± 0.14 OUR AVERAGE					

$0.94 \pm 0.12 \pm 0.10$

¹ WIECHCZYN...15 BELL $e^+e^- \rightarrow \Upsilon(4S)$

$1.10 \pm 0.26 \pm 0.20$

¹ AUBERT 08G BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<40	90	² ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----	----	-----------------------	-----	-----	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 7.3 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^- \pi^+ K^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)]$ assuming $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^{*-} \pi^+ K^0)/\Gamma_{\text{total}}$		Γ_{141}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	

< 1.10

90

¹ AUBERT 08G BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<25	90	² ALBRECHT	93E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----	----	-----------------------	-----	-----	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ALBRECHT 93E reports $< 4.2 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*-} \pi^+ K^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^- K^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.71 \pm 0.31 \pm 0.34$	¹ AAIJ	12AX LHCB	pp at 7 TeV

¹ AAIJ 12AX reports $[\Gamma(B^0 \rightarrow D_s^- K^+ \pi^+ \pi^-)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow D_s^- K^+ \pi^+ \pi^-)] = 0.54 \pm 0.07 \pm 0.07$ which we multiply by our best value $B(B_s^0 \rightarrow D_s^- K^+ \pi^+ \pi^-) = (3.2 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D_s^- \pi^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{143}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.0 \times 10^{-3}$	90	¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 5.0 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^- \pi^+ K^*(892)^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(D_s^{*-} \pi^+ K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-3}$	90	¹ ALBRECHT	93E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ ALBRECHT 93E reports $< 2.7 \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow D_s^{*-} \pi^+ K^*(892)^0)/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)]$ assuming $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$, which we rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 4.5 \times 10^{-2}$.

$\Gamma(\bar{D}^0 K^0)/\Gamma_{\text{total}}$ Γ_{145}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
5.5 ± 0.4 OUR AVERAGE			
$5.6 \pm 0.5 \pm 0.2$	KUMAR	23 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$5.3 \pm 0.7 \pm 0.3$	¹ AUBERT,B	06L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.0^{+1.3}_{-1.2} \pm 0.6$	¹ KROKOVNY	03 BELL	Repl. by KUMAR 23
-----------------------------	-----------------------	---------	-------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$88 \pm 15 \pm 9$	¹ AUBERT	06A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^0 K^+ \pi^-)/\Gamma(\bar{D}^0 \pi^+ \pi^-)$ Γ_{146}/Γ_{47}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.106 \pm 0.007 \pm 0.008$	AAIJ	13AQ LHCB	pp at 7 TeV

$\Gamma(\bar{D}^0 K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.5 ± 0.6 OUR AVERAGE			
$5.4 \pm 0.3 \pm 1.1$	^{1,2} AAIJ	15X	LHCB pp at 7, 8 TeV
$4.0 \pm 0.7 \pm 0.3$	³ AUBERT,B	06L	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$4.8^{+1.1}_{-1.0} \pm 0.5$	³ KROKOVNY	03	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.7 \pm 0.9 \pm 0.6$	³ AUBERT	06A	BABR Repl. by AUBERT,B 06L
¹ AAIJ 15X reports $(5.13 \pm 0.20 \pm 0.15 \pm 0.24 \pm 0.60) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow \bar{D}^0 K^*(892)^0)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-)]$ assuming $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (9.2 \pm 0.6 \pm 0.7 \pm 0.6) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (8.8 \pm 1.7) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
² Measured via amplitude analysis of $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$, which excludes contribution from decay via $D^*(2010)^-$ resonance.			
³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			

 $\Gamma(\bar{D}^0 K^*(1410)^0)/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.7 \times 10^{-5}$	90	¹ AAIJ	15X	LHCB pp at 7, 8 TeV
¹ Measured via amplitude analysis of $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$, which excludes contribution from decay via $D^*(2010)^-$ resonance.				

 $\Gamma(\bar{D}^0 K_0^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.7 \pm 0.7 \pm 0.1$	^{1,2} AAIJ	15X	LHCB pp at 7, 8 TeV
¹ AAIJ 15X reports $(0.71 \pm 0.27 \pm 0.33 \pm 0.47 \pm 0.08) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow \bar{D}^0 K_0^*(1430)^0)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-)]$ assuming $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (9.2 \pm 0.6 \pm 0.7 \pm 0.6) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (8.8 \pm 1.7) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
² Measured via amplitude analysis of $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$, which excludes contribution from decay via $D^*(2010)^-$ resonance.			

 $\Gamma(\bar{D}^0 K_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$2.1 \pm 0.8 \pm 0.4$	^{1,2} AAIJ	15X	LHCB pp at 7, 8 TeV
¹ AAIJ 15X reports $(2.04 \pm 0.45 \pm 0.30 \pm 0.54 \pm 0.25) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow \bar{D}^0 K_2^*(1430)^0)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-)]$ assuming $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (9.2 \pm 0.6 \pm 0.7 \pm 0.6) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (8.8 \pm 1.7) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
² Measured via amplitude analysis of $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$, which excludes contribution from decay via $D^*(2010)^-$ resonance.			

 $\Gamma(D_0^{*-}(2300) K^+, D_0^{*-} \rightarrow \bar{D}^0 \pi^-)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.8 \pm 0.4$	^{1,2} AAIJ	15X	LHCB pp at 7, 8 TeV

- ¹ AAIJ 15X reports $(1.77 \pm 0.26 \pm 0.19 \pm 0.67 \pm 0.20) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow D_0^{*}(2300)^- K^+, D_0^{*-} \rightarrow \bar{D}^0 \pi^-)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-)]$ assuming $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (9.2 \pm 0.6 \pm 0.7 \pm 0.6) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (8.8 \pm 1.7) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² Measured via amplitude analysis of $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$, which excludes contribution from decay via $D^*(2010)^-$ resonance.

$\Gamma(D_2^{*}(2460)^- K^+, D_2^{*-} \rightarrow \bar{D}^0 \pi^-)/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
20.3 ± 3.5 OUR AVERAGE			
22 ± 2 ± 4	^{1,2} AAIJ	15X LHCB	pp at 7, 8 TeV
18.3 $\pm 4.0 \pm 3.1$	³ AUBERT	06A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ AAIJ 15X reports $(2.12 \pm 0.10 \pm 0.11 \pm 0.11 \pm 0.25) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow D_2^{*}(2460)^- K^+, D_2^{*-} \rightarrow \bar{D}^0 \pi^-)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-)]$ assuming $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (9.2 \pm 0.6 \pm 0.7 \pm 0.6) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow \bar{D}^0 K^+ \pi^-) = (8.8 \pm 1.7) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² Measured via amplitude analysis of $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$, which excludes contribution from decay via $D^*(2010)^-$ resonance.
- ³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D_3^{*}(2760)^- K^+, D_3^{*-} \rightarrow \bar{D}^0 \pi^-)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.10 \times 10^{-5}$	90	¹ AAIJ	15X LHCB	pp at 7, 8 TeV

- ¹ Measured via amplitude analysis of $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$, which excludes contribution from decay via $D^*(2010)^-$ resonance.

$\Gamma(\bar{D}^0 K^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 37	90	¹ AUBERT	06A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma([K^+ K^-]_D K^*(892)^0)/\Gamma(\bar{D}^0 K^*(892)^0)$ $\Gamma_{155}/\Gamma_{147}$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.811 \pm 0.057 \pm 0.017$	AAIJ	24M LHCB	pp at 7, 8, 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.92 ± 0.10 ± 0.02	AAIJ	19N LHCB	Repl. by AAIJ 24M
1.05 $^{+0.17}_{-0.15}$ ± 0.04	AAIJ	14BN LHCB	Repl. by AAIJ 16S
1.36 $^{+0.37}_{-0.32}$ ± 0.07	AAIJ	13L LHCB	Repl. by AAIJ 14BN

$\Gamma([\pi^+ \pi^-]_D K^*(892)^0)/\Gamma(\bar{D}^0 K^*(892)^0)$ $\Gamma_{156}/\Gamma_{147}$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.104 \pm 0.111 \pm 0.026$	AAIJ	24M LHCB	pp at 7, 8, 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.32 ± 0.19 ± 0.03	AAIJ	19N LHCB	Repl. by AAIJ 24M
1.21 $^{+0.28}_{-0.25}$ ± 0.05	AAIJ	14BN LHCB	Repl. by AAIJ 16S

$$\Gamma([\pi^+ K^-]_D K^*(892)^0)/\Gamma([K^+ \pi^-]_D K^*(892)^0) \quad \Gamma_{157}/\Gamma_{158}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.080 ± 0.015 ± 0.002	AAIJ	19N	LHCB pp at 7, 8, 13 TeV

$$\Gamma([\pi^+ \pi^- \pi^+ \pi^-]_D K^{*0})/\Gamma(\bar{D}^0 K^*(892)^0) \quad \Gamma_{159}/\Gamma_{147}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.882 ± 0.086 ± 0.033	AAIJ	24M	LHCB pp at 7, 8, 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.01 ± 0.16 ± 0.04	AAIJ	19N	LHCB Repl. by AAIJ 24M
--------------------	------	-----	------------------------

$$\Gamma([\pi^+ K^- \pi^+ \pi^-]_D K^{*0})/\Gamma([K^+ \pi^- \pi^+ \pi^-]_D K^{*0}) \quad \Gamma_{160}/\Gamma_{161}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.073 ± 0.018 ± 0.002	AAIJ	19N	LHCB pp at 7, 8, 13 TeV

$$\Gamma(\bar{D}^0 \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{162}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
2.67 ± 0.09 OUR AVERAGE				
2.70 ± 0.06 ± 0.10		BLOOMFIELD 22	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
2.69 ± 0.09 ± 0.13		¹ LEES	11M	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
2.25 ± 0.14 ± 0.35		¹ BLYTH	06	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
2.74 $^{+0.36}_{-0.32}$ ± 0.55		¹ COAN	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.9 ± 0.2 ± 0.3		¹ AUBERT	04B	BABR Repl. by LEES 11M
3.1 ± 0.4 ± 0.5		¹ ABE	02J	BELL Repl. by BLYTH 06
< 1.2	90	² NEMAT1	98	CLE2 Repl. by COAN 02
< 4.8	90	³ ALAM	94	CLE2 Repl. by NEMAT1 98

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMAT1 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$$\Gamma(\bar{D}^0 \rho^0)/\Gamma_{\text{total}} \quad \Gamma_{163}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
3.21 ± 0.21 OUR AVERAGE				
3.21 ± 0.10 ± 0.21		¹ AAIJ	15Y	LHCB pp at 7, 8 TeV
3.19 ± 0.20 ± 0.45		^{2,3} KUZMIN	07	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.9 ± 1.0 ± 0.4		² SATPATHY	03	BELL Repl. by KUZMIN 07
< 3.9	90	⁴ NEMAT1	98	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
< 5.5	90	⁵ ALAM	94	CLE2 Repl. by NEMAT1 98
< 6.0	90	⁶ BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
< 27.0	90	⁷ ALBRECHT	88K	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Measured using isobar formalism in the decay chain $B^0 \rightarrow \bar{D}^0 \rho(770)$, $\rho \rightarrow \pi^+ \pi^-$ assuming $B(\rho(770) \rightarrow \pi^+ \pi^-) = 1$. The second uncertainty combines in quadrature all systematic uncertainties quoted in the paper.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Our second uncertainty combines systematics and model errors quoted in the paper.

⁴ NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

⁵ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

⁶ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

⁷ ALBRECHT 88K reports < 0.003 assuming $B^0 \bar{B}^0 : B^+ B^-$ production ratio is 45:55. We rescale to 50%.

$\Gamma(\bar{D}^0 f_2)/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.56 ± 0.21 OUR AVERAGE			
1.68 ± 0.11 ± 0.21	¹ AAIJ	15Y	LHCB pp at 7, 8 TeV
1.20 ± 0.18 ± 0.38	^{2,3} KUZMIN	07	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Result obtained using the isobar formalism. The second uncertainty combines in quadrature all systematic uncertainties quoted in the paper. Measured in the decay chain $B^0 \rightarrow \bar{D}^0 f_2(1270)$, $f_2 \rightarrow \pi^+ \pi^-$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Our second uncertainty combines systematics and model errors quoted in the paper.

$\Gamma(\bar{D}^0 \eta)/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
2.56 ± 0.12 OUR AVERAGE				
2.66 ± 0.12 ± 0.21		KUMAR	23	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
2.53 ± 0.09 ± 0.11		¹ LEES	11M	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.77 ± 0.16 ± 0.21		¹ BLYTH	06	BELL Repl. by KUMAR 23
2.5 ± 0.2 ± 0.3		¹ AUBERT	04B	BABR Repl. by LEES 11M
1.4 $^{+0.5}_{-0.4}$ ± 0.3		¹ ABE	02J	BELL Repl. by BLYTH 06
<1.3	90	² NEMATI	98	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
<6.8	90	³ ALAM	94	CLE2 Repl. by NEMATI 98

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\bar{D}^0 \eta')/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.38 ± 0.16 OUR AVERAGE		Error includes scale factor of 1.3.		
1.48 ± 0.13 ± 0.07		¹ LEES	11M	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
1.14 ± 0.20 $^{+0.10}_{-0.13}$		¹ SCHUMANN	05	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.7 ± 0.4 ± 0.2		¹ AUBERT	04B	BABR Repl. by LEES 11M
<9.4	90	² NEMATI	98	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

<8.6 90 ³ ALAM 94 CLE2 Repl. by NEMATI 98

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\overline{D}^0 \eta')/\Gamma(\overline{D}^0 \eta)$

$\Gamma_{166}/\Gamma_{165}$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.54 \pm 0.07 \pm 0.01$	LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.7 \pm 0.2 \pm 0.1$	AUBERT	04B BABR	Repl. by LEES 11M

$\Gamma(\overline{D}^0 \omega)/\Gamma_{\text{total}}$

Γ_{167}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
2.54 ± 0.16 OUR AVERAGE				
$2.75 \pm 0.72 \pm 0.35$		¹ AAIJ	15Y LHCb	pp at 7, 8 TeV
$2.57 \pm 0.11 \pm 0.14$		² LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$2.37 \pm 0.23 \pm 0.28$		² BLYTH	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$3.0 \pm 0.3 \pm 0.4$		² AUBERT	04B BABR	Repl. by LEES 11M
$1.8 \pm 0.5 \pm 0.4$		² ABE	02J BELL	Repl. by BLYTH 06
<5.1	90	³ NEMATI	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<6.3	90	⁴ ALAM	94 CLE2	Repl. by NEMATI 98

¹ Result obtained using the isobar model. The second uncertainty combines in quadrature all systematic uncertainties quoted in the paper.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ NEMATI 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\overline{D}^0 \phi)/\Gamma_{\text{total}}$

Γ_{168}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
$7.7 \pm 2.1 \pm 1.0$		¹ AAIJ	23AZ LHCb	pp at 7, 8, 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 23	95	AAIJ	18AY LHCb	Repl. by AAIJ 23AZ
<116	90	² AUBERT	07A0 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ The last uncertainty includes the uncertainties of the branching fractions $B(B^0 \rightarrow \overline{D}^0 K^+ K^-)$ and $B(\phi \rightarrow K^+ K^-)$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^0 K^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{169}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<19	90	¹ AUBERT	06A BABR	Repl. by AUBERT 09AE

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^0 K^+ \pi^-)/\Gamma(\bar{D}^0 K^+ \pi^-)$ $\Gamma_{169}/\Gamma_{146}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.060 ± 0.034 OUR AVERAGE

0.045 ^{+0.056+0.028} _{-0.050-0.018}	1,2 NEGISHI	12	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
---	-------------	----	------	-----------------------------------

0.068 ± 0.042	3 AUBERT	09AE	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
---------------	----------	------	------	-----------------------------------

¹ Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.² Uses $D^0 \rightarrow K^- \pi^+$ mode. Restricts $K^+ \pi^-$ mass within ± 50 MeV of the nominal K^{*0} mass. Corresponds to the upper limit, < 0.16 at 95% CL.³ Reports a signal at the level of 2.5 standard deviations after combining results from $D^0 \rightarrow K^+ \pi^-$, $K^+ \pi^- \pi^0$, and $K^+ \pi^- \pi^+ \pi^-$. $\Gamma(D^0 K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{170}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.1	90	1 AUBERT,B	06L	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	------------	-----	------	-----------------------------------

< 1.8	90	1 KROKOVNY	03	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	------------	----	------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(D^0 K^*(892)^0)/\Gamma(\bar{D}^0 K^*(892)^0)$ $\Gamma_{170}/\Gamma_{147}$ "OUR EVALUATION" is derived from $r_{B^0}(B^0 \rightarrow DK^{*0})$ data block listed in "CP violation parameters" section.

VALUE (units 10^{-2})	DOCUMENT ID
--------------------------	-------------

6.6^{+1.1}_{-1.2} OUR EVALUATION (Produced by HFLAV) $\Gamma(\bar{D}^{*0} \gamma)/\Gamma_{\text{total}}$ Γ_{171}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 2.5 \times 10^{-5}$	90	1 AUBERT,B	05Q	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
---	----	------------	-----	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 5.0 \times 10^{-5}$	90	1 ARTUSO	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
------------------------	----	----------	----	------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. $\Gamma(\bar{D}^{*0}(2007)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{172}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

2.2 ± 0.6 OUR AVERAGE Error includes scale factor of 2.6. See the ideogram below.

3.05 ± 0.14 ± 0.28	1 LEES	11M	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------------------	--------	-----	------	-----------------------------------

1.39 ± 0.18 ± 0.26	1 BLYTH	06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
--------------------	---------	----	------	-----------------------------------

2.20 ^{+0.59} _{-0.52} ± 0.79	1 COAN	02	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
---	--------	----	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.9 ± 0.4 ± 0.5	1 AUBERT	04B	BABR	Repl. by LEES 11M
-----------------	----------	-----	------	-------------------


2.7 ^{+0.8} _{-0.7} ± 0.5	1 ABE	02J	BELL	Repl. by BLYTH 06
---	-------	-----	------	-------------------

< 4.4	90	2 NEMAT1	98	CLE2	Repl. by COAN 02
---------	----	----------	----	------	------------------

< 9.7	90	3 ALAM	94	CLE2	Repl. by NEMAT1 98
---------	----	--------	----	------	--------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.² NEMAT1 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.



$\Gamma(\bar{D}^0 \pi^0) / \Gamma(\bar{D}^*(2007)^0 \pi^0)$

$\Gamma_{162} / \Gamma_{172}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.90±0.08 OUR AVERAGE			
0.88±0.05±0.06	LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
1.62±0.23±0.35	BLYTH	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.0 ±0.1 ±0.2	AUBERT	04B BABR	Repl. by LEES 11M

$\Gamma(\bar{D}^*(2007)^0 \rho^0) / \Gamma_{\text{total}}$

Γ_{173} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.1 × 10⁻⁴	90	¹ SATPATHY	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.00056	90	² NEMAT1	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<0.00117	90	³ ALAM	94 CLE2	Repl. by NEMAT1 98

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMAT1 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(\bar{D}^*(2007)^0 \eta) / \Gamma_{\text{total}}$ Γ_{174} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
2.3 $\pm$ 0.6 OUR AVERAGE		Error includes scale factor of 2.8.		
2.69 $\pm$ 0.14 $\pm$ 0.23		¹ LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
1.40 $\pm$ 0.28 $\pm$ 0.26		¹ BLYTH	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.6 $\pm$ 0.4 $\pm$ 0.4		¹ AUBERT	04B BABR	Repl. by LEES 11M
<4.6	90	¹ ABE	02J BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
<2.6	90	² NEMAT1	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<6.9	90	³ ALAM	94 CLE2	Repl. by NEMAT1 98

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMAT1 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0) / B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-) / B(D^0 \rightarrow K^- \pi^+)$.

 $\Gamma(\bar{D}^0 \eta) / \Gamma(\bar{D}^*(2007)^0 \eta)$ $\Gamma_{165} / \Gamma_{174}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.99 $\pm$ 0.10 OUR AVERAGE			
0.97 $\pm$ 0.07 $\pm$ 0.07	LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
1.27 $\pm$ 0.29 $\pm$ 0.25	BLYTH	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.9 $\pm$ 0.2 $\pm$ 0.1	AUBERT	04B BABR	Repl. by LEES 11M

 $\Gamma(\bar{D}^*(2007)^0 \eta') / \Gamma(\bar{D}^*(2007)^0 \eta)$ $\Gamma_{175} / \Gamma_{174}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.61 $\pm$ 0.14 $\pm$ 0.02	LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.5 $\pm$ 0.3 $\pm$ 0.1	AUBERT	04B BABR	Repl. by LEES 11M

 $\Gamma(\bar{D}^*(2007)^0 \eta') / \Gamma_{\text{total}}$ Γ_{175} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.40 $\pm$ 0.22 OUR AVERAGE				
1.48 $\pm$ 0.22 $\pm$ 0.13		¹ LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
1.21 $\pm$ 0.34 $\pm$ 0.22		¹ SCHUMANN	05 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.3 $\pm$ 0.7 $\pm$ 0.2		^{1,2} AUBERT	04B BABR	Repl. by LEES 11M
<14	90	BRANDENB...	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<19	90	³ NEMAT1	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<27	90	⁴ ALAM	94 CLE2	Repl. by NEMAT1 98

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Reports an upper limit $< 2.6 \times 10^{-4}$ at 90% CL.

³ NEMAT1 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

⁴ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0) / B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-) / B(D^0 \rightarrow K^- \pi^+)$.

$$\Gamma(\bar{D}^0 \eta') / \Gamma(\bar{D}^*(2007)^0 \eta') \quad \Gamma_{166} / \Gamma_{175}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.96 ± 0.18 ± 0.06	LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.3 ± 0.8 ± 0.2	AUBERT	04B BABR	Repl. by LEES 11M

$$\Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{176} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
(6.2 ± 1.2 ± 1.8) × 10⁻⁴	1,2 SATPATHY	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² No assumption about the intermediate mechanism is made in the analysis.

$$\Gamma(\bar{D}^*(2007)^0 K^+ \pi^-) / \Gamma(\bar{D}^*(2007)^0 \pi^+ \pi^-) \quad \Gamma_{177} / \Gamma_{176}$$

VALUE (units 10 ⁻²)	DOCUMENT ID	TECN	COMMENT
8.36 ± 0.43 ± 0.61	AAIJ	22N LHCB	pp at 13 TeV

$$\Gamma(\bar{D}^*(2007)^0 K^0) / \Gamma_{\text{total}} \quad \Gamma_{178} / \Gamma$$

VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT
3.6 ± 1.2 ± 0.3		¹ AUBERT,B	06L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.6 90 ¹ KROKOVNY 03 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\bar{D}^*(2007)^0 K^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{179} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.9 × 10⁻⁵	90	¹ KROKOVNY	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\bar{D}^*(2007)^0 \phi) / \Gamma_{\text{total}} \quad \Gamma_{180} / \Gamma$$

VALUE (units 10 ⁻⁶)	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.5 ± 0.3	¹ AAIJ	23AZ LHCB	pp at 7, 8, 13 TeV

¹ The last uncertainty includes the uncertainties of the branching fraction $B(B^0 \rightarrow \bar{D}^0 K^+ K^-)$ and $B(\phi \rightarrow K^+ K^-)$.

$$\Gamma(D^*(2007)^0 K^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{181} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.0 × 10⁻⁵	90	¹ KROKOVNY	03 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{182} / \Gamma$$

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
2.7 ± 0.5 OUR AVERAGE			
2.60 ± 0.47 ± 0.37	¹ MAJUMDER	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
3.0 ± 0.7 ± 0.6	¹ EDWARDS	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^*(2007)^0 \pi^+ \pi^+ \pi^- \pi^-) / \Gamma(D^*(2010)^- \pi^+ \pi^+ \pi^- \pi^0) \quad \Gamma_{182} / \Gamma_{67}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.04 ± 0.02	¹ EDWARDS	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(D^*(2010)^+ D^*(2010)^-) / \Gamma_{\text{total}} \quad \Gamma_{183} / \Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
8.0 ± 0.6 OUR AVERAGE				
7.82 ± 0.38 ± 0.63		¹ KRONENBIT...12	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
8.1 ± 0.6 ± 1.0		¹ AUBERT,B	06A BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
9.9 $^{+4.2}_{-3.3}$ ± 1.2		¹ LIPELES	00 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.1 ± 0.8 ± 1.1		¹ MIYAKE	05 BELL	Repl. by KRONENBIT-TER 12
8.3 ± 1.6 ± 1.2		^{1,2} AUBERT	02M BABR	Repl. by AUBERT,B 06B
6.2 $^{+4.0}_{-2.9}$ ± 1.0		³ ARTUSO	99 CLE2	Repl. by LIPELES 00
<61	90	⁴ BARATE	98Q ALEP	$e^+ e^- \rightarrow Z$
<22	90	⁵ ASNER	97 CLE2	Repl. by ARTUSO 99

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AUBERT 02M also assumes the measured CP -odd fraction of the final states is $0.22 \pm 0.18 \pm 0.03$.

³ ARTUSO 99 uses $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = (48 \pm 4)\%$.

⁴ BARATE 98Q (ALEPH) observes 2 events with an expected background of 0.10 ± 0.03 which corresponds to a branching ratio of $(2.3^{+1.9}_{-1.2} \pm 0.4) \times 10^{-3}$.

⁵ ASNER 97 at CLEO observes 1 event with an expected background of 0.022 ± 0.011 . This corresponds to a branching ratio of $(5.3^{+7.1}_{-3.7} \pm 1.0) \times 10^{-4}$.

$$\Gamma(\bar{D}^*(2007)^0 \omega) / \Gamma_{\text{total}} \quad \Gamma_{184} / \Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
3.6 ± 1.1 OUR AVERAGE				Error includes scale factor of 3.1.
4.55 ± 0.24 ± 0.39		¹ LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
2.29 ± 0.39 ± 0.40		¹ BLYTH	06 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2 ± 0.7 ± 0.9	90	¹ AUBERT	04B BABR	Repl. by LEES 11M
< 7.9	90	¹ ABE	02J BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
< 7.4	90	² NEMAT1	98 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
<21	90	³ ALAM	94 CLE2	Repl. by NEMAT1 98

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² NEMAT1 98 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the PDG 96 values for D^0 , D^{*0} , η , η' , and ω branching fractions.

³ ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0) / B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- 2\pi^+ \pi^-) / B(D^0 \rightarrow K^- \pi^+)$.

$$\Gamma(\bar{D}^0 \omega) / \Gamma(\bar{D}^*(2007)^0 \omega) \quad \Gamma_{167} / \Gamma_{184}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.58 ± 0.06 OUR AVERAGE			
0.56 ± 0.04 ± 0.04	LEES	11M BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

1.04±0.20±0.17 BLYTH 06 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.7 ±0.1 ±0.1 AUBERT 04B BABR Repl. by LEES 11M

$\Gamma(D^*(2010)^+ D^-)/\Gamma_{\text{total}}$ Γ_{185}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
6.1±1.5 OUR AVERAGE		Error includes scale factor of 1.6.		
5.7±0.7±0.7		¹ AUBERT,B	06A BABR	$e^+e^- \rightarrow \Upsilon(4S)$
11.7±2.6 ^{+2.2} _{-2.5}		^{1,2} ABE	02Q BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.8±1.0±1.3		¹ AUBERT	03J BABR	Repl. by AUBERT,B 06B
14.8±3.8 ^{+2.8} _{-3.1}		^{1,3} ABE	02Q BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 6.3	90	¹ LIPELES	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<56	90	BARATE	98Q ALEP	$e^+e^- \rightarrow Z$
<18	90	ASNER	97 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The measurement is performed using fully reconstructed D^* and D^+ decays.

³ The measurement is performed using a partial reconstruction technique for the D^* and fully reconstructed D^+ decays as a cross check.

$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$ Γ_{186}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.9	90	¹ AUBERT,B	06A BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<270	90	BARATE	98Q ALEP	$e^+e^- \rightarrow Z$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- D^0 K^+)/\Gamma_{\text{total}}$ Γ_{187}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.07±0.07±0.09	¹ DEL-AMO-SA...11B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.7 ±0.3 ±0.3	¹ AUBERT	03X BABR	Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- D^*(2007)^0 K^+)/\Gamma_{\text{total}}$ Γ_{188}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.46±0.18±0.37	¹ DEL-AMO-SA...11B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.6 ±0.7 ±0.7	¹ AUBERT	03X BABR	Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^*(2010)^- D^0 K^+)/\Gamma_{\text{total}}$ Γ_{189}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.47±0.10±0.18	¹ DEL-AMO-SA...11B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.1 $^{+0.4}_{-0.3} \pm 0.4$ ¹ AUBERT 03X BABR Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^{*(2010)-} D^0 K^+)/\Gamma(D^- D^0 K^+)$ $\Gamma_{189}/\Gamma_{187}$

VALUE	DOCUMENT ID	TECN	COMMENT
1.754 ± 0.028 ± 0.038	¹ AAIJ	20AN LHCb	pp at 7, 8, 13 TeV

¹ Uses $D^+ \rightarrow K^- \pi^+ \pi^+$, $D^0 \rightarrow K^- \pi^+$ and $D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-$ decays.

$\Gamma(D^{*(2010)-} D^{*(2007)^0} K^+)/\Gamma_{\text{total}}$ Γ_{190}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
10.6 ± 0.33 ± 0.86	¹ DEL-AMO-SA...11B	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.8 ± 1.0 ± 1.7 ¹ AUBERT 03X BABR Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^- D^+ K^0)/\Gamma_{\text{total}}$ Γ_{191}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
0.75 ± 0.12 ± 0.12		¹ DEL-AMO-SA...11B	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7 90 ¹ AUBERT 03X BABR Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$[\Gamma(D^{*(2010)-} D^+ K^0) + \Gamma(D^- D^{*(2010)+} K^0)]/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.41 ± 0.36 ± 0.39	¹ DEL-AMO-SA...11B	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.5 ± 1.2 ± 1.0 ¹ AUBERT 03X BABR Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(D^{*(2010)-} D^{*(2010)+} K^0)/\Gamma_{\text{total}}$ Γ_{193}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.7 OUR AVERAGE			

8.26 ± 0.43 ± 0.67 ¹ DEL-AMO-SA...11B BABR $e^+ e^- \rightarrow \Upsilon(4S)$

6.8 ± 0.8 ± 1.4 ^{1,2} DALSENO 07 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

8.8 ± 0.8 ± 1.4 ^{1,2} AUBERT,B 06Q BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.8 $^{+1.5}_{-1.4} \pm 1.3$ ¹ AUBERT 03X BABR Repl. by AUBERT,B 06Q

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The result is rescaled by a factor of 2 to convert from K_S^0 to K^0 .

$\Gamma(D^{*-} D_{s1}(2536)^+, D_{s1}^+ \rightarrow D^{*+} K^0)/\Gamma_{\text{total}}$ Γ_{194}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.0 ± 2.4 OUR AVERAGE			
$7.6^{+4.8+1.6}_{-4.2-1.4}$	1,2 DALSENO	07 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$8.2 \pm 2.6 \pm 1.2$	1,2 AUBERT,B	06Q BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² The result is rescaled by a factor of 2 to convert from K_S^0 to K^0 .

 $\Gamma(\bar{D}^0 D^0 K^0)/\Gamma_{\text{total}}$ Γ_{195}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$0.27 \pm 0.10 \pm 0.05$		¹ DEL-AMO-SA...11B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.4	90	¹ AUBERT	03X BABR	Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(D^0 \bar{D}^0 K^+ \pi^-)/\Gamma(D^{*}(2010)^- D^0 K^+)$ $\Gamma_{196}/\Gamma_{189}$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$14.2 \pm 1.1 \pm 1.0$	¹ AAIJ	20AG LHCB	pp at 7, 8, and 13 TeV
¹ AAIJ 20AG excluded contributions from $B^0 \rightarrow D^{*-} D^0 K^+$ transitions with $D^{*-} \rightarrow \bar{D}^0 \pi^-$.			

 $[\Gamma(\bar{D}^0 D^{*}(2007)^0 K^0) + \Gamma(\bar{D}^{*}(2007)^0 D^0 K^0)]/\Gamma_{\text{total}}$ Γ_{197}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$1.08 \pm 0.32 \pm 0.36$		¹ DEL-AMO-SA...11B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3.7	90	¹ AUBERT	03X BABR	Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\bar{D}^{*}(2007)^0 D^{*}(2007)^0 K^0)/\Gamma_{\text{total}}$ Γ_{198}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
$2.40 \pm 0.55 \pm 0.67$		¹ DEL-AMO-SA...11B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<6.6	90	¹ AUBERT	03X BABR	Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma((\bar{D} + \bar{D}^*)(D + D^*)K)/\Gamma_{\text{total}}$ Γ_{199}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$3.68 \pm 0.10 \pm 0.24$	¹ DEL-AMO-SA...11B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$4.3 \pm 0.3 \pm 0.6$	¹ AUBERT	03X BABR	Repl. by DEL-AMO-SANCHEZ 11B

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta_c K^0)/\Gamma_{\text{total}}$ Γ_{200}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.90 ± 0.11 OUR AVERAGE			
$1.04^{+0.18}_{-0.15} \pm 0.13$	¹ CHILIKIN	19	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$0.62^{+0.21}_{-0.20} \pm 0.05$	^{2,3} AUBERT	07AV	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$0.91 \pm 0.16 \pm 0.05$	^{2,4} AUBERT,B	04B	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$1.23 \pm 0.23^{+0.40}_{-0.41}$	² FANG	03	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$1.09^{+0.55}_{-0.42} \pm 0.33$	⁵ EDWARDS	01	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ CHILIKIN 19 reports $[\Gamma(B^0 \rightarrow \eta_c K^0)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow p\bar{p}\pi^+\pi^-)] = (38.0^{+6.4+1.3}_{-2.9-4.7}) \times 10^{-7}$ which we divide by our best value $B(\eta_c(1S) \rightarrow p\bar{p}\pi^+\pi^-) = (3.7 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AUBERT 07AV reports $[\Gamma(B^0 \rightarrow \eta_c K^0)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow p\bar{p})] = (0.83^{+0.28}_{-0.26} \pm 0.05) \times 10^{-6}$ which we divide by our best value $B(\eta_c(1S) \rightarrow p\bar{p}) = (1.33 \pm 0.11) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AUBERT,B 04B reports $[\Gamma(B^0 \rightarrow \eta_c K^0)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow K\bar{K}\pi)] = (0.0648 \pm 0.0085 \pm 0.0071) \times 10^{-3}$ which we divide by our best value $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = (7.1 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma\eta_c)$ in those modes have been accounted for.

$\Gamma(\eta_c K^0)/\Gamma(J/\psi(1S)K^0)$ $\Gamma_{200}/\Gamma_{214}$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.39 \pm 0.20 \pm 0.45$	¹ AUBERT,B	04B	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses BABAR measurement of $B(B^0 \rightarrow J/\psi K^0) = (8.5 \pm 0.5 \pm 0.6) \times 10^{-4}$.

$\Gamma(\eta_c(1S)K^+\pi^-)/\Gamma(J/\psi(1S)K^+\pi^-)$ $\Gamma_{201}/\Gamma_{215}$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.57 \pm 0.03 \pm 0.05$	¹ AAIJ	18AN	LHCB pp at 7, 8, 13 TeV

¹ AAIJ 18AN reports $[\Gamma(B^0 \rightarrow \eta_c(1S)K^+\pi^-)/\Gamma(B^0 \rightarrow J/\psi(1S)K^+\pi^-)] \times [B(\eta_c(1S) \rightarrow p\bar{p})] / [B(J/\psi(1S) \rightarrow p\bar{p})] = 0.357 \pm 0.015 \pm 0.008$ which we multiply or divide by our best values $B(\eta_c(1S) \rightarrow p\bar{p}) = (1.33 \pm 0.11) \times 10^{-3}$, $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.120 \pm 0.029) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\eta_c(1S)K^*(1410)^0)/\Gamma(\eta_c(1S)K^+\pi^-)$ $\Gamma_{204}/\Gamma_{201}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$32 \pm 24 \pm 6$	¹ AAIJ	18AN	LHCB pp at 7, 8, 13 TeV

¹ AAIJ 18AN reports $[\Gamma(B^0 \rightarrow \eta_c(1S)K^*(1410)^0)/\Gamma(B^0 \rightarrow \eta_c(1S)K^+\pi^-)] \times [B(K^*(1410) \rightarrow K\pi)] = 0.021 \pm 0.011 \pm 0.011$ which we divide by our best value $B(K^*(1410) \rightarrow K\pi) = (6.6 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\eta_c(1S)K^+\pi^-(NR))/\Gamma(\eta_c(1S)K^+\pi^-) \quad \Gamma_{202}/\Gamma_{201}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$10.3 \pm 1.4^{+1.0}_{-1.2}$	AAIJ	18AN LHCB	pp at 7, 8, 13 TeV

$$\Gamma(\eta_c(1S)K_0^*(1430)^0)/\Gamma(\eta_c(1S)K^+\pi^-) \quad \Gamma_{205}/\Gamma_{201}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$27 \pm 5 \pm 3$	¹ AAIIJ	18AN LHCB	pp at 7, 8, 13 TeV

¹ AAIIJ 18AN reports $[\Gamma(B^0 \rightarrow \eta_c(1S)K_0^*(1430)^0)/\Gamma(B^0 \rightarrow \eta_c(1S)K^+\pi^-)] \times [B(K_0^*(1430) \rightarrow K\pi)] = 0.253 \pm 0.035^{+0.035}_{-0.028}$ which we divide by our best value $B(K_0^*(1430) \rightarrow K\pi) = (93 \pm 10) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\eta_c(1S)K_2^*(1430)^0)/\Gamma(\eta_c(1S)K^+\pi^-) \quad \Gamma_{206}/\Gamma_{201}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$8.2^{+3.6}_{-4.4} \pm 0.2$	¹ AAIIJ	18AN LHCB	pp at 7, 8, 13 TeV

¹ AAIIJ 18AN reports $[\Gamma(B^0 \rightarrow \eta_c(1S)K_2^*(1430)^0)/\Gamma(B^0 \rightarrow \eta_c(1S)K^+\pi^-)] \times [B(K_2^*(1430) \rightarrow K\pi)] = 0.041 \pm 0.015^{+0.010}_{-0.016}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\eta_c(1S)K^*(1680)^0)/\Gamma(\eta_c(1S)K^+\pi^-) \quad \Gamma_{207}/\Gamma_{201}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$5.7^{+6.5}_{-6.8} \pm 0.4$	¹ AAIIJ	18AN LHCB	pp at 7, 8, 13 TeV

¹ AAIIJ 18AN reports $[\Gamma(B^0 \rightarrow \eta_c(1S)K^*(1680)^0)/\Gamma(B^0 \rightarrow \eta_c(1S)K^+\pi^-)] \times [B(K^*(1680) \rightarrow K\pi)] = 0.022 \pm 0.020^{+0.015}_{-0.017}$ which we divide by our best value $B(K^*(1680) \rightarrow K\pi) = (38.7 \pm 2.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\eta_c(1S)K_0^*(1950)^0)/\Gamma(\eta_c(1S)K^+\pi^-) \quad \Gamma_{208}/\Gamma_{201}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$7^{+4}_{-6} \pm 2$	¹ AAIIJ	18AN LHCB	pp at 7, 8, 13 TeV

¹ AAIIJ 18AN reports $[\Gamma(B^0 \rightarrow \eta_c(1S)K_0^*(1950)^0)/\Gamma(B^0 \rightarrow \eta_c(1S)K^+\pi^-)] \times [B(K_0^*(1950) \rightarrow K^-\pi^+)] = 0.038 \pm 0.018^{+0.014}_{-0.025}$ which we divide by our best value $B(K_0^*(1950) \rightarrow K^-\pi^+) = (52 \pm 14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(T_{c\bar{c}}(4100)^-K^+, T_{c\bar{c}}^- \rightarrow \eta_c\pi^-)/\Gamma(\eta_c(1S)K^+\pi^-) \quad \Gamma_{203}/\Gamma_{201}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$3.3 \pm 1.1^{+1.2}_{-1.1}$	AAIJ	18AN LHCB	pp at 7, 8, 13 TeV

$\Gamma(\eta_c K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{209}/Γ

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

5.3 $^{+0.8}_{-0.9}$ OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.

4.42 ± 0.24 $^{+0.54}_{-0.66}$	¹ AAIJ	18AN LHCB	$p\bar{p}$ at 7, 8, 13 TeV
$6.8 \pm 0.9 \pm 0.4$	^{2,3} AUBERT	08AB BABR	$e^+e^- \rightarrow \Upsilon(4S)$
7.7 $^{+2.4}_{-2.2}$ ± 0.7	^{4,5} AUBERT	07AV BABR	$e^+e^- \rightarrow \Upsilon(4S)$
16.2 ± 3.2 $^{+5.5}_{-6.0}$	⁵ FANG	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AAIJ 18AN reports $B(B^0 \rightarrow \eta_c K^*(892)^0, K^*(892)^0 \rightarrow K^+\pi^-) = (2.95 \pm 0.16^{+0.36}_{-0.44}) \times 10^{-4}$ using the fitted fraction of $0.514 \pm 0.019^{+0.017}_{-0.048}$ from Dalitz decay of $B(B^0 \rightarrow \eta_c K^+\pi^-) = (5.73 \pm 0.24 \pm 0.67) \times 10^{-4}$ and corrected for $B(K^*(892)^0 \rightarrow K^+\pi^-) = 2/3$.


² AUBERT 08AB reports $[\Gamma(B^0 \rightarrow \eta_c K^*(892)^0)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \eta_c K^+)] = 0.62 \pm 0.06 \pm 0.05$ which we multiply by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.10 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Uses the production ratio of $(B^+B^-)/(B^0\bar{B}^0) = 1.026 \pm 0.032$ at $\Upsilon(4S)$.

⁴ AUBERT 07AV reports $[\Gamma(B^0 \rightarrow \eta_c K^*(892)^0)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow p\bar{p})] = (1.03^{+0.27}_{-0.24} \pm 0.17) \times 10^{-6}$ which we divide by our best value $B(\eta_c(1S) \rightarrow p\bar{p}) = (1.33 \pm 0.11) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

WEIGHTED AVERAGE
5.3+0.8-0.9 (Error scaled by 1.7)



$$\Gamma(\eta_c(2S) K_S^0, \eta_c \rightarrow p \bar{p} \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{210} / \Gamma$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$4.2^{+1.4+0.3}_{-1.2-0.3}$	CHILIKIN	19	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$$\Gamma(\eta_c(2S) K^{*0}) / \Gamma_{\text{total}} \quad \Gamma_{211} / \Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.9	90	¹ AUBERT	08AB BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses the production ratio of $(B^+ B^-) / (B^0 \bar{B}^0) = 1.026 \pm 0.032$ at $\Upsilon(4S)$.

$$\Gamma(h_c(1P) K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{212} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-5}$	CHILIKIN	19	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$$\Gamma(B^0 \rightarrow h_c(1P) K^{*0}) / \Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \quad \Gamma_{213} / \Gamma \times \Gamma_{30}^{h_c(1P)} / \Gamma_{h_c(1P)}$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	¹ AUBERT	08AB BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses the production ratio of $(B^+ B^-) / (B^0 \bar{B}^0) = 1.026 \pm 0.032$ at $\Upsilon(4S)$.

$$\Gamma(\eta_c K^{*(892)^0}) / \Gamma(\eta_c K^0) \quad \Gamma_{209} / \Gamma_{200}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.33 \pm 0.36^{+0.24}_{-0.33}$	FANG	03	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

$$\Gamma(J/\psi(1S) K^0) / \Gamma_{\text{total}} \quad \Gamma_{214} / \Gamma$$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
8.91 ± 0.21 OUR FIT					
8.91 ± 0.21 OUR AVERAGE					
9.02 ± 0.10 ± 0.26			¹ CHOUDHURY	21	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
8.1 ± 0.9 ± 0.6			² CHILIKIN	19	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
8.8 $^{+1.4}_{-1.3}$ ± 0.1			^{3,4} AUBERT	07AV BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
8.69 ± 0.22 ± 0.30			⁴ AUBERT	05J	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
9.5 ± 0.8 ± 0.6			⁴ AVERY	00	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
11.5 ± 2.3 ± 1.7			⁵ ABE	96H	CDF $p \bar{p}$ at 1.8 TeV
6.93 ± 4.07 ± 0.04			⁶ BORTOLETTO	092	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
9.24 ± 7.21 ± 0.05			⁷ ALBRECHT	90J	ARG $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.9 ± 0.4 ± 0.9			⁴ ABE	03B	BELL Repl. by CHOUDHURY 21
8.3 ± 0.4 ± 0.5			⁴ AUBERT	02	BABR Repl. by AUBERT 05J
8.5 $^{+1.4}_{-1.2}$ ± 0.6			⁴ JESSOP	97	CLE2 Repl. by AVERY 00
7.5 ± 2.4 ± 0.8			⁶ ALAM	94	CLE2 Sup. by JESSOP 97
<50	90		ALAM	86	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

¹ CHOUDHURY 21 uses the relative production fraction of charged (f^{+-}) to neutral (f^{00}) B mesons at $\Upsilon(4S)$ value of $f^{+-} / f^{00} = 1.058 \pm 0.024$.

² CHILIKIN 19 reports $[\Gamma(B^0 \rightarrow J/\psi(1S)K^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow p\bar{p}\pi^+\pi^-)] = (48.6^{+4.6+2.4}_{-4.4-2.6}) \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}\pi^+\pi^-) = (6.0 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AUBERT 07AV reports $[\Gamma(B^0 \rightarrow J/\psi(1S)K^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow p\bar{p})] = (1.87^{+0.28}_{-0.26} \pm 0.07) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.120 \pm 0.029) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁵ ABE 96H assumes that $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.

⁶ BORTOLETTO 92 reports $(6 \pm 3 \pm 2) \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)K^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁷ ALBRECHT 90J reports $(8 \pm 6 \pm 2) \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)K^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{215}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
1.15 ± 0.05 OUR AVERAGE				
1.15 $\pm 0.01 \pm 0.05$		CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow J/\psi K^- \pi^+$
1.16 $\pm 0.56 \pm 0.01$		¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.079 ± 0.011		² AUBERT 09AA	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
<1.3	90	³ ALBRECHT 87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<6.3	90	GILES 84	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ BORTOLETTO 92 reports $(1.0 \pm 0.4 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)K^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Does not report systematic uncertainties.

³ ALBRECHT 87D assume $B^+B^-/B^0\bar{B}^0$ ratio is 55/45. $K\pi$ system is specifically selected as nonresonant.

$\Gamma(J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{216}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.27 ± 0.05 OUR FIT				
1.28 ± 0.05 OUR AVERAGE				
1.19 $\pm 0.01 \pm 0.08$		CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow J/\psi K^- \pi^+$
1.33 $^{+0.22}_{-0.21} \pm 0.02$		^{1,2} AUBERT 07AV	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

1.309 ± 0.026 ± 0.077		² AUBERT	05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
1.29 ± 0.05 ± 0.13		² ABE	02N	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
1.74 ± 0.20 ± 0.18		³ ABE	98O	CDF	$p\bar{p}$ 1.8 TeV
1.32 ± 0.17 ± 0.17		⁴ JESSOP	97	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
1.27 ± 0.65 ± 0.01		⁵ BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
1.27 ± 0.60 ± 0.01	6	⁶ ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
4.04 ± 1.81 ± 0.02	5	⁷ BEBEK	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.24 ± 0.05 ± 0.09		² AUBERT	02	BABR	Repl. by AUBERT 05J
1.36 ± 0.27 ± 0.22		⁸ ABE	96H	CDF	Sup. by ABE 98O
1.69 ± 0.31 ± 0.18	29	⁹ ALAM	94	CLE2	Sup. by JESSOP 97
		¹⁰ ALBRECHT	94G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
4.0 ± 0.30		¹¹ ALBAJAR	91E	UA1	$E_{\text{cm}}^{p\bar{p}} = 630$ GeV
3.3 ± 0.18	5	¹² ALBRECHT	87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
4.1 ± 0.18	5	¹³ ALAM	86	CLEO	Repl. by BEBEK 87

¹ AUBERT 07AV reports $[\Gamma(B^0 \rightarrow J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow p\bar{p})] = (2.82^{+0.30+0.36}_{-0.28-0.35}) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow p\bar{p}) = (2.120 \pm 0.029) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ABE 98O reports $[B(B^0 \rightarrow J/\psi(1S)K^*(892)^0)]/[B(B^+ \rightarrow J/\psi(1S)K^+)] = 1.76 \pm 0.14 \pm 0.15$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁵ BORTOLETTO 92 reports $(1.1 \pm 0.5 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁶ ALBRECHT 90J reports $(1.1 \pm 0.5 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁷ BEBEK 87 reports $(3.5 \pm 1.6 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)K^*(892)^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$, which we rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BORTOLETTO 92 to use the same assumptions.

⁸ ABE 96H assumes that $B(B^+ \rightarrow J/\psi K^+) = (1.02 \pm 0.14) \times 10^{-3}$.

⁹ The neutral and charged B events together are predominantly longitudinally polarized, $\Gamma_L/\Gamma = 0.080 \pm 0.08 \pm 0.05$. This can be compared with a prediction using HQET, 0.73 (KRAMER 92). This polarization indicates that the $B \rightarrow \psi K^*$ decay is dominated by the $CP = -1$ CP eigenstate. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

¹⁰ ALBRECHT 94G measures the polarization in the vector-vector decay to be predominantly longitudinal, $\Gamma_T/\Gamma = 0.03 \pm 0.16 \pm 0.15$ making the neutral decay a CP eigenstate when the K^*0 decays through $K_S^0 \pi^0$.

¹¹ ALBAJAR 91E assumes B_d^0 production fraction of 36%.

¹² ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

¹³ ALAM 86 assumes $B^\pm/B^0$ ratio is 60/40. The observation of the decay $B^+ \rightarrow J/\psi K^*(892)^+$ (HAAS 85) has been retracted in this paper.

$\Gamma(J/\psi(1S)K^*(892)^0)/\Gamma(J/\psi(1S)K^0)$

$\Gamma_{216}/\Gamma_{214}$

VALUE	DOCUMENT ID	TECN	COMMENT
1.50 ± 0.09 OUR AVERAGE			
1.51 ± 0.05 ± 0.08	AUBERT	05J	BABR $e^+e^- \rightarrow \Upsilon(4S)$
1.39 ± 0.36 ± 0.10	ABE	96Q	CDF $p\bar{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.49 ± 0.10 ± 0.08	¹ AUBERT	02	BABR Repl. by AUBERT 05J
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			

$\Gamma(J/\psi(1S)\eta K_S^0)/\Gamma_{\text{total}}$

Γ_{217}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
5.4 ± 0.9 OUR AVERAGE			
5.22 ± 0.78 ± 0.49	¹ IWASHITA	14	BELL $e^+e^- \rightarrow \Upsilon(4S)$
8.4 ± 2.6 ± 2.7	¹ AUBERT	04Y	BABR $e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			

$\Gamma(J/\psi(1S)\eta' K_S^0)/\Gamma_{\text{total}}$

Γ_{218}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	90	¹ XIE	07	BELL $e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				

$\Gamma(J/\psi(1S)\omega K^0)/\Gamma_{\text{total}}$

Γ_{220}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.3 ± 0.3	¹ DEL-AMO-SA..10B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.1 ± 0.6 ± 0.3	¹ AUBERT	08W	BABR Repl. by DEL-AMO-SANCHEZ 10B
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			

$\Gamma(\chi_{c0}(3915), \chi_{c0} \rightarrow J/\psi\omega)/\Gamma_{\text{total}}$

Γ_{221}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
2.1 ± 0.9 ± 0.3	¹ DEL-AMO-SA..10B	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.3 ^{+1.3} _{-1.1} ± 0.2	^{1,2} AUBERT	08W	BABR Repl. by DEL-AMO-SANCHEZ 10B
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			
² Corresponds to upper limit of 3.9×10^{-5} at 90% CL.			

$\Gamma(J/\psi(1S)\phi K^0)/\Gamma_{\text{total}}$

Γ_{219}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.9 ± 1.0 OUR AVERAGE	Error	includes scale factor of 1.3.	
4.43 ± 0.76 ± 0.19	LEES	15	BABR $e^+e^- \rightarrow \Upsilon(4S)$

$10.2 \pm 3.8 \pm 1.0$	¹ AUBERT	030	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$8.8^{+3.5}_{-3.0} \pm 1.3$	² ANASTASSOV 00	CLE2		$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ANASTASSOV 00 finds 10 events on a background of 0.5 ± 0.2 . Assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$, a uniform Dalitz plot distribution, isotropic $J/\psi(1S)$ and ϕ decays, and $B(B^+ \rightarrow J/\psi(1S)\phi K^+) = B(B^0 \rightarrow J/\psi(1S)\phi K^0)$.

$\Gamma(J/\psi(1S)K(1270)^0)/\Gamma_{\text{total}}$ Γ_{222}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.30 \pm 0.34 \pm 0.32$	¹ ABE	01L	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the PDG value of $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.00 \pm 0.10) \times 10^{-3}$.

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{223}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
1.65 ± 0.08 OUR AVERAGE				
$1.65 \pm 0.10 \pm 0.09$		¹ AAIJ	24N	LHCB pp at 7, 8, 13 TeV
$1.62 \pm 0.11 \pm 0.06$		² PAL	18	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$1.69 \pm 0.14 \pm 0.07$		² AUBERT	08AU	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$2.5^{+1.1}_{-0.9} \pm 0.2$		² AVERY	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.94 \pm 0.22 \pm 0.17$		² AUBERT,B	06B	BABR	Repl. by AUBERT 08AU
$2.3 \pm 0.5 \pm 0.2$		² ABE	03B	BELL	Repl. by PAL 18
$2.0 \pm 0.6 \pm 0.2$		² AUBERT	02	BABR	Repl. by AUBERT,B 06B
< 32	90	³ ACCIARRI	97C	L3	
< 5.8	90	BISHAI	96	CLE2	Sup. by AVERY 00
< 690	90	² ALEXANDER	95	CLE2	Sup. by BISHAI 96

¹ AAIJ 24N reports $[\Gamma(B^0 \rightarrow J/\psi(1S)\pi^0)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow J/\psi(1S)K^*(892)^+)] = (1.153 \pm 0.053 \pm 0.048) \times 10^{-2}$ which we multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^*(892)^+) = (1.43 \pm 0.08) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ ACCIARRI 97C assumes B^0 production fraction $(39.5 \pm 4.0)\%$ and B_S $(12.0 \pm 3.0)\%$.

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ Γ_{224}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
10.8 ± 2.3 OUR AVERAGE				Error includes scale factor of 1.5.
$7.3 \pm 2.5 \pm 1.3$		¹ AAIJ	15D	LHCB pp at 7, 8 TeV
$12.3^{+1.8}_{-1.7} \pm 0.7$		^{2,3} CHANG	12	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.5 \pm 1.7 \pm 0.8$		³ CHANG	07A	BELL	Repl. by CHANG 12
< 27	90	³ AUBERT	030	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 1200	90	⁴ ACCIARRI	97C	L3	

¹ AAIJ 15D reports $[\Gamma(B^0 \rightarrow J/\psi(1S)\eta)/\Gamma_{\text{total}}] / [B(B_S^0 \rightarrow J/\psi(1S)\eta)] = (1.85 \pm 0.61 \pm 0.14) \times 10^{-2}$ which we multiply by our best value $B(B_S^0 \rightarrow J/\psi(1S)\eta) =$

$(4.0 \pm 0.7) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Reconstructs η in $\gamma\gamma$ and $\pi^+\pi^-\pi^0$ decays.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ ACCIARRI 97C assumes B^0 production fraction $(39.5 \pm 4.0\%)$ and B_S $(12.0 \pm 3.0\%)$.

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{225}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
3.99 ± 0.15 OUR AVERAGE			
$3.98 \pm 0.14 \pm 0.07$	^{1,2} AAIJ	13M LHCb	pp at 7 TeV
$4.6 \pm 0.7 \pm 0.6$	³ AUBERT	03B BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AAIJ 13M reports $(3.97 \pm 0.09 \pm 0.11 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow J/\psi(1S)K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.018 \pm 0.042) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AAIJ 13M does not report correlations between various measurements of the $J/\psi\pi\pi$ final state.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)\pi^+\pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{226}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ AUBERT	07AC BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S)f_0(500), f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_{227}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.8 \pm 0.5^{+1.1}_{-1.5}$	¹ AAIJ	14X LHCb	pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.4^{+2.5}_{-1.1} \pm 0.2$	^{2,3} AAIJ	13M LHCb	Repl. by AAIJ 14X
-----------------------------	---------------------	----------	-------------------

¹ AAIJ 14X uses Dalitz plot analysis of $B^0 \rightarrow J/\psi\pi^+\pi^-$.

² AAIJ 13M reports $(6.4 \pm 0.8^{+2.4}_{-0.8}) \times 10^{-6}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)f_0(500), f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(B^0 \rightarrow J/\psi(1S)\pi^+\pi^-) = (3.97 \pm 0.09 \pm 0.11 \pm 0.16) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow J/\psi(1S)\pi^+\pi^-) = (3.99 \pm 0.15) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AAIJ 13M does not report correlations between various measurements of the $J/\psi\pi\pi$ final state. Measured in Dalitz plot like analysis of $B^0 \rightarrow J/\psi\pi^+\pi^-$.

$\Gamma(J/\psi(1S)f_2)/\Gamma_{\text{total}}$ Γ_{228}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$0.33^{+0.05}_{-0.06}$ OUR AVERAGE				Error includes scale factor of 1.5.

$0.30 \pm 0.03^{+0.02}_{-0.03}$	¹ AAIJ	14X LHCb	pp at 7, 8 TeV
---------------------------------	-------------------	----------	------------------

$0.42 \pm 0.06 \pm 0.02$	^{2,3} AAIJ	13M LHCb	pp at 7 TeV
--------------------------	---------------------	----------	---------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.5	90	^{4,5} AUBERT	07AC BABR	$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	-----------------------	-----------	-----------------------------------

¹ AAIJ 14X uses Dalitz plot analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$.

² AAIJ 13M reports $[\Gamma(B^0 \rightarrow J/\psi(1S) f_2)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (3.5 \pm 0.4 \pm 0.4) \times 10^{-6}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S) f_2)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] / [B(B^0 \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(B^0 \rightarrow J/\psi(1S) \pi^+ \pi^-) = (3.97 \pm 0.09 \pm 0.11 \pm 0.16) \times 10^{-5}$, which we rescale to our best values $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$, $B(B^0 \rightarrow J/\psi(1S) \pi^+ \pi^-) = (3.99 \pm 0.15) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

³ AAIJ 13M does not report correlations between various measurements of the $J/\psi \pi \pi$ final state. Measured in Dalitz plot like analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$.

⁴ AUBERT 07AC reports $[\Gamma(B^0 \rightarrow J/\psi(1S) f_2)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] < 0.46 \times 10^{-5}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = 84.3 \times 10^{-2}$.

⁵ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(J/\psi(1S) \rho^0)/\Gamma_{\text{total}}$ Γ_{229}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$2.55^{+0.18}_{-0.16}$ OUR AVERAGE

$2.50 \pm 0.10^{+0.18}_{-0.15}$	¹ AAIJ	14X	LHCB	pp at 7, 8 TeV
---------------------------------	-------------------	-----	------	------------------

$2.7 \pm 0.3 \pm 0.2$	² AUBERT	07AC	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	---------------------	------	------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.51^{+0.22}_{-0.23} \pm 0.10$	^{3,4} AAIJ	13M	LHCB	Repl. by AAIJ 14X
---------------------------------	---------------------	-----	------	-------------------

$1.6 \pm 0.6 \pm 0.4$	² AUBERT	03B	BABR	Repl. by AUBERT 07AC
-----------------------	---------------------	-----	------	----------------------

< 25	90	BISHAI	96	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
--------	----	--------	----	---

¹ AAIJ 14X uses Dalitz plot analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$. We assume $B(\rho(770)^0 \rightarrow \pi^+ \pi^-) = 100\%$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AAIJ 13M reports $(2.49^{+0.20+0.16}_{-0.13-0.23}) \times 10^{-5}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S) \rho^0)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(B^0 \rightarrow J/\psi(1S) \pi^+ \pi^-) = (3.97 \pm 0.09 \pm 0.11 \pm 0.16) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow J/\psi(1S) \pi^+ \pi^-) = (3.99 \pm 0.15) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AAIJ 13M does not report correlations between various measurements of the $J/\psi \pi \pi$ final state. Measured in Dalitz plot like analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$. Assumes $B(\rho(770)^0 \rightarrow \pi\pi) = 100\%$.

$\Gamma(J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{230}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 1.1 \times 10^{-6}$	90	¹ AAIJ	13M	LHCB pp at 7 TeV
---	----	-------------------	-----	--------------------

¹ AAIJ 13M does not provide correlations between various measurements of the $J/\psi \pi^+ \pi^-$ final state. The measurements were obtained from a Dalitz plot like analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$. Also reports $\Gamma(J/\psi(1S) f_0(980), f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}} = (6.1^{+3.1+1.7}_{-2.0-1.4}) \times 10^{-6}$.

$\Gamma(J/\psi(1S)\rho(1450)^0, \rho^0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_{231}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

 $2.9^{+1.6}_{-0.7}$ OUR AVERAGE

$4.6 \pm 1.1 \pm 1.9$	¹ AAIJ	14X	LHCB	pp at 7, 8 TeV
$2.1^{+2.4}_{-0.7} \pm 0.1$	^{2,3} AAIJ	13M	LHCB	pp at 7 TeV

¹ AAIJ 14X uses Dalitz plot analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$.

² AAIJ 13M reports $(2.1^{+1.0+2.2}_{-0.6-0.4}) \times 10^{-6}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)\rho(1450)^0, \rho^0 \rightarrow \pi\pi)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(B^0 \rightarrow J/\psi(1S)\pi^+\pi^-) = (3.97 \pm 0.09 \pm 0.11 \pm 0.16) \times 10^{-5}$, which we rescale to our best value $B(B^0 \rightarrow J/\psi(1S)\pi^+\pi^-) = (3.99 \pm 0.15) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ AAIJ 13M does not report correlations between various measurements of the $J/\psi \pi \pi$ final state. Measured in Dalitz plot like analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$.

 $\Gamma(J/\psi \rho(1700)^0, \rho^0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{232}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.0 \pm 0.5 \pm 1.2$	¹ AAIJ	14X	LHCB	pp at 7, 8 TeV
---	-------------------	-----	------	------------------

¹ AAIJ 14X uses Dalitz plot analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$.

 $\Gamma(J/\psi(1S)\omega)/\Gamma_{\text{total}}$ Γ_{233}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$1.8^{+0.7}_{-0.5} \pm 0.1$		¹ AAIJ	14X	LHCB	pp at 7, 8 TeV
---	--	-------------------	-----	------	------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<27 90 BISHAI 96 CLE2 $e^+e^- \rightarrow \gamma(4S)$

¹ AAIJ 14X reports $[\Gamma(B^0 \rightarrow J/\psi(1S)\omega)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-)] = (2.7^{+0.8+0.7}_{-0.6-0.5}) \times 10^{-7}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-) = (1.53 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(J/\psi(1S)\omega)/\Gamma(J/\psi(1S)\rho^0)$ $\Gamma_{233}/\Gamma_{229}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.61^{+0.39}_{-0.21} \pm 0.05$	^{1,2} AAIJ	13M	LHCB	pp at 7 TeV
---	---------------------	-----	------	---------------

¹ AAIJ 13M reports $0.61^{+0.24+0.31}_{-0.14-0.16}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)\omega)/\Gamma(B^0 \rightarrow J/\psi(1S)\rho^0)] \times [B(\omega(782) \rightarrow \pi^+\pi^-)]$ assuming $B(\omega(782) \rightarrow \pi^+\pi^-) = (1.53^{+0.11}_{-0.13}) \times 10^{-2}$, which we rescale to our best value $B(\omega(782) \rightarrow \pi^+\pi^-) = (1.53 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AAIJ 13M does not report correlations between various measurements of the $J/\psi \pi \pi$ final state. Measured in Dalitz plot like analysis of $B^0 \rightarrow J/\psi \pi^+ \pi^-$. Assumes $B(\rho(770)^0 \rightarrow \pi\pi) = 100\%$.

 $\Gamma(J/\psi(1S)\omega)/\Gamma(J/\psi(1S)\rho^0)$ $\Gamma_{233}/\Gamma_{229}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

$0.89 \pm 0.19^{+0.07}_{-0.13}$	AAIJ	13A	LHCB	pp at 7 TeV
---	------	-----	------	---------------

$\Gamma(J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$ Γ_{234}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$2.53 \pm 0.35 \pm 0.05$	¹ AAIJ	13BT LHCB	pp at 7 TeV

¹ AAIJ 13BT reports $(2.53 \pm 0.31 \pm 0.19) \times 10^{-6}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow J/\psi(1S)K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.018 \pm 0.042) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(J/\psi(1S)a_0(980), a_0 \rightarrow K^+K^-)/\Gamma_{\text{total}}$ Γ_{235}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.470 \pm 0.331 \pm 0.072$	¹ AAIJ	13BT LHCB	pp at 7 TeV

¹ AAIJ 13BT uses $B(\bar{B}^0 \rightarrow J/\psi K^+K^-) = (2.53 \pm 0.31 \pm 0.19) \times 10^{-6}$ to derive this result. It also reports the equivalent upper limit of $< 9.0 \times 10^{-7}$ at 90% CL.

 $\Gamma(J/\psi(1S)\phi)/\Gamma_{\text{total}}$ Γ_{236}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.1 \times 10^{-7}$	90	AAIJ	21K LHCB	pp at 7, 8, 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 10.1 \times 10^{-7}$	90	LEES	15 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$< 1.9 \times 10^{-7}$	90	¹ AAIJ	13BT LHCB	pp at 7 TeV
$< 9.4 \times 10^{-7}$	90	² LIU	08I BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$< 9.2 \times 10^{-6}$	90	² AUBERT	03O BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AAIJ 13BT uses $B(B^0 \rightarrow J/\psi(1S)K^+K^-) = (2.53 \pm 0.31 \pm 0.19) \times 10^{-6}$ and $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ to obtain this result.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(J/\psi(1S)\eta'(958))/\Gamma_{\text{total}}$ Γ_{237}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$7.6 \pm 2.2 \pm 1.0$		¹ AAIJ	15D LHCB	pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 7.4	90	^{2,3} CHANG	12 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 63	90	³ AUBERT	03O BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AAIJ 15D reports $[\Gamma(B^0 \rightarrow J/\psi(1S)\eta'(958))/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow J/\psi(1S)\eta')] = (2.28 \pm 0.65 \pm 0.16) \times 10^{-2}$ which we multiply by our best value $B(B_s^0 \rightarrow J/\psi(1S)\eta') = (3.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Reconstructs $\eta'(985)$ in $(\eta\pi^+\pi^-)$ and $\rho(770)$ decays.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\eta'(958))$ $\Gamma_{224}/\Gamma_{237}$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.111 \pm 0.475 \pm 0.062$	¹ AAIJ	15D LHCB	pp at 7, 8 TeV

¹ Uses $J/\psi \rightarrow \mu^+\mu^-$, $\eta' \rightarrow \rho^0\gamma$, and $\eta' \rightarrow \eta\pi^+\pi^-$ decays.

$$\Gamma(J/\psi(1S)K^0\pi^+\pi^-)/\Gamma(J/\psi(1S)K^0) \quad \Gamma_{238}/\Gamma_{214}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.50 ± 0.04 OUR AVERAGE			
0.493 ± 0.034 ± 0.027	AAIJ	14L	LHCB pp at 7 TeV
1.24 ± 0.40 ± 0.15	AFFOLDER	02B	CDF $p\bar{p}$ 1.8 TeV

$$\Gamma(J/\psi(1S)K^0K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_{240}/\Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
25 ± 7 OUR AVERAGE			Error includes scale factor of 1.8.
34.9 ± 6.7 ± 1.5	LEES	15	BABR $e^+e^- \rightarrow \gamma(4S)$
20.2 ± 4.3 ± 1.9	¹ AAIJ	14L	LHCB pp at 7 TeV

¹ Measured with $B(B^0 \rightarrow J/\psi K_S^0 K^+ K^-) / B(B^0 \rightarrow J/\psi K_S^0)$ using PDG 12 for the involved branching fractions.

$$\Gamma(J/\psi(1S)K^0K^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{239}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 21 × 10⁻⁶	90	¹ AAIJ	14L	LHCB pp at 7 TeV

¹ Measured with $B(B^0 \rightarrow J/\psi K_S^0 K^\pm \pi^\mp) / B(B^0 \rightarrow J/\psi K_S^0 \pi^+ \pi^-)$ using PDG 12 values for the involved branching fractions.

$$\Gamma(J/\psi(1S)K^0\rho^0)/\Gamma_{\text{total}} \quad \Gamma_{242}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
5.4 ± 2.9 ± 0.9	¹ AFFOLDER	02B	CDF $p\bar{p}$ 1.8 TeV

¹ Uses $B^0 \rightarrow J/\psi(1S)K_S^0$ decay as a reference and $B(B^0 \rightarrow J/\psi(1S)K^0) = 8.3 \times 10^{-4}$.

$$\Gamma(J/\psi(1S)K^*(892)^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{243}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
7.7 ± 4.1 ± 1.3	¹ AFFOLDER	02B	CDF $p\bar{p}$ 1.8 TeV

¹ Uses $B^0 \rightarrow J/\psi(1S)K_S^0$ decay as a reference and $B(B^0 \rightarrow J/\psi(1S)K^0) = 8.3 \times 10^{-4}$.

$$\Gamma(J/\psi(1S)\pi^+\pi^-\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{244}/\Gamma_{225}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.361 ± 0.017 ± 0.021	¹ AAIJ	14Y	LHCB pp at 7, 8 TeV

¹ Excludes contributions from $\psi(2S)$ and $\chi_{c1}(3872)$ decaying to $J/\psi(1S)\pi^+\pi^-$.

$$\Gamma(J/\psi(1S)f_1(1285))/\Gamma_{\text{total}} \quad \Gamma_{245}/\Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
8.4^{+2.1}_{-2.0} ± 0.5	¹ AAIJ	14Y	LHCB pp at 7, 8 TeV

¹ AAIJ 14Y reports $(8.37 \pm 1.95^{+0.71}_{-0.66} \pm 0.35) \times 10^{-6}$ from a measurement of $[\Gamma(B^0 \rightarrow J/\psi(1S)f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow 2\pi^+2\pi^-)]$ assuming $B(f_1(1285) \rightarrow 2\pi^+2\pi^-) = 0.11^{+0.007}_{-0.006}$, which we rescale to our best value $B(f_1(1285) \rightarrow 2\pi^+2\pi^-) = (10.9 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(J/\psi(1S)K^*(892)^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{246}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$6.6 \pm 1.9 \pm 1.1$		¹ AFFOLDER	02B CDF	$p\bar{p}$ 1.8 TeV

¹ Uses $B^0 \rightarrow J/\psi(1S)K^*(892)^0$ decay as a reference and $B(B^0 \rightarrow J/\psi(1S)K^0) = 12.4 \times 10^{-4}$.

 $\Gamma(\eta_{c2}(1D)K_S^0, \eta_{c2} \rightarrow h_c\gamma)/\Gamma_{\text{total}}$ Γ_{247}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.5 \times 10^{-5}$	90	CHILIKIN	20 BELL	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(\eta_{c2}(1D)\pi^-K^+, \eta_{c2} \rightarrow h_c\gamma)/\Gamma_{\text{total}}$ Γ_{248}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.0 \times 10^{-4}$	90	CHILIKIN	20 BELL	$e^+e^- \rightarrow \gamma(4S)$

 $\Gamma(\chi_{c1}(3872)K^0)/\Gamma_{\text{total}}$ Γ_{251}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
1.1 ± 0.4 OUR AVERAGE				
$2.5^{+0.9}_{-0.8} \pm 0.8$		¹ HIRATA	23 BELL	$e^+e^- \rightarrow \gamma(4S)$
$0.93 \pm 0.27 \pm 0.29$		² CHOI	11 BELL	$e^+e^- \rightarrow \gamma(4S)$
$1.2 \pm 0.6^{+0.4}_{-0.5}$		³ DEL-AMO-SA..10B	BABR	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.5	90	^{4,5} BHARDWAJ	11 BELL	$e^+e^- \rightarrow \gamma(4S)$
$2.1 \pm 1.0 \pm 0.7$		^{5,6} AUSHEV	10 BELL	$e^+e^- \rightarrow \gamma(4S)$
< 5	90	^{7,8} AUBERT	09B BABR	$e^+e^- \rightarrow \gamma(4S)$
$4.8 \pm 2.4 \pm 1.7$		^{5,7,9} AUBERT	08B BABR	$e^+e^- \rightarrow \gamma(4S)$
< 1.4	90	¹⁰ AUBERT	08Y BABR	$e^+e^- \rightarrow \gamma(4S)$
< 2.4	90	¹¹ AUBERT	06 BABR	Repl. by AUBERT 08Y
$3.0 \pm 1.4^{+1.6}_{-1.5}$		^{12,13} GOKHROO	06 BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ HIRATA 23 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872)K^0)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = 1.34^{+0.47+0.10}_{-0.40-0.12}$ which we multiply by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CHOI 11 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872)K^0)/\Gamma_{\text{total}}] / [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = 0.50 \pm 0.14 \pm 0.04$ which we multiply by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ DEL-AMO-SANCHEZ 10B reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872)K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))] = (6 \pm 3 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S)) = (5.0 \pm 1.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ BHARDWAJ 11 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872)K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \gamma J/\psi)] < 2.4 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \gamma J/\psi) = 10 \times 10^{-3}$.

⁵ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

⁶ AUSHEV 10 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872)K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \bar{D}^{*0}D^0)] = (0.97 \pm 0.46 \pm 0.13) \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow$

$\bar{D}^{*0} D^0 = (46 \pm 16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷ Uses $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.6 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = (48.4 \pm 0.6)\%$.

⁸ AUBERT 09B reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \gamma J/\psi)] < 4.9 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \gamma J/\psi) = 10 \times 10^{-3}$.

⁹ AUBERT 08B reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)] = (2.22 \pm 1.05 \pm 0.42) \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0) = (46 \pm 16) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰ AUBERT 08Y reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 6.0 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 4.3 \times 10^{-2}$.

¹¹ AUBERT 06 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 10.3 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 4.3 \times 10^{-2}$.

¹² GOKHROO 06 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)] = (1.66 \pm 0.70^{+0.32}_{-0.37}) \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0) = (55 \pm 28) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹³ Measure the near-threshold enhancements in the $(D^0 \bar{D}^0 \pi^0)$ system at a mass $3875.2 \pm 0.7^{+0.3}_{-1.6} \pm 0.8$ MeV/ c^2 .

$\Gamma(\chi_{c1}(3872)^- K^+)/\Gamma_{\text{total}}$ Γ_{249}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5 \times 10^{-4}$	90	¹ AUBERT	06E	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Perform measurements of absolute branching fractions using a missing mass technique.

$\Gamma(\chi_{c1}(3872)^- K^+, \chi_{c1}(3872)^- \rightarrow J/\psi(1S) \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{250}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.2	90	^{1,2} CHOI	11	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.4	90	^{2,3} AUBERT	05B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
---------	----	-----------------------	-----	---

¹ Assumes $\pi^+ \pi^0$ originates from ρ^+ .

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ The isovector- X hypothesis is excluded with a likelihood test at 1×10^{-4} level.

$\Gamma(\chi_{c1}(3872) K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{252}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$0.92 \pm 0.35 \pm 0.29$		¹ BALA	15	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.9	90	^{2,3} AUBERT	09B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
		^{3,4} AUBERT	09B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ BALA 15 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^*(892)^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] = (4.0 \pm 1.5 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = (4.3 \pm 1.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² AUBERT 09B reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^*(892)^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \gamma J/\psi)] < 2.8 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \gamma J/\psi) = 10 \times 10^{-3}$.

³ Uses $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.6 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = (48.4 \pm 0.6)\%$.

⁴ AUBERT 09B reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^*(892)^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \gamma \psi(2S))] < 4.4 \times 10^{-6}$ at 90% CL.

$\Gamma(\chi_{c1}(3872) K^+ \pi^-)/\Gamma_{\text{total}}$

Γ_{253}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.8 \pm 0.3 \pm 0.6$	1,2 BALA 15	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ BALA 15 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) K^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = (4.3 \pm 1.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1}(3872) \gamma)/\Gamma_{\text{total}}$

Γ_{254}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.2 \times 10^{-5}$	90	1,2 CHOU 19	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at $\Upsilon(4S)$.

² CHOU 19 reports $[\Gamma(B^0 \rightarrow \chi_{c1}(3872) \gamma)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 5.1 \times 10^{-7}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 4.3 \times 10^{-2}$.

$\Gamma(T_{c\bar{c}1}(4430)^\pm K^\mp, T_{c\bar{c}1}^\pm \rightarrow \psi(2S) \pi^\pm)/\Gamma_{\text{total}}$

Γ_{255}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$6.0^{+1.7+2.5}_{-2.0-1.4}$		CHILIKIN 13	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.1	95	¹ AUBERT	09AA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$3.2^{+1.8+5.3}_{-0.9-1.6}$		¹ MIZUK	09 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$4.1 \pm 1.0 \pm 1.4$		1,2 CHOI	08 BELL	Repl. by MIZUK 09

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Establishes the $(Z_c(4430))^+$ with a significance of 6.5 sigma. Needs confirmation.

$\Gamma(T_{c\bar{c}1}(4430)^\pm K^\mp, T_{c\bar{c}1}^\pm \rightarrow J/\psi \pi^\pm)/\Gamma_{\text{total}}$

Γ_{256}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$5.4^{+4.0+1.1}_{-1.0-0.6}$		CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4	95	¹ AUBERT	09AA BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	---------------------	-----------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(T_{c\bar{c}1}(3900)^\pm K^\mp, T_{c\bar{c}1}^\pm \rightarrow J/\psi \pi^\pm)/\Gamma_{\text{total}}$

Γ_{257}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$< 9 \times 10^{-7}$	CHILIKIN 14	BELL	$\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

$$\Gamma(T_{c\bar{c}1}(4200)^{\pm} K^{\mp}, T_{c\bar{c}1}^{\pm} \rightarrow J/\psi \pi^{\pm})/\Gamma_{\text{total}} \quad \Gamma_{258}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$2.2^{+0.7+1.1}_{-0.5-0.6}$		CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

$$\Gamma(T_{c\bar{c}1}(3900)^{\pm} K^{\mp}, T_{c\bar{c}1}^{\pm} \rightarrow J/\psi \pi^{\pm})/\Gamma(J/\psi(1S)K^{*}(892)^0) \quad \Gamma_{257}/\Gamma_{216}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-2}$	90	ABAZOV	18B	D0 $p\bar{p}$ at 1.96 TeV

$$\Gamma(J/\psi(1S)p\bar{p})/\Gamma_{\text{total}} \quad \Gamma_{259}/\Gamma$$

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
$4.51 \pm 0.40 \pm 0.44$		¹ AAIJ	19U	LHCB pp at 7, 8, 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.2	90	² AAIJ	13Z	LHCB Repl. by AAIJ 19U
< 8.3	90	³ XIE	05	BELL $e^+e^- \rightarrow \Upsilon(4S)$
< 19	90	³ AUBERT	03K	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Measured relative to $B_S^0 \rightarrow J/\psi \phi$ assuming $B(B_S^0 \rightarrow J/\psi \phi) = (10.5 \pm 0.13 \pm 0.64) \times 10^{-4}$ and taking into account small K^+K^- S -wave contribution. Measurement assumes $f_s/f_d = 0.259 \pm 0.015$ for 7, 8 TeV data and f_s/f_d multiplied by 1.068 ± 0.046 for 13 TeV data.

² Uses $B(B_S^0 \rightarrow J/\psi(1S)\pi^+\pi^-) = (1.98 \pm 0.20) \times 10^{-4}$.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(J/\psi(1S)\gamma)/\Gamma_{\text{total}} \quad \Gamma_{260}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	¹ AAIJ	15BB	LHCB pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	² AUBERT,B	04T	BABR $e^+e^- \rightarrow \Upsilon(4S)$
--------	----	-----------------------	-----	--

¹ Branching fractions of normalization modes $B^0 \rightarrow J/\psi \gamma X$ taken from PDG 14. Uses $f_s/f_d = 0.259 \pm 0.015$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(J/\psi \mu^+ \mu^-, J/\psi \rightarrow \mu^+ \mu^-)/\Gamma_{\text{total}} \quad \Gamma_{261}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-9}$	95	AAIJ	22Q	LHCB pp at 7, 8, 13 TeV

$$\Gamma(J/\psi(1S)\bar{D}^0)/\Gamma_{\text{total}} \quad \Gamma_{262}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-7}$	90	AAIJ	24F	LHCB pp at 7, 8, 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.3 \times 10^{-5}$	90	¹ AUBERT	05U	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$<2.0 \times 10^{-5}$	90	¹ ZHANG	05B	BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\psi(2S)\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{263}/\Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.17 \pm 0.17 \pm 0.08$	¹ CHOBANOVA 16	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\psi(2S)K^0)/\Gamma_{\text{total}}$ Γ_{264}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
5.8 ± 0.5 OUR FIT				
5.8 ± 0.5 OUR AVERAGE				
4.7 ± 0.7 ± 0.7		¹ AAIJ	14L LHCb	pp at 7 TeV
6.46 ± 0.65 ± 0.51		² AUBERT	05J BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.7 ± 1.1		² ABE	03B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
5.0 ± 1.1 ± 0.6		² RICHICHI	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.9 ± 1.1 ± 1.1		² AUBERT	02 BABR	Repl. by AUBERT 05J
< 8	90	² ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 15	90	² BORTOLETTO	92 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 28	90	² ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Measured with $B(B^0 \rightarrow \psi(2S)K_S^0) \times B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) / B(B^0 \rightarrow J/\psi K_S^0)$ using PDG 12 values for the involved branching fractions.				
² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				

 $\Gamma(\psi(2S)K^0\pi^+\pi^-)/\Gamma(\psi(2S)K^0)$ $\Gamma_{265}/\Gamma_{264}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.480 ± 0.013 ± 0.032	TUMASYAN	22Al CMS	pp at 13 TeV

 $\Gamma(\psi(2S)K^0)/\Gamma(J/\psi(1S)K^0)$ $\Gamma_{264}/\Gamma_{214}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.82 ± 0.13 ± 0.12	¹ AUBERT	02 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.			

 $\Gamma(\psi(3770)K^0, \psi \rightarrow \bar{D}^0 D^0)/\Gamma_{\text{total}}$ Γ_{266}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.23	90	¹ AUBERT	08B BABR	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				

 $\Gamma(\psi(3770)K^0, \psi \rightarrow D^- D^+)/\Gamma_{\text{total}}$ Γ_{267}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.88	90	¹ AUBERT	08B BABR	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.				

 $\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ $\Gamma_{268}/\Gamma_{225}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.56 ± 0.07 ± 0.05	¹ AAIJ	13AA LHCb	pp at 7 TeV
¹ Assuming lepton universality for dimuon decay modes of J/ψ and $\psi(2S)$ mesons, the ratio $B(J/\psi \rightarrow \mu^+ \mu^-)/B(\psi(2S) \rightarrow \mu^+ \mu^-) = B(J/\psi \rightarrow e^+ e^-)/B(\psi(2S) \rightarrow e^+ e^-) = 7.69 \pm 0.19$ was used.			

 $\Gamma(\psi(2S)K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{269}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
5.80 ± 0.39		^{1,2} CHILIKIN	13 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.57 ± 0.16		³ AUBERT	09AA BABR	$e^+e^- \rightarrow \Upsilon(4S)$

$5.68 \pm 0.13 \pm 0.42$		² MIZUK	09	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
<10	90	² ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
¹ Combines measurements with $\psi(2S) \rightarrow \ell^+\ell^-$ with measurement from MIZUK 09 which uses $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$.					
² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.					
³ Does not report systematic uncertainties.					

$\Gamma(\psi(2S)K^*(892)^0)/\Gamma_{\text{total}}$	Γ_{270}/Γ
VALUE (units 10^{-4})	CL%
5.9 ± 0.4 OUR FIT	

$6.0^{+0.5}_{-0.7}$ OUR AVERAGE Error includes scale factor of 1.1.

$5.55^{+0.22+0.41}_{-0.23-0.84}$		¹ CHILIKIN	13	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$6.49 \pm 0.59 \pm 0.97$		¹ AUBERT	05J	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$7.6 \pm 1.1 \pm 1.0$		¹ RICHICHI	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
$9.0 \pm 2.2 \pm 0.9$		² ABE	980	CDF	$p\bar{p}$ 1.8 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$5.52^{+0.35+0.53}_{-0.32-0.58}$		¹ MIZUK	09	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
<19	90	¹ ALAM	94	CLE2	Repl. by RICHICHI 01
14	$\pm 8 \pm 4$	¹ BORTOLETTO	92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<23	90	¹ ALBRECHT	90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 980 reports $[B(B^0 \rightarrow \psi(2S)K^*(892)^0)]/[B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.908 \pm 0.194 \pm 0.10$. We multiply by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (9.9 \pm 1.0) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\psi(2S)K^*(892)^0)/\Gamma(J/\psi(1S)K^*(892)^0)$	$\Gamma_{270}/\Gamma_{216}$
VALUE	DOCUMENT ID
$0.487 \pm 0.018 \pm 0.014$	^{1,2} AAIJ
	12L LHCb pp at 7 TeV

¹ AAIJ 12L reports $0.476 \pm 0.014 \pm 0.010 \pm 0.012$ from a measurement of $[\Gamma(B^0 \rightarrow \psi(2S)K^*(892)^0)/\Gamma(B^0 \rightarrow J/\psi(1S)K^*(892)^0)] \times [B(J/\psi(1S) \rightarrow e^+e^-)] / [B(\psi(2S) \rightarrow e^+e^-)]$ assuming $B(J/\psi(1S) \rightarrow e^+e^-) = (5.94 \pm 0.06) \times 10^{-2}$, $B(\psi(2S) \rightarrow e^+e^-) = (7.72 \pm 0.17) \times 10^{-3}$, which we rescale to our best values $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032) \times 10^{-2}$, $B(\psi(2S) \rightarrow e^+e^-) = (7.94 \pm 0.22) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² Assumes $B(J/\psi \rightarrow \mu^+\mu^-) / B(\psi(2S) \rightarrow \mu^+\mu^-) = B(J/\psi \rightarrow e^+e^-) / B(\psi(2S) \rightarrow e^+e^-) = 7.69 \pm 0.19$.

$\Gamma(\psi(2S)K^*(892)^0)/\Gamma(\psi(2S)K^0)$	$\Gamma_{270}/\Gamma_{264}$
VALUE	DOCUMENT ID
1.02 ± 0.10 OUR FIT	
$1.00 \pm 0.14 \pm 0.09$	AUBERT
	05J BABR $e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\chi_{c0}K^0)/\Gamma_{\text{total}}$	Γ_{271}/Γ
VALUE (units 10^{-6})	CL%
$195^{+42}_{-36} \pm 11$	¹ AAIJ
	18F LHCb pp at 7, 8 TeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

$145^{+103}_{-85} \pm 8$	2,3 LEES	12I BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$148 \pm 30 \pm 13$	2,4 LEES	12O BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$142^{+55}_{-44} \pm 22$	2,5 AUBERT	09AU BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 113	90 ⁵ GARMASH	07 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 1240	90 ² AUBERT	05K BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 500	90 ⁶ EDWARDS	01 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ AAIJ 18F uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used. We compute $B(B^0 \rightarrow \chi_{c0} K^0)$ using the PDG value $B(\chi_{c0} \rightarrow \pi\pi) = (8.51 \pm 0.33) \times 10^{-3}$ and 2/3 for the $\pi^+ \pi^-$ fraction. Our first error is their experiment's error and the second error is systematic error from using our best value.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ LEES 12I reports $[\Gamma(B^0 \rightarrow \chi_{c0} K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)] = (0.46^{+0.25}_{-0.17} \pm 0.21) \times 10^{-6}$ which we divide by our best value $B(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0) = (3.18 \pm 0.19) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Measured in the $B^0 \rightarrow K_S^0 K^+ K^-$ decay.

⁵ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

⁶ EDWARDS 01 assumes equal production of B^0 and B^+ at the $\Upsilon(4S)$. The correlated uncertainties (28.3)% from $B(J/\psi(1S) \rightarrow \gamma \eta_c)$ in those modes have been accounted for.

$\Gamma(\chi_{c0} K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{272}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
$1.7 \pm 0.3 \pm 0.2$		¹ AUBERT	08BD BABR	$e^+e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •

< 7.7	90	¹ AUBERT	05K BABR	Repl. by AUBERT 08BD
---------	----	---------------------	----------	----------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1} \pi^0)/\Gamma_{\text{total}}$ Γ_{273}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.12 \pm 0.25 \pm 0.12$	¹ KUMAR	08 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\chi_{c1} K^0)/\Gamma_{\text{total}}$ Γ_{274}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
3.95 ± 0.27 OUR AVERAGE				
15 $^{+5}_{-4} \pm 6$		¹ CHILIKIN	19 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$3.78^{+0.17}_{-0.16} \pm 0.33$		² BHARDWAJ	11 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$4.2 \pm 0.3 \pm 0.3$		³ AUBERT	09B BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$3.1^{+1.5}_{-1.1} \pm 0.1$		⁴ AVERY	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •

$3.51 \pm 0.33 \pm 0.45$	² SONI	06 BELL	Repl. by BHARDWAJ 11
--------------------------	-------------------	---------	----------------------

- 4.53 ± 0.41 ± 0.51 ² AUBERT 05J BABR Repl. by AUBERT 09B
 4.3 ± 1.4 ± 0.2 ⁵ AUBERT 02 BABR Repl. by AUBERT 05J
^{<27} 90 ² ALAM 94 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
- ¹ CHILIKIN 19 reports $[\Gamma(B^0 \rightarrow \chi_{c1} K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^+\pi^-)] = (7.4^{+2.4+0.6}_{-2.0-0.4}) \times 10^{-7}$ which we divide by our best value $B(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^+\pi^-) = (5.0 \pm 1.9) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ³ Uses $\chi_{c1,2} \rightarrow J/\psi\gamma$. Assumes $B(\Upsilon(4S) \rightarrow B^+B^-) = (51.6 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = (48.4 \pm 0.6)\%$.
- ⁴ AVERY 00 reports $(3.9^{+1.9}_{-1.3} \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow \chi_{c1} K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ⁵ AUBERT 02 reports $(5.4 \pm 1.4 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow \chi_{c1} K^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\chi_{c1} K^0)/\Gamma(J/\psi(1S) K^0)$ $\Gamma_{274}/\Gamma_{214}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.53±0.16±0.02	¹ AUBERT 02 BABR		$e^+e^- \rightarrow \Upsilon(4S)$

- ¹ AUBERT 02 reports $0.66 \pm 0.11 \pm 0.17$ from a measurement of $[\Gamma(B^0 \rightarrow \chi_{c1} K^0)/\Gamma(B^0 \rightarrow J/\psi(1S) K^0)] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\chi_{c1} \pi^- K^+)/\Gamma_{\text{total}}$ Γ_{275}/Γ

<i>VALUE</i> (units 10^{-4})	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>
4.97±0.12±0.28	¹ BHARDWAJ 16	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.83 ± 0.10 ± 0.39	¹ MIZUK 08 BELL	Repl. by BHARDWAJ 16
--------------------	------------------------------------	----------------------

- ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\chi_{c1} \pi^- K^+)/\Gamma(J/\psi(1S) K^+ \pi^-)$ $\Gamma_{275}/\Gamma_{215}$

<i>VALUE</i>	<i>DOCUMENT ID</i>	<i>TECN</i>
0.476±0.021±0.018	¹ LEES 12B	BABR

- ¹ LEES 12B reports $0.474 \pm 0.013 \pm 0.026$ from a measurement of $[\Gamma(B^0 \rightarrow \chi_{c1} \pi^- K^+)/\Gamma(B^0 \rightarrow J/\psi(1S) K^+ \pi^-)] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.4 \pm 1.5) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\chi_{c1} K^*(892)^0)/\Gamma_{\text{total}} \quad \Gamma_{276}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

2.38 $\pm$ 0.19 OUR FIT Error includes scale factor of 1.2.

2.22 $^{+0.40}_{-0.31}$ OUR AVERAGE Error includes scale factor of 1.6.

2.5 $\pm$ 0.2 $\pm$ 0.2 ¹ AUBERT 09B BABR $e^+e^- \rightarrow \Upsilon(4S)$

1.73 $^{+0.15+0.34}_{-0.12-0.22}$ ² MIZUK 08 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.14 $\pm$ 0.34 $\pm$ 0.72 ² SONI 06 BELL Repl. by MIZUK 08

3.27 $\pm$ 0.42 $\pm$ 0.64 ² AUBERT 05J BABR Repl. by AUBERT 09B

3.8 $\pm$ 1.3 $\pm$ 0.1 ³ AUBERT 02 BABR Repl. by AUBERT 05J

<21 90 ⁴ ALAM 94 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses $\chi_{c1,2} \rightarrow J/\psi\gamma$. Assumes $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.6 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = (48.4 \pm 0.6)\%$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ AUBERT 02 reports $(4.8 \pm 1.4 \pm 0.9) \times 10^{-4}$ from a measurement of $[\Gamma(B^0 \rightarrow \chi_{c1} K^*(892)^0)/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.273 \pm 0.016$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\chi_{c1} K^*(892)^0)/\Gamma(J/\psi(1S) K^*(892)^0) \quad \Gamma_{276}/\Gamma_{216}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

18.8 $\pm$ 1.5 OUR FIT Error includes scale factor of 1.1.

19.8 $\pm$ 1.1 $\pm$ 1.5 ¹ AAIJ 13AC LHCB pp at 7 TeV

¹ Uses $B(\chi_{c1} \rightarrow J/\psi\gamma) = (34.4 \pm 1.5)\%$.

$$\Gamma(\chi_{c1} K^*(892)^0)/\Gamma(\chi_{c1} K^0) \quad \Gamma_{276}/\Gamma_{274}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.72 $\pm$ 0.11 $\pm$ 0.12 AUBERT 05J BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89 $\pm$ 0.34 $\pm$ 0.17 ¹ AUBERT 02 BABR Repl. by AUBERT 05J

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(T_{c\bar{c}}(4050)^- K^+, T_{c\bar{c}}^- \rightarrow \chi_{c1} \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{277}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

3.0 $^{+1.5+3.7}_{-0.8-1.6}$ ¹ MIZUK 08 BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.8 90 ^{1,2} LEES 12B BABR

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses $\chi_{c1} \rightarrow J/\psi\gamma$ mode. Uses $\chi_{c1} \rightarrow J/\psi\gamma$ mode. Finds a good description of the data without this $B^0 \rightarrow X(4051)^+ K^-$ decay mode in a fit.

$$\Gamma(T_{c\bar{c}}(4250)^- K^+, T_{c\bar{c}}^- \rightarrow \chi_{c1} \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{278} / \Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
$4.0^{+2.3+19.7}_{-0.9-0.5}$		¹ MIZUK	08	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.0 90 ^{1,2} LEES 12B BABR

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses $\chi_{c1} \rightarrow J/\psi \gamma$ mode. Finds a good description of the data without this $B^0 \rightarrow X(4248)^+ K^-$ decay mode in a fit.

$$\Gamma(\chi_{c1} \pi^+ \pi^- K^0) / \Gamma_{\text{total}} \quad \Gamma_{279} / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.16 \pm 0.35 \pm 0.32$	¹ BHARDWAJ	16	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\chi_{c1} \pi^- \pi^0 K^+) / \Gamma_{\text{total}} \quad \Gamma_{280} / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.52 \pm 0.52 \pm 0.24$	¹ BHARDWAJ	16	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\chi_{c2} K^0) / \Gamma_{\text{total}} \quad \Gamma_{281} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-5}$	90	¹ BHARDWAJ	11	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<2.8 \times 10^{-5}$	90	² AUBERT	09B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
$<2.6 \times 10^{-5}$	90	¹ SONI	06	BELL Repl. by BHARDWAJ 11
$<4.1 \times 10^{-5}$	90	¹ AUBERT	05K	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses $\chi_{c1,2} \rightarrow J/\psi \gamma$. Assumes $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.6 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = (48.4 \pm 0.6)\%$.

$$\Gamma(\chi_{c2} K^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{282} / \Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
4.9 ± 1.2 OUR FIT	Error includes scale factor of 1.1.			
$6.6 \pm 1.8 \pm 0.5$		¹ AUBERT	09B	BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7.1 90 ² SONI 06 BELL $e^+ e^- \rightarrow \Upsilon(4S)$
 <3.6 90 ² AUBERT 05K BABR Repl. by AUBERT 09B

¹ Uses $\chi_{c1,2} \rightarrow J/\psi \gamma$. Assumes $B(\Upsilon(4S) \rightarrow B^+ B^-) = (51.6 \pm 0.6)\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = (48.4 \pm 0.6)\%$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(\chi_{c2} K^*(892)^0) / \Gamma(\chi_{c1} K^*(892)^0) \quad \Gamma_{282} / \Gamma_{276}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
20 ± 5 OUR FIT	Error includes scale factor of 1.1.		
$17.1 \pm 5.0 \pm 2.0$	¹ AAIJ	13AC	LHCB pp at 7 TeV

¹ Uses $B(\chi_{c1} \rightarrow J/\psi \gamma) / B(\chi_{c2} \rightarrow J/\psi \gamma) = 1.76 \pm 0.11$.

$\Gamma(\chi_{c2}\pi^-K^+)/\Gamma_{\text{total}}$ Γ_{283}/Γ

VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT
$0.72 \pm 0.09 \pm 0.05$		¹ BHARDWAJ 16	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\chi_{c2}\pi^+\pi^-K^0)/\Gamma_{\text{total}}$ Γ_{284}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.70 \times 10^{-4}$	90	¹ BHARDWAJ 16	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\chi_{c2}\pi^-\pi^0K^+)/\Gamma_{\text{total}}$ Γ_{285}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.74 \times 10^{-4}$	90	¹ BHARDWAJ 16	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

 $\Gamma(\psi(4660)K^0, \psi \rightarrow \Lambda_c^+ \Lambda_c^-)/\Gamma_{\text{total}}$ Γ_{286}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.3 \times 10^{-4}$	90	¹ LI 18D	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes $B(\gamma(4S) \rightarrow B^0 \bar{B}^0) = 48.6 \pm 0.6\%$ and $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = 6.23 \pm 0.33\%$.

 $\Gamma(\psi(4230)^0 K^0, \psi^0 \rightarrow J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{287}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 17 \times 10^{-6}$	90	¹ GARG 19	BELL	$e^+e^- \rightarrow \gamma(4S)$

¹ Assumes equal production of B^0 and B^+ at the $\gamma(4S)$.

 $\Gamma(K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{288}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
20.0 ± 0.4 OUR FIT				
20.0 ± 0.4 OUR AVERAGE				
$20.67 \pm 0.37 \pm 0.62$		ADACHI 24	BELL	$e^+e^- \rightarrow \gamma(4S)$
$20.00 \pm 0.34 \pm 0.60$		¹ DUH 13	BELL	$e^+e^- \rightarrow \gamma(4S)$
$19.1 \pm 0.6 \pm 0.6$		¹ AUBERT 07B	BABR	$e^+e^- \rightarrow \gamma(4S)$
$18.0 \begin{smallmatrix} + 2.3 \\ - 2.1 \end{smallmatrix} \begin{smallmatrix} + 1.2 \\ - 0.9 \end{smallmatrix}$		¹ BORNHEIM 03	CLE2	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$19.9 \pm 0.4 \pm 0.8$		¹ LIN 07A	BELL	Repl. by DUH 13
$18.5 \pm 1.0 \pm 0.7$		¹ CHAO 04	BELL	Repl. by LIN 07A
$17.9 \pm 0.9 \pm 0.7$		¹ AUBERT 02Q	BABR	Repl. by AUBERT 07B
$22.5 \pm 1.9 \pm 1.8$		¹ CASEY 02	BELL	Repl. by CHAO 04
$19.3 \begin{smallmatrix} + 3.4 \\ - 3.2 \end{smallmatrix} \begin{smallmatrix} + 1.5 \\ - 0.6 \end{smallmatrix}$		¹ ABE 01H	BELL	Repl. by CASEY 02
$16.7 \pm 1.6 \pm 1.3$		¹ AUBERT 01E	BABR	Repl. by AUBERT 02Q
< 66	90	² ABE 00C	SLD	$e^+e^- \rightarrow Z$
$17.2 \begin{smallmatrix} + 2.5 \\ - 2.4 \end{smallmatrix} \pm 1.2$		¹ CRONIN-HEN..00	CLE2	Repl. by BORNHEIM 03
$15 \begin{smallmatrix} + 5. \\ - 4 \end{smallmatrix} \pm 1.4$		GODANG 98	CLE2	Repl. by CRONIN-HENNESSY 00

24	$+17$ -11	± 2	³ ADAM	96D	DLPH	$e^+e^- \rightarrow Z$
< 17		90	ASNER	96	CLE2	Sup. by ADAM 96D
< 30		90	⁴ BUSKULIC	96V	ALEP	$e^+e^- \rightarrow Z$
< 90		90	⁵ ABREU	95N	DLPH	Sup. by ADAM 96D
< 81		90	⁶ AKERS	94L	OPAL	$e^+e^- \rightarrow Z$
< 26		90	⁷ BATTLE	93	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 180		90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 90		90	⁸ AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 320		90	AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

³ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

⁴ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

⁵ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

⁶ Assumes $B(Z \rightarrow b\bar{b}) = 0.217$ and B_d^0 (B_s^0) fraction 39.5% (12%).

⁷ BATTLE 93 assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

⁸ Assumes the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

$\Gamma(K^+\pi^-)/\Gamma(K^0\pi^0)$ $\Gamma_{288}/\Gamma_{289}$

VALUE	DOCUMENT ID	TECN	COMMENT
2.16$\pm$0.16$\pm$0.16	LIN	07A	BELL $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.20 $^{+0.50+0.22}_{-0.58-0.32}$	¹ ABE	01H	BELL Repl. by LIN 07A
-----------------------------------	------------------	-----	-----------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$[\Gamma(K^+\pi^-) + \Gamma(\pi^+\pi^-)]/\Gamma_{\text{total}}$ $(\Gamma_{288} + \Gamma_{423})/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
19$\pm$ 6 OUR AVERAGE				

28 $^{+15}_{-10} \pm 20$		¹ ADAM	96D	DLPH $e^+e^- \rightarrow Z$
18 $^{+6+3}_{-5-4}$	17.2	ASNER	96	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

24 $^{+8}_{-7} \pm 2$		² BATTLE	93	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	--	---------------------	----	--

¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

² BATTLE 93 assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(K^0\pi^0)/\Gamma_{\text{total}}$ Γ_{289}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
10.1 $\pm$ 0.4 OUR AVERAGE				

10.73 $\pm$ 0.63 $\pm$ 0.62		¹ ADACHI	24	BELL $e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	--	---------------------	----	--

$9.68 \pm 0.46 \pm 0.50$	² DUH	13	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$10.1 \pm 0.6 \pm 0.4$	² LEES	13D	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$12.8 \begin{smallmatrix} +4.0 & +1.7 \\ -3.3 & -1.4 \end{smallmatrix}$	² BORNHEIM	03	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$8.7 \pm 0.5 \pm 0.6$	² FUJIKAWA	10A	BELL	Repl. by DUH 13
$10.3 \pm 0.7 \pm 0.6$	² AUBERT	08E	BABR	Repl. by LEES 13D
$9.2 \pm 0.7 \pm 0.6$	² LIN	07A	BELL	Repl. by FUJIKAWA 10A
$11.4 \pm 0.9 \pm 0.6$	² AUBERT	05Y	BABR	Repl. by AUBERT 08E
$11.4 \pm 1.7 \pm 0.8$	² AUBERT	04M	BABR	Repl. by AUBERT 05Y
$11.7 \pm 2.3 \begin{smallmatrix} +1.2 \\ -1.3 \end{smallmatrix}$	² CHAO	04	BELL	Repl. by LIN 07A
$8.0 \begin{smallmatrix} +3.3 \\ -3.1 \end{smallmatrix} \pm 1.6$	² CASEY	02	BELL	Repl. by CHAO 04
$16.0 \begin{smallmatrix} +7.2 & +2.5 \\ -5.9 & -2.7 \end{smallmatrix}$	² ABE	01H	BELL	Repl. by CASEY 02
$8.2 \begin{smallmatrix} +3.1 \\ -2.7 \end{smallmatrix} \pm 1.2$	² AUBERT	01E	BABR	Repl. by AUBERT 04M
$14.6 \begin{smallmatrix} +5.9 & +2.4 \\ -5.1 & -3.3 \end{smallmatrix}$	² CRONIN-HEN..00	CLE2		Repl. by BORNHEIM 03
<41	90	GODANG	98	CLE2 Repl. by CRONIN-HENNESSY 00
<40	90	ASNER	96	CLE2 Rep. by GODANG 98

¹ This is the combined result of this analysis (ADACHI 24) and the updated analysis of ADACHI 23E. The individual results are $(10.40 \pm 0.66 \pm 0.60) \times 10^{-6}$ and $(11.15 \pm 0.68 \pm 0.62) \times 10^{-6}$, respectively.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K^0)/\Gamma_{\text{total}}$				Γ_{290}/Γ
VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT	
66 ± 4 OUR AVERAGE	Error includes scale factor of 1.4.			
$68.5 \pm 2.2 \pm 3.1$	¹ AUBERT	09AV	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$58.9 \begin{smallmatrix} +3.6 \\ -3.5 \end{smallmatrix} \pm 4.3$	¹ SCHUEMANN	06	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$89 \begin{smallmatrix} +18 \\ -16 \end{smallmatrix} \pm 9$	¹ RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$66.6 \pm 2.6 \pm 2.8$	¹ AUBERT	07AE	BABR	Repl. by AUBERT 09AV
$67.4 \pm 3.3 \pm 3.2$	¹ AUBERT	05M	BABR	AUBERT 07AE
$60.6 \pm 5.6 \pm 4.6$	¹ AUBERT	03W	BABR	Repl. by AUBERT 05M
$55 \begin{smallmatrix} +19 \\ -16 \end{smallmatrix} \pm 8$	¹ ABE	01M	BELL	Repl. by SCHUEMANN 06
$42 \begin{smallmatrix} +13 \\ -11 \end{smallmatrix} \pm 4$	¹ AUBERT	01G	BABR	Repl. by AUBERT 03W
$47 \begin{smallmatrix} +27 \\ -20 \end{smallmatrix} \pm 9$	BEHRENS	98	CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K^*(892)^0)/\Gamma_{\text{total}}$				Γ_{291}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE				
$2.6 \pm 0.7 \pm 0.2$		¹ SATO	14	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$3.1 \begin{smallmatrix} +0.9 \\ -0.8 \end{smallmatrix} \pm 0.3$		¹ DEL-AMO-SA..10A	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.8 \pm 1.1 \pm 0.5$		¹ AUBERT	07E	BABR	Repl. by DEL-AMO-SANCHEZ 10A
< 2.6	90	¹ SCHUEMANN	07	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 7.6	90	¹ AUBERT,B	04D	BABR	Repl. by AUBERT 07E
< 24	90	¹ RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 39	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K_0^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{292}/Γ

VALUE (units 10^{-6})		DOCUMENT ID	TECN	COMMENT
$6.3 \pm 1.3 \pm 0.9$		¹ DEL-AMO-SA...10A	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta' K_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{293}/Γ

VALUE (units 10^{-6})		DOCUMENT ID	TECN	COMMENT
$13.7^{+3.0}_{-2.9} \pm 1.2$		¹ DEL-AMO-SA...10A	BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^0)/\Gamma_{\text{total}}$ Γ_{294}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$1.23^{+0.27}_{-0.24}$ OUR AVERAGE				
$1.27^{+0.33}_{-0.29} \pm 0.08$		¹ HOI	12	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$1.15^{+0.43}_{-0.38} \pm 0.09$		¹ AUBERT	09AV	BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.9	90	¹ CHANG	07B	BELL	Repl. by HOI 12
< 2.9	90	¹ AUBERT,B	06V	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 2.5	90	¹ AUBERT,B	05K	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 2.0	90	¹ CHANG	05A	BELL	Repl. by CHANG 07B
< 5.2	90	¹ AUBERT	04H	BABR	Repl. by AUBERT,B 05K
< 9.3	90	¹ RICHICHI	00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 33	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K^*(892)^0)/\Gamma_{\text{total}}$ Γ_{295}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
15.9 ± 1.0 OUR AVERAGE				
$15.2 \pm 1.2 \pm 1.0$		¹ WANG	07B	BELL $e^+e^- \rightarrow \Upsilon(4S)$
$16.5 \pm 1.1 \pm 0.8$		¹ AUBERT,B	06H	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$13.8^{+5.5}_{-4.6} \pm 1.6$		¹ RICHICHI	00	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$18.6 \pm 2.3 \pm 1.2$		¹ AUBERT,B	04D	BABR	Repl. by AUBERT,B 06H
< 30	90	BEHRENS	98	CLE2	Repl. by RICHICHI 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\eta K_0^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{296}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$11.0 \pm 1.6 \pm 1.5$	¹ AUBERT,B	06H BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\eta K_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{297}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$9.6 \pm 1.8 \pm 1.1$	¹ AUBERT,B	06H BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\omega K^0)/\Gamma_{\text{total}}$ Γ_{298}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
4.8 ± 0.4 OUR AVERAGE				
$4.5 \pm 0.4 \pm 0.3$		¹ CHOBANOVA 14	BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$5.4 \pm 0.8 \pm 0.3$		¹ AUBERT	07AE BABR	$e^+e^- \rightarrow \Upsilon(4S)$
$10.0^{+5.4}_{-4.2} \pm 1.4$		¹ JESSOP 00	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.2 \pm 1.0 \pm 0.4$		¹ AUBERT,B	06E BABR	Repl. by AUBERT 07AE
$4.4^{+0.8}_{-0.7} \pm 0.4$		¹ JEN	06 BELL	Repl. by CHOBANOVA 14
$5.9^{+1.6}_{-1.3} \pm 0.5$		¹ AUBERT	04H BABR	Repl. by AUBERT,B 06E
$4.0^{+1.9}_{-1.6} \pm 0.5$		¹ WANG	04A BELL	Repl. by JEN 06
<13	90	¹ AUBERT	01G BABR	Repl. by AUBERT 04H
<57	90	¹ BERGFELD	98 CLE2	Repl. by JESSOP 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(a_0(980)^0 K^0, a_0^0 \rightarrow \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{299}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<7.8	90	¹ AUBERT,BE	04 BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of charged and neutral B mesons at $\Upsilon(4S)$.

 $\Gamma(b_1^0 K^0, b_1^0 \rightarrow \omega \pi^0)/\Gamma_{\text{total}}$ Γ_{300}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<7.8	90	¹ AUBERT	08AG BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(a_0(980)^\pm K^\mp, a_0^\pm \rightarrow \eta \pi^\pm)/\Gamma_{\text{total}}$ Γ_{301}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	90	¹ AUBERT	07Y BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	¹ AUBERT,BE	04 BABR	Repl. by AUBERT 07Y
------	----	------------------------	---------	---------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(b_1^- K^+, b_1^- \rightarrow \omega \pi^-)/\Gamma_{\text{total}}$ Γ_{302}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$7.4 \pm 1.0 \pm 1.0$	¹ AUBERT	07BI BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(b_1^0 K^{*0}, b_1^0 \rightarrow \omega \pi^0)/\Gamma_{\text{total}}$ Γ_{303}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.0 \times 10^{-6}$	90	¹ AUBERT	09AF BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(b_1^- K^{*+}, b_1^- \rightarrow \omega \pi^-)/\Gamma_{\text{total}}$ Γ_{304}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.0 \times 10^{-6}$	90	¹ AUBERT	09AF BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(a_0(1450)^\pm K^\mp, a_0^\pm \rightarrow \eta \pi^\pm)/\Gamma_{\text{total}}$ Γ_{305}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	¹ AUBERT	07Y BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_S^0 X^0(\text{Familon}))/\Gamma_{\text{total}}$ Γ_{306}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<53	90	¹ AMMAR	01B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ AMMAR 01B searched for the two-body decay of the B meson to a massless neutral feebly-interacting particle X^0 such as the familon, the Nambu-Goldstone boson associated with a spontaneously broken global family symmetry.

$\Gamma(\omega K^{*}(892)^0)/\Gamma_{\text{total}}$ Γ_{307}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
2.0 ± 0.5 OUR AVERAGE				
$2.2 \pm 0.6 \pm 0.2$		¹ AUBERT	09H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$1.8 \pm 0.7 \pm 0.3$		¹ GOLDENZWE..08	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.2	90	¹ AUBERT,B	06T BABR	Repl. by AUBERT 09H
< 6.0	90	¹ AUBERT	05O BABR	Repl. by AUBERT,B 06T
< 23	90	¹ BERGFELD	98 CLE2	

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega(K\pi)_0^{*0})/\Gamma_{\text{total}}$ Γ_{308}/Γ

$(K\pi)_0^{*0}$ is the total S-wave composed of $K_0^{*}(1430)$ and nonresonant that are described using LASS shape.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$18.4 \pm 1.8 \pm 1.7$	¹ AUBERT	09H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega K_0^{*}(1430)^0)/\Gamma_{\text{total}}$ Γ_{309}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$16.0 \pm 1.6 \pm 3.0$	¹ AUBERT	09H BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(\omega K_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{310}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
10.1$\pm$2.0$\pm$1.1	¹ AUBERT	09H BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\omega K^+\pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{311}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
5.1$\pm$0.7$\pm$0.7	^{1,2} GOLDENZWE..08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² For the $K\pi$ mass range 0.755–1.250 GeV/ c^2 , excluding $K^*(892)$.

 $\Gamma(K^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{312}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
37.8$\pm$3.2 OUR AVERAGE				
38.5 $\pm$ 1.0 $\pm$ 3.9		^{1,2} LEES	11 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
36.6 $^{+4.2}_{-4.3}$ $\pm$ 3.0		¹ CHANG	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

35.7 $^{+2.6}_{-1.5}$ $\pm$ 2.2		¹ AUBERT	08AQ BABR	Repl. by LEES 11
<40	90	¹ ECKHART	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

 $\Gamma(K^+\rho^-)/\Gamma_{\text{total}}$ Γ_{313}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
7.0$\pm$0.9 OUR AVERAGE				
6.6 $\pm$ 0.5 $\pm$ 0.8		^{1,2} LEES	11 BABR	$e^+e^- \rightarrow \Upsilon(4S)$
15.1 $^{+3.4+2.4}_{-3.3-2.6}$		¹ CHANG	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.0 $^{+0.8}_{-1.3}$ $\pm$ 0.6		¹ AUBERT	08AQ BABR	Repl. by LEES 11
7.3 $^{+1.3}_{-1.2}$ $\pm$ 1.3		¹ AUBERT	03T BABR	Repl. by AUBERT 08AQ
<32	90	¹ JESSOP	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
<35	90	ASNER	96 CLE2	Repl. by JESSOP 00

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

 $\Gamma(K^+\rho(1450)^-)/\Gamma_{\text{total}}$ Γ_{314}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
2.4$\pm$1.0$\pm$0.6		^{1,2} LEES	11 BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	¹ AUBERT	08AQ BABR	Repl. by LEES 11
------	----	---------------------	-----------	------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

$\Gamma(K^+\rho(1700)^-)/\Gamma_{\text{total}}$ Γ_{315}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$0.6 \pm 0.6 \pm 0.4$ 1,2 LEES 11 BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.1 90 1 AUBERT 08AQ BABR Repl. by LEES 11

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

 $\Gamma((K^+\pi^-\pi^0)_{\text{nonresonant}})/\Gamma_{\text{total}}$ Γ_{316}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$2.8 \pm 0.5 \pm 0.4$ 1,2 LEES 11 BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.4 \pm 0.9 \pm 0.5$ 1 AUBERT 08AQ BABR Repl. by LEES 11

<9.4 90 1 CHANG 04 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays. The quoted value is only for the flat part of the non-resonant component.

 $\Gamma((K\pi)_0^{*+}\pi^-, (K\pi)_0^{*+} \rightarrow K^+\pi^0)/\Gamma_{\text{total}}$ Γ_{317}/Γ

$(K\pi)_0^{*+}$ is the total S-wave composed of $K_0^*(1430)$ and nonresonant that are described using LASS shape.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$34.2 \pm 2.4 \pm 4.1$ 1,2 LEES 11 BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.4^{+1.1+2.3}_{-1.3-2.1}$ 1 AUBERT 08AQ BABR Repl. by LEES 11

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

 $\Gamma((K\pi)_0^{*0}\pi^0, (K\pi)_0^{*0} \rightarrow K^+\pi^-)/\Gamma_{\text{total}}$ Γ_{318}/Γ

$(K\pi)_0^{*0}$ is the total S-wave composed of $K_0^*(1430)$ and nonresonant that are described using LASS shape.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$8.6 \pm 1.1 \pm 1.3$ 1,2 LEES 11 BABR $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$8.7^{+1.1+2.8}_{-0.9-2.6}$ 1 AUBERT 08AQ BABR Repl. by LEES 11

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

 $\Gamma(K_2^*(1430)^0\pi^0)/\Gamma_{\text{total}}$ Γ_{319}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<4.0 90 1 AUBERT 08AQ BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(1680)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{320}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<7.5	90	¹ AUBERT	08AQ BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K_x^{*0} \pi^0)/\Gamma_{\text{total}}$ Γ_{321}/Γ

K_x^{*0} stands for the possible candidates of $K^*(1410)$, $K_0^*(1430)$ and $K_2^*(1430)$.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.1^{+1.6+0.5}_{-1.5-0.6}$	¹ CHANG	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{322}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
49.7 ± 1.8 OUR FIT				
49.6 ± 2.0 OUR AVERAGE				
$50.2 \pm 1.5 \pm 1.8$		¹ AUBERT	09AU BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$47.5 \pm 2.4 \pm 3.7$		² GARMASH	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
$50^{+10}_{-9} \pm 7$		¹ ECKHART	02 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$43.0 \pm 2.3 \pm 2.3$		¹ AUBERT	06I BABR	Repl. by AUBERT 09AU
$43.7 \pm 3.8 \pm 3.4$		¹ AUBERT,B	04O BABR	Repl. by AUBERT 06I
$45.4 \pm 5.2 \pm 5.9$		¹ GARMASH	04 BELL	Repl. by GARMASH 07
<440	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.


 $\Gamma(K^0 \pi^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{323}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$13.9^{+2.6}_{-1.8}$ OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
$12.1 \pm 0.6 \pm 2.9$	¹ AAIJ	18F LHCB	pp at 7, 8 TeV
$11.1^{+2.5}_{-1.0} \pm 0.9$	² AUBERT	09AU BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$19.9 \pm 2.5^{+1.7}_{-2.0}$	³ GARMASH	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.




$\Gamma(K^0 \rho^0) / \Gamma_{\text{total}}$		Γ_{324} / Γ	
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN COMMENT
3.4 ±1.1 OUR AVERAGE Error includes scale factor of 2.3. See the ideogram below.			
1.89 ^{+0.55} _{-0.79} ±0.40		¹ AAIJ 18F LHCb	pp at 7, 8 TeV
4.4 ^{+0.7} _{-0.6} ±0.3		² AUBERT 09AU BABR	$e^+e^- \rightarrow \Upsilon(4S)$
6.1 ±1.0 ^{+1.1} _{-1.2}		³ GARMASH 07 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.9 ±0.8 ±0.9		² AUBERT 07F BABR	Repl. by AUBERT 09AU
< 39	90	ASNER 96 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<320	90	ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<500	90	⁴ AVERY 89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

³ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

⁴ AVERY 89B reports $< 5.8 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.



$\Gamma(K^*(892)^+\pi^-)/\Gamma_{\text{total}}$				Γ_{325}/Γ	
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
7.5 $\pm$0.4 OUR AVERAGE					
7.02 $\pm$ 0.30 $\pm$ 0.45		¹ AAIJ	18F LHCb	pp at 7, 8 TeV	
8.0 $\pm$ 1.1 $\pm$ 0.8		^{2,3} LEES	11 BABR	$e^+e^- \rightarrow \Upsilon(4S)$	
8.3 $\begin{smallmatrix} +0.9 \\ -0.8 \end{smallmatrix}$ $\pm$ 0.8		^{3,4} AUBERT	09AU BABR	$e^+e^- \rightarrow \Upsilon(4S)$	
8.4 $\pm$ 1.1 $\begin{smallmatrix} +1.0 \\ -0.9 \end{smallmatrix}$		⁴ GARMASH	07 BELL	$e^+e^- \rightarrow \Upsilon(4S)$	
16 $\begin{smallmatrix} +6 \\ -5 \end{smallmatrix}$ $\pm$ 2		³ ECKHART	02 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
12.6 $\begin{smallmatrix} +2.7 \\ -1.6 \end{smallmatrix}$ $\pm$ 0.9		^{2,3} AUBERT	08AQ BABR	Repl. by LEES 11	
11.0 $\pm$ 1.5 $\pm$ 0.71		³ AUBERT	06I BABR	Repl. by AUBERT 09AU	
12.9 $\pm$ 2.4 $\pm$ 1.4		³ AUBERT,B	04O BABR	Repl. by AUBERT 06I	
14.8 $\begin{smallmatrix} +4.6 \\ -4.4 \end{smallmatrix}$ $\begin{smallmatrix} +2.8 \\ -1.3 \end{smallmatrix}$		³ CHANG	04 BELL	Repl. by GARMASH 07	
< 72	90	ASNER	96 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$	
<620	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
<380	90	⁵ AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
<560	90	⁶ AVERY	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+ \pi^- \pi^0$ decays.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

⁵ AVERY 89B reports $< 4.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

⁶ AVERY 87 reports $< 7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K_0^*(1430)^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{326}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
33 ± 7 OUR AVERAGE	Error includes scale factor of 2.0.		
29.9 $^{+2.3}_{-1.7} \pm 3.6$	^{1,2} AUBERT	09AU BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
49.7 ± 3.8 $^{+6.8}_{-8.2}$	² GARMASH	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

$\Gamma(K_x^{*+} \pi^-)/\Gamma_{\text{total}}$ Γ_{327}/Γ

K_x^{*+} stands for the possible candidates of $K^*(1410)$, $K_0^*(1430)$ and $K_2^*(1430)$.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
5.1 ± 1.5 $^{+0.6}_{-0.7}$	¹ CHANG	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(1410)^+ \pi^-, K^{*+} \rightarrow K^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{328}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.8	90	¹ GARMASH	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.

$\Gamma((K\pi)_0^{*+} \pi^-, (K\pi)_0^{*+} \rightarrow K^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{329}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
16.2 $\pm 0.69 \pm 1.15$	¹ AAIJ	18F LHCb	pp at 7, 8 TeV

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. $(K\pi)_0^{*+}$ is the S-wave component of $K^0 \pi^+$. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

$\Gamma(f_0(980) K^0, f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{330}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			


9.23 $\pm 0.40 \pm 0.79$	¹ AAIJ	18F LHCb	pp at 7, 8 TeV
6.9 $\pm 0.8 \pm 0.6$	² AUBERT	09AU BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
7.6 ± 1.7 $^{+0.9}_{-1.3}$	³ GARMASH	07 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.5 $\pm 0.7 \pm 0.6$	² AUBERT	06I BABR	Repl. by AUBERT 09AU
<360	90	⁴ AVERY	89B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$

- ¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.
- ² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
- ³ Uses Dalitz plot analysis of the $B^0 \rightarrow K^0 \pi^+ \pi^-$ final state decays.
- ⁴ AVERY 89B reports $< 4.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

WEIGHTED AVERAGE
 8.1 ± 0.8 (Error scaled by 1.3)



$\Gamma(K^0 f_0(500))/\Gamma_{\text{total}}$

Γ_{331}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.16^{+0.20}_{-0.04} \pm 0.15$	¹ AAIJ	18F LHCb	pp at 7, 8 TeV

- ¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

$\Gamma(K^0 f_0(1500))/\Gamma_{\text{total}}$

Γ_{332}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$1.29 \pm 0.27 \pm 0.70$	¹ AAIJ	18F LHCb	pp at 7, 8 TeV

- ¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

$\Gamma(f_2(1270)K^0)/\Gamma_{\text{total}}$ Γ_{333}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$2.7^{+1.0}_{-0.8} \pm 0.9$		¹ AUBERT	09AU BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.5 90 ² GARMASH 07 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² GARMASH 07 reports $B(B^0 \rightarrow f_2(1270)K^0) \times B(f_2(1270) \rightarrow \pi^+\pi^-) < 1.4 \times 10^{-6}$ using Dalitz plot analysis. We compute $B(B^0 \rightarrow f_2(1270)K^0)$ using the PDG value $B(f_2(1270) \rightarrow \pi\pi) = 84.3 \times 10^{-2}$ and 2/3 for the $\pi^+\pi^-$ fraction.

$\Gamma(f_x(1300)K^0, f_x \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{334}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$1.81^{+0.55}_{-0.45} \pm 0.48$		¹ AUBERT	09AU BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^*(892)^0\pi^0)/\Gamma_{\text{total}}$ Γ_{335}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$3.3 \pm 0.5 \pm 0.4$		^{1,2} LEES	11 BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.6 \pm 0.7 \pm 0.4$ ^{1,2} AUBERT 08AQ BABR Repl. by LEES 11
 < 3.5 90 ² CHANG 04 BELL $e^+e^- \rightarrow \Upsilon(4S)$
 < 3.6 90 JESSOP 00 CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
 < 28 90 ASNER 96 CLE2 Repl. by JESSOP 00

¹ Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_2^*(1430)^+\pi^-)/\Gamma_{\text{total}}$ Γ_{336}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$3.65^{+0.15}_{-0.12} \pm 0.31$		¹ AAIJ	18F LHCb	pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 16.2 90 ^{2,3} AUBERT 08AQ BABR $e^+e^- \rightarrow \Upsilon(4S)$
 < 6 90 ⁴ GARMASH 07 BELL $e^+e^- \rightarrow \Upsilon(4S)$
 < 18 90 ³ GARMASH 04 BELL Repl. by GARMASH 07
 < 2600 90 ALBRECHT 91B ARG $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0\pi^+\pi^-$ final state decays. We compute $B(B^0 \rightarrow K_2^*(1430)^+\pi^-)$ using the PDG 18 value $B(K_2^*(1430) \rightarrow K\pi) = 49.9 \times 10^{-2}$ and 2/3 for the $K^0\pi^+$ fraction. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0\pi^+\pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+\pi^-\pi^0$ decays.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ GARMASH 07 reports $B(B^0 \rightarrow K_2^*(1430)^+\pi^-) \times B(K_2^+ \rightarrow K^0\pi^+) < 2.1 \times 10^{-6}$ using Dalitz plot analysis. We compute $B(B^0 \rightarrow K_2^*(1430)^+\pi^-)$ using the PDG value $B(K_2^*(1430) \rightarrow K\pi) = 49.9 \times 10^{-2}$ and 2/3 for the $K^0\pi^+$ fraction.

$\Gamma(K^*(1680)^+\pi^-)/\Gamma_{\text{total}}$ Γ_{337}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$14.1 \pm 0.58 \pm 0.84$ ¹ AAIJ 18F LHCb pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<25 90 ^{2,3} AUBERT 08AQ BABR $e^+e^- \rightarrow \Upsilon(4S)$

<10 90 ⁴ GARMASH 07 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ final state decays. We compute $B(B^0 \rightarrow K_2^{*+}(1430) \pi^-)$ using the PDG 18 value $B(K_2^{*+}(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ and 2/3 for the $K^0 \pi^+$ fraction. For the branching fraction of the reference mode, the PDG 18 average $B(B^0 \rightarrow K_S^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ is used.

² Uses Dalitz plot analysis of $B^0 \rightarrow K^+ \pi^- \pi^0$ decays.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ GARMASH 07 reports $B(B^0 \rightarrow K^*(1680)^+ \pi^-) \times B(K^{*+} \rightarrow K^0 \pi^+) < 2.6 \times 10^{-6}$ using Dalitz plot analysis. We compute $B(B^0 \rightarrow K^*(1680)^+ \pi^-)$ using the PDG value $B(K^*(1680) \rightarrow K \pi) = 38.7 \times 10^{-2}$ and 2/3 for the $K^0 \pi^+$ fraction.

 $\Gamma(K^+ \pi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{338}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 2.3 \times 10^{-4}$ 90 ¹ ADAM 96D DLPH $e^+e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1 $\times 10^{-4}$ 90 ² ABREU 95N DLPH Sup. by ADAM 96D

¹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

² Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

 $\Gamma(\rho^0 K^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{339}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.8 \pm 0.5 \pm 0.5$ ^{1,2} KYEONG 09 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Required $0.75 < m_{K^+ \pi^-} < 1.20 \text{ GeV}/c^2$.

 $\Gamma(f_0(980) K^+ \pi^-, f_0 \rightarrow \pi \pi)/\Gamma_{\text{total}}$ Γ_{340}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.4 \pm 0.4^{+0.3}_{-0.4}$ ^{1,2} KYEONG 09 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Required $0.75 < m_{K^+ K^-} < 1.2 \text{ GeV}/c^2$.

 $\Gamma(K^+ \pi^- \pi^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{341}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 2.1 \times 10^{-6}$ 90 ^{1,2} KYEONG 09 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Required $0.55 < m_{\pi^+ \pi^-} < 1.42$ and $0.75 < m_{K^+ \pi^-} < 1.20 \text{ GeV}/c^2$.

$$\Gamma(K^*(892)^0 \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{342} / \Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

54.5 ± 2.9 ± 4.3¹ AUBERT 07AS BABR $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.5^{+1.1+0.9}_{-1.0-1.6}^{1,2} KYEONG 09 BELL $e^+ e^- \rightarrow \Upsilon(4S)$ <1400 90 ALBRECHT 91E ARG $e^+ e^- \rightarrow \Upsilon(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.² Required $0.55 < m_{\pi^+ \pi^-} < 1.42 \text{ GeV}/c^2$.
$$\Gamma(K^*(892)^0 \rho^0) / \Gamma_{\text{total}} \quad \Gamma_{343} / \Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

3.9 ± 1.3 OUR AVERAGE

Error includes scale factor of 1.9.

5.1 ± 0.6^{+0.6}_{-0.8}¹ LEES 12K BABR $e^+ e^- \rightarrow \Upsilon(4S)$ 2.1^{+0.8+0.9}_{-0.7-0.5}¹ KYEONG 09 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.6 ± 0.9 ± 1.3

¹ AUBERT,B 06G BABR Repl. by LEES 12K< 34 90 ² GODANG 02 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$ <286 90 ³ ABE 00C SLD $e^+ e^- \rightarrow Z$ <460 90 ALBRECHT 91B ARG $e^+ e^- \rightarrow \Upsilon(4S)$ <580 90 ⁴ AVERY 89B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$ <960 90 ⁵ AVERY 87 CLEO $e^+ e^- \rightarrow \Upsilon(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.² Assumes a helicity 00 configuration. For a helicity 11 configuration, the limit decreases to 2.4×10^{-5} .³ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.⁴ AVERY 89B reports $< 6.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.⁵ AVERY 87 reports $< 1.2 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.
$$\Gamma(K^*(892)^0 f_0(980), f_0 \rightarrow \pi\pi) / \Gamma_{\text{total}} \quad \Gamma_{344} / \Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

3.9^{+2.1}_{-1.8} OUR AVERAGE

Error includes scale factor of 3.9.

5.7 ± 0.6 ± 0.4

¹ LEES 12K BABR $e^+ e^- \rightarrow \Upsilon(4S)$ 1.4^{+0.6+0.6}_{-0.5-0.4}^{1,2} KYEONG 09 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.3 90 ¹ AUBERT,B 06G BABR $e^+ e^- \rightarrow \Upsilon(4S)$ <170 90 ³ AVERY 89B CLEO $e^+ e^- \rightarrow \Upsilon(4S)$ ¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.² The upper limit is 2.2×10^{-6} at 90% CL.³ AVERY 89B reports $< 2.0 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K_1(1270)^+\pi^-)/\Gamma_{\text{total}}$ Γ_{345}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-5}$	90	¹ AUBERT	10D BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K_1(1400)^+\pi^-)/\Gamma_{\text{total}}$ Γ_{346}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7 \times 10^{-5}$	90	¹ AUBERT	10D BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------------	----	----------	---------	-----------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(a_1(1260)^-K^+)/\Gamma_{\text{total}}$ Γ_{347}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$16.3 \pm 2.9 \pm 2.3$		^{1,2} AUBERT	08F BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<230	90	³ ADAM	96D DLPH	$e^+e^- \rightarrow Z$
<390	90	⁴ ABREU	95N DLPH	Sup. by ADAM 96D

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Assumes $a_1^\pm$ decays only to 3π and $B(a_1^\pm \rightarrow \pi^\pm \pi^\mp \pi^\pm) = 0.5$.

³ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

⁴ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

 $\Gamma(K^*(892)^+\rho^-)/\Gamma_{\text{total}}$ Γ_{348}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$10.3 \pm 2.3 \pm 1.3$		¹ LEES	12K BABR	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12.0	90	¹ AUBERT,B	06G BABR	Repl. by LEES 12K
---------	----	-----------------------	----------	-------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K_0^*(1430)^+\rho^-)/\Gamma_{\text{total}}$ Γ_{349}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$28 \pm 10 \pm 6$	¹ LEES	12K BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K_1(1400)^0\rho^0)/\Gamma_{\text{total}}$ Γ_{350}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(K_0^*(1430)^0\rho^0)/\Gamma_{\text{total}}$ Γ_{351}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$27 \pm 4 \pm 4$	¹ LEES	12K BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K_0^*(1430)^0 f_0(980), f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_{352}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$2.7 \pm 0.7 \pm 0.6$	¹ LEES	12K BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K_2^*(1430)^0 f_0(980), f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_{353}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.6 \pm 1.7 \pm 1.0$	¹ LEES	12K BABR	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{354}/Γ

VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
$7.80 \pm 1.27 \pm 0.84$		¹ AAIJ	17G LHCB	pp at 7 and 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10 ± 8 ± 4		^{2,3} DUH	13 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
12 ± 8 ± 2		⁴ AAIJ	12AR LHCB	Repl. by AAIJ 17G
23 ± 10 ± 10		⁵ AALTONEN	12L CDF	$p\bar{p}$ at 1.96 TeV
< 70	90	⁶ AALTONEN	09C CDF	Repl. by AALTONEN 12L
< 50	90	³ AUBERT	07B BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 41	90	³ LIN	07 BELL	Repl. by DUH 13
< 180	90	⁷ ABULENCIA,A	06D CDF	Repl. by AALTONEN 09C
< 37	90	ABE	05G BELL	Repl. by LIN 07
< 70	90	CHAO	04 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 80	90	³ BORNHEIM	03 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 60	90	³ AUBERT	02Q BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 90	90	³ CASEY	02 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 270	90	³ ABE	01H BELL	$e^+e^- \rightarrow \Upsilon(4S)$
< 250	90	³ AUBERT	01E BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 6600	90	⁸ ABE	00C SLD	$e^+e^- \rightarrow Z$
< 190	90	³ CRONIN-HEN.	00 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 430	90	GODANG	98 CLE2	Repl. by CRONIN-HENNESSY 00
< 4600		⁹ ADAM	96D DLPH	$e^+e^- \rightarrow Z$
< 400	90	ASNER	96 CLE2	Repl. by GODANG 98
< 1800	90	¹⁰ BUSKULIC	96V ALEP	$e^+e^- \rightarrow Z$
< 12000	90	¹¹ ABREU	95N DLPH	Sup. by ADAM 96D
< 700	90	³ BATTLE	93 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Supersedes results of AAIJ 12AR.

² DUH 13 reports also for the same data $B(B^0 \rightarrow K^+K^-) < 0.20 \times 10^{-6}$ at 90% CL.

³ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

⁴ AAIJ 12AR reports $[\Gamma(B^0 \rightarrow K^+K^-)/\Gamma_{\text{total}}] / [B(B_s^0 \rightarrow K^+K^-)] / [\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0)] = 0.018^{+0.008}_{-0.007} \pm 0.009$ which we multiply by our best values $B(B_s^0 \rightarrow K^+K^-) = (2.72 \pm 0.23) \times 10^{-5}$, $\Gamma(\bar{b} \rightarrow B_s^0)/\Gamma(\bar{b} \rightarrow B^0) = 0.246 \pm 0.023$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

⁵ Reported a central value of $(0.23 \pm 0.10 \pm 0.10) \times 10^{-6}$ using $B(B^0 \rightarrow K^+ \pi^-) = (19.4 \pm 0.6) \times 10^{-6}$.

⁶ Obtains this result from $B(K^+ K^-)/B(K^+ \pi^-) = 0.020 \pm 0.008 \pm 0.006$, assuming $B(B^0 \rightarrow K^+ \pi^-) = (19.4 \pm 0.6) \times 10^{-6}$.

⁷ ABULENCIA, A 06D obtains this from $\Gamma(K^+ K^-)/\Gamma(K^+ \pi^-) < 0.10$ at 90% CL, assuming $B(B^0 \rightarrow K^+ \pi^-) = (18.9 \pm 0.7) \times 10^{-6}$.

⁸ ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.

⁹ ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

¹⁰ BUSKULIC 96V assumes PDG 96 production fractions for B^0 , B^+ , B_s , b baryons.

¹¹ Assumes a B^0 , B^- production fraction of 0.39 and a B_s production fraction of 0.12. Contributions from B^0 and B_s decays cannot be separated. Limits are given for the weighted average of the decay rates for the two neutral B mesons.

$\Gamma(K^0 \bar{K}^0)/\Gamma_{\text{total}}$

Γ_{355}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
1.21 ± 0.16 OUR AVERAGE				
1.26 ± 0.19 ± 0.05		¹ DUH	13	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
1.08 ± 0.28 ± 0.11		¹ AUBERT, BE	06C	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.87 $^{+0.25}_{-0.20}$ ± 0.09		¹ LIN	07	BELL Repl. by DUH 13
0.8 ± 0.3 ± 0.9		¹ ABE	05G	BELL Repl. by LIN 07
1.19 $^{+0.40}_{-0.35}$ ± 0.13		¹ AUBERT, BE	05E	BABR Repl. by AUBERT, BE 06C
< 1.8	90	¹ AUBERT	04M	BABR $e^+ e^- \rightarrow \Upsilon(4S)$
< 1.5	90	¹ CHAO	04	BELL Repl. by ABE 05G
< 3.3	90	¹ BORNHEIM	03	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
< 4.1	90	¹ CASEY	02	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
< 17	90	GODANG	98	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(K^0 \bar{K}^0)/\Gamma(K^0 \phi)$

$\Gamma_{355}/\Gamma_{364}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.08 ± 0.02	¹ AAIJ	20F	LHCB pp at 7, 8, 13 TeV

¹ Observed signal with a significance of 3.5 σ .

$\Gamma(K^0 K^- \pi^+)/\Gamma_{\text{total}}$

Γ_{356}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
6.7 ± 0.5 OUR FIT				
7.0 ± 0.6 OUR AVERAGE				
7.2 ± 0.7 ± 0.3		¹ LAI	19	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
6.4 ± 1.0 ± 0.6		¹ DEL-AMO-SA..10E	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 18	90	¹ GARMASH	04	BELL $e^+ e^- \rightarrow \Upsilon(4S)$
< 21	90	¹ ECKHART	02	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}} \quad \Gamma_{357}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.4 \times 10^{-6}$	90	AAIJ	14BMLHCB	pp at 7 TeV

$$\Gamma(K^0 K^- \pi^+)/\Gamma(K^0 \pi^+ \pi^-) \quad \Gamma_{356}/\Gamma_{322}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.134 ± 0.011 OUR FIT

0.123 ± 0.009 ± 0.015 AAIJ 17BP LHCB pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.128 ± 0.017 ± 0.009 AAIJ 13BP LHCB Repl. by AAIJ 17BP

$$[\Gamma(\bar{K}^{*0} K^0) + \Gamma(K^{*0} \bar{K}^0)]/\Gamma_{\text{total}} \quad \Gamma_{358}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.96 90 ¹ AAIJ 16 LHCB pp at 7 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9 90 ² AUBERT, BE 06N BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes $B(B^0 \rightarrow K^0 \pi^+ \pi^-) = (4.96 \pm 0.20) \times 10^{-5}$.

² Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{359}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

2.17 ± 0.60 ± 0.24 ¹ GAUR 13 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19 90 ¹ ECKHART 02 CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(K_S^0 K_S^0 \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{360}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.9 × 10⁻⁶ 90 ¹ AUBERT 09AD BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(K_S^0 K_S^0 \eta)/\Gamma_{\text{total}} \quad \Gamma_{361}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.0 × 10⁻⁶ 90 ¹ AUBERT 09AD BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(K_S^0 K_S^0 \eta')/\Gamma_{\text{total}} \quad \Gamma_{362}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10⁻⁶ 90 ¹ AUBERT 09AD BABR $e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$$\Gamma(K^0 K^+ K^-)/\Gamma_{\text{total}} \quad \Gamma_{363}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

26.8 ± 1.1 OUR FIT

26.6 ± 1.2 OUR AVERAGE

26.5 ± 0.9 ± 0.8 ^{1,2} LEES 120 BABR $e^+ e^- \rightarrow \Upsilon(4S)$

28.3 ± 3.3 ± 4.0 ¹ GARMASH 04 BELL $e^+ e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •

$23.8 \pm 2.0 \pm 1.6$		¹ AUBERT,B	04V	BABR	Repl. by LEES 120
< 1300	90	ALBRECHT	91E	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² All intermediate charmonium and charm resonances are removed, except of χ_{c0} .

$\Gamma(K^0 K^+ K^-)/\Gamma(K^0 \pi^+ \pi^-)$ $\Gamma_{363}/\Gamma_{322}$

VALUE		DOCUMENT ID	TECN	COMMENT
0.539 ± 0.025 OUR FIT				
$0.549 \pm 0.018 \pm 0.033$		AAIJ	17BP	LHCB pp at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.385 \pm 0.031 \pm 0.023$		AAIJ	13BP	LHCB Repl. by AAIJ 17BP

$\Gamma(K^0 \phi)/\Gamma_{\text{total}}$ Γ_{364}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
7.3 ± 0.7 OUR AVERAGE				
$7.1 \pm 0.6^{+0.4}_{-0.3}$		¹ LEES	120	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$9.0^{+2.2}_{-1.8} \pm 0.7$		¹ CHEN	03B	BELL $e^+e^- \rightarrow \Upsilon(4S)$

- • • We do not use the following data for averages, fits, limits, etc. • • •

$8.4^{+1.5}_{-1.3} \pm 0.5$		¹ AUBERT	04A	BABR	Repl. by LEES 120
$8.1^{+3.1}_{-2.5} \pm 0.8$		¹ AUBERT	01D	BABR	$e^+e^- \rightarrow \Upsilon(4S)$
< 12.3	90	¹ BRIERE	01	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 31	90	¹ BERGFELD	98	CLE2	
< 88	90	ASNER	96	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
< 720	90	ALBRECHT	91B	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 420	90	² AVERY	89B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 1000	90	³ AVERY	87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² AVERY 89B reports $< 4.9 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

³ AVERY 87 reports $< 1.3 \times 10^{-3}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(f_0(980) K^0, f_0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}$ Γ_{365}/Γ

VALUE (units 10^{-6})		DOCUMENT ID	TECN	COMMENT
$7.0^{+2.6}_{-1.8} \pm 2.4$		¹ LEES	120	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_0(1500) K^0)/\Gamma_{\text{total}}$ Γ_{366}/Γ

VALUE (units 10^{-6})		DOCUMENT ID	TECN	COMMENT
$13.3^{+5.8}_{-4.4} \pm 3.2$		¹ LEES	120	BABR $e^+e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

$\Gamma(f_2'(1525)^0 K^0)/\Gamma_{\text{total}}$ Γ_{367}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.29^{+0.27}_{-0.18} \pm 0.36$	¹ LEES	120 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(f_0(1710) K^0, f_0 \rightarrow K^+ K^-)/\Gamma_{\text{total}}$ Γ_{368}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 0.7 \pm 0.5$	¹ LEES	120 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^0 K^+ K^- \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{369}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$33 \pm 5 \pm 9$	¹ LEES	120 BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{370}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
6.0 $\pm$ 0.5 OUR AVERAGE	Error includes scale factor of 1.1.		
$6.19 \pm 0.48 \pm 0.19$	¹ LEES	12i BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
$4.2^{+1.6}_{-1.3} \pm 0.8$	¹ GARMASH	04 BELL	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.9^{+0.9}_{-0.8} \pm 0.6$	¹ AUBERT,B	05 BABR	Repl. by LEES 12i
-----------------------------	-----------------------	---------	-------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(f_0(980) K^0, f_0 \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{371}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$2.7^{+1.3}_{-1.2} \pm 1.3$	^{1,2} LEES	12i BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decay.

 $\Gamma(f_0(1710) K^0, f_0 \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{372}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.50^{+0.46}_{-0.24} \pm 0.11$	^{1,2} LEES	12i BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decay.

 $\Gamma(f_2(2010) K^0, f_2 \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{373}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.54^{+0.21}_{-0.20} \pm 0.52$	^{1,2} LEES	12i BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decay.

$\Gamma(K_S^0 K_S^0 K_S^0 \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{374}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$13.3^{+2.2}_{-2.3} \pm 2.2$	1,2 LEES	12I BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² Uses Dalitz plot analysis of the $B^0 \rightarrow K_S^0 K_S^0 K_S^0$ decay.

 $\Gamma(K_S^0 K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{375}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	¹ AUBERT,B	06R BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^*(892)^0 K^+ K^-)/\Gamma_{\text{total}}$ Γ_{376}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
$27.5 \pm 1.3 \pm 2.2$		¹ AUBERT	07AS BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<610	90	ALBRECHT	91E ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
--------	----	----------	---------	------------------------------------

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^*(892)^0 \phi)/\Gamma_{\text{total}}$ Γ_{377}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
10.0 ± 0.5 OUR FIT				
10.0 ± 0.5 OUR AVERAGE				

$10.4 \pm 0.5 \pm 0.6$	¹ PRIM	13	BELL	$e^+ e^- \rightarrow \Upsilon(4S)$
------------------------	-------------------	----	------	------------------------------------

$9.7 \pm 0.5 \pm 0.5$	¹ AUBERT	08BG	BABR	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------	---------------------	------	------	------------------------------------

$11.5^{+4.5+1.8}_{-3.7-1.7}$	¹ BRIERE	01	CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
------------------------------	---------------------	----	------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9.2 \pm 0.7 \pm 0.6$	¹ AUBERT	07D	BABR	Repl. by AUBERT 08BG
-----------------------	---------------------	-----	------	----------------------

$9.2 \pm 0.9 \pm 0.5$	¹ AUBERT,B	04W	BABR	Repl. by AUBERT 07D
-----------------------	-----------------------	-----	------	---------------------

$11.2 \pm 1.3 \pm 0.8$	¹ AUBERT	03V	BABR	Repl. by AUBERT,B 04W
------------------------	---------------------	-----	------	-----------------------

$10.0^{+1.6+0.7}_{-1.5-0.8}$	¹ CHEN	03B	BELL	Repl. by PRIM 13
------------------------------	-------------------	-----	------	------------------

$8.7^{+2.5}_{-2.1} \pm 1.1$	¹ AUBERT	01D	BABR	Repl. by AUBERT 03V
-----------------------------	---------------------	-----	------	---------------------

<384	90	² ABE	00C	SLD $e^+ e^- \rightarrow Z$
--------	----	------------------	-----	-----------------------------

<21	90	¹ BERGFELD	98	CLE2
-------	----	-----------------------	----	------

<43	90	ASNER	96	CLE2 $e^+ e^- \rightarrow \Upsilon(4S)$
-------	----	-------	----	---

<320	90	ALBRECHT	91B	ARG $e^+ e^- \rightarrow \Upsilon(4S)$
--------	----	----------	-----	--

<380	90	³ AVERY	89B	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
--------	----	--------------------	-----	---


<380	90	⁴ AVERY	87	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$
--------	----	--------------------	----	---

¹ Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

² ABE 00C assumes $B(Z \rightarrow b\bar{b}) = (21.7 \pm 0.1)\%$ and the B fractions $f_{B^0} = f_{B^+} = (39.7^{+1.8}_{-2.2})\%$ and $f_{B_s} = (10.5^{+1.8}_{-2.2})\%$.


³ AVERY 89B reports $< 4.4 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

⁴ AVERY 87 reports $< 4.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 40% to $B^0 \bar{B}^0$. We rescale to 50%.



$\phi_{\parallel}$ in $B^0 \rightarrow J/\psi K^{*0}$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
-2.92 ± 0.04 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
$-2.94 \pm 0.02 \pm 0.03$	AAIJ	13AT LHCb	pp at 7 TeV
$-2.69 \pm 0.08 \pm 0.11$	¹ ABAZOV	09E D0	$p\bar{p}$ at 1.96 TeV
$-2.93 \pm 0.08 \pm 0.04$	² AUBERT	07AD BABR	$e^+e^- \rightarrow \Upsilon(4S)$




¹ Obtained $\phi_{\parallel}$ as $\delta_2 - \delta_1$, assuming they are uncorrelated.

² Obtained by combining the B^0 and B^+ modes.

$\phi_{\perp}$ in $B^0 \rightarrow J/\psi K^{*0}$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
2.96 ± 0.05 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.		
$2.94 \pm 0.02 \pm 0.02$	AAIJ	13AT LHCb	pp at 7 TeV
$3.21 \pm 0.06 \pm 0.06$	ABAZOV	09E D0	$p\bar{p}$ at 1.96 TeV
$2.91 \pm 0.05 \pm 0.03$	¹ AUBERT	07AD BABR	$e^+e^- \rightarrow \gamma(4S)$

¹ Obtained by combining the B^0 and B^+ modes.



Γ_L/Γ in $B^0 \rightarrow \psi(2S) K^{*}(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.463^{+0.028}_{-0.040}$ OUR AVERAGE			
$0.455^{+0.031+0.014}_{-0.029-0.049}$	CHILIKIN	13 BELL	$e^+e^- \rightarrow \gamma(4S)$
$0.48 \pm 0.05 \pm 0.02$	¹ AUBERT	07AD BABR	$e^+e^- \rightarrow \gamma(4S)$
$0.45 \pm 0.11 \pm 0.04$	² RICHICHI	01 CLE2	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.448^{+0.040+0.040}_{-0.027-0.053}$	MIZUK	09 BELL	$e^+e^- \rightarrow \gamma(4S)$
---------------------------------------	-------	---------	---------------------------------

¹ Obtained by combining the B^0 and B^+ modes.

² Averages between charged and neutral B mesons.

- ¹ Obtained by amplitude analysis of $\bar{B}^0 \rightarrow D^{*-} \omega \pi^+$. The second uncertainty combines in quadrature experimental systematic and model uncertainties.
² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.

 Γ_L/Γ in $B^0 \rightarrow \bar{D}_1(2420)^0 \omega$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$67.1 \pm 11.7^{+2.3}_{-5.0}$	1,2 MATVIENKO	15	BELL $e^+e^- \rightarrow \Upsilon(4S)$

- ¹ Obtained by amplitude analysis of $\bar{B}^0 \rightarrow D^{*-} \omega \pi^+$. The second uncertainty combines in quadrature experimental systematic and model uncertainties.
² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.

 Γ_L/Γ in $B^0 \rightarrow \bar{D}_2^*(2460)^0 \omega$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
$76.0^{+18.3+3.5}_{-8.5-2.8}$	1,2 MATVIENKO	15	BELL $e^+e^- \rightarrow \Upsilon(4S)$

- ¹ Obtained by amplitude analysis of $\bar{B}^0 \rightarrow D^{*-} \omega \pi^+$. The second uncertainty combines in quadrature experimental systematic and model uncertainties.
² Assumes equal production of B^0 and B^+ at $\Upsilon(4S)$.

 Γ_L/Γ in $B^0 \rightarrow D^{*-} \omega \pi^+$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.654 \pm 0.042 \pm 0.016$	¹ AUBERT	06L	BABR $e^+e^- \rightarrow \Upsilon(4S)$

- ¹ Invariant mass of the $[\omega\pi]$ system is restricted in the region 1.1 and 1.9 GeV.

 Γ_L/Γ in $B^0 \rightarrow \omega K^{*0}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.69 ± 0.11 OUR AVERAGE			
$0.68 \pm 0.17 \pm 0.16$	AAIJ	19J	LHCB pp at 7, 8 TeV
$0.72 \pm 0.14 \pm 0.02$	AUBERT	09H	BABR $e^+e^- \rightarrow \Upsilon(4S)$
$0.56 \pm 0.29^{+0.18}_{-0.08}$	GOLDENZWE..08	BELL	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma_{\perp}/\Gamma$ in $B^0 \rightarrow \omega K^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.10 \pm 0.09 \pm 0.09$	AAIJ	19J	LHCB pp at 7, 8 TeV

 A_{CP}^0 in $B^0 \rightarrow \omega K^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.13 \pm 0.27 \pm 0.13$	AAIJ	19J	LHCB pp at 7, 8 TeV

 $A_{CP}^{\perp}$ in $B^0 \rightarrow \omega K^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.3 \pm 0.8 \pm 0.4$	AAIJ	19J	LHCB pp at 7, 8 TeV

 $A_{CP}^{\parallel}$ in $B^0 \rightarrow \omega K^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.26 \pm 0.55 \pm 0.22$	AAIJ	19J	LHCB pp at 7, 8 TeV


 ϕ_0 in $B^0 \rightarrow \omega K^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
$-0.86 \pm 0.29 \pm 0.71$	AAIJ	19J	LHCB pp at 7, 8 TeV

$\phi_{\parallel}$ in $B^0 \rightarrow \phi K^*(892)^0$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
2.43 ±0.11 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.		
2.562±0.069±0.040	AAIJ	14AMLHCB	$p\bar{p}$ at 7 TeV
2.23 ±0.10 ±0.02	PRIM	13 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
2.40 ±0.13 ±0.08	AUBERT	08BG BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.31 ±0.14 ±0.08	AUBERT	07D BABR	Repl. by AUBERT 08BG
2.40 $^{+0.28}_{-0.24}$ ±0.07	¹ CHEN	05A BELL	Repl. by PRIM 13
2.34 $^{+0.23}_{-0.20}$ ±0.05	AUBERT,B	04W BABR	Repl. by AUBERT 07D


¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.



$\phi_{\perp}$ in $B^0 \rightarrow \phi K^*(892)^0$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
2.53 ±0.09 OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.		
2.633±0.062±0.037	AAIJ	14AMLHCB	$p\bar{p}$ at 7 TeV
2.37 ±0.10 ±0.04	PRIM	13 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
2.35 ±0.13 ±0.09	AUBERT	08BG BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.24 ±0.15 ±0.09	AUBERT	07D BABR	Repl. by AUBERT 08BG
2.51 ±0.25 ±0.06	¹ CHEN	05A BELL	Repl. by PRIM 13
2.47 ±0.25 ±0.05	AUBERT,B	04W BABR	Repl. by AUBERT 07D

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.



$\delta_0(B^0 \rightarrow \phi K^*(892)^0)$

VALUE (rad)	DOCUMENT ID	TECN	COMMENT
2.88 ± 0.10 OUR AVERAGE			
$2.91 \pm 0.10 \pm 0.08$	PRIM	13 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$2.82 \pm 0.15 \pm 0.09$	AUBERT	08BG BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2.78 \pm 0.17 \pm 0.09$	AUBERT	07D BABR	Repl. by AUBERT 08BG

A_{CP}^0 in $B^0 \rightarrow \phi K^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.007 ± 0.030 OUR AVERAGE			
$-0.003 \pm 0.038 \pm 0.005$	AAIJ	14AMLHCB	pp at 7 TeV
$-0.030 \pm 0.061 \pm 0.007$	PRIM	13 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$0.01 \pm 0.07 \pm 0.02$	AUBERT	08BG BABR	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$-0.03 \pm 0.08 \pm 0.02$	AUBERT	07D BABR	Repl. by AUBERT 08BG
$0.13 \pm 0.12 \pm 0.04$	¹ CHEN	05A BELL	Repl. by PRIM 13
$-0.06 \pm 0.10 \pm 0.01$	AUBERT,B	04W BABR	Repl. by AUBERT 07D

¹ This quantity was recalculated by the BELLE authors from numbers in the original paper.

$A_{CP}^{\perp}$ in $B^0 \rightarrow \phi K^*(892)^0$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02 ± 0.06 OUR AVERAGE			
$0.047 \pm 0.074 \pm 0.009$	AAIJ	14AMLHCB	pp at 7 TeV
$-0.14 \pm 0.11 \pm 0.01$	PRIM	13 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
$-0.04 \pm 0.15 \pm 0.06$	AUBERT	08BG BABR	$e^+e^- \rightarrow \Upsilon(4S)$

⁸ ALBRECHT 87I is inclusive measurement with like-sign dileptons, with tagged B decays plus leptons, and one fully reconstructed event. Measures $r=0.21 \pm 0.08$. We convert to χ for comparison. Superseded by ALBRECHT 92L.

⁹ BEAN 87B measured $r < 0.24$; we converted to χ .

¹⁰ Same-sign dilepton events. Limit assumes semileptonic BR for B^+ and B^0 equal. If $B^0/B^\pm$ ratio < 0.58 , no limit exists. The limit was corrected in BEAN 87B from $r < 0.30$ to $r < 0.37$. We converted this limit to χ .

$$\Delta m_{B^0} = m_{B_H^0} - m_{B_L^0}$$


Δm_{B^0} is a measure of 2π times the B^0 - $\bar{B}^0$ oscillation frequency in time-dependent mixing experiments.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The averaging/rescaling procedure takes into account correlations between the measurements and includes Δm_d calculated from χ_d measured at $\Upsilon(4S)$.

VALUE ($10^{12} \hbar s^{-1}$)	DOCUMENT ID	TECN	COMMENT
0.5069$\pm$0.0019 OUR EVALUATION (Produced by HFLAV)			
0.516 $\pm$ 0.008 $\pm$ 0.005	¹ ABUDINEN	23D BELL	$e^+e^- \rightarrow \Upsilon(4S)$
0.5050 $\pm$ 0.0021 $\pm$ 0.0010	² AAIJ	16AV LHCb	pp at 7, 8 TeV
0.503 $\pm$ 0.011 $\pm$ 0.013	³ AAIJ	13CF LHCb	pp at 7 TeV
0.5156 $\pm$ 0.0051 $\pm$ 0.0033	⁴ AAIJ	13F LHCb	pp at 7 TeV
0.499 $\pm$ 0.032 $\pm$ 0.003	⁵ AAIJ	12I LHCb	pp at 7 TeV
0.506 $\pm$ 0.020 $\pm$ 0.016	⁶ ABAZOV	06W D0	$p\bar{p}$ at 1.96 TeV
0.511 $\pm$ 0.007 $^{+0.007}_{-0.006}$	⁷ AUBERT	06G BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.511 $\pm$ 0.005 $\pm$ 0.006	⁸ ABE	05B BELL	$e^+e^- \rightarrow \Upsilon(4S)$
0.531 $\pm$ 0.025 $\pm$ 0.007	⁹ ABDALLAH	03B DLPH	$e^+e^- \rightarrow Z$
0.492 $\pm$ 0.018 $\pm$ 0.013	¹⁰ AUBERT	03C BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.503 $\pm$ 0.008 $\pm$ 0.010	¹¹ HASTINGS	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
0.509 $\pm$ 0.017 $\pm$ 0.020	¹² ZHENG	03 BELL	$e^+e^- \rightarrow \Upsilon(4S)$
0.516 $\pm$ 0.016 $\pm$ 0.010	¹³ AUBERT	02I BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.493 $\pm$ 0.012 $\pm$ 0.009	¹⁴ AUBERT	02J BABR	$e^+e^- \rightarrow \Upsilon(4S)$
0.497 $\pm$ 0.024 $\pm$ 0.025	¹⁵ ABBIENDI,G	00B OPAL	$e^+e^- \rightarrow Z$
0.503 $\pm$ 0.064 $\pm$ 0.071	¹⁶ ABE	99K CDF	$p\bar{p}$ at 1.8 TeV
0.500 $\pm$ 0.052 $\pm$ 0.043	¹⁷ ABE	99Q CDF	$p\bar{p}$ at 1.8 TeV
0.516 $\pm$ 0.099 $^{+0.029}_{-0.035}$	¹⁸ AFFOLDER	99C CDF	$p\bar{p}$ at 1.8 TeV
0.471 $^{+0.078}_{-0.068}$ $^{+0.033}_{-0.034}$	¹⁹ ABE	98C CDF	$p\bar{p}$ at 1.8 TeV
0.458 $\pm$ 0.046 $\pm$ 0.032	²⁰ ACCIARRI	98D L3	$e^+e^- \rightarrow Z$
0.437 $\pm$ 0.043 $\pm$ 0.044	²¹ ACCIARRI	98D L3	$e^+e^- \rightarrow Z$
0.472 $\pm$ 0.049 $\pm$ 0.053	²² ACCIARRI	98D L3	$e^+e^- \rightarrow Z$
0.523 $\pm$ 0.072 $\pm$ 0.043	²³ ABREU	97N DLPH	$e^+e^- \rightarrow Z$
0.493 $\pm$ 0.042 $\pm$ 0.027	²¹ ABREU	97N DLPH	$e^+e^- \rightarrow Z$
0.499 $\pm$ 0.053 $\pm$ 0.015	²⁴ ABREU	97N DLPH	$e^+e^- \rightarrow Z$
0.480 $\pm$ 0.040 $\pm$ 0.051	²⁰ ABREU	97N DLPH	$e^+e^- \rightarrow Z$
0.444 $\pm$ 0.029 $^{+0.020}_{-0.017}$	²¹ ACKERSTAFF	97U OPAL	$e^+e^- \rightarrow Z$
0.430 $\pm$ 0.043 $^{+0.028}_{-0.030}$	²⁰ ACKERSTAFF	97V OPAL	$e^+e^- \rightarrow Z$
0.482 $\pm$ 0.044 $\pm$ 0.024	²⁵ BUSKULIC	97D ALEP	$e^+e^- \rightarrow Z$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.52±0.25±0.19 AAIJ 12U LHCb Repl. by AAIJ 13Y



$B(B^0 \rightarrow K^{*0} \ell^+ \ell^-) (14.18 < q^2 < 16.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.09±0.10 OUR AVERAGE	Error includes scale factor of 1.1.		
1.22±0.11±0.09	KHACHATRY...16D	CMS	pp at 8 TeV
1.02±0.11 ^{+0.11} _{-0.15}	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
0.84 ^{+0.16} _{-0.15} ±0.09	CHATRCHYAN 13BL	CMS	pp at 7 TeV
1.34±0.26±0.08	AALTONEN 11AI	CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.15±0.20±0.09	AAIJ	12U LHCb	Repl. by AAIJ 13Y

$B(B^0 \rightarrow K^{*0} \ell^+ \ell^-) (16.0 < q^2 < 19.0 \text{ GeV}^2/c^4)$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
1.27±0.09 OUR AVERAGE			
1.26±0.09±0.09	KHACHATRY...16D	CMS	pp at 8 TeV
1.23±0.12 ^{+0.15} _{-0.18}	AAIJ	13Y LHCb	pp at 7 TeV, $K^{*0} \mu^+ \mu^-$
1.56±0.18±0.15	CHATRCHYAN 13BL	CMS	pp at 7 TeV
0.97±0.26±0.07	AALTONEN 11AI	CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.50±0.24±0.15	AAIJ	12U LHCb	Repl. by AAIJ 13Y

AUBERT	09A	PR D79 012002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AA	PR D79 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AC	PR D79 112009	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AD	PR D80 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AE	PR D80 031102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AF	PR D80 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AG	PR D80 051105	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AL	PR D80 092007	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AO	PRL 103 211802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AU	PR D80 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09AV	PR D80 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09C	PR D79 032002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09G	PRL 102 141802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09H	PR D79 052005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09I	PR D79 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09K	PR D79 072009	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09S	PR D79 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09T	PRL 102 091803	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		EPAPS Document No. E-PRLTAO-102-060910		(BABAR Collab.)
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHANG	09	PR D79 052006	Y.-W. Chang <i>et al.</i>	(BELLE Collab.)
DALSENO	09	PR D79 072004	J. Dalseno <i>et al.</i>	(BELLE Collab.)
KYEONG	09	PR D80 051103	S.-H. Kyeong <i>et al.</i>	(BELLE Collab.)
MIZUK	09	PR D80 031104	R. Mizuk <i>et al.</i>	(BELLE Collab.)
VERVINK	09	PR D80 111104	K. Vervink <i>et al.</i>	(BELLE Collab.)
WEI	09A	PRL 103 171801	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
Also		EPAPS Supplement	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
ADACHI	08	PR D77 091101	I. Adachi <i>et al.</i>	(BELLE Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AC	PR D77 071102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AD	PR D77 091104	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AF	PR D78 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AG	PR D78 011104	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AH	PR D78 011107	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AJ	PR D78 032005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AP	PR D78 051103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AQ	PR D78 052005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AU	PRL 101 021801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08AV	PRL 101 081801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BA	PR D78 071102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BB	PR D78 071104	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BC	PR D78 072007	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BD	PR D78 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BG	PR D78 092008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BH	PR D78 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BK	PRL 101 201801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BL	PRL 101 261802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BN	PR D78 112003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08C	PR D77 011104	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08E	PR D77 012003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08F	PRL 100 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08G	PRL 100 171803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08H	PR D77 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08I	PRL 100 081801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08N	PRL 100 021801	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D79 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08P	PR D77 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08Q	PRL 100 151802	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08R	PR D77 032002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08W	PRL 101 082001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHEN	08C	PRL 100 251801	J.-H. Chen <i>et al.</i>	(BELLE Collab.)
CHIANG	08	PR D78 111102	C.C. Chiang <i>et al.</i>	(BELLE Collab.)
CHOI	08	PRL 100 142001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
GOLDENZWE...	08	PRL 101 231801	P. Goldenzweig <i>et al.</i>	(BELLE Collab.)
KIM	08	PL B669 287	H.O. Kim <i>et al.</i>	(BELLE Collab.)
KUMAR	08	PR D78 091104	R. Kumar <i>et al.</i>	(BELLE Collab.)
KUSAKA	08	PR D77 072001	A. Kusaka <i>et al.</i>	(BELLE Collab.)
LEE	08A	PR D77 071101	S.E. Lee <i>et al.</i>	(BELLE Collab.)

