



$I(J^P) = 0(0^-)$   
 $I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

**$B_c^+$  MASS**

| VALUE (MeV)                                                                   | DOCUMENT ID             | TECN      | COMMENT                |
|-------------------------------------------------------------------------------|-------------------------|-----------|------------------------|
| <b>6274.47 ± 0.27 ± 0.17</b>                                                  | <sup>1</sup> AAIJ       | 20R LHCB  | $pp$ at 7, 8, 13 TeV   |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                         |           |                        |
| 6274.28 ± 1.40 ± 0.32                                                         | <sup>2</sup> AAIJ       | 17L LHCB  | Repl. by AAIJ 20R      |
| 6274.0 ± 1.8 ± 0.4                                                            | <sup>3</sup> AAIJ       | 14AQ LHCB | Repl. by AAIJ 20R      |
| 6276.28 ± 1.44 ± 0.36                                                         | <sup>4</sup> AAIJ       | 13AS LHCB | Repl. by AAIJ 20R      |
| 6273.7 ± 1.3 ± 1.6                                                            | <sup>5</sup> AAIJ       | 12AV LHCB | Repl. by AAIJ 20R      |
| 6275.6 ± 2.9 ± 2.5                                                            | <sup>6</sup> AALTONEN   | 08M CDF   | $p\bar{p}$ at 1.96 TeV |
| 6300 ± 14 ± 5                                                                 | <sup>6</sup> ABAZOV     | 08T D0    | $p\bar{p}$ at 1.96 TeV |
| 6285.7 ± 5.3 ± 1.2                                                            | <sup>6</sup> ABULENCIA  | 06C CDF   | Repl. by AALTONEN 08M  |
| 6400 ± 390 ± 130                                                              | <sup>7</sup> ABE        | 98M CDF   | $p\bar{p}$ at 1.8 TeV  |
| 6320 ± 60                                                                     | <sup>8</sup> ACKERSTAFF | 98O OPAL  | $e^+e^- \rightarrow Z$ |

- <sup>1</sup> AAIJ 20R uses the  $B_c^+ \rightarrow J/\psi \pi^+, J/\psi \pi^+ \pi^- \pi^+, J/\psi p\bar{p} \pi^+, J/\psi D_s^+, J/\psi D^0 K^+$  and  $B_s^0 \pi^+$  modes.
- <sup>2</sup> Measured using  $B_c^+ \rightarrow J/\psi D^0 K^+$  decays.
- <sup>3</sup> Uses  $B_c^+ \rightarrow J/\psi p\bar{p} \pi^+$  decays.
- <sup>4</sup> AAIJ 13AS uses the  $B_c^+ \rightarrow J/\psi D_s^+$ .
- <sup>5</sup> AAIJ 12AV uses the  $B_c^+ \rightarrow J/\psi \pi^+$  mode and also measures the mass difference  $M(B_c^+) - M(B^+) = 994.6 \pm 1.3 \pm 0.6 \text{ MeV}/c^2$ .
- <sup>6</sup> Measured using a fully reconstructed decay mode of  $B_c \rightarrow J/\psi \pi$ .
- <sup>7</sup> ABE 98M observed  $20.4^{+6.2}_{-5.5}$  events in the  $B_c^+ \rightarrow J/\psi(1S) \ell \nu_\ell$  with a significance of  $> 4.8$  standard deviations. The mass value is estimated from  $m(J/\psi(1S) \ell)$ .
- <sup>8</sup> ACKERSTAFF 98O observed 2 candidate events in the  $B_c^+ \rightarrow J/\psi(1S) \pi^+$  channel with an estimated background of  $0.63 \pm 0.20$  events.

**$m_{B_c^+} - m_{B_s^0}$**

| VALUE (MeV)                                                                                                                                                             | DOCUMENT ID       | TECN     | COMMENT              |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------|----------|----------------------|
| <b>907.75 ± 0.37 ± 0.27</b>                                                                                                                                             | <sup>1</sup> AAIJ | 20R LHCB | $pp$ at 7, 8, 13 TeV |
| <sup>1</sup> AAIJ 20R uses the $B_c^+ \rightarrow J/\psi \pi^+, J/\psi \pi^+ \pi^- \pi^+, J/\psi p\bar{p} \pi^+, J/\psi D_s^+, J/\psi D^0 K^+$ and $B_s^0 \pi^+$ modes. |                   |          |                      |

**$B_c^+$  MEAN LIFE**

| VALUE ( $10^{-12} \text{ s}$ )      | DOCUMENT ID           | TECN     | COMMENT          |
|-------------------------------------|-----------------------|----------|------------------|
| <b>0.510 ± 0.009 OUR EVALUATION</b> | (Produced by HFLAV)   |          |                  |
| <b>0.510 ± 0.009 OUR AVERAGE</b>    |                       |          |                  |
| 0.541 ± 0.026 ± 0.014               | <sup>1</sup> SIRUNYAN | 18BY CMS | $pp$ at 8 TeV    |
| 0.5134 ± 0.0110 ± 0.0057            | <sup>2,3</sup> AAIJ   | 15G LHCB | $pp$ at 7, 8 TeV |
| 0.509 ± 0.008 ± 0.012               | <sup>4</sup> AAIJ     | 14G LHCB | $pp$ at 8 TeV    |

|                                           |                        |     |     |                        |
|-------------------------------------------|------------------------|-----|-----|------------------------|
| 0.452 $\pm 0.048$ $\pm 0.027$             | <sup>3</sup> AALTONEN  | 13  | CDF | $p\bar{p}$ at 1.96 TeV |
| 0.448 $\pm 0.038$ $\pm 0.032$<br>$-0.036$ | <sup>5</sup> ABAZOV    | 09H | D0  | $p\bar{p}$ at 1.96 TeV |
| 0.463 $\pm 0.073$ $\pm 0.036$<br>$-0.065$ | <sup>5</sup> ABULENCIA | 06O | CDF | $p\bar{p}$ at 1.96 TeV |
| 0.46 $\pm 0.18$ $\pm 0.03$<br>$-0.16$     | <sup>5</sup> ABE       | 98M | CDF | $p\bar{p}$ 1.8 TeV     |

<sup>1</sup> The lifetime is measured using the decays  $B_c^+ \rightarrow J/\psi \pi^+$  and  $B^+ \rightarrow J/\psi K^+$ .

<sup>2</sup> Also measures the width difference  $\Delta\Gamma = \Gamma_{B_c^+} - \Gamma_{B^+} = 4.46 \pm 0.14 \pm 0.07 \text{ mm}^{-1} \text{ c}$ .

<sup>3</sup> Uses fully reconstructed  $B_c^+ \rightarrow J/\psi \pi^+$  decays.

<sup>4</sup> Measured using  $B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu X$  decays.

<sup>5</sup> The lifetime is measured from the  $J/\psi e$  decay vertices.

## $B_c^+$ DECAY MODES $\times B(\bar{b} \rightarrow B_c)$

The following quantities are not pure branching ratios; rather the fractions  $\Gamma_i/\Gamma \times B(\bar{b} \rightarrow B_c)$ .  $B_c^-$  modes are charge conjugates of the modes below.

| Mode                                                | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level |
|-----------------------------------------------------|--------------------------------|------------------|
| $\Gamma_1$ $J/\psi(1S)\ell^+\nu_\ell$ anything      | seen                           |                  |
| $\Gamma_2$ $J/\psi(1S)\mu^+\nu_\mu$                 | seen                           |                  |
| $\Gamma_3$ $J/\psi(1S)\tau^+\nu_\tau$               | seen                           |                  |
| $\Gamma_4$ $J/\psi(1S)\pi^+$                        | seen                           |                  |
| $\Gamma_5$ $J/\psi(1S)K^+$                          | seen                           |                  |
| $\Gamma_6$ $J/\psi(1S)\pi^+\pi^0$                   |                                |                  |
| $\Gamma_7$ $J/\psi(1S)\pi^+\pi^+\pi^-$              | seen                           |                  |
| $\Gamma_8$ $J/\psi(1S)K^+\pi^-\pi^+$                |                                |                  |
| $\Gamma_9$ $J/\psi(1S)K^+K^-\pi^+$                  |                                |                  |
| $\Gamma_{10}$ $J/\psi(1S)a_1(1260)$                 | not seen                       |                  |
| $\Gamma_{11}$ $J/\psi(1S)K^+K^-\pi^+$               | seen                           |                  |
| $\Gamma_{12}$ $J/\psi(1S)\pi^+\pi^+\pi^+\pi^-\pi^-$ | seen                           |                  |
| $\Gamma_{13}$ $\psi(2S)\pi^+$                       | seen                           |                  |
| $\Gamma_{14}$ $\psi(2S)\pi^+\pi^-\pi^+$             |                                |                  |
| $\Gamma_{15}$ $\psi(2S)K^+K^-\pi^+$                 |                                |                  |
| $\Gamma_{16}$ $J/\psi(1S)D^0K^+$                    | seen                           |                  |
| $\Gamma_{17}$ $J/\psi(1S)D^*(2007)^0K^+$            | seen                           |                  |
| $\Gamma_{18}$ $J/\psi(1S)D^*(2010)^+K^{*0}$         | seen                           |                  |
| $\Gamma_{19}$ $J/\psi(1S)D^+K^{*0}$                 | seen                           |                  |
| $\Gamma_{20}$ $J/\psi(1S)D_s^+$                     | seen                           |                  |
| $\Gamma_{21}$ $J/\psi(1S)D_s^{*+}$                  | seen                           |                  |
| $\Gamma_{22}$ $J/\psi(1S)p\bar{p}\pi^+$             | seen                           |                  |
| $\Gamma_{23}$ $\chi_{c0}\pi^+$                      |                                |                  |
| $\Gamma_{24}$ $\chi_{c1}\pi^+$                      |                                |                  |

|               |                                                                |                        |     |
|---------------|----------------------------------------------------------------|------------------------|-----|
| $\Gamma_{25}$ | $\chi_{c2} \pi^+$                                              |                        |     |
| $\Gamma_{26}$ | $p \bar{p} \pi^+$                                              | not seen               |     |
| $\Gamma_{27}$ | $D^0 K^+$                                                      | seen                   |     |
| $\Gamma_{28}$ | $D^0 \pi^+$                                                    | not seen               |     |
| $\Gamma_{29}$ | $D^{*0} \pi^+$                                                 | not seen               |     |
| $\Gamma_{30}$ | $D^{*0} K^+$                                                   | not seen               |     |
| $\Gamma_{31}$ | $D_s^+ \bar{D}^0$                                              | $< 7.2 \times 10^{-4}$ | 90% |
| $\Gamma_{32}$ | $D_s^+ D^0$                                                    | $< 3.0 \times 10^{-4}$ | 90% |
| $\Gamma_{33}$ | $D_s^+ \bar{D}^0$                                              | $< 1.9 \times 10^{-4}$ | 90% |
| $\Gamma_{34}$ | $D^+ D^0$                                                      | $< 1.4 \times 10^{-4}$ | 90% |
| $\Gamma_{35}$ | $D_s^{*+} \bar{D}^0$                                           | $< 5.3 \times 10^{-4}$ | 90% |
| $\Gamma_{36}$ | $D_s^+ \bar{D}^*(2007)^0$                                      | $< 4.6 \times 10^{-4}$ | 90% |
| $\Gamma_{37}$ | $D_s^{*+} D^0$                                                 | $< 9 \times 10^{-4}$   | 90% |
| $\Gamma_{38}$ | $D_s^+ D^*(2007)^0$                                            | $< 6.6 \times 10^{-4}$ | 90% |
| $\Gamma_{39}$ | $D^*(2010)^+ \bar{D}^0$                                        | $< 3.8 \times 10^{-4}$ | 90% |
| $\Gamma_{40}$ | $D^*(2010)^+ \bar{D}^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma$ | not seen               |     |
| $\Gamma_{41}$ | $D^+ \bar{D}^*(2007)^0$                                        | $< 6.5 \times 10^{-4}$ | 90% |
| $\Gamma_{42}$ | $D^*(2007)^+ D^0$                                              | $< 2.0 \times 10^{-4}$ | 90% |
| $\Gamma_{43}$ | $D^*(2010)^+ D^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma$       | not seen               |     |
| $\Gamma_{44}$ | $D^+ D^*(2007)^0$                                              | $< 3.7 \times 10^{-4}$ | 90% |
| $\Gamma_{45}$ | $D_s^{*+} \bar{D}^*(2007)^0$                                   | $< 1.3 \times 10^{-3}$ | 90% |
| $\Gamma_{46}$ | $D_s^{*+} D^*(2007)^0$                                         | $< 1.3 \times 10^{-3}$ | 90% |
| $\Gamma_{47}$ | $D^*(2010)^+ \bar{D}^*(2007)^0$                                | $< 1.0 \times 10^{-3}$ | 90% |
| $\Gamma_{48}$ | $D^*(2010)^+ D^*(2007)^0$                                      | $< 7.7 \times 10^{-4}$ | 90% |
| $\Gamma_{49}$ | $D^+ K^{*0}$                                                   | not seen               |     |
| $\Gamma_{50}$ | $D^+ \bar{K}^{*0}$                                             | not seen               |     |
| $\Gamma_{51}$ | $D_s^+ K^{*0}$                                                 | not seen               |     |
| $\Gamma_{52}$ | $D_s^+ \bar{K}^{*0}$                                           | not seen               |     |
| $\Gamma_{53}$ | $D_s^+ \phi$                                                   | not seen               |     |
| $\Gamma_{54}$ | $K^+ K^0$                                                      | not seen               |     |
| $\Gamma_{55}$ | $B_s^0 \pi^+ / B(\bar{b} \rightarrow B_s)$                     | seen                   |     |
| $\Gamma_{56}$ | $B_s^0 \pi^+$                                                  |                        |     |
| $\Gamma_{57}$ | $\pi^+ \mu^+ \mu^-$                                            |                        |     |
| $\Gamma_{58}$ | $D_s^+ \mu^+ \mu^-$                                            |                        |     |

### $B_c^+$ BRANCHING RATIOS

| $\Gamma(J/\psi(1S) \ell^+ \nu_\ell \text{ anything}) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ |     |                                     | $\Gamma_1 / \Gamma \times B$ |                         |  |
|-----------------------------------------------------------------------------------------------------------------|-----|-------------------------------------|------------------------------|-------------------------|--|
| VALUE (units $10^{-5}$ )                                                                                        | CL% | DOCUMENT ID                         | TECN                         | COMMENT                 |  |
| <b><math>8.2 \pm 1.3</math> OUR AVERAGE</b>                                                                     |     | Error includes scale factor of 1.4. |                              |                         |  |
| $8.8 \pm 1.0 \pm 0.2$                                                                                           |     | <sup>1,2</sup> AALTONEN             | 16A CDF                      | $p \bar{p}$ at 1.96 TeV |  |
| $5.2^{+2.4}_{-2.1}$                                                                                             |     | <sup>3</sup> ABE                    | 98M CDF                      | $p \bar{p}$ 1.8 TeV     |  |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|     |    |                         |     |      |                        |
|-----|----|-------------------------|-----|------|------------------------|
| <16 | 90 | <sup>4</sup> ACKERSTAFF | 98O | OPAL | $e^+e^- \rightarrow Z$ |
| <19 | 90 | <sup>5</sup> ABREU      | 97E | DLPH | $e^+e^- \rightarrow Z$ |
| <12 | 90 | <sup>6</sup> BARATE     | 97H | ALEP | $e^+e^- \rightarrow Z$ |

<sup>1</sup> AALTONEN 16A reports  $[\Gamma(B_c^+ \rightarrow J/\psi(1S)\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.211 \pm 0.012^{+0.021}_{-0.020}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = (40.8 \pm 0.7) \times 10^{-2}$ ,  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>2</sup> AALTONEN 16A also measures the cross-section  $\sigma(B_c) \times B(B_c \rightarrow J/\psi\mu\nu_\mu) = 0.60 \pm 0.09$  nb and estimates the total cross-section  $\sigma(B_c)$  to be in the range  $25 \pm 4$  to  $52 \pm 8$  nb for  $p_T(B_c) > 6$  GeV/c and  $|y(B_c)| < 1$ .

<sup>3</sup> ABE 98M result is derived from the measurement of  $[\sigma(B_c) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell)] / [\sigma(B^+) \times B(B^+ \rightarrow J/\psi(1S)K^+)] = 0.132^{+0.041}_{-0.037}(\text{stat}) \pm 0.031(\text{sys})^{+0.032}_{-0.020}(\text{lifetime})$  by using PDG 98 values of  $B(b \rightarrow B^+)$  and  $B(B^+ \rightarrow J/\psi(1S)K^+)$ .

<sup>4</sup> ACKERSTAFF 98O reports  $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 6.95 \times 10^{-5}$  at 90%CL. We rescale to our PDG 98 values of  $B(Z \rightarrow b\bar{b})$ .

<sup>5</sup> ABREU 97E value listed is for an assumed  $\tau_{B_c} = 0.4$  ps and improves to  $1.6 \times 10^{-4}$  for  $\tau_{B_c} = 1.4$  ps.

<sup>6</sup> BARATE 97H reports  $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S)\ell\nu_\ell) < 5.2 \times 10^{-5}$  at 90%CL. We rescale to our PDG 96 values of  $B(Z \rightarrow b\bar{b})$ . A  $B_c^+ \rightarrow J/\psi(1S)\mu^+\nu_\mu$  candidate event is found, compared to all the known background sources  $2 \times 10^{-3}$ , which gives  $m_{B_c} = 5.96^{+0.25}_{-0.19}$  GeV and  $\tau_{B_c} = 1.77 \pm 0.17$  ps.

### $\Gamma(J/\psi(1S)\tau^+\nu_\tau)/\Gamma(J/\psi(1S)\mu^+\nu_\mu)$ $\Gamma_3/\Gamma_2$

| VALUE                 | DOCUMENT ID       | TECN | COMMENT               |
|-----------------------|-------------------|------|-----------------------|
| <b>0.71±0.17±0.18</b> | <sup>1</sup> AAIJ | 18C  | LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 18C uses  $\tau^+ \rightarrow \mu^+\nu_\mu\bar{\nu}_\tau$  mode to obtain the ratio value.

### $\Gamma(J/\psi(1S)\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_4/\Gamma \times B$

| VALUE | CL% | DOCUMENT ID                   | TECN | COMMENT                    |
|-------|-----|-------------------------------|------|----------------------------|
| seen  |     | <sup>1</sup> AABOUD           | 21   | ATLS $pp$ at 8 TeV         |
| seen  |     | <sup>2</sup> AAIJ             | 15M  | LHCB $pp$ at 8 TeV         |
| seen  |     | <sup>3</sup> KHACHATRY...15AA | CMS  | $pp$ at 7 TeV              |
| seen  |     | AALTONEN                      | 13   | CDF $p\bar{p}$ at 1.96 TeV |
| seen  |     | <sup>4</sup> AAIJ             | 12AV | LHCB $pp$ at 7 TeV         |
| seen  |     | AALTONEN                      | 08M  | CDF $p\bar{p}$ at 1.96 TeV |
| seen  |     | ABAZOV                        | 08T  | D0 $p\bar{p}$ at 1.96 TeV  |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                         |    |                         |     |      |                        |
|-------------------------|----|-------------------------|-----|------|------------------------|
| <2.4 × 10 <sup>-4</sup> | 90 | <sup>5</sup> ACKERSTAFF | 98O | OPAL | $e^+e^- \rightarrow Z$ |
| <3.4 × 10 <sup>-4</sup> | 90 | <sup>6</sup> ABREU      | 97E | DLPH | $e^+e^- \rightarrow Z$ |
| <8.2 × 10 <sup>-5</sup> | 90 | <sup>7</sup> BARATE     | 97H | ALEP | $e^+e^- \rightarrow Z$ |
| <2.0 × 10 <sup>-5</sup> | 95 | <sup>8</sup> ABE        | 96R | CDF  | $p\bar{p}$ 1.8 TeV     |

- <sup>1</sup> AABOUD 21 reports a measurement of  $B(B_c^+ \rightarrow J/\psi \pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.34 \pm 0.04_{-0.02}^{+0.06} \pm 0.01) \%$ , at  $p_T > 13$  GeV and  $|y| < 2.3$ .
- <sup>2</sup> AAIJ 15M reports a measurement of  $B(B_c^+ \rightarrow J/\psi \pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.683 \pm 0.018 \pm 0.009) \%$  at  $p_T(B) < 20$  GeV and  $2.0 < y(B) < 4.5$ .
- <sup>3</sup> KHACHATRYAN 15AA reports a measurement of  $B(B_c^+ \rightarrow J/\psi \pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.48 \pm 0.05 \pm 0.03 \pm 0.05) \%$ , at  $p_T > 15$  GeV and  $|\eta(B)| < 1.6$ .
- <sup>4</sup> AAIJ 12AV reports a measurement of  $B(B_c^+ \rightarrow J/\psi \pi^+) / B(B^+ \rightarrow J/\psi K^+) \cdot f_c/f_u = (0.68 \pm 0.10 \pm 0.03 \pm 0.05) \%$  at  $p_T(B) > 4$  GeV and  $2.5 < \eta(B) < 4.5$ .
- <sup>5</sup> ACKERSTAFF 98O reports  $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S) \pi^+) < 1.06 \times 10^{-4}$  at 90%CL. We rescale to our PDG 98 values of  $B(Z \rightarrow b\bar{b})$ .
- <sup>6</sup> ABREU 97E value listed is for an assumed  $\tau_{B_c} = 0.4$  ps and improves to  $2.7 \times 10^{-4}$  for  $\tau_{B_c} = 1.4$  ps.
- <sup>7</sup> BARATE 97H reports  $B(Z \rightarrow B_c X) / B(Z \rightarrow qq) \cdot B(B_c \rightarrow J/\psi(1S) \pi) < 3.6 \times 10^{-5}$  at 90%CL. We rescale to our PDG 96 values of  $B(Z \rightarrow b\bar{b})$ .
- <sup>8</sup> ABE 96R reports  $B(b \rightarrow B_c X) / B(b \rightarrow B^+ X) \cdot B(B_c^+ \rightarrow J/\psi(1S) \pi^+) / B(B^+ \rightarrow J/\psi(1S) K^+) < 0.053$  at 95%CL for  $\tau_{B_c} = 0.8$  ps. It changes from 0.15 to 0.04 for  $0.17 \text{ ps} < \tau_{B_c} < 1.6$  ps. We rescale to our PDG 96 values of  $B(b \rightarrow B^+) = 0.378 \pm 0.022$  and  $B(B^+ \rightarrow J/\psi(1S) K^+) = 0.00101 \pm 0.00014$ .

### $\Gamma(J/\psi(1S)\pi^+)/\Gamma(J/\psi(1S)\mu^+\nu_\mu)$ $\Gamma_4/\Gamma_2$

| VALUE (units $10^{-2}$ )                   | DOCUMENT ID       | TECN     | COMMENT       |
|--------------------------------------------|-------------------|----------|---------------|
| <b><math>4.69 \pm 0.28 \pm 0.46</math></b> | <sup>1</sup> AAIJ | 14W LHCb | $pp$ at 7 TeV |

- <sup>1</sup> AAIJ 14W reports also a measurement  $B(B_c^+ \rightarrow J/\psi \pi^+) / B(B_c^+ \rightarrow J/\psi \mu^+ \nu_\mu) = 0.271 \pm 0.016 \pm 0.016$  in the region  $m_{J/\psi \mu^+} > 5.3$  GeV.

### $\Gamma(J/\psi(1S)K^+)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_5/\Gamma_4$

| VALUE                                         | EVTs | DOCUMENT ID | TECN      | COMMENT          |
|-----------------------------------------------|------|-------------|-----------|------------------|
| <b><math>0.079 \pm 0.007 \pm 0.003</math></b> |      | AAIJ        | 16AF LHCb | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                             |    |      |           |                    |
|-----------------------------|----|------|-----------|--------------------|
| $0.069 \pm 0.019 \pm 0.005$ | 50 | AAIJ | 13BY LHCb | Repl. by AAIJ 16AF |
|-----------------------------|----|------|-----------|--------------------|

### $\Gamma(J/\psi(1S)\pi^+\pi^0)/\Gamma(J/\psi(1S)\pi^+)$ $\Gamma_6/\Gamma_4$

| VALUE                                      | DOCUMENT ID | TECN     | COMMENT              |
|--------------------------------------------|-------------|----------|----------------------|
| <b><math>2.80 \pm 0.15 \pm 0.19</math></b> | AAIJ        | 24L LHCb | $pp$ at 7, 8, 13 TeV |

### $\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ $\Gamma_7/\Gamma \times B$

| VALUE | CL% | DOCUMENT ID | TECN     | COMMENT       |
|-------|-----|-------------|----------|---------------|
| seen  |     | AAIJ        | 12Y LHCb | $pp$ at 7 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                        |    |                    |          |                        |
|------------------------|----|--------------------|----------|------------------------|
| $< 5.7 \times 10^{-4}$ | 90 | <sup>1</sup> ABREU | 97E DLPH | $e^+e^- \rightarrow Z$ |
|------------------------|----|--------------------|----------|------------------------|

- <sup>1</sup> ABREU 97E value listed is independent of  $0.4 \text{ ps} < \tau_{B_c} < 1.4$  ps.

$$\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_7/\Gamma_4$$

| VALUE                                                | DOCUMENT ID      | TECN     | COMMENT       |
|------------------------------------------------------|------------------|----------|---------------|
| <b>2.4 ± 0.4 OUR AVERAGE</b>                         |                  |          |               |
| 2.55 ± 0.80 ± 0.33 <sup>+0.04</sup> <sub>-0.01</sub> | KHACHATRY...15AA | CMS      | $pp$ at 7 TeV |
| 2.41 ± 0.30 ± 0.33                                   | AAIJ             | 12Y LHCb | $pp$ at 7 TeV |

$$\Gamma(J/\psi(1S)a_1(1260))/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{10}/\Gamma \times B$$

| VALUE                             | CL% | DOCUMENT ID                 | TECN | COMMENT                |
|-----------------------------------|-----|-----------------------------|------|------------------------|
| <b>&lt; 1.2 × 10<sup>-3</sup></b> | 90  | <sup>1</sup> ACKERSTAFF 98O | OPAL | $e^+e^- \rightarrow Z$ |

<sup>1</sup> ACKERSTAFF 98O reports  $B(Z \rightarrow B_c X)/B(Z \rightarrow qq) \times B(B_c \rightarrow J/\psi(1S)a_1(1260)) < 5.29 \times 10^{-4}$  at 90%CL. We rescale to our PDG 98 values of  $B(Z \rightarrow b\bar{b})$ .

$$\Gamma(J/\psi(1S)K^+K^-\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{11}/\Gamma \times B$$

| VALUE       | DOCUMENT ID       | TECN      | COMMENT          |
|-------------|-------------------|-----------|------------------|
| <b>seen</b> | <sup>1</sup> AAIJ | 13CA LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> A signal yield of  $78 \pm 14$  decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of  $3 \text{ fb}^{-1}$  data.

$$\Gamma(J/\psi(1S)K^+K^-\pi^+)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{11}/\Gamma_4$$

| VALUE                     | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------|-------------------|-----------|------------------|
| <b>0.53 ± 0.10 ± 0.05</b> | <sup>1</sup> AAIJ | 13CA LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> A signal yield of  $78 \pm 14$  decays is reported with a significance of 6.2 standard deviations using an integrated luminosity of  $3 \text{ fb}^{-1}$  data.

$$\Gamma(J/\psi(1S)\pi^+\pi^+\pi^+\pi^-\pi^-)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{12}/\Gamma_4$$

| VALUE                     | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------------|-------------------|----------|------------------|
| <b>1.74 ± 0.44 ± 0.24</b> | <sup>1</sup> AAIJ | 14P LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> A signal yield of  $32 \pm 8$  decays is reported with a significance of 4.5 standard deviations.

$$\Gamma(J/\psi(1S)D_s^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{20}/\Gamma \times B$$

| VALUE (units 10 <sup>-6</sup> ) | DOCUMENT ID       | TECN     | COMMENT              |
|---------------------------------|-------------------|----------|----------------------|
| <b>6.7 ± 0.8 ± 0.1</b>          | <sup>1</sup> AAIJ | 24F LHCb | $pp$ at 7, 8, 13 TeV |

<sup>1</sup> AAIJ 24F reports  $[\Gamma(B_c^+ \rightarrow J/\psi(1S)D_s^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] = (1.63 \pm 0.15 \pm 0.13) \times 10^{-5}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(J/\psi(1S)D_s^{*+})/\Gamma(J/\psi(1S)D_s^+) \quad \Gamma_{21}/\Gamma_{20}$$

| VALUE                     | DOCUMENT ID | TECN     | COMMENT              |
|---------------------------|-------------|----------|----------------------|
| <b>1.91 ± 0.20 ± 0.07</b> | AAIJ        | 24F LHCb | $pp$ at 7, 8, 13 TeV |

$$\Gamma(\psi(2S)\pi^+)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{13}/\Gamma_4$$

| VALUE                        | DOCUMENT ID         | TECN     | COMMENT                 |
|------------------------------|---------------------|----------|-------------------------|
| <b>0.254 ± 0.018 ± 0.006</b> | <sup>1,2</sup> AAIJ | 24W LHCb | $pp$ at 7, 8 and 13 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                               |                     |           |                    |
|-------------------------------|---------------------|-----------|--------------------|
| 0.268 ± 0.032 ± 0.007 ± 0.006 | <sup>3,4</sup> AAIJ | 15AY LHCb | $pp$ at 7, 8 TeV   |
| 0.250 ± 0.068 ± 0.014 ± 0.006 | <sup>3</sup> AAIJ   | 13AMLHCb  | Repl. by AAIJ 15AY |

<sup>1</sup> The last uncertainty includes the uncertainties on the branching fractions of the leptonic  $J/\psi$  and  $\psi(2S)$  decays ( $\pm 0.005$ ).

<sup>2</sup> Supersedes AAIJ 15AY.

<sup>3</sup> The last uncertainty is due to the uncertainty of the  $B(\psi(2S) \rightarrow \mu^+ \mu^-)/B(J/\psi \rightarrow \mu^+ \mu^-)$  ratio measurement.

<sup>4</sup> Replaced by AAIJ 24W.

| $\Gamma(\psi(2S)\pi^+)/\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)$ <span style="float:right"><math>\Gamma_{13}/\Gamma_7</math></span>           |             |      |                           |
|----------------------------------------------------------------------------------------------------------------------------------------|-------------|------|---------------------------|
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $(3.5 \pm 0.6 \pm 0.2) \times 10^{-2}$                                                                                                 | AAIJ        | 22P  | LHCB $pp$ at 7, 8, 13 TeV |
| $\Gamma(\psi(2S)K^+K^-\pi^+)/\Gamma(J/\psi(1S)K^+K^-\pi^+)$ <span style="float:right"><math>\Gamma_{15}/\Gamma_{11}</math></span>      |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $(3.7 \pm 1.2 \pm 0.1) \times 10^{-2}$                                                                                                 | AAIJ        | 22P  | LHCB $pp$ at 7, 8, 13 TeV |
| $\Gamma(J/\psi(1S)K^+K^-K^+)/\Gamma(J/\psi(1S)K^+K^-\pi^+)$ <span style="float:right"><math>\Gamma_9/\Gamma_{11}</math></span>         |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $(7.0 \pm 1.8 \pm 0.2) \times 10^{-2}$                                                                                                 | AAIJ        | 22P  | LHCB $pp$ at 7, 8, 13 TeV |
| $\Gamma(J/\psi(1S)K^+\pi^-\pi^+)/\Gamma(J/\psi(1S)K^+K^-\pi^+)$ <span style="float:right"><math>\Gamma_8/\Gamma_{11}</math></span>     |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $0.35 \pm 0.06 \pm 0.01$                                                                                                               | AAIJ        | 22P  | LHCB $pp$ at 7, 8, 13 TeV |
| $\Gamma(J/\psi(1S)K^+K^-\pi^+)/\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)$ <span style="float:right"><math>\Gamma_{11}/\Gamma_7</math></span>   |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $0.185 \pm 0.013 \pm 0.006$                                                                                                            | AAIJ        | 22P  | LHCB $pp$ at 7, 8, 13 TeV |
| $\Gamma(\psi(2S)\pi^+\pi^-\pi^+)/\Gamma(J/\psi(1S)\pi^+\pi^+\pi^-)$ <span style="float:right"><math>\Gamma_{14}/\Gamma_7</math></span> |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $(1.9 \pm 0.4 \pm 0.1) \times 10^{-2}$                                                                                                 | AAIJ        | 22P  | LHCB $pp$ at 7, 8, 13 TeV |
| $\Gamma(J/\psi(1S)D^0K^+)/\Gamma(J/\psi(1S)\pi^+)$ <span style="float:right"><math>\Gamma_{16}/\Gamma_4</math></span>                  |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $0.432 \pm 0.136 \pm 0.028$                                                                                                            | AAIJ        | 17L  | LHCB $pp$ at 7, 8 TeV     |
| $\Gamma(J/\psi(1S)D^*(2007)^0K^+)/\Gamma(J/\psi(1S)D^0K^+)$ <span style="float:right"><math>\Gamma_{17}/\Gamma_{16}</math></span>      |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $5.1 \pm 1.8 \pm 0.4$                                                                                                                  | AAIJ        | 17L  | LHCB $pp$ at 7, 8 TeV     |
| $\Gamma(J/\psi(1S)D^*(2010)^+K^{*0})/\Gamma(J/\psi(1S)D^0K^+)$ <span style="float:right"><math>\Gamma_{18}/\Gamma_{16}</math></span>   |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $2.10 \pm 1.08 \pm 0.34$                                                                                                               | AAIJ        | 17L  | LHCB $pp$ at 7, 8 TeV     |
| $\Gamma(J/\psi(1S)D^+K^{*0})/\Gamma(J/\psi(1S)D^0K^+)$ <span style="float:right"><math>\Gamma_{19}/\Gamma_{16}</math></span>           |             |      |                           |
| VALUE                                                                                                                                  | DOCUMENT ID | TECN | COMMENT                   |
| $0.63 \pm 0.39 \pm 0.08$                                                                                                               | AAIJ        | 17L  | LHCB $pp$ at 7, 8 TeV     |

$$\Gamma(J/\psi(1S)D_s^+)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{20}/\Gamma_4$$

| VALUE                                                                         | DOCUMENT ID      | TECN      | COMMENT          |
|-------------------------------------------------------------------------------|------------------|-----------|------------------|
| <b><math>2.8 \pm 0.4</math> OUR AVERAGE</b>                                   |                  |           |                  |
| $2.76 \pm 0.33 \pm 0.33$                                                      | <sup>1</sup> AAD | 220 ATLS  | $pp$ at 13 TeV   |
| $2.90 \pm 0.57 \pm 0.24$                                                      | AAIJ             | 13AS LHCB | $pp$ at 7, 8 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                  |           |                  |
| $3.8 \pm 1.1 \pm 0.4$                                                         | AAD              | 16H ATLS  | $pp$ at 7, 8 TeV |

<sup>1</sup>Supersedes the measurement of AAD 16H.

$$\Gamma(J/\psi(1S)D_s^{*+})/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{21}/\Gamma_4$$

| VALUE                                                                         | DOCUMENT ID | TECN     | COMMENT          |
|-------------------------------------------------------------------------------|-------------|----------|------------------|
| <b><math>5.33 \pm 0.61 \pm 0.74</math></b>                                    | AAD         | 220 ATLS | $pp$ at 13 TeV   |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |             |          |                  |
| $10.4 \pm 3.1 \pm 1.6$                                                        | AAD         | 16H ATLS | Repl. by AAD 220 |

$$\Gamma(J/\psi(1S)D_s^{*+})/\Gamma(J/\psi(1S)D_s^+) \quad \Gamma_{21}/\Gamma_{20}$$

| VALUE                                                                         | DOCUMENT ID      | TECN      | COMMENT          |
|-------------------------------------------------------------------------------|------------------|-----------|------------------|
| <b><math>2.00 \pm 0.23</math> OUR AVERAGE</b>                                 |                  |           |                  |
| $1.93 \pm 0.24 \pm 0.09$                                                      | <sup>1</sup> AAD | 220 ATLS  | $pp$ at 13 TeV   |
| $2.37 \pm 0.56 \pm 0.10$                                                      | AAIJ             | 13AS LHCB | $pp$ at 7, 8 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • |                  |           |                  |
| $2.8 \begin{smallmatrix} +1.2 \\ -0.8 \end{smallmatrix} \pm 0.3$              | AAD              | 16H ATLS  | $pp$ at 7, 8 TeV |

<sup>1</sup>Supersedes the measurement of AAD 16H.

$$\Gamma(J/\psi(1S)p\bar{p}\pi^+)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{22}/\Gamma_4$$

| VALUE                                                                            | DOCUMENT ID | TECN      | COMMENT          |
|----------------------------------------------------------------------------------|-------------|-----------|------------------|
| <b><math>0.143 \begin{smallmatrix} +0.041 \\ -0.036 \end{smallmatrix}</math></b> | AAIJ        | 14AQ LHCB | $pp$ at 7, 8 TeV |

$$\Gamma(\chi_{c0}\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{23}/\Gamma \times B$$

| VALUE (units $10^{-6}$ )                                                            | DOCUMENT ID         | TECN      | COMMENT             |
|-------------------------------------------------------------------------------------|---------------------|-----------|---------------------|
| <b><math>24.0 \begin{smallmatrix} +8.6 \\ -7.6 \end{smallmatrix} \pm 0.4</math></b> | <sup>1,2</sup> AAIJ | 16AT LHCB | $pp$ at 7 and 8 TeV |

<sup>1</sup>AAIJ 16AT reports  $[\Gamma(B_c^+ \rightarrow \chi_{c0}\pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] \times [\Gamma(\bar{b} \rightarrow B^+)/\Gamma_{\text{total}}] = (9.8 \begin{smallmatrix} +3.4 \\ -3.0 \end{smallmatrix} \pm 0.8) \times 10^{-6}$  which we divide by our best value  $\Gamma(\bar{b} \rightarrow B^+)/\Gamma_{\text{total}} = 0.408 \pm 0.007$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup>The significance of the observed signal is 4.0 standard deviations.

$$\Gamma(\chi_{c1}\pi^+)/\Gamma(\chi_{c2}\pi^+) \quad \Gamma_{24}/\Gamma_{25}$$

| VALUE                        | CL% | DOCUMENT ID | TECN     | COMMENT                 |
|------------------------------|-----|-------------|----------|-------------------------|
| <b><math>&lt;0.49</math></b> | 90  | AAIJ        | 24i LHCB | $pp$ at 7, 8 and 13 TeV |

$$\Gamma(\chi_{c2}\pi^+)/\Gamma(J/\psi(1S)\pi^+) \quad \Gamma_{25}/\Gamma_4$$

| VALUE                                       | DOCUMENT ID       | TECN     | COMMENT                 |
|---------------------------------------------|-------------------|----------|-------------------------|
| <b><math>0.37 \pm 0.06 \pm 0.022</math></b> | <sup>1</sup> AAIJ | 24i LHCB | $pp$ at 7, 8 and 13 TeV |

<sup>1</sup>The last uncertainty includes the knowledge of the  $\chi_{c2} \rightarrow J/\psi\gamma$  branching fraction ( $\pm 0.01$ ).



$$\Gamma(p\bar{p}\pi^+)/\Gamma_{\text{total}} \quad \Gamma_{26}/\Gamma$$

| VALUE           | DOCUMENT ID       | TECN     | COMMENT          |
|-----------------|-------------------|----------|------------------|
| <b>not seen</b> | <sup>1</sup> AAIJ | 16K LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> Measures the ratio  $(f_c/f_u) \times B(B_c^+ \rightarrow p\bar{p}\pi^+) < 3.6 \times 10^{-8}$  at 95% CL, in the region  $m(p\bar{p}) < 2.85 \text{ GeV}/c^2$ , where  $f_c$  ( $f_u$ ) represents the fragmentation fraction of the  $b$ -quark into the  $B_c^+$  ( $B_u^+$ ) meson.

$$\Gamma(D^0 K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{27}/\Gamma \times B$$

| VALUE (units $10^{-7}$ )                      | DOCUMENT ID       | TECN      | COMMENT          |
|-----------------------------------------------|-------------------|-----------|------------------|
| <b><math>3.8^{+1.2}_{-1.0} \pm 0.1</math></b> | <sup>1</sup> AAIJ | 17AG LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^0 K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] = (9.3^{+2.8}_{-2.5} \pm 0.6) \times 10^{-7}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(D^0 \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{28}/\Gamma \times B$$

| VALUE                                       | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|---------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt; 1.6 \times 10^{-7}</math></b> | 95  | <sup>1</sup> AAIJ | 17AG LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^0 \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 3.9 \times 10^{-7}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(D^{*0} \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{29}/\Gamma \times B$$

| VALUE                                     | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|-------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt; 4 \times 10^{-7}</math></b> | 95  | <sup>1</sup> AAIJ | 17AG LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^{*0} \pi^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(D^{*0} K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{30}/\Gamma \times B$$

| VALUE                                     | CL% | DOCUMENT ID       | TECN      | COMMENT          |
|-------------------------------------------|-----|-------------------|-----------|------------------|
| <b><math>&lt; 4 \times 10^{-7}</math></b> | 95  | <sup>1</sup> AAIJ | 17AG LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 17AG reports  $[\Gamma(B_c^+ \rightarrow D^{*0} K^+)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(D_s^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{31}/\Gamma \times B$$

| VALUE                                       | CL% | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------------------------------|-----|-------------------|----------|------------------|
| <b><math>&lt; 1.4 \times 10^{-7}</math></b> | 90  | <sup>1</sup> AAIJ | 18P LHCB | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 0.9 \times 10^{-3}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D_s^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{32}/\Gamma \times B$$

| VALUE               | CL% | DOCUMENT ID       | TECN     | COMMENT          |
|---------------------|-----|-------------------|----------|------------------|
| $<6 \times 10^{-8}$ | 90  | <sup>1</sup> AAIJ | 18P LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 3.7 \times 10^{-4}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{33}/\Gamma \times B$$

| VALUE                 | CL% | DOCUMENT ID       | TECN     | COMMENT          |
|-----------------------|-----|-------------------|----------|------------------|
| $<3.0 \times 10^{-6}$ | 90  | <sup>1</sup> AAIJ | 18P LHCb | $pp$ at 7, 8 TeV |

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.9 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{34}/\Gamma \times B$$

| VALUE                 | CL% | DOCUMENT ID       | TECN     | COMMENT            |
|-----------------------|-----|-------------------|----------|--------------------|
| $<1.9 \times 10^{-6}$ | 90  | <sup>1</sup> AAIJ | 18P LHCb | Repl. by AAIJ 21AF |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D^+ D^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.2 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$[\Gamma(D_s^{*+} \bar{D}^0) + \Gamma(D_s^+ \bar{D}^{*0}(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad (\Gamma_{35} + \Gamma_{36})/\Gamma \times B$$

| VALUE               | CL% | DOCUMENT ID       | TECN     | COMMENT            |
|---------------------|-----|-------------------|----------|--------------------|
| $<4 \times 10^{-7}$ | 90  | <sup>1</sup> AAIJ | 18P LHCb | Repl. by AAIJ 21AF |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^{*+} \bar{D}^0) + \Gamma(B_c^+ \rightarrow D_s^+ \bar{D}^{*0}(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 2.8 \times 10^{-3}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$[\Gamma(D_s^{*+} D^0) + \Gamma(D_s^+ D^{*0}(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad (\Gamma_{37} + \Gamma_{38})/\Gamma \times B$$

| VALUE               | CL% | DOCUMENT ID       | TECN     | COMMENT            |
|---------------------|-----|-------------------|----------|--------------------|
| $<5 \times 10^{-7}$ | 90  | <sup>1</sup> AAIJ | 18P LHCb | Repl. by AAIJ 21AF |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D_s^{*+} D^0) + \Gamma(B_c^+ \rightarrow D_s^+ D^{*0}(2007)^0)]/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 3.0 \times 10^{-3}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D^{*0}(2010)^+ \bar{D}^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{39}/\Gamma \times B$$

| VALUE                 | CL% | DOCUMENT ID         | TECN     | COMMENT                 |
|-----------------------|-----|---------------------|----------|-------------------------|
| $<6.2 \times 10^{-3}$ | 90  | <sup>1</sup> BARATE | 98Q ALEP | $e^+ e^- \rightarrow Z$ |

<sup>1</sup> BARATE 98Q reports  $B(Z \rightarrow B_c X) \times B(B_c \rightarrow D^{*0}(2010)^+ \bar{D}^0) < 1.9 \times 10^{-3}$  at 90%CL. We rescale to our PDG 98 values of  $B(Z \rightarrow b \bar{b})$ .

$$\left[ \Gamma(D^*(2010)^+ \bar{D}^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma) + \Gamma(D^+ \bar{D}^*(2007)^0) \right] / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad (\Gamma_{40} + \Gamma_{41}) / \Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 9 \times 10^{-6}$       90      <sup>1</sup> AAIJ      18P    LHCB    Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $\left[ \Gamma(B_c^+ \rightarrow D^*(2010)^+ \bar{D}^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma) + \Gamma(B_c^+ \rightarrow D^+ \bar{D}^*(2007)^0) \right] / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)]$   
 $< 5.5 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  
 $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\left[ \Gamma(D^*(2010)^+ D^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma) + \Gamma(D^+ D^*(2007)^0) \right] / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad (\Gamma_{43} + \Gamma_{44}) / \Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.4 \times 10^{-6}$       90      <sup>1</sup> AAIJ      18P    LHCB    Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $\left[ \Gamma(B_c^+ \rightarrow D^*(2010)^+ D^0, D^{*+} \rightarrow D^+ \pi^0 / \gamma) + \Gamma(B_c^+ \rightarrow D^+ D^*(2007)^0) \right] / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)]$   
 $< 2.2 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  
 $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D_s^{*+} \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{45} / \Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.7 \times 10^{-6}$       90      <sup>1</sup> AAIJ      18P    LHCB    Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $\Gamma(B_c^+ \rightarrow D_s^{*+} \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.1 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D_s^{*+} D^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{46} / \Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.1 \times 10^{-6}$       90      <sup>1</sup> AAIJ      18P    LHCB    Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $\Gamma(B_c^+ \rightarrow D_s^{*+} D^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 2.0 \times 10^{-2}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D^*(2010)^+ \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{47} / \Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.0 \times 10^{-4}$       90      <sup>1</sup> AAIJ      18P    LHCB    Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $\Gamma(B_c^+ \rightarrow D^*(2010)^+ \bar{D}^*(2007)^0) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 6.5 \times 10^{-1}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D^*(2010)^+ D^*(2007)^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{48}/\Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.0 \times 10^{-5}$  90 <sup>1</sup> AAIJ 18P LHCB Repl. by AAIJ 21AF

<sup>1</sup> AAIJ 18P reports  $[\Gamma(B_c^+ \rightarrow D^*(2010)^+ D^*(2007)^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] / [B(B^+ \rightarrow \bar{D}^0 D^+)] < 1.3 \times 10^{-1}$  which we multiply by our best values  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ ,  $B(B^+ \rightarrow \bar{D}^0 D^+) = 3.8 \times 10^{-4}$ .

$$\Gamma(D^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{49}/\Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

$< 2.0 \times 10^{-7}$  90 <sup>1</sup> AAIJ 13R LHCB  $pp$  at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.5 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(D^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{50}/\Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

$< 1.6 \times 10^{-7}$  90 <sup>1</sup> AAIJ 13R LHCB  $pp$  at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.4 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(D_s^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{51}/\Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

$< 2.9 \times 10^{-7}$  90 <sup>1</sup> AAIJ 13R LHCB  $pp$  at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D_s^+ K^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.7 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(D_s^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{52}/\Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

$< 4 \times 10^{-7}$  90 <sup>1</sup> AAIJ 13R LHCB  $pp$  at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D_s^+ \bar{K}^{*0})/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 1.1 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(D_s^+ \phi)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{53}/\Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

$< 3.3 \times 10^{-7}$  90 <sup>1</sup> AAIJ 13R LHCB  $pp$  at 7 TeV

<sup>1</sup> AAIJ 13R reports  $[\Gamma(B_c^+ \rightarrow D_s^+ \phi)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 0.8 \times 10^{-6}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

$$\Gamma(K^+ K^0)/\Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c) \quad \Gamma_{54}/\Gamma \times B$$

| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
|-------|-----|-------------|------|---------|
|-------|-----|-------------|------|---------|

$< 4.6 \times 10^{-7}$  90 <sup>1</sup> AAIJ 13BS LHCB  $pp$  at 7 TeV

<sup>1</sup> Derived from  $\Gamma(K^+ K^0)/\Gamma \times B(\bar{b} \rightarrow B_c) / (B(B^+ \rightarrow K^0 \pi^+) B(\bar{b} \rightarrow B^+)) < 5.8\%$  at 90% CL using normalization mode  $B(B^+ \rightarrow K^0 \pi^+) = (23.97 \pm 0.53 \pm 0.71) \times 10^{-6}$  and assuming a  $B$  production ratio  $f(\bar{b} \rightarrow B_u^+) = 0.33$ .

| $\Gamma(B_s^0 \pi^+ / B(\bar{b} \rightarrow B_s)) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$ | $\Gamma_{55} / \Gamma \times B$              |
|--------------------------------------------------------------------------------------------------------------|----------------------------------------------|
| VALUE (units $10^{-3}$ )                                                                                     | DOCUMENT ID TECN COMMENT                     |
| $2.37 \pm 0.31 \pm 0.11^{+0.17}_{-0.13}$                                                                     | <sup>1</sup> AAIJ 13BU LHCB $pp$ at 7, 8 TeV |

<sup>1</sup> The last uncertainty is due to the uncertainty of the  $B_c^+$  lifetime measurement.

| $\Gamma(D_s^+ \bar{D}^0) / \Gamma_{\text{total}}$ | $\Gamma_{31} / \Gamma$                     |
|---------------------------------------------------|--------------------------------------------|
| VALUE CL%                                         | DOCUMENT ID TECN COMMENT                   |
| $< 7.2 \times 10^{-4}$ 90                         | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

| $\Gamma(D_s^+ D^0) / \Gamma_{\text{total}}$ | $\Gamma_{32} / \Gamma$                     |
|---------------------------------------------|--------------------------------------------|
| VALUE CL%                                   | DOCUMENT ID TECN COMMENT                   |
| $< 3.0 \times 10^{-4}$ 90                   | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

| $\Gamma(D^+ \bar{D}^0) / \Gamma_{\text{total}}$ | $\Gamma_{33} / \Gamma$                     |
|-------------------------------------------------|--------------------------------------------|
| VALUE CL%                                       | DOCUMENT ID TECN COMMENT                   |
| $< 1.9 \times 10^{-4}$ 90                       | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

| $\Gamma(D^+ D^0) / \Gamma_{\text{total}}$ | $\Gamma_{34} / \Gamma$                     |
|-------------------------------------------|--------------------------------------------|
| VALUE CL%                                 | DOCUMENT ID TECN COMMENT                   |
| $< 1.4 \times 10^{-4}$ 90                 | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

| $\Gamma(D_s^{*+} \bar{D}^0) / \Gamma_{\text{total}}$ | $\Gamma_{35} / \Gamma$                     |
|------------------------------------------------------|--------------------------------------------|
| VALUE CL%                                            | DOCUMENT ID TECN COMMENT                   |
| $< 5.3 \times 10^{-4}$ 90                            | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

| $\Gamma(D_s^+ \bar{D}^{*(2007)0}) / \Gamma_{\text{total}}$ | $\Gamma_{36} / \Gamma$                     |
|------------------------------------------------------------|--------------------------------------------|
| VALUE CL%                                                  | DOCUMENT ID TECN COMMENT                   |
| $< 4.6 \times 10^{-4}$ 90                                  | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

| $\Gamma(D_s^{*+} D^0) / \Gamma_{\text{total}}$ | $\Gamma_{37} / \Gamma$                     |
|------------------------------------------------|--------------------------------------------|
| VALUE CL%                                      | DOCUMENT ID TECN COMMENT                   |
| $< 0.9 \times 10^{-3}$ 90                      | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

| $\Gamma(D_s^+ D^{*(2007)0}) / \Gamma_{\text{total}}$ | $\Gamma_{38} / \Gamma$                     |
|------------------------------------------------------|--------------------------------------------|
| VALUE CL%                                            | DOCUMENT ID TECN COMMENT                   |
| $< 6.6 \times 10^{-4}$ 90                            | <sup>1</sup> AAIJ 21AF LHCB $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c) / B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19AI.

$\Gamma(D^*(2010)^+\bar{D}^0)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<3.8 \times 10^{-4}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(D^*(2007)^+D^0)/\Gamma_{\text{total}}$   $\Gamma_{42}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<2.0 \times 10^{-4}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(D^+\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<6.5 \times 10^{-4}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(D^+D^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<3.7 \times 10^{-4}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(D_s^{*+}\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_{45}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<1.3 \times 10^{-3}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(D_s^{*+}D^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_{46}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<1.3 \times 10^{-3}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(D^*(2010)^+\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<1.0 \times 10^{-3}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(D^*(2010)^+D^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$ 

| VALUE                 | CL% | DOCUMENT ID       | TECN      | COMMENT        |
|-----------------------|-----|-------------------|-----------|----------------|
| $<7.7 \times 10^{-4}$ | 90  | <sup>1</sup> AAIJ | 21AF LHCB | $pp$ at 13 TeV |

<sup>1</sup> Uses  $B(\bar{b} \rightarrow B_c)/B(\bar{b} \rightarrow B^+) = 0.76\%$  determined by AAIJ 19Al.

 $\Gamma(B_s^0\pi^+)/\Gamma(J/\psi(1S)\pi^+)$   $\Gamma_{56}/\Gamma_4$ 

| VALUE               | DOCUMENT ID       | TECN     | COMMENT        |
|---------------------|-------------------|----------|----------------|
| $91 \pm 10 \pm 8.5$ | <sup>1</sup> AAIJ | 23M LHCB | $pp$ at 13 TeV |

<sup>1</sup> The  $B_s^0$  mesons are reconstructed via the decays  $B_s^0 \rightarrow J/\psi\phi$  and  $B_s^0 \rightarrow D_s^- \pi^+$ . The third uncertainty includes systematic ( $\pm 8$ ) and imprecise knowledge of the branching fractions ( $\pm 3$ ).

$$\Gamma(\pi^+ \mu^+ \mu^-) / \Gamma(J/\psi(1S) \pi^+)$$

$$\Gamma_{57}/\Gamma_4$$

| VALUE                 | CL% | DOCUMENT ID | TECN     | COMMENT                 |
|-----------------------|-----|-------------|----------|-------------------------|
| $<2.1 \times 10^{-4}$ | 90  | AAIJ        | 24W LHCB | $pp$ at 7, 8 and 13 TeV |

$$\Gamma(D_s^+ \mu^+ \mu^-) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)$$

$$\Gamma_{58}/\Gamma \times B$$

| VALUE                 | CL% | DOCUMENT ID       | TECN     | COMMENT              |
|-----------------------|-----|-------------------|----------|----------------------|
| $<3.1 \times 10^{-8}$ | 90  | <sup>1</sup> AAIJ | 24F LHCB | $pp$ at 7, 8, 13 TeV |

<sup>1</sup> AAIJ 24F reports  $[\Gamma(B_c^+ \rightarrow D_s^+ \mu^+ \mu^-) / \Gamma_{\text{total}} \times B(\bar{b} \rightarrow B_c)] / [B(\bar{b} \rightarrow B^+)] < 7.5 \times 10^{-8}$  which we multiply by our best value  $B(\bar{b} \rightarrow B^+) = 40.8 \times 10^{-2}$ .

## POLARIZATION IN $B_c^+$ DECAY

In decays involving two vector mesons, one can distinguish among the states in which meson polarizations are both longitudinal ( $L$ ) or both are transverse and parallel ( $\parallel$ ) or perpendicular ( $\perp$ ) to each other with the parameters  $\Gamma_L/\Gamma$ ,  $\Gamma_\perp/\Gamma$ , and the relative phases  $\phi_\parallel$  and  $\phi_\perp$ . See the definitions in the note on “Polarization in  $B$  Decays” review in the  $B^0$  Particle Listings.

$$\Gamma_L/\Gamma \text{ in } B_c^+ \rightarrow J/\psi D_s^{*+}$$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------|-------------|------|---------|
|-------|-------------|------|---------|

**0.34 $\pm$ 0.09 OUR AVERAGE**

|                            |                    |           |                  |
|----------------------------|--------------------|-----------|------------------|
| 0.30 $\pm$ 0.10 $\pm$ 0.04 | <sup>1,2</sup> AAD | 22O ATLS  | $pp$ at 13 TeV   |
| 0.48 $\pm$ 0.20            | <sup>3</sup> AAIJ  | 13AS LHCB | $pp$ at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

|                 |                  |          |                  |
|-----------------|------------------|----------|------------------|
| 0.62 $\pm$ 0.24 | <sup>4</sup> AAD | 16H ATLS | $pp$ at 7, 8 TeV |
|-----------------|------------------|----------|------------------|

<sup>1</sup> Supersedes the measurement of AAD 16H.

<sup>2</sup> AAD 22O measures  $1 - \Gamma_L/\Gamma = 0.70 \pm 0.10 \pm 0.04$ .

<sup>3</sup> AAIJ 13AS measures  $1 - \Gamma_L/\Gamma = 0.52 \pm 0.20$ .

<sup>4</sup> AAD 16H measures  $1 - \Gamma_L/\Gamma = 0.38 \pm 0.24$ .

$$A_P(B_c^+)$$

$$A_P(B_c^+) = [\sigma(B_c^-) - \sigma(B_c^+)] / [\sigma(B_c^-) + \sigma(B_c^+)]$$

| VALUE (units $10^{-2}$ ) | DOCUMENT ID | TECN | COMMENT |
|--------------------------|-------------|------|---------|
|--------------------------|-------------|------|---------|

**−1.0 $\pm$ 1.0 OUR AVERAGE**

|                          |                   |           |                |
|--------------------------|-------------------|-----------|----------------|
| −2.5 $\pm$ 2.1 $\pm$ 0.5 | <sup>1</sup> AAIJ | 19AI LHCB | $pp$ at 7 TeV  |
| −0.5 $\pm$ 1.1 $\pm$ 0.4 | <sup>1</sup> AAIJ | 19AI LHCB | $pp$ at 13 TeV |

<sup>1</sup> Measured using  $B_c^+$  semileptonic decays.

**$B_c^+$  REFERENCES**

|              |      |                        |                              |                  |
|--------------|------|------------------------|------------------------------|------------------|
| AAIJ         | 24F  | JHEP 2402 032          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 24I  | JHEP 2402 173          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 24L  | JHEP 2404 151          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 24W  | EPJ C84 468            | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 23M  | JHEP 2307 066          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAD          | 22O  | JHEP 2208 087          | G. Aad <i>et al.</i>         | (ATLAS Collab.)  |
| AAIJ         | 22P  | JHEP 2201 065          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AABOUD       | 21   | PR D104 012010         | M. Aaboud <i>et al.</i>      | (ATLAS Collab.)  |
| AAIJ         | 21AF | JHEP 2112 117          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 20R  | JHEP 2007 123          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 19AI | PR D100 112006         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 18C  | PRL 120 121801         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 18P  | NP B930 563            | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| SIRUNYAN     | 18BY | EPJ C78 457            | A.M. Sirunyan <i>et al.</i>  | (CMS Collab.)    |
| AAIJ         | 17AG | PRL 118 111803         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 17L  | PR D95 032005          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAD          | 16H  | EPJ C76 4              | G. Aad <i>et al.</i>         | (ATLAS Collab.)  |
| AAIJ         | 16AF | JHEP 1609 153          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 16AT | PR D94 091102          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 16K  | PL B759 313            | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AALTONEN     | 16A  | PR D93 052001          | T. Aaltonen <i>et al.</i>    | (CDF Collab.)    |
| AAIJ         | 15AY | PR D92 072007          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 15G  | PL B742 29             | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 15M  | PRL 114 132001         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| KHACHATRY... | 15AA | JHEP 1501 063          | V. Khachatryan <i>et al.</i> | (CMS Collab.)    |
| AAIJ         | 14AQ | PRL 113 152003         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 14G  | EPJ C74 2839           | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 14P  | JHEP 1405 148          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 14W  | PR D90 032009          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 13AM | PR D87 071103          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 13AS | PR D87 112012          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| Also         |      | PR D89 019901 (errat.) | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 13BS | PL B726 646            | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 13BU | PRL 111 181801         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 13BY | JHEP 1309 075          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 13CA | JHEP 1311 094          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 13R  | JHEP 1302 043          | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AALTONEN     | 13   | PR D87 011101          | T. Aaltonen <i>et al.</i>    | (CDF Collab.)    |
| AAIJ         | 12AV | PRL 109 232001         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| AAIJ         | 12Y  | PRL 108 251802         | R. Aaij <i>et al.</i>        | (LHCb Collab.)   |
| ABAZOV       | 09H  | PRL 102 092001         | V.M. Abazov <i>et al.</i>    | (D0 Collab.)     |
| AALTONEN     | 08M  | PRL 100 182002         | T. Aaltonen <i>et al.</i>    | (CDF Collab.)    |
| ABAZOV       | 08T  | PRL 101 012001         | V.M. Abazov <i>et al.</i>    | (D0 Collab.)     |
| ABULENCIA    | 06C  | PRL 96 082002          | A. Abulencia <i>et al.</i>   | (CDF Collab.)    |
| ABULENCIA    | 06O  | PRL 97 012002          | A. Abulencia <i>et al.</i>   | (CDF Collab.)    |
| ABE          | 98M  | PRL 81 2432            | F. Abe <i>et al.</i>         | (CDF Collab.)    |
| Also         |      | PR D58 112004          | F. Abe <i>et al.</i>         | (CDF Collab.)    |
| ACKERSTAFF   | 98O  | PL B420 157            | K. Akerstaff <i>et al.</i>   | (OPAL Collab.)   |
| BARATE       | 98Q  | EPJ C4 387             | R. Barate <i>et al.</i>      | (ALEPH Collab.)  |
| PDG          | 98   | EPJ C3 1               | C. Caso <i>et al.</i>        | (PDG Collab.)    |
| ABREU        | 97E  | PL B398 207            | P. Abreu <i>et al.</i>       | (DELPHI Collab.) |
| BARATE       | 97H  | PL B402 213            | R. Barate <i>et al.</i>      | (ALEPH Collab.)  |
| ABE          | 96R  | PRL 77 5176            | F. Abe <i>et al.</i>         | (CDF Collab.)    |
| PDG          | 96   | PR D54 1               | R. M. Barnett <i>et al.</i>  | (PDG Collab.)    |