

$K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

 $K_4^*(2045)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2048^{+8}_{-9} OUR AVERAGE		Error includes scale factor of 1.1.			
$2090 \pm 9^{+11}_{-29}$	183k	ABLIKIM	19AQ BES	$\pm$	$J/\psi \rightarrow K^+ K^- \pi^0$
$2062 \pm 14 \pm 13$		¹ ASTON	86 LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
2039 ± 10	400	^{2,3} CLELAND	82 SPEC	$\pm$	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$
2070^{+100}_{-40}		⁴ ASTON	81C LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2079 ± 7	431	TORRES	86 MPSF		$400 pA \rightarrow 4KX$
2088 ± 20	650	BAUBILLIER	82 HBC	—	$8.25 K^- p \rightarrow K_S^0 \pi^- p$
2115 ± 46	488	CARMONY	77 HBC	0	$9 K^+ d \rightarrow K^+ \pi's X$
¹ From a fit to all moments.					
² From a fit to 8 moments.					
³ Number of events evaluated by us.					
⁴ From energy-independent partial-wave analysis.					

 $K_4^*(2045)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
199^{+27}_{-19} OUR AVERAGE					
$201 \pm 19^{+57}_{-17}$	183k	ABLIKIM	19AQ BES	$\pm$	$J/\psi \rightarrow K^+ K^- \pi^0$
$221 \pm 48 \pm 27$		⁵ ASTON	86 LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
189 ± 35	400	^{6,7} CLELAND	82 SPEC	$\pm$	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
61 ± 58	431	TORRES	86 MPSF		$400 pA \rightarrow 4KX$
170^{+100}_{-50}	650	BAUBILLIER	82 HBC	—	$8.25 K^- p \rightarrow K_S^0 \pi^- p$
240^{+500}_{-100}		⁸ ASTON	81C LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
300 ± 200		CARMONY	77 HBC	0	$9 K^+ d \rightarrow K^+ \pi's X$
⁵ From a fit to all moments.					
⁶ From a fit to 8 moments.					
⁷ Number of events evaluated by us.					
⁸ From energy-independent partial-wave analysis.					

$K_4^*(2045)$ DECAY MODES

Mode		Fraction (Γ_i/Γ)
Γ_1	$K\pi$	$(9.9\pm 1.2)\%$
Γ_2	$K^*(892)\pi\pi$	$(9\pm 5)\%$
Γ_3	$K^*(892)\pi\pi\pi$	$(7\pm 5)\%$
Γ_4	$\rho K\pi$	$(5.7\pm 3.2)\%$
Γ_5	$\omega K\pi$	$(5.0\pm 3.0)\%$
Γ_6	$\phi K\pi$	$(2.8\pm 1.4)\%$
Γ_7	$\phi K^*(892)$	$(1.4\pm 0.7)\%$

$K_4^*(2045)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.099 ± 0.012	ASTON	88	LASS	0	$11\ K^-p\rightarrow K^-\pi^+n$
$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$					Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.89 ± 0.53	BAUBILLIER	82	HBC	—	$8.25\ K^-p\rightarrow pK_S^03\pi$
$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$					Γ_3/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.75 ± 0.49	BAUBILLIER	82	HBC	—	$8.25\ K^-p\rightarrow pK_S^03\pi$
$\Gamma(\rho K\pi)/\Gamma(K\pi)$					Γ_4/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.58 ± 0.32	BAUBILLIER	82	HBC	—	$8.25\ K^-p\rightarrow pK_S^03\pi$
$\Gamma(\omega K\pi)/\Gamma(K\pi)$					Γ_5/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.50 ± 0.30	BAUBILLIER	82	HBC	—	$8.25\ K^-p\rightarrow pK_S^03\pi$
$\Gamma(\phi K\pi)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.028 ± 0.014	⁹ TORRES	86	MPSF	400 pA $\rightarrow$ 4K X	
$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.014 ± 0.007	⁹ TORRES	86	MPSF	400 pA $\rightarrow$ 4K X	
⁹ Error determination is model dependent.					

$K_4^*(2045)$ REFERENCES

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
TORRES	86	PR D34 707	S. Torres <i>et al.</i>	(VPI, ARIZ, FNAL, FSU+)
BAUBILLIER	82	PL 118B 447	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
CARMONY	77	PR D16 1251	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)
