$$T_{b\overline{s}}(5568)^+$$

$$I(J^{P}) = 1(?^{?})$$

OMITTED FROM SUMMARY TABLE was  $X(5568)^{\pm}$ 

Seen as a peak in the  $B_s \pi^{\pm}$  mass spectrum with a significance of more than  $3\sigma$  by ABAZOV 16E and ABAZOV 18A in inclusive  $p\overline{p}$  collisions at 1.96 TeV. Not seen by AAIJ 16AI, AABOUD 18L, AALTONEN 18A, and SIRUNYAN 18J. Needs confirmation.

## *Т<sub>b</sub>*(5568)<sup>+</sup> MASS

VALUE (MeV)	EVTS	DOCUMENT I	0	TECN	COMMENT	
$5566.9^{+3.2}_{-3.1}^{+0.6}_{-1.2}$	278	<sup>1</sup> ABAZOV	18A	D0	$p \overline{p} \rightarrow B^0_s \pi^{\pm} X$	
ullet $ullet$ $ullet$ We do not use the following data for averages, fits, limits, etc. $ullet$ $ullet$						
$5567.8 {\pm} 2.9 {+} 0.9 {-} {1.9}$	133	<sup>2</sup> ABAZOV	16E	D0	$p \overline{p} \rightarrow B^0_s \pi^{\pm} X$	
<sup>1</sup> From the combined analysis of $B_s^0 \to J/\psi \phi$ and $B_s^0 \to D_s^{\pm} \mu^{\mp} X$ decays. <sup>2</sup> Assumes $T_{b\overline{s}}(5568)^{\pm} \to B_s \pi^{\pm}$ decay. If $T_{b\overline{s}}(5568)^{\pm} \to B_s^* \pi^{\pm}$ decay is assumed, the mass shifts upward by 49 MeV.						

## *Т<sub>b</sub>*(5568)<sup>+</sup> WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	COMMENT
$18.6^{+7.9+3.5}_{-6.1-3.8}$	278	<sup>1</sup> ABAZOV	18A	D0	$p\overline{p} \rightarrow B_{s}\pi^{\pm}X$
ullet $ullet$ $ullet$ We do not use the following data for averages, fits, limits, etc. $ullet$ $ullet$					
$21.9\!\pm\!6.4^{+5.0}_{-2.5}$	133	ABAZOV	16E	D0	$p\overline{p} \rightarrow B_{s}\pi^{\pm}X$
$^1$ From the combined analysis of $B^0_{m{s}} o \ J/\psi\phi$ and $B^0_{m{s}} o \ D^\pm_{m{s}}\mu^\mp X$ decays.					

# T<sub>bs</sub>(5568)<sup>+</sup> DECAY MODES

	Mode	Fraction $(\Gamma_i/\Gamma)$
$\Gamma_1$	$B_s \pi^+$	seen

#### Tbs(5568)<sup>+</sup> BRANCHING RATIOS

$\Gamma(B_s \pi^+) / \Gamma_{total}$						$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID		TECN	COMMENT	
seen	145	<sup>1</sup> ABAZOV	18A	D0	$p\overline{p} \rightarrow B_s^0 \pi^{\pm} X$	
seen	133	<sup>2</sup> ABAZOV	16E	D0	$p \overline{p} \rightarrow B_s^{0} \pi^{\pm} X$	

https://pdg.lbl.gov

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen			$pp \rightarrow B_{s}^{0}\pi^{\pm}X$
not seen			$p\overline{p} \rightarrow B_{s}^{0}\pi^{\pm}X$
not seen	<sup>5</sup> SIRUNYAN	18J CMS	$pp \rightarrow B_{s}^{0}\pi^{\pm}X$
not seen	<sup>6</sup> AAIJ	16AI LHCB	$pp \rightarrow B_s^0 \pi^{\pm} X$

<sup>1</sup>With  $B_s$  mesons reconstructed in decays to  $D_s^{\pm} \mu^{\mp} X$ .

- <sup>2</sup> Seen in  $p\overline{p}$  collisions at 1.96 TeV at a rate of  $(8.6 \pm 1.9 \pm 1.4)\%$  relative to inclusive  $B_s$  production in the kinematic region  $10 < p_T(B_s) < 30$  GeV/c, with  $B_s$  mesons reconstructed in decays to  $J/\psi\phi$ . An alternative possibility,  $T_{b\overline{s}}(5568)^{\pm} \rightarrow B_s^* \pi^{\pm}$  with a missing  $\gamma$ , could not be ruled out.
- <sup>3</sup>Not seen in 24.4 fb<sup>-1</sup> of pp collision data at  $\sqrt{s} = 7$  and 8 TeV with  $B_s$  mesons reconstructed in decays to  $J/\psi\phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^{\pm} \rightarrow B_s \pi^{\pm}$  relative to inclusive  $B_s$  production is less than 1.5% at  $p_T(B_s) > 10$  GeV/c and less than 1.6% at  $p_T(B_s) > 15$  GeV/c at 95% CL.
- <sup>4</sup> Not seen in 9.6 fb<sup>-1</sup> of  $p\overline{p}$  collision data at  $\sqrt{s} = 1.96$  TeV with  $B_s$  mesons reconstructed in decays to  $J/\psi \phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^{\pm} \rightarrow B_s \pi^{\pm}$  relative to inclusive  $B_s$  production is less than 6.7% at 95% CL.
- <sup>5</sup> Not seen in 19.7 fb<sup>-1</sup> of pp collisions data at  $\sqrt{s} = 8$  TeV with  $B_s$  mesons reconstructed in decays to  $J/\psi\phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^{\pm} \rightarrow B_s \pi^{\pm}$  relative to inclusive  $B_s$  production is less than 1.1% at  $p_T(B_s)$ > 10 GeV/c and less than 1.0% at  $p_T(B_s) > 15$  GeV/c at 95%CL.

<sup>6</sup> Not seen in 3 fb<sup>-1</sup> of pp collision data at  $\sqrt{s} = 7$  and 8 TeV in a scan over the  $T_{b\overline{s}}(5568)$  mass and width, with  $B_s$  mesons reconstructed in decays to  $D_s^- \pi^+$  or  $J/\psi\phi$ . An upper limit on the production rate times branching fraction for  $T_{b\overline{s}}(5568)^{\pm} \rightarrow B_s \pi^{\pm}$  relative to inclusive  $B_s$  production is less than 2.1% at  $p_T(B_s) > 10$  GeV/c at 90% CL.

### T<sub>bs</sub>(5568)<sup>+</sup> REFERENCES

AABOUD AALTONEN ABAZOV SIRUNYAN AAIJ ABAZOV	18A 18A 18J 16AI	PRL 120 202007 PRL 120 202006 PR D97 092004 PRL 120 202005 PRL 117 152003 PRL 117 022003	M. Aaboud <i>et al.</i> T. Aaltonen <i>et al.</i> V.M. Abazov <i>et al.</i> A.M. Sirunyan <i>et al.</i> R. Aaij <i>et al.</i> V.M. Abazov <i>et al.</i>	(ATLAS Collab.) (CDF Collab.) (D0 Collab.) (CMS Collab.) (LHCb Collab.) (D0 Collab.)
ABAZOV	16E	PRL 117 022003	V.M. Abazov et al.	(D0 Collab.)