

X(1835)

$$I^G(J^{PC}) = 0^+(0^-\,+)$$

OMITTED FROM SUMMARY TABLE

Could be a superposition of two states, one appearing as threshold enhancement in  $p\bar{p}$  the other one with a lower mass at 1835 MeV. Coupled-channel analyses with more sophisticated model are needed.

X(1835) MASS									
VALUE (MeV)			EVTS	DOCUMENT ID		TECN	COMMENT		
1831.8 <sup>+</sup> <sub>-</sub> 4.0 <sub>2.6</sub> OUR AVERAGE									
1832.5 ± 3.1 ± 2.5			21k	<sup>1</sup> ABLIKIM	24B	BES3	J/ψ → γ3(π <sup>+</sup> π <sup>-</sup> )		
1825.3 ± 2.4 <sup>+</sup> <sub>-</sub> 17.3 <sub>2.4</sub>				<sup>2</sup> ABLIKIM	16J	BES3	J/ψ → γπ <sup>+</sup> π <sup>-</sup> η'		
1844 ± 9 <sup>+</sup> <sub>-</sub> 16 <sub>25</sub>				<sup>3</sup> ABLIKIM	15T	BES3	J/ψ → γK <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> η		
• • • We do not use the following data for averages, fits, limits, etc. • • •									
1818 ± 9 ± 2.5			37k	<sup>4</sup> ABLIKIM	24B	BES3	J/ψ → γ3(π <sup>+</sup> π <sup>-</sup> )		
1839 ± 26 ± 26				<sup>5</sup> ABLIKIM	18I	BES3	J/ψ → γγϕ(1020)		
1909.5 ± 15.9 <sup>+</sup> <sub>-</sub> 9.4 <sub>27.5</sub>				<sup>6</sup> ABLIKIM	16J	BES3	J/ψ → γπ <sup>+</sup> π <sup>-</sup> η'		
1842.2 ± 4.2 <sup>+</sup> <sub>-</sub> 7.1 <sub>2.6</sub>			0.6k	<sup>7</sup> ABLIKIM	13U	BES3	J/ψ → γ3(π <sup>+</sup> π <sup>-</sup> )		
1832 <sup>+</sup> <sub>-</sub> 19 <sub>5</sub> ± 26				<sup>8</sup> ABLIKIM	12D	BES3	J/ψ → γp $\bar{p}$		
1836.5 ± 3.0 <sup>+</sup> <sub>-</sub> 5.6 <sub>2.1</sub>			4265	<sup>9</sup> ABLIKIM	11C	BES3	J/ψ → γπ <sup>+</sup> π <sup>-</sup> η'		
1877.3 ± 6.3 <sup>+</sup> <sub>-</sub> 3.4 <sub>7.4</sub>				<sup>10</sup> ABLIKIM	11J	BES3	J/ψ → ω(ηπ <sup>+</sup> π <sup>-</sup> )		
1837 <sup>+</sup> <sub>-</sub> 10 <sub>12</sub> <sup>+</sup> <sub>-</sub> 9 <sub>7</sub>			231	<sup>11,12</sup> ALEXANDER	10	CLEO	J/ψ → γp $\bar{p}$		
1833.7 ± 6.1 ± 2.7			264	ABLIKIM	05R	BES2	J/ψ → γπ <sup>+</sup> π <sup>-</sup> η'		
1831 ± 7				<sup>12,13</sup> ABLIKIM	05R	BES2	J/ψ → γp $\bar{p}$		
1859 <sup>+</sup> <sub>-</sub> 3 <sub>10</sub> <sup>+</sup> <sub>-</sub> 5 <sub>25</sub>				<sup>12</sup> BAI	03F	BES2	J/ψ → γp $\bar{p}$		

<sup>1</sup> From a fit of the measured  $3(\pi^+\pi^-)$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold through interference with a second previously unseen narrow resonance near 1880 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known background contributions. A second solution of the fit gives 37k events.

<sup>2</sup> From a fit of the measured  $\pi^+\pi^-\eta'$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold through interference with a second previously unseen narrow resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors.

<sup>3</sup> Decay dominated by  $f_0(980)\eta$  hence  $I^G(J^{PC}) = 0^+(0^-\,+)$ .

<sup>4</sup> From a fit of the measured  $3(\pi^+\pi^-)$  lineshape to a Flatte formula that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold. The fit also includes known background contributions.

<sup>5</sup> From a fit to  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^-\,+$ . Other  $J^{PC}$  not excluded.

<sup>6</sup> Pole mass from a fit of the measured  $\pi^+\pi^-\eta'$  lineshape to a Flatte formula that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold; the fit also includes

known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner function ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit.

<sup>7</sup> Superseded by ABLIKIM 24B.

<sup>8</sup> From the fit including final state interaction effects in isospin 0  $S$ -wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.

<sup>9</sup> From a fit of the  $\pi^+\pi^-\eta'$  mass distribution to a combination of  $\gamma f_1(1510)$ ,  $\gamma X(1835)$ , and two states  $\gamma X(2120)$  and  $\gamma \eta(2370)$ , for  $M(\pi^+\pi^-\eta') < 2.8$  GeV, and accounting for backgrounds from non- $\eta'$  events and  $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$ .

<sup>10</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$  with  $B(J/\psi \rightarrow \omega X \rightarrow \omega a_0(980)^\pm (\rightarrow \eta\pi^\pm)\pi^\mp) = (1.50 \pm 0.26^{+0.72}_{-0.36}) \times 10^{-4}$ . Not seen in  $J/\psi(1S) \rightarrow \omega K^+K^-\eta$  by ABLIKIM 24BQ with 90% CL upper limit of  $9.55 \times 10^{-7}$ . This state may be also due to  $\eta_2(1870)$  or to a combination of  $X(1835)$  and  $\eta_2(1870)$ .

<sup>11</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma X(1835)$ ,  $\gamma R$  with  $M(R) = 2100$  MeV and  $\Gamma(R) = 160$  MeV, and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV.

<sup>12</sup> Evidence for a threshold enhancement in the  $p\bar{p}$  mass spectrum was also reported by ABE 02K, AUBERT, B 05L, and WANG 05A in  $B^+ \rightarrow p\bar{p}K^+$ , WANG 05A in  $B^0 \rightarrow p\bar{p}K_S^0$ , ABE 02W in  $\bar{B}^0 \rightarrow p\bar{p}D^0$ , DEL-AMO-SANCHEZ 12 in  $B \rightarrow D(D^*)p\bar{p}(\pi)$ , and WEI 08 in  $B^+ \rightarrow p\bar{p}\pi^+$  decays. Not seen by ATHAR 06 in  $\Upsilon(1S) \rightarrow p\bar{p}\gamma$ .

<sup>13</sup> From the fit including final state interaction effects in isospin 0  $S$ -wave according to SIBIRTSEV 05A. Systematic errors not estimated.

## X(1835) WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>120 ± 70</b>	<b>OUR AVERAGE</b>		Error includes scale factor of 8.8.		
80.7 ± 5.2 ± 7.7		21k	<sup>1</sup> ABLIKIM	24B BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
245.2 ± 13.1 <sup>+</sup> <sub>-4.6</sub>			<sup>2</sup> ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma \pi^+\pi^-\eta'$
192 <sup>+</sup> <sub>-17</sub> <sup>+</sup> <sub>-43</sub>			<sup>3</sup> ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
175 ± 57 ± 25			<sup>4</sup> ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
273.5 ± 21.4 <sup>+</sup> <sub>-64.0</sub>			<sup>5</sup> ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma \pi^+\pi^-\eta'$
83 ± 14 ± 11		0.6k	<sup>6</sup> ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
< 76	90		<sup>7</sup> ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
190 ± 9 <sup>+</sup> <sub>-36</sub>		4265	<sup>8</sup> ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+\pi^-\eta'$
57 ± 12 <sup>+</sup> <sub>-4</sub>			<sup>9</sup> ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
0 <sup>+</sup> <sub>-0</sub>		231	<sup>10,11</sup> ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
67.7 ± 20.3 ± 7.7		264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma \pi^+\pi^-\eta'$
< 153	90		<sup>11,12</sup> ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p\bar{p}$
< 30			<sup>11</sup> BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

<sup>1</sup> From a fit of the measured  $3(\pi^+\pi^-)$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold through interference with a second previously unseen narrow resonance near 1880 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known background contributions. A second solution of the fit gives 37k events.

<sup>2</sup> From a fit of the measured  $\pi^+\pi^-\eta'$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold through interference with a second previously unseen narrow

resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors.

<sup>3</sup> Decay dominated by  $f_0(980)\eta$  hence  $I^G(J^{PC}) = 0^+(0^-+)$ .

<sup>4</sup> From a fit to  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^-+$ . Other  $J^{PC}$  not excluded.

<sup>5</sup> Pole width from a fit of the measured  $\pi^+\pi^-\eta'$  lineshape to a Flatte formula that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold; the fit also includes known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner function ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit.

<sup>6</sup> Superseded by ABLIKIM 24B.

<sup>7</sup> From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.

<sup>8</sup> From a fit of the  $\pi^+\pi^-\eta'$  mass distribution to a combination of  $\gamma f_1(1510)$ ,  $\gamma X(1835)$ , and two states  $\gamma X(2120)$  and  $\gamma\eta(2370)$ , for  $M(\pi^+\pi^-\eta') < 2.8$  GeV, and accounting for backgrounds from non- $\eta'$  events and  $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$ .

<sup>9</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$  with  $B(J/\psi \rightarrow \omega X \rightarrow \omega a_0(980)^\pm (\rightarrow \eta\pi^\pm)\pi^\mp) = (1.50 \pm 0.26^{+0.72}_{-0.36}) \times 10^{-4}$ . Not seen in  $J/\psi(1S) \rightarrow \omega K^+K^-\eta$  by ABLIKIM 24BQ with 90% CL upper limit of  $9.55 \times 10^{-7}$ . This state may be also due to  $\eta_2(1870)$  or to a combination of  $X(1835)$  and  $\eta_2(1870)$ .

<sup>10</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma X(1835)$ ,  $\gamma R$  with  $M(R) = 2100$  MeV and  $\Gamma(R) = 160$  MeV, and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV.

<sup>11</sup> Evidence for a threshold enhancement in the  $p\bar{p}$  mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in  $B^+ \rightarrow p\bar{p}K^+$ , WANG 05A in  $B^0 \rightarrow p\bar{p}K_S^0$ , ABE 02W in  $\bar{B}^0 \rightarrow p\bar{p}D^0$ , DEL-AMO-SANCHEZ 12 in  $B \rightarrow D(D^*)p\bar{p}(\pi)$ , and WEI 08 in  $B^+ \rightarrow p\bar{p}\pi^+$  decays. Not seen by ATHAR 06 in  $\Upsilon(1S) \rightarrow p\bar{p}\gamma$ .

<sup>12</sup> From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Systematic errors not estimated.

### X(1835) DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $p\bar{p}$	seen
$\Gamma_2$ $\eta'\pi^+\pi^-$	seen
$\Gamma_3$ $\gamma\gamma$	not seen
$\Gamma_4$ $K_S^0 K_S^0 \eta$	seen
$\Gamma_5$ $\gamma\phi(1020)$	possibly seen
$\Gamma_6$ $3(\pi^+\pi^-)$	seen

### X(1835) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta'\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_3/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<35.6	90	<sup>1</sup> ZHANG	12A	BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
<83	90	<sup>2</sup> ZHANG	12A	BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

<sup>1</sup> From a two-resonance fit and constructive interference of the  $\eta(1760)$  and  $X(1835)$ , a significance of  $2.8\sigma$ .

<sup>2</sup> From a two-resonance fit and destructive interference of the  $\eta(1760)$  and  $X(1835)$ , a significance of  $2.8\sigma$ .

**X(1835) BRANCHING RATIOS**

$\Gamma(p\bar{p})/\Gamma(\eta'\pi^+\pi^-)$

$\Gamma_1/\Gamma_2$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.333	ABLIKIM	05R	BES2 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

$\Gamma(\eta'\pi^+\pi^-)/\Gamma(K_S^0 K_S^0 \eta)$

$\Gamma_2/\Gamma_4$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$6.7 \pm 1.8$	<sup>1</sup> ABLIKIM	15T	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

<sup>1</sup>Using results from ABLIKIM 05R.

$\Gamma(\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> ABLIKIM	16J	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

<sup>1</sup> ABLIKIM 16J quotes  $B(J/\psi \rightarrow \gamma X(1835)) \times B(X(1835) \rightarrow \pi^+\pi^-\eta') = (3.93 \pm 0.38^{+0.31}_{-0.84}) \times 10^{-4}$  from a fit of the measured  $\pi^+\pi^-\eta'$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the X(1835); fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

$\Gamma(\gamma\phi(1020))/\Gamma_{\text{total}}$

$\Gamma_5/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	<sup>1</sup> ABLIKIM	18i	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$

<sup>1</sup> Seen as a peak in  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^{-+}$ . Other  $J^{PC}$  not excluded.

$\Gamma(\gamma\gamma)/\Gamma(\eta'\pi^+\pi^-)$

$\Gamma_3/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.80 \times 10^{-3}$	90	<sup>1</sup> ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma$

<sup>1</sup>Using results from ABLIKIM 16J.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$

$\Gamma_6/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	21k	<sup>1</sup> ABLIKIM	24B	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<b>seen</b>	0.6k	<sup>2</sup> ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
-------------	------	----------------------	-----	--

<sup>1</sup> ABLIKIM 24B quotes  $B(J/\psi \rightarrow \gamma X(1835)) \times B(X(1835) \rightarrow 3(\pi^+\pi^-)) = (1.19 \pm 0.30 \pm 0.15) \times 10^{-5}$  for constructive interference and  $(2.07 \pm 0.50 \pm 0.36) \times 10^{-5}$  for destructive interference from a fit of the measured  $3(\pi^+\pi^-)$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold through interference with a second narrow resonance near 1880 MeV. The solution for destructive interference gives 37k events.

<sup>2</sup>Superseded by ABLIKIM 24B.

**X(1835) REFERENCES**

ABLIKIM	24B	PRL 132 151901	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BQ	PR D110 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.) JPC
DEL-AMO-SA...	12	PR D85 092017	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10G	CP C34 421	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
WEI	08	PL B659 80	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT,B	05L	PR D72 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)

---