

$\chi_{b2}(3P)$

$I^G(J^{PC}) = 0^+(2^{++})$   
 $J$  needs confirmation.

Observed in the radiative decay to  $\Upsilon(3S)$ , therefore  $C = +$ .

$\chi_{b2}(3P)$  MASS

| VALUE (MeV)                                                                                                                                                                                                                                                               | DOCUMENT ID           | TECN     | COMMENT                               |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------|----------|---------------------------------------|
| <b>10524.02±0.57±0.53</b>                                                                                                                                                                                                                                                 | <sup>1</sup> SIRUNYAN | 18N CMS  | $pp \rightarrow \gamma \mu^+ \mu^- X$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • •                                                                                                                                                                                             |                       |          |                                       |
| 10530 ±5 ±9                                                                                                                                                                                                                                                               | <sup>2</sup> AAD      | 12A ATLS | $pp \rightarrow \gamma \mu^+ \mu^- X$ |
| <sup>1</sup> Systematic error includes an additional 0.5 MeV for the uncertainty on the $\Upsilon(3S)$ mass. Also measures $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$ MeV. A total of 372 $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ events was observed. |                       |          |                                       |
| <sup>2</sup> The mass barycenter of the merged lineshapes from the $J = 1$ and 2 states.                                                                                                                                                                                  |                       |          |                                       |

$\chi_{b2}(3P)$  DECAY MODES

| Mode                                | Fraction ( $\Gamma_i/\Gamma$ ) |
|-------------------------------------|--------------------------------|
| $\Gamma_1 \quad \Upsilon(3S)\gamma$ | seen                           |

$\chi_{b2}(3P)$  BRANCHING RATIOS

| $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$ |             |         |                                       | $\Gamma_1/\Gamma$ |
|----------------------------------------------------|-------------|---------|---------------------------------------|-------------------|
| VALUE                                              | DOCUMENT ID | TECN    | COMMENT                               |                   |
| <b>seen</b>                                        | SIRUNYAN    | 18N CMS | $pp \rightarrow \gamma \mu^+ \mu^- X$ |                   |

$\chi_{b2}(3P)$  REFERENCES

|          |     |                |                             |                 |
|----------|-----|----------------|-----------------------------|-----------------|
| SIRUNYAN | 18N | PRL 121 092002 | A.M. Sirunyan <i>et al.</i> | (CMS Collab.)   |
| AAD      | 12A | PRL 108 152001 | G. Aad <i>et al.</i>        | (ATLAS Collab.) |