

$\chi_{c0}(1P)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

$\chi_{c0}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3414.71± 0.30 OUR AVERAGE				
3413.0 ± 1.9 ±0.6	933	¹ AAIJ	17BB LHCb	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3414.2 ± 0.5 ±2.3	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
3406 ± 7 ±6	230	² ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
3414.21± 0.39±0.27		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
3414.7 + 0.7 - 0.6 ±0.2		³ ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
3415.5 ± 0.4 ±0.4	392	⁴ BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
3417.4 + 1.8 - 1.9 ±0.2		³ AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
3414.1 ± 0.6 ±0.8		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3417.8 ± 0.4 ±4		³ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3416 ± 3 ±4		⁵ TANENBAUM	78 MRK1	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3414.6 ± 1.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
3416.5 ± 3.0		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c0}$
3422 ±10		⁵ BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3415 ± 9		⁵ BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$

¹ From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c0}(1P)$ fixed to the PDG 16 value.
² From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.
³ Using mass of $\psi(2S) = 3686.0$ MeV.
⁴ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.
⁵ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

$\chi_{c0}(1P)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.9±0.6 OUR FIT	Error includes scale factor of 1.1.			
10.5±0.8 OUR AVERAGE	Error includes scale factor of 1.1.			
10.6±1.9±2.6	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
12.6 +1.5+0.9 -1.6-1.1		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
8.6 +1.7 -1.3 ±0.1		ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
9.7±1.0	392	¹ BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
16.6 +5.2 -3.7 ±0.1		AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
14.3±2.0±3.0		BAI	98I BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
13.5±3.3±4.2		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X, \gamma\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
13.2±2.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ Recalculated by ANDREOTTI 05A. **$\chi_{c0}(1P)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic decays		
Γ_1 $2(\pi^+\pi^-)$	$(2.18 \pm 0.11) \%$	S=1.2
Γ_2 $\rho^0\pi^+\pi^-$	$(8.5 \pm 2.7) \times 10^{-3}$	
Γ_3 $\rho^0\rho^0$		
Γ_4 $f_0(980)f_0(980)$	$(6.7 \pm 2.1) \times 10^{-4}$	
Γ_5 $\pi^+\pi^-\pi^0\pi^0$	$(3.3 \pm 0.4) \%$	
Γ_6 $\rho^+\pi^-\pi^0 + \text{c.c.}$	$(2.9 \pm 0.4) \%$	
Γ_7 $4\pi^0$	$(3.3 \pm 0.4) \times 10^{-3}$	
Γ_8 $\pi^+\pi^-K^+K^-$	$(1.81 \pm 0.16) \%$	S=1.2
Γ_9 $K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-$	$(9.9 \pm 4.0 \pm 2.8) \times 10^{-4}$	
Γ_{10} $K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	$(8.0 \pm 2.0 \pm 2.4) \times 10^{-4}$	
Γ_{11} $K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	$(6.3 \pm 1.9) \times 10^{-3}$	
Γ_{12} $K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	$< 2.7 \times 10^{-3}$	CL=90%
Γ_{13} $f_0(980)f_0(980)$	$(1.6 \pm 1.0 \pm 0.9) \times 10^{-4}$	
Γ_{14} $f_0(980)f_0(2200)$	$(8.0 \pm 2.0 \pm 2.5) \times 10^{-4}$	
Γ_{15} $f_0(1370)f_0(1370)$	$< 2.7 \times 10^{-4}$	CL=90%
Γ_{16} $f_0(1370)f_0(1500)$	$< 1.7 \times 10^{-4}$	CL=90%
Γ_{17} $f_0(1370)f_0(1710)$	$(6.7 \pm 3.5 \pm 2.3) \times 10^{-4}$	
Γ_{18} $f_0(1500)f_0(1370)$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{19} $f_0(1500)f_0(1500)$	$< 5 \times 10^{-5}$	CL=90%
Γ_{20} $f_0(1500)f_0(1710)$	$< 7 \times 10^{-5}$	CL=90%
Γ_{21} $K^+K^-\pi^+\pi^-\pi^0$	$(8.6 \pm 0.9) \times 10^{-3}$	
Γ_{22} $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(4.2 \pm 0.4) \times 10^{-3}$	
Γ_{23} $K^+K^-\pi^0\pi^0$	$(5.6 \pm 0.9) \times 10^{-3}$	
Γ_{24} $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(2.50 \pm 0.33) \%$	
Γ_{25} $\rho^+K^-K^0 + \text{c.c.}$	$(1.21 \pm 0.21) \%$	
Γ_{26} $K^*(892)^-K^+\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(4.6 \pm 1.2) \times 10^{-3}$	
Γ_{27} $K_S^0 K_S^0 \pi^+ \pi^-$	$(5.7 \pm 1.1) \times 10^{-3}$	
Γ_{28} $K^+K^-\eta\pi^0$	$(3.0 \pm 0.7) \times 10^{-3}$	
Γ_{29} $3(\pi^+\pi^-)$	$(1.96 \pm 0.22) \%$	S=3.4
Γ_{30} $K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(7.4 \pm 1.6) \times 10^{-3}$	

Γ_{31}	$K^*(892)^0 \bar{K}^*(892)^0$	$(1.7 \pm 0.6) \times 10^{-3}$	
Γ_{32}	$\pi\pi$	$(8.6 \pm 0.4) \times 10^{-3}$	S=1.2
Γ_{33}	$\pi^0\eta$	$< 1.8 \times 10^{-4}$	
Γ_{34}	$\pi^0\eta'$	$< 1.1 \times 10^{-3}$	
Γ_{35}	$\pi^0\eta_c$	$< 1.6 \times 10^{-3}$	CL=90%
Γ_{36}	$\eta\eta$	$(3.02 \pm 0.25) \times 10^{-3}$	S=1.3
Γ_{37}	$\eta\eta'$	$(9.1 \pm 1.1) \times 10^{-5}$	
Γ_{38}	$\eta'\eta'$	$(2.18 \pm 0.12) \times 10^{-3}$	
Γ_{39}	$\omega\omega$	$(9.7 \pm 1.1) \times 10^{-4}$	
Γ_{40}	$\omega\phi$	$(1.42 \pm 0.13) \times 10^{-4}$	
Γ_{41}	$\omega K^+ K^-$	$(1.94 \pm 0.21) \times 10^{-3}$	
Γ_{42}	$K^+ K^-$	$(6.07 \pm 0.33) \times 10^{-3}$	S=1.1
Γ_{43}	$K_S^0 K_S^0$	$(3.18 \pm 0.19) \times 10^{-3}$	S=1.1
Γ_{44}	$\pi^+ \pi^- \eta$	$< 2.0 \times 10^{-4}$	CL=90%
Γ_{45}	$\pi^+ \pi^- \eta'$	$< 4 \times 10^{-4}$	CL=90%
Γ_{46}	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$< 9 \times 10^{-5}$	CL=90%
Γ_{47}	$K^+ K^- \pi^0$	$< 6 \times 10^{-5}$	CL=90%
Γ_{48}	$K^+ K^- \eta$	$< 2.3 \times 10^{-4}$	CL=90%
Γ_{49}	$K^+ K^- K_S^0 K_S^0$	$(1.4 \pm 0.5) \times 10^{-3}$	
Γ_{50}	$K_S^0 K_S^0 K_S^0 K_S^0$	$(5.8 \pm 0.5) \times 10^{-4}$	
Γ_{51}	$K^+ K^- K^+ K^-$	$(2.8 \pm 0.4) \times 10^{-3}$	S=1.5
Γ_{52}	$K^+ K^- \phi$	$(9.7 \pm 2.5) \times 10^{-4}$	
Γ_{53}	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(3.7 \pm 0.6) \times 10^{-3}$	
Γ_{54}	$K^+ K^- \pi^0 \phi$	$(1.90 \pm 0.35) \times 10^{-3}$	
Γ_{55}	$3(K^+ K^-)$	$(1.08 \pm 0.22) \times 10^{-5}$	
Γ_{56}	$\phi \pi^+ \pi^- \pi^0$	$(1.18 \pm 0.15) \times 10^{-3}$	
Γ_{57}	$\phi\phi$	$(8.48 \pm 0.31) \times 10^{-4}$	
Γ_{58}	$\phi\phi\eta$	$(8.4 \pm 1.0) \times 10^{-4}$	
Γ_{59}	$p\bar{p}$	$(2.21 \pm 0.14) \times 10^{-4}$	S=1.6
Γ_{60}	$p\bar{p}\pi^0$	$(7.0 \pm 0.7) \times 10^{-4}$	S=1.3
Γ_{61}	$p\bar{p}\eta$	$(3.6 \pm 0.4) \times 10^{-4}$	
Γ_{62}	$p\bar{p}\omega$	$(5.3 \pm 0.6) \times 10^{-4}$	
Γ_{63}	$p\bar{p}\phi$	$(6.0 \pm 1.4) \times 10^{-5}$	
Γ_{64}	$p\bar{p}\pi^+ \pi^-$	$(2.1 \pm 0.7) \times 10^{-3}$	S=1.4
Γ_{65}	$p\bar{p}\pi^0 \pi^0$	$(1.04 \pm 0.28) \times 10^{-3}$	
Γ_{66}	$p\bar{p}K^+ K^- (\text{non-resonant})$	$(1.22 \pm 0.26) \times 10^{-4}$	
Γ_{67}	$p\bar{p}K_S^0 K_S^0$	$< 8.8 \times 10^{-4}$	CL=90%
Γ_{68}	$p\bar{p}K_S^0 K^- \pi^+ + \text{c.c.}$	$(2.6 \pm 0.4) \times 10^{-5}$	
Γ_{69}	$p\bar{n}\pi^-$	$(1.27 \pm 0.11) \times 10^{-3}$	
Γ_{70}	$\bar{p}n\pi^+$	$(1.37 \pm 0.12) \times 10^{-3}$	
Γ_{71}	$p\bar{n}\pi^- \pi^0$	$(2.35 \pm 0.21) \times 10^{-3}$	
Γ_{72}	$\bar{p}n\pi^+ \pi^0$	$(2.22 \pm 0.19) \times 10^{-3}$	
Γ_{73}	$\Lambda\bar{\Lambda}$	$(3.61 \pm 0.16) \times 10^{-4}$	S=1.1
Γ_{74}	$\Lambda\bar{\Lambda}\pi^+ \pi^-$	$(1.18 \pm 0.13) \times 10^{-3}$	

Γ_{75}	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	< 5	$\times 10^{-4}$	CL=90%
Γ_{76}	$\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$	< 5	$\times 10^{-4}$	CL=90%
Γ_{77}	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	< 5	$\times 10^{-4}$	CL=90%
Γ_{78}	$\Lambda\bar{\Lambda}\eta$	(2.3 ± 0.4)	$\times 10^{-4}$	
Γ_{79}	$\Lambda\bar{\Lambda}\omega$	(2.38 ± 0.34)	$\times 10^{-4}$	
Γ_{80}	$\Lambda\bar{\Lambda}\phi$	(3.0 ± 1.3)	$\times 10^{-5}$	
Γ_{81}	$K^+\bar{p}\Lambda + \text{c.c.}$	(1.25 ± 0.12)	$\times 10^{-3}$	S=1.3
Γ_{82}	$nK_S^0\bar{\Lambda} + \text{c.c.}$	(6.7 ± 0.5)	$\times 10^{-4}$	
Γ_{83}	$K^*(892)^+\bar{p}\Lambda + \text{c.c.}$	(4.8 ± 0.9)	$\times 10^{-4}$	
Γ_{84}	$K^+\bar{p}\Lambda(1520) + \text{c.c.}$	(3.0 ± 0.8)	$\times 10^{-4}$	
Γ_{85}	$\bar{p}\Lambda(1520)K_S^0\pi^+ + \text{c.c.}$	(1.6 ± 0.7)	$\times 10^{-5}$	
Γ_{86}	$\Lambda(1520)\bar{\Lambda}(1520)$	(3.1 ± 1.2)	$\times 10^{-4}$	
Γ_{87}	$\Sigma^0\bar{\Sigma}^0$	(4.70 ± 0.32)	$\times 10^{-4}$	
Γ_{88}	$\Sigma^+\bar{p}K_S^0 + \text{c.c.}$	(3.54 ± 0.27)	$\times 10^{-4}$	
Γ_{89}	$\Sigma^0\bar{p}K^+ + \text{c.c.}$	(3.05 ± 0.20)	$\times 10^{-4}$	
Γ_{90}	$\Sigma^+\bar{\Sigma}^-$	(4.7 ± 0.8)	$\times 10^{-4}$	S=2.6
Γ_{91}	$\Sigma^+\bar{\Sigma}^-\eta$	(1.27 ± 0.24)	$\times 10^{-4}$	
Γ_{92}	$\Sigma^-\bar{\Sigma}^+$	(5.2 ± 0.5)	$\times 10^{-4}$	
Γ_{93}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	(1.6 ± 0.6)	$\times 10^{-4}$	
Γ_{94}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	(2.3 ± 0.7)	$\times 10^{-4}$	
Γ_{95}	$K^-\Lambda\Xi^+ + \text{c.c.}$	(1.95 ± 0.35)	$\times 10^{-4}$	
Γ_{96}	$\Xi^0\Xi^0$	(4.5 ± 0.5)	$\times 10^{-4}$	S=1.7
Γ_{97}	$\Xi^-\Xi^+$	(4.48 ± 0.20)	$\times 10^{-4}$	
Γ_{98}	$\Omega^-\bar{\Omega}^+$	(3.5 ± 0.6)	$\times 10^{-5}$	
Γ_{99}	$\eta_c\pi^+\pi^-$	< 7	$\times 10^{-4}$	CL=90%

Radiative decays

Γ_{100}	$\gamma J/\psi(1S)$	$(1.41 \pm 0.09) \%$		S=1.7
Γ_{101}	$\gamma\rho^0$	< 9	$\times 10^{-6}$	CL=90%
Γ_{102}	$\gamma\omega$	< 8	$\times 10^{-6}$	CL=90%
Γ_{103}	$\gamma\phi$	< 6	$\times 10^{-6}$	CL=90%
Γ_{104}	$\gamma\gamma$	(2.06 ± 0.10)	$\times 10^{-4}$	S=1.1
Γ_{105}	$e^+e^- J/\psi(1S)$	(1.34 ± 0.30)	$\times 10^{-4}$	
Γ_{106}	$\mu^+\mu^- J/\psi(1S)$	< 1.9	$\times 10^{-5}$	CL=90%

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 88 branching ratios uses 255 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 393.1$ for 206 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_2	15									
x_8	10	1								
x_{30}	4	1	33							
x_{32}	21	3	9	3						
x_{36}	13	2	5	1	13					
x_{42}	19	3	8	3	20	12				
x_{43}	19	3	8	3	19	11	17			
x_{51}	7	1	5	3	6	3	6	5		
x_{57}	5	1	3	1	4	3	4	4	2	
x_{59}	6	1	2	1	0	1	5	5	2	1
x_{73}	24	4	9	3	24	15	22	21	6	5
x_{100}	6	1	2	1	8	5	6	5	1	1
x_{104}	11	2	-4	-4	14	9	11	12	-1	1
Γ	-19	-3	-21	-11	-16	-8	-16	-14	-12	-7
	x_1	x_2	x_8	x_{30}	x_{32}	x_{36}	x_{42}	x_{43}	x_{51}	x_{57}

x_{73}	7			
x_{100}	-37	7		
x_{104}	3	17	8	
Γ	-3	-15	-5	-43
	x_{59}	x_{73}	x_{100}	x_{104}

$\chi_{c0}(1P)$ PARTIAL WIDTHS

$\chi_{c0}(1P) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$					$\Gamma_1 \Gamma_{104} / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
48.7 ± 3.5 OUR FIT	Error includes scale factor of 1.2.					
49 ± 10 OUR AVERAGE	Error includes scale factor of 1.8.					
44.7 ± 3.6 ± 4.9	3.6k	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+\pi^-)$	
75 ± 13 ± 8		EISENSTEIN	01	CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c0}$	

$$\Gamma(\rho^0 \rho^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_3 \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	<252	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+ \pi^-)$
-----	----	------	--------	----	--

$$\Gamma(\pi^+ \pi^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_8 \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

41 ± 4 OUR FIT Error includes scale factor of 1.1.

38.8 ± 3.7 ± 4.7		1.7k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+ K^- \pi^+ \pi^-$
-------------------------	--	------	--------	----	---

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{21} \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

26 ± 4 ± 4		1094	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-------------------	--	------	-----------------	------	--

$$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{30} \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

17 ± 4 OUR FIT

16.7 ± 6.1 ± 3.0		495 ± 182	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+ K^- \pi^+ \pi^-$
-------------------------	--	-----------	--------	----	---

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{31} \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90	<148	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+ K^- \pi^+ \pi^-$
----	----	------	--------	----	---

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{32} \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

19.1 ± 1.5 OUR FIT Error includes scale factor of 1.2.

23 ± 5 OUR AVERAGE

29.7 ^{+17.4} _{-12.0} ± 4.8		103 ⁺⁶⁰ ₋₄₂	¹ UEHARA	09	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
--	--	-----------------------------------	---------------------	----	---

22.7 ± 3.2 ± 3.5		129 ± 18	² NAKAZAWA	05	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
------------------	--	----------	-----------------------	----	---

¹We multiplied the measurement by 3 to convert from $\pi^0 \pi^0$ to $\pi\pi$. Interference with the continuum included.

²We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{36} \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

9.4 ± 2.3 ± 1.2		22	¹ UEHARA	10A	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$
------------------------	--	----	---------------------	-----	--

¹Interference with the continuum not included.

$$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{39} \Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.9	90		¹ LIU	12B	BELL $\gamma\gamma \rightarrow 2(\pi^+ \pi^- \pi^0)$
------	----	--	------------------	-----	--

¹Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

$$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{40}\Gamma_{104}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.34	90	¹ LIU	12B	BELL $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-------	----	------------------	-----	---

¹ Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

$$\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_{104}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

13.6±1.1 OUR FIT Error includes scale factor of 1.1.

14.3±1.6±2.3	153 ± 17	NAKAZAWA 05	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
---------------------	----------	-------------	------	--

$$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{43}\Gamma_{104}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

7.1 ±0.6 OUR FIT Error includes scale factor of 1.1.

8.7 ±1.7 ±0.9	266	¹ UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
----------------------	-----	---------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.00±0.65±0.71	134 ± 12	CHEN	07B	BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c0}$
----------------	----------	------	-----	--

¹ Supersedes CHEN 07B.

$$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_{104}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

6.3±1.0 OUR FIT Error includes scale factor of 1.4.

7.9±1.3±1.1	215 ± 36	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+ K^-)$
--------------------	----------	--------	----	--

$$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{57}\Gamma_{104}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.89±0.13 OUR FIT Error includes scale factor of 1.1.

1.72±0.33±0.14	56 ± 11	¹ LIU	12B	BELL $\gamma\gamma \rightarrow 2(K^+ K^-)$
-----------------------	---------	------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ±0.9 ±0.4	23.6 ± 9.6	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+ K^-)$
---------------	------------	--------	----	--

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

$\chi_{c0}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-)) \quad \Gamma_2/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.39±0.12 OUR FIT

0.39±0.12	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$
------------------	--------------	------	---

$$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.14±0.09±0.05	1946	¹ ABLIKIM	24BT	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
----------------	------	----------------------	------	--

¹ ABLIKIM 24BT reports $2.127 \pm 0.002 \pm 0.101\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.7 \pm 2.1 \pm 0.2$	36 ± 9	¹ ABLIKIM	04G BES	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

¹ ABLIKIM 04G reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (6.5 \pm 1.6 \pm 1.3) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.3 \pm 0.4 \pm 0.1$	1751.4	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $3.54 \pm 0.10 \pm 0.43 \pm 0.18$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.9 \pm 0.4 \pm 0.1$	1358.5	^{1,2} HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $3.04 \pm 0.18 \pm 0.42 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.3 \pm 0.4 \pm 0.1$	3296	¹ ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ABLIKIM 11A reports $(3.34 \pm 0.06 \pm 0.44) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma(\pi^+\pi^-K^+K^-)$ Γ_{30}/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.09 OUR FIT			
0.41 ± 0.10	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

 $\Gamma(K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.9^{+3.6}_{-2.8} \pm 0.2$	83	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+\pi^-K^+K^-$

¹ ABLIKIM 05Q reports $(10.44 \pm 2.37^{+3.05}_{-1.90}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$8.0^{+2.0}_{-2.4} \pm 0.2$	62	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $(8.49 \pm 1.66^{+1.32}_{-1.99}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$6.3 \pm 1.9 \pm 0.1$	68	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $(6.66 \pm 1.31^{+1.60}_{-1.51}) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The measurement assumes $B(K_1(1270) \rightarrow K \rho(770)) = 42 \pm 6\%$.

$\Gamma(K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 2.7	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $< 2.85 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$. The measurement assumes $B(K_1(1400) \rightarrow K^*(892)\pi) = 94 \pm 6\%$.

$\Gamma(f_0(980) f_0(980))/\Gamma_{\text{total}}$					Γ_{13}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$16.3^{+10.5}_{-9.0} \pm 0.4$	28	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980) f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.59 \pm 0.50^{+0.89}_{-0.72}) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. One of the $f_0(980)$ mesons is identified via decay to $\pi^+ \pi^-$ while the other via $K^+ K^-$ decay.

$\Gamma(f_0(980)f_0(2200))/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.0^{+2.0}_{-2.5} \pm 0.2$	77	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

¹ ABLIKIM 05Q reports $(8.42 \pm 1.42^{+1.65}_{-2.29}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(2200))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(980) \rightarrow \pi^+ \pi^-$ and $f_0(2200) \rightarrow K^+ K^-$ decays.

 $\Gamma(f_0(1370)f_0(1370))/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

¹ ABLIKIM 05Q reports $< 2.9 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$. One of the $f_0(1370)$ mesons is identified via decay to $\pi^+ \pi^-$ while the other via $K^+ K^-$ decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

 $\Gamma(f_0(1370)f_0(1500))/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

¹ ABLIKIM 05Q reports $< 1.8 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+ \pi^-$ and $f_0(1500) \rightarrow K^+ K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

 $\Gamma(f_0(1370)f_0(1710))/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.7^{+3.5}_{-2.3} \pm 0.2$	61	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

¹ ABLIKIM 05Q reports $(7.12 \pm 1.85^{+3.28}_{-1.68}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+ \pi^-$ and $f_0(1710) \rightarrow K^+ K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

$\Gamma(f_0(1500)f_0(1370))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
¹ ABLIKIM 05Q reports $< 1.4 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+ \pi^-$ and $f_0(1370) \rightarrow K^+ K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

 $\Gamma(f_0(1500)f_0(1500))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
¹ ABLIKIM 05Q reports $< 0.55 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$. One of the $f_0(1500)$ is identified via decay to $\pi^+ \pi^-$ while the other via $K^+ K^-$ decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

 $\Gamma(f_0(1500)f_0(1710))/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
¹ ABLIKIM 05Q reports $< 0.73 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+ \pi^-$ and $f_0(1710) \rightarrow K^+ K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.61 \pm 0.13 \pm 0.94$	9.0k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.				

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.22 \pm 0.10 \pm 0.43$	2.7k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.				

 $\Gamma(K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.56 \pm 0.09 \pm 0.01$	213.5	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

^1HE 08B reports $0.59 \pm 0.05 \pm 0.08 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.50 \pm 0.33 \pm 0.06$	401.7	^1HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

^1HE 08B reports $2.64 \pm 0.15 \pm 0.31 \pm 0.14$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.21 \pm 0.21 \pm 0.03$	179.7	^1HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

^1HE 08B reports $1.28 \pm 0.16 \pm 0.15 \pm 0.07$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.46 \pm 0.12 \pm 0.01$	64.1	^1HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

^1HE 08B reports $0.49 \pm 0.10 \pm 0.07 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.7 \pm 1.1 \pm 0.1$	152 ± 14	$^1\text{ABLIKIM}$	050 BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$

$^1\text{ABLIKIM}$ 050 reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ $= (0.558 \pm 0.051 \pm 0.089) \times 10^{-3}$ which we divide by our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.30 \pm 0.07 \pm 0.01$	56.4	^1HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹HE 08B reports $0.32 \pm 0.05 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3(\pi^+ \pi^-)) / \Gamma_{\text{total}}$


Γ_{29} / Γ

VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT
19.6±2.2 OUR AVERAGE	Error includes scale factor of 3.4. See the ideogram below.			
20.9±0.5±0.5	145K	¹ ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
11.7±1.0±1.9		² BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
12.5±2.9±0.5		² TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ABLIKIM 22Q reports $(2.080 \pm 0.006 \pm 0.068) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow 3(\pi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

WEIGHTED AVERAGE
19.6±2.2 (Error scaled by 3.4)



$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.72^{+0.60}_{-0.55} \pm 0.04$	64	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.57 \pm 0.40 \pm 0.04$ 30 ± 6 ^{2,3} ABLIKIM 04H BES Repl. by ABLIKIM 05Q

¹ ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(0.168 \pm 0.035^{+0.047}_{-0.040}) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

³ ABLIKIM 04H reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(1.53 \pm 0.29 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-3}$	90	¹ ABLIKIM	15N BES3	$\psi(2S) e^+ e^- \rightarrow \gamma \pi^0 \eta_c$

¹ Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$.

 $\Gamma(\eta \eta')/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$9.1 \pm 1.1 \pm 0.2$	85		¹ ABLIKIM	17AI BES3	$\psi(2S) \rightarrow \gamma \eta' \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 24 90 35 ± 13 ² ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \eta' \eta$

< 50 90 ³ ADAMS 07 CLEO $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ABLIKIM 17AI reports $(8.92 \pm 0.84 \pm 0.65) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ASNER 09 reports $< 0.25 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

³ Superseded by ASNER 09. ADAMS 07 reports $< 0.5 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

 $\Gamma(\eta' \eta')/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.18 ± 0.12 OUR AVERAGE				

$2.24 \pm 0.13 \pm 0.05$ 2.5k ¹ ABLIKIM 17AI BES3 $\psi(2S) \rightarrow \gamma \eta' \eta'$

$2.01 \pm 0.21 \pm 0.05$ 0.4k ² ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \eta' \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.61 \pm 0.41 \pm 0.04$ 23 ³ ADAMS 07 CLEO $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ABLIKIM 17Al reports $(2.19 \pm 0.03 \pm 0.14) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ASNER 09 reports $(2.12 \pm 0.13 \pm 0.21) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by ASNER 09. ADAMS 07 reports $(1.7 \pm 0.4 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.0922 \pm 0.0011 \pm 0.0046$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$

Γ_{39}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97±0.11 OUR AVERAGE				
$0.94 \pm 0.11 \pm 0.02$	991	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
$2.18 \pm 0.66 \pm 0.05$	38.1 ± 9.6	² ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow \gamma 6\pi$

¹ ABLIKIM 11K reports $(0.95 \pm 0.03 \pm 0.11) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 05N reports $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(0.212 \pm 0.053 \pm 0.037) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\phi)/\Gamma_{\text{total}}$

Γ_{40}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.42±0.13±0.03				
	486	¹ ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.18 \pm 0.22 \pm 0.03$	76	^{2,3} ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
--------------------------	----	------------------------	----------	---------------------------------------

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(13.83 \pm 0.70 \pm 1.01) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 11K reports $(1.2 \pm 0.1 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by ABLIKIM 19J.

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.94 \pm 0.06 \pm 0.20$	1.4k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.

$\Gamma(\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.0 90 ² ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ATHAR 07 reports $< 0.21 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

² ABLIKIM 06R reports $< 1.1 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $< 0.38 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.09	90	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7 90 ^{2,3} ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$

<0.7 90 ^{3,4} BAI 99B BES $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ATHAR 07 reports $< 0.10 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

² ABLIKIM 06R reports $< 0.70 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

³ We have multiplied the $K_S^0 K^+ \pi^-$ measurement by a factor of 2 to convert to $K^0 K^+ \pi^-$.

⁴ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	¹ ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
¹ ATHAR 07 reports $< 0.06 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.				

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.23	90	¹ ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
¹ ATHAR 07 reports $< 0.24 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.				

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.48 \pm 0.03$	16.8 ± 4.8	¹ ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
¹ ABLIKIM 050 reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.138 \pm 0.039 \pm 0.025) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.8 \pm 0.5 \pm 0.1$	319	¹ ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$
¹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (5.64 \pm 0.33 \pm 0.37) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..				

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.97 \pm 0.25 \pm 0.02$	38	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
¹ ABLIKIM 06T reports $(1.03 \pm 0.22 \pm 0.15) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$3.68 \pm 0.30 \pm 0.50$	ABLIKIM 15M	BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(K^+ K^- \pi^0 \phi) / \Gamma_{\text{total}}$ Γ_{54} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$1.90 \pm 0.14 \pm 0.32$	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(\phi \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{56} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.18 \pm 0.07 \pm 0.13$	538	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.

$\Gamma(\phi \phi) / \Gamma_{\text{total}}$ Γ_{57} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.848 ± 0.031 OUR FIT				
$0.859 \pm 0.027 \pm 0.020$	2701	¹ ABLIKIM	23N BES3	$\psi(2S) \rightarrow \gamma$ hadrons

¹ Measured using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ and $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ from PDG 22.

$\Gamma(\phi \phi \eta) / \Gamma_{\text{total}}$ Γ_{58} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.4 \pm 1.0 \pm 0.2$	186.6	¹ ABLIKIM	20B BES3	$\psi(2S) \rightarrow \gamma \phi \phi \eta$

¹ ABLIKIM 20B reports $(8.41 \pm 0.74 \pm 0.62) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \phi \phi \eta) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p \bar{p} \pi^0) / \Gamma_{\text{total}}$ Γ_{60} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.70 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.3.		
$0.73 \pm 0.06 \pm 0.02$	¹ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p \bar{p} X$
$0.56 \pm 0.12 \pm 0.01$	² ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(7.76 \pm 0.37 \pm 0.51 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p \bar{p} \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.59 \pm 0.10 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p \bar{p} \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p \bar{p} \eta) / \Gamma_{\text{total}}$ Γ_{61} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.36 ± 0.04 OUR AVERAGE			
$0.35 \pm 0.04 \pm 0.01$	¹ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p \bar{p} X$
$0.37 \pm 0.11 \pm 0.01$	² ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(3.73 \pm 0.38 \pm 0.28 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.39 \pm 0.11 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$

Γ_{62}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$0.53 \pm 0.06 \pm 0.01$	¹ ONYISI 10	CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$

¹ ONYISI 10 reports $(5.57 \pm 0.48 \pm 0.42 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$

Γ_{63}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.0 \pm 1.4 \pm 0.1$	42 ± 8	¹ ABLIKIM 11F	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $(6.12 \pm 1.18 \pm 0.86) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

Γ_{64}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.1 ± 0.7	OUR EVALUATION		Error includes scale factor of 1.4. Treating systematic error as correlated.

2.1 ± 1.0 OUR AVERAGE Error includes scale factor of 2.0.

$1.57 \pm 0.21 \pm 0.53$ ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c0}$

$4.20 \pm 1.15 \pm 0.18$ ¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$

Γ_{65}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.104 \pm 0.028 \pm 0.002$	39.5	¹ HE 08B	CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹ HE 08B reports $0.11 \pm 0.02 \pm 0.02 \pm 0.01\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.22 \pm 0.26 \pm 0.03$	48 ± 8	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $(1.24 \pm 0.20 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8.8	90	¹ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$

¹ Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.7 ± 1.1 OUR AVERAGE				
$12.9 \pm 1.1 \pm 0.3$	5150	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
$11.3 \pm 3.1 \pm 0.3$		² ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^-X$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ = $(1.26 \pm 0.02 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06I reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ = $(1.10 \pm 0.24 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$13.7 \pm 1.2 \pm 0.3$	5808	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ = $(1.34 \pm 0.03 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$23.5 \pm 2.0 \pm 0.5$	2480	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ = $(2.29 \pm 0.08 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$22.2 \pm 1.8 \pm 0.5$	2757	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+\pi^0$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (2.16 \pm 0.07 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$118 \pm 12 \pm 3$		426	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<400 90 ² ABLIKIM 06D BES2 $\psi(2S) \rightarrow \chi_{c0} \gamma$

¹ ABLIKIM 12I reports $(119.0 \pm 6.4 \pm 11.4) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda} \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.2 \pm 0.5)\%$

$\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

¹ ABLIKIM 12I reports $< 54 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda} \pi^+ \pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Lambda} \pi^-$

¹ ABLIKIM 12I reports $< 55 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Lambda} \pi^+$

¹ ABLIKIM 12I reports $< 50 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.25±0.12 OUR AVERAGE		Error includes scale factor of 1.3.		
1.31±0.09±0.03	9k	^{1,2} ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$
1.01±0.19±0.02		³ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ABLIKIM 13D reports $(1.32 \pm 0.03 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.

³ ATHAR 07 reports $(1.07 \pm 0.17 \pm 0.12) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.8±0.9±0.1	254	¹ ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+} \bar{p} \Lambda$

¹ ABLIKIM 19AU reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.7 \pm 0.7 \pm 0.5) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+\bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0±0.7±0.1	62 ± 12	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

¹ ABLIKIM 11F reports $(3.00 \pm 0.58 \pm 0.50) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±0.5±0.2	1284	¹ ABLIKIM	21AV BES3	$\psi(2S) \rightarrow \gamma n K_S^0 \bar{\Lambda} + \text{c.c.}$

¹ ABLIKIM 21AV reports $(6.65 \pm 0.26 \pm 0.41) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.0979 \pm 0.0020$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Also uses $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = (63.9 \pm 0.5)\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$.

$\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.1 \pm 1.2 \pm 0.1$	28 ± 10	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

¹ ABLIKIM 11F reports $(3.18 \pm 1.11 \pm 0.53) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.70 ± 0.32 OUR AVERAGE				

$4.84 \pm 0.34 \pm 0.11$	1046	¹ ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
--------------------------	------	----------------------	----------	---

$4.2 \pm 0.7 \pm 0.1$	78 ± 10	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
-----------------------	-------------	-------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.7 \pm 0.5 \pm 0.1$	243	^{3,4} ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
-----------------------	-----	------------------------	----------	---

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(4.72 \pm 0.18 \pm 0.28) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² NAIK 08 reports $(4.41 \pm 0.56 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 13H reports $(4.78 \pm 0.34 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Superseded by ABLIKIM 18V

 $\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.7 ± 0.8 OUR AVERAGE	Error includes scale factor of 2.6.			

$5.12 \pm 0.35 \pm 0.12$	747	¹ ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
--------------------------	-----	----------------------	----------	---

$3.1 \pm 0.7 \pm 0.1$	39 ± 7	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
-----------------------	------------	-------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.5 \pm 0.5 \pm 0.1$	148	^{3,4} ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
-----------------------	-----	------------------------	----------	---

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(4.99 \pm 0.24 \pm 0.24) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² NAIK 08 reports $(3.25 \pm 0.57 \pm 0.43) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 13H reports $(4.54 \pm 0.42 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Superseded by ABLIKIM 18V

$\Gamma(\Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}$

Γ_{91}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
12.7$\pm$2.4$\pm$0.3	74	¹ ABLIKIM	24CA BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}(1P)$

¹ ABLIKIM 24CA reports $(12.6 \pm 2.0 \pm 1.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$

Γ_{92}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.2$\pm$0.5$\pm$0.1	2143	¹ ABLIKIM	20I BES3	$\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

¹ ABLIKIM 20I reports $(5.13 \pm 0.24 \pm 0.41) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

Γ_{93}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.3$\pm$5.9$\pm$0.4	27	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

¹ ABLIKIM 12I reports $(16.4 \pm 5.7 \pm 1.6) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$

Γ_{94}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
23$\pm$7$\pm$1	33	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

¹ ABLIKIM 12I reports $(23.5 \pm 6.2 \pm 2.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.95 \pm 0.35 \pm 0.04$	57	¹ ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \Xi^+ + \text{c.c.}$
¹ ABLIKIM 15I reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.90 \pm 0.30 \pm 0.16) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.5 ± 0.5 OUR AVERAGE	Error includes scale factor of 1.7.			
$4.69 \pm 0.31 \pm 0.11$	1741	¹ ABLIKIM	22O BES3	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$
$3.2 \pm 0.8 \pm 0.1$	23.3 ± 4.9	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$
¹ ABLIKIM 22O reports $(4.67 \pm 0.19 \pm 0.26) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² NAIK 08 reports $(3.34 \pm 0.70 \pm 0.48) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.48 ± 0.20 OUR AVERAGE					
$4.45 \pm 0.18 \pm 0.10$		4932	¹ ABLIKIM	22O BES3	$\psi(2S) \rightarrow \gamma \Xi^- \Xi^+$
$4.9 \pm 0.7 \pm 0.1$		95	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^- \Xi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<10.3		90	³ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0} \gamma$
¹ ABLIKIM 22O reports $(4.43 \pm 0.08 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² NAIK 08 reports $(5.14 \pm 0.60 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
³ Using $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.2 \pm 0.5)\%$					

 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.51 \pm 0.54 \pm 0.29$	284	ABLIKIM	23T BES3	$\chi_{cJ} \rightarrow \Omega^- \bar{\Omega}^+$

$$\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}} \qquad \Gamma_{99}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7 \times 10^{-4}$	90	^{1,2} ABLIKIM 13B	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 41 \times 10^{-4}$	90	^{1,3} ABLIKIM 13B	BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
-----------------------	----	----------------------------	------	---

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.

² From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

³ From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}} \qquad \Gamma_{59}/\Gamma \times \Gamma_{32}/\Gamma$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

18.9 ± 1.6 OUR FIT Error includes scale factor of 1.5.

15.3 ± 2.4 ± 0.8	¹ ANDREOTTI 03	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0 \pi^0$
-------------------------	---------------------------	------	--

¹ We have multiplied $B(p\bar{p}) \cdot B(\pi^0 \pi^0)$ measurement by 3 to obtain $B(p\bar{p}) \cdot B(\pi\pi)$.

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0 \eta)/\Gamma_{\text{total}} \qquad \Gamma_{59}/\Gamma \times \Gamma_{33}/\Gamma$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

< 0.4	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$
-----------------	---------------	------	-----------------------------------

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0 \eta')/\Gamma_{\text{total}} \qquad \Gamma_{59}/\Gamma \times \Gamma_{34}/\Gamma$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

< 2.5	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$
-----------------	---------------	------	-----------------------------------

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}} \qquad \Gamma_{59}/\Gamma \times \Gamma_{36}/\Gamma$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

6.7 ± 0.7 OUR FIT Error includes scale factor of 1.4.

4.0 ± 1.2^{+0.5}_{-0.3}	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \eta\eta$
--	---------------	------	---------------------------------

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta')/\Gamma_{\text{total}} \qquad \Gamma_{59}/\Gamma \times \Gamma_{37}/\Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.1^{+2.3}_{-1.5}$	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$
---------------------	---------------	------	-----------------------------------

RADIATIVE DECAYS

$$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}} \qquad \Gamma_{101}/\Gamma$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 9	90	1.2 ± 4.5	¹ BENNETT 08A	CLEO	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
---------------	----	---------------	--------------------------	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 10	90	6 ± 12	² ABLIKIM 11E	BES3	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
--------	----	------------	--------------------------	------	---

¹ BENNETT 08A reports $< 9.6 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

² ABLIKIM 11E reports $< 10.5 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$ Γ_{102}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

< 8	90	0.0 $\pm$ 2.8	¹ BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$
---------------	----	---------------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13	90	5 $\pm$ 11	² ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
---------------	----	------------	----------------------	----------	---

¹ BENNETT 08A reports $< 8.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

² ABLIKIM 11E reports $< 12.9 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{103}/Γ

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

< 6	90	0.1 $\pm$ 1.6	¹ BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
---------------	----	---------------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	15 $\pm$ 7	² ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
---------------	----	------------	----------------------	----------	---

¹ BENNETT 08A reports $< 6.4 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

² ABLIKIM 11E reports $< 16.2 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(e^+e^-J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{105}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.55 $\pm$ 0.33 $\pm$ 0.03	56	^{1,2} ABLIKIM	17I BES3	$\psi(2S) \rightarrow \gamma e^+e^-J/\psi$
----------------------------	----	------------------------	----------	--

¹ ABLIKIM 17I reports $(1.51 \pm 0.30 \pm 0.13) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow e^+e^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.22) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Not independent from other measurements reported by ABLIKIM 17I

$\Gamma(e^+e^-J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$ $\Gamma_{105}/\Gamma_{100}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

9.5 $\pm$ 1.9 $\pm$ 0.7	56	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$
--	----	----------------------	----------	--

¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(\mu^+\mu^-J/\psi(1S))/\Gamma(e^+e^-J/\psi(1S))$ $\Gamma_{106}/\Gamma_{105}$

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	-------------	--------------------	-------------	----------------

<0.14	90	<9.5	ABLIKIM	19Z BES3	$\psi(2S) \rightarrow \gamma\chi_c \rightarrow$ $\gamma(\mu^+\mu^-J/\psi)$
-----------------	----	------	---------	----------	---

$$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S)) \quad \Gamma_{104}/\Gamma_{100}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.46 $\pm$ 0.13 OUR FIT Error includes scale factor of 1.6.**2.0 $\pm$ 0.4 OUR AVERAGE**2.2 $\pm$ 0.4 $^{+0.1}_{-0.2}$ ¹ ANDREOTTI 04 E835 $p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$ 1.45 $\pm$ 0.74 ² AMBROGIANI 00B E835 $\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$ ¹ The values of $B(p\bar{p})B(\gamma\gamma)$ and $B(\gamma\gamma)B(\gamma J/\psi)$ measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.² Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.
$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_{59}/\Gamma \times \Gamma_{100}/\Gamma$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

31.2 $\pm$ 2.2 OUR FIT Error includes scale factor of 1.4.**28.2 $\pm$ 2.1 OUR AVERAGE**28.0 $\pm$ 1.9 $\pm$ 1.3 392 ^{1,2,3} BAGNASCO 02 E835 $\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$ 29.3 $^{+5.7}_{-4.7}$ $\pm$ 1.5 89 ^{1,2} AMBROGIANI 99B $\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$ ¹ Values in $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$ and $(\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$ are not independent. The latter is used in the fit since it is less correlated to the total width.² Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.³ Recalculated by ANDREOTTI 05A. **$\chi_{c0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS**

$$\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \quad \Gamma_1/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

6.12 $\pm$ 0.26 OUR FIT Error includes scale factor of 1.1.**6.9 $\pm$ 2.4 OUR AVERAGE** Error includes scale factor of 3.8.4.4 $\pm$ 0.1 $\pm$ 0.9 ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c0}$ 9.3 $\pm$ 0.9 ² TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c0}$ ¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow 2\pi^+2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].² The value $B(\psi(1S) \rightarrow \gamma\chi_{c0}) \times B(\chi_{c0} \rightarrow 2\pi^+2\pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
$$\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.12 $\pm$ 0.09 OUR FIT Error includes scale factor of 1.1.**2.082 $\pm$ 0.002 $\pm$ 0.089** 1946 ¹ ABLIKIM 24BT BES3 $\psi(2S) \rightarrow \gamma\chi_{c0}$ ¹ Calculated by us. The value given here is derived from the value of $B(\chi_{c0} \rightarrow 2(\pi^+\pi^-))$ reported in ABLIKIM 24BT using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.79 \pm 0.20)\%$ [PDG 22].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_8/\Gamma \times \Gamma_{181}^{\psi(2S)}}{\Gamma_{181}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.77 ± 0.15 OUR FIT	Error includes scale factor of 1.1.		
1.64 ± 0.05 ± 0.2	ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_8/\Gamma \times \Gamma_{181}^{\psi(2S)}}{\Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.1 ± 0.4 OUR FIT	Error includes scale factor of 1.1.		
5.8 ± 1.6 OUR AVERAGE	Error includes scale factor of 2.3.		
4.22 ± 0.20 ± 0.97	BAI	99B	BES $\psi(2S) \rightarrow \gamma \chi_{c0}$
7.4 ± 1.0	¹ TANENBAUM	78	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi \pi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{32}/\Gamma \times \Gamma_{181}^{\psi(2S)}}{\Gamma_{181}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.33 ± 0.34 OUR FIT	Error includes scale factor of 1.2.			
8.80 ± 0.34 OUR AVERAGE				
9.11 ± 0.08 ± 0.65	17k	¹ ABLIKIM	10A	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
8.81 ± 0.11 ± 0.43	8.9k	² ASNER	09	CLEO $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
8.13 ± 0.19 ± 0.89	2.8k	³ ASNER	09	CLEO $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c0} \rightarrow \pi^0 \pi^0) = (3.23 \pm 0.03 \pm 0.23 \pm 0.14) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.

² Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^+ \pi^-) = (6.37 \pm 0.08 \pm 0.31 \pm 0.32) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi \pi$.

³ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^0 \pi^0) = (2.94 \pm 0.07 \pm 0.32 \pm 0.15) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi \pi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{32}/\Gamma \times \Gamma_{181}^{\psi(2S)}}{\Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
24.0 ± 1.0 OUR FIT	Error includes scale factor of 1.2.			
20.7 ± 1.7 OUR AVERAGE				
23.9 ± 2.7 ± 4.1	97 ± 11	¹ BAI	03C	BES $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow \gamma \pi^0 \pi^0$
20.2 ± 1.1 ± 1.5	720 ± 32	² BAI	98I	BES $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow \gamma \pi^+ \pi^-$

¹ We have multiplied $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.

² Calculated by us. The value for $B(\chi_{c0} \rightarrow \pi^+ \pi^-)$ reported in BAI 98I is derived using $B(\psi' \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi' \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi \pi$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}} \quad \Gamma_{36}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.94±0.22 OUR FIT		Error includes scale factor of 1.2.		

3.12±0.19 OUR AVERAGE

3.23±0.09±0.23 2132 ¹ ABLIKIM 10A BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

2.93±0.12±0.29 0.9k ² ASNER 09 CLEO $\psi(2S) \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.86±0.46±0.37 48 ³ ADAMS 07 CLEO $\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c0} \rightarrow \eta\eta) = (3.44 \pm 0.10 \pm 0.24 \pm 0.13) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$.

² Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \eta\eta) = (3.18 \pm 0.13 \pm 0.31 \pm 0.16) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

³ Superseded by ASNER 09. Calculated by us. The value of $B(\chi_{c0}(1P) \rightarrow \eta\eta)$ reported by ADAMS 07 was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46)\%$ (ATHAR 04).

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{36}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.85 ±0.06 OUR FIT	Error includes scale factor of 1.2.		

0.578±0.241±0.158 BAI 03C BES $\psi(2S) \rightarrow \gamma\eta\eta$

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}} \quad \Gamma_{42}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.92±0.28 OUR FIT				

5.97±0.07±0.32 8.1k ¹ ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K^+K^-$

¹ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow K^+K^-) = (6.47 \pm 0.08 \pm 0.35 \pm 0.32) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \quad \Gamma_{42}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.71±0.08 OUR FIT				

1.63±0.10±0.15 774 ± 38 ¹ BAI 98I BES $\psi(2S) \rightarrow \gamma K^+K^-$

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow K^+K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}} \quad \Gamma_{43}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.10±0.16 OUR FIT				

3.18±0.17 OUR AVERAGE

3.22±0.07±0.17 2.1k ¹ ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

3.02±0.19±0.33 322 ABLIKIM 05O BES2 $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow K_S^0 K_S^0) = (3.49 \pm 0.08 \pm 0.18 \pm 0.17) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{43}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.9±0.5 OUR FIT			
5.6±0.8±1.3	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{51}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8 ±0.4 OUR FIT	Error includes scale factor of 1.5.			
3.20±0.11±0.41	278	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma_{51}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.0±1.2 OUR FIT	Error includes scale factor of 1.5.		
6.1±0.8±0.9	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow 3(K^+ K^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{55}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±1.8±1.1	37.7 ± 6.2	¹ ABLIKIM	24P BES3	$e^+ e^- \rightarrow \psi(2S)$

¹ Systematic error derived by us, based on the text.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{57}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.827±0.033 OUR FIT				
0.78 ±0.08 OUR AVERAGE				
0.77 ±0.03 ±0.08	612	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma \text{ hadrons}$
0.86 ±0.19 ±0.12	26	² ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31)\%$.

² Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{57}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

2.38±0.10 OUR FIT**2.6 ±1.0 ±1.1**¹ BAI

99B

BES

 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{59}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6})

EVTS

DOCUMENT ID

TECN

COMMENT

21.6±1.3 OUR FIT Error includes scale factor of 1.5.**23.7±1.0 OUR AVERAGE**

23.7±0.8±0.9

1222

ABLIKIM

13v

BES3

 $\psi(2S) \rightarrow \gamma p\bar{p}$

23.7±1.4±1.4

383 ± 22

¹ NAIK

08

CLEO

 $\psi(2S) \rightarrow \gamma p\bar{p}$ 23.6^{+3.7}_{-3.4}±3.489.5⁺¹⁴₋₁₃

BAI

04F

BES

 $\psi(2S) \rightarrow \gamma\chi_{c0}(1P) \rightarrow \gamma\bar{p}p$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow p\bar{p}) = (25.7 \pm 1.5 \pm 1.5 \pm 1.3) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{59}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5})

DOCUMENT ID

TECN

COMMENT

6.2±0.4 OUR FIT Error includes scale factor of 1.5.**4.6±1.9**¹ BAI

98I

BES

 $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\bar{p}p$

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{68}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6})

EVTS

DOCUMENT ID

TECN

COMMENT

2.55±0.26±0.31

173

ABLIKIM

24BX

BES3

 $\psi(2S) \rightarrow \gamma\chi_{c0}$

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}\Lambda(1520)K_S^0 \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{85}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6})

EVTS

DOCUMENT ID

TECN

COMMENT

1.57^{+0.66}_{-0.62}±0.22

27

ABLIKIM

24BX

BES3

 $\psi(2S) \rightarrow \gamma\chi_{c0}$

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \times \Gamma_{73} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

35.2±1.3 OUR FIT**35.1±1.4 OUR AVERAGE** Error includes scale factor of 1.1.35.6±1.0±1.0 1486 ABLIKIM 21L BES3 $\psi(2S) \rightarrow \gamma p \pi^- \bar{p} \pi^+$ 31.2±3.3±2.0 131 ¹ NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

32.0±1.9±2.2 369 ^{2,3} ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$ ¹ Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow \Lambda \bar{\Lambda}) = (33.8 \pm 3.6 \pm 2.2 \pm 1.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.² Superseded by ABLIKIM 21L³ Calculated by us. ABLIKIM 13H reports $B(\chi_{c0} \rightarrow \Lambda \bar{\Lambda}) = (33.3 \pm 2.0 \pm 2.6) \times 10^{-5}$ from a measurement of $B(\chi_{c0} \rightarrow \Lambda \bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma \chi_{c0})$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.62 \pm 0.31)\%$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \times \Gamma_{73} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

10.1±0.4 OUR FIT**13.0^{+3.6}_{-3.5}±2.5** 15.2^{+4.2}_{-4.0} ¹ BAI 03E BES $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$ ¹ BAI 03E reports $[B(\chi_{c0} \rightarrow \Lambda \bar{\Lambda}) B(\psi(2S) \rightarrow \gamma \chi_{c0}) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p \bar{p})] = (2.45^{+0.68}_{-0.65} \pm 0.46)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p \bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda} \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \times \Gamma_{78} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.26±0.30±0.20 67 ABLIKIM 22AO BES3 $\psi(2S) \rightarrow \gamma p \pi^- \bar{p} \pi^+ \gamma \gamma$

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda} \omega) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \times \Gamma_{79} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

23.2±2.2±2.4 316 ± 30 ¹ ABLIKIM 24BE BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$ ¹ Calculated by us. The authors report $B(\chi_{c0} \rightarrow \Lambda \bar{\Lambda} \omega)$ obtained from a product using PDG 22 value of $B(\psi(2S) \rightarrow \gamma \chi_{c0})$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda} \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \times \Gamma_{80} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.92±1.22±0.19 7.2 ABLIKIM 24AC BES3 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$$\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \\ \Gamma_{88} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.45 ± 0.17 ± 0.19	493	¹ ABLIKIM	19BB BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_c^0 \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (3.52 \pm 0.19 \pm 0.21) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_c^0) = (9.79 \pm 0.20)\%$ and other branching fractions from PDG 18.

$$\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \\ \Gamma_{89} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.97 ± 0.12 ± 0.14	871	¹ ABLIKIM	20AE BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+ + \text{c.c.}$

¹ Calculated by us. ABLIKIM 20AE reports $B(\chi_c^0 \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (3.03 \pm 0.12 \pm 0.15) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_c^0) = (9.79 \pm 0.20)\%$ and other branching fractions from PDG 20.

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}} \\ \Gamma_{100} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.137 ± 0.009 OUR FIT	Error includes scale factor of 1.7.			

0.147 ± 0.029 OUR AVERAGE Error includes scale factor of 4.6.

0.158 ± 0.003 ± 0.006	4.8k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
0.024 ± 0.015 ± 0.205	12k	ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
0.069 ± 0.018		² OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.4 ± 0.3		³ BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.16 ± 0.11		³ BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma \chi_{c0}$
3.3 ± 1.7		⁴ BIDDICK	77 CNTR	$e^+ e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.151 ± 0.003 ± 0.010	4.3k	⁵ ABLIKIM	120 BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.125 ± 0.007 ± 0.013	560	⁶ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.18 ± 0.01 ± 0.02	172	⁷ ADAM	05A CLEO	Repl. by MENDEZ 08

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{100}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}{\Gamma_{104}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.396±0.025 OUR FIT	Error includes scale factor of 1.7.			
0.358±0.020±0.037	560	MENDEZ	08	CLEO $\psi(2S) \rightarrow \gamma \chi_{c0}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.55 ±0.04 ±0.06	172	¹ ADAM	05A	CLEO Repl. by MENDEZ 08

¹ Not independent from other values reported by ADAM 05A.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma \gamma)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{104}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}{\Gamma_{104}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00±0.09 OUR FIT	Error includes scale factor of 1.1.			
1.95±0.09 OUR AVERAGE				
1.93±0.08±0.05	3.5k	ABLIKIM	17AE	BES3 $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$
2.17±0.32±0.10	0.2k	ECKLUND	08A	CLEO $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$
3.7 ±1.8 ±1.0		LEE	85	CBAL $\psi(2S) \rightarrow \gamma \chi_{c0}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.17±0.17±0.12	0.8k	¹ ABLIKIM	12A	BES3 $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$

¹ Superseded by ABLIKIM 17AE.

$\chi_{c0}(1P)$ REFERENCES

ABLIKIM	24AC	PR D110 032016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BE	PR D110 032022	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BT	PR D110 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BX	PR D110 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CA	PR D110 112013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24P	PR D109 072016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23N	JHEP 2305 069	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23T	PR D107 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AO	PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22O	JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ABLIKIM	21AV	JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AE	PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20I	PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AI	PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ANDREOTTI	05C	PR D72 112002	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	04	PL B584 16	M. Andreotti <i>et al.</i>	(E835 Collab.)
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ANDREOTTI	03	PRL 91 091801	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
BAGNASCO	02	PL B533 237	S. Bagnasco <i>et al.</i>	(FNAL E835 Collab.)
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
AMBROGIANI	99B	PRL 83 2902	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)
Also		Private Comm.	G. Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)