

$\chi_{c1}(3872)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

also known as $X(3872)$

This state shows properties different from a conventional $q\bar{q}$ state. A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^-J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^-J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow \chi_{c1}(3872)K^+$ decays, where $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to $< 4\%$ at 95% CL.

See the review on “Spectroscopy of Mesons Containing Two Heavy Quarks.”

$\chi_{c1}(3872)$ T-MATRIX POLE $\sqrt{s}$

Note that $\Gamma = -2\text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$(3871.70 \pm 0.15^{+0.07}_{-0.08}) - i(0.19 \pm 0.08^{+0.14}_{-0.19})$	¹ ABLIKIM	24C BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

¹From simultaneous line shape fits of $e^+e^- \rightarrow \gamma\chi_{c1}(3872) \rightarrow \gamma[D^0\bar{D}^0\pi^0]$ and $\gamma[J/\psi\pi^+\pi^-]$. The most prominent pole is reported at $7.04 \pm 0.15^{+0.07}_{-0.08}$ MeV above the $D^0\bar{D}^0\pi^0$ threshold of 3864.66 MeV on the first sheet with respect to the $D^{*0}\bar{D}^0$ channel. The uncertainty in the D^0 width is included in the uncertainty of the pole mass.

$\chi_{c1}(3872)$ MASS FROM $J/\psi X$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3871.64 ± 0.06	OUR AVERAGE			
$3870.2 \pm 0.7 \pm 0.3$	24.6	ABLIKIM	23W BES3	$e^+e^- \rightarrow J/\psi(1S)\pi^+\pi^-\omega$
$3871.64 \pm 0.06 \pm 0.01$	19.8k	¹ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
$3871.9 \pm 0.7 \pm 0.2$	20	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
$3871.95 \pm 0.48 \pm 0.12$	0.6k	AAIJ	12H LHCb	$pp \rightarrow J/\psi\pi^+\pi^-X$
$3871.85 \pm 0.27 \pm 0.19$	170	² CHOI	11 BELL	$B \rightarrow K\pi^+\pi^-J/\psi$

3873	$+1.8$ -1.6	± 1.3	27	³ DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3871.61	± 0.16	± 0.19	6k	^{3,4} AALTONEN	09AU CDF2	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
3871.4	± 0.6	± 0.1	93.4	AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
3868.7	± 1.5	± 0.4	9.4	AUBERT	08Y BABR	$B^0 \rightarrow K_S^0 J/\psi \pi^+ \pi^-$
3871.8	± 3.1	± 3.0	522	^{3,5} ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
3871.57	± 0.09		155	⁶ AAIJ	23AP LHCb	$B_s^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$
3871.695	± 0.067	± 0.068	15.6k	⁷ AAIJ	20AD LHCb	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
3871.59	± 0.06	± 0.03	4.2k	⁸ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
3873.3	± 1.1	± 1.0	45	⁹ ABLIKIM	19V BES	$e^+ e^- \rightarrow \gamma \omega J/\psi$
3860.0	± 10.4		13.6	^{3,10} AGHASYAN	18A COMP	$\gamma^* N \rightarrow X \pi^\pm N'$
3868.6	± 1.2	± 0.2	8	¹¹ AUBERT	06 BABR	$B^0 \rightarrow K_S^0 J/\psi \pi^+ \pi^-$
3871.3	± 0.6	± 0.1	61	¹¹ AUBERT	06 BABR	$B^- \rightarrow K^- J/\psi \pi^+ \pi^-$
3873.4	± 1.4		25	¹² AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
3871.3	± 0.7	± 0.4	730	^{3,13} ACOSTA	04 CDF2	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
3872.0	± 0.6	± 0.5	36	¹⁴ CHOI	03 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
3836	± 13		58	^{3,15} ANTONIAZZI	94 E705	$300 \pi^\pm \text{Li} \rightarrow J/\psi \pi^+ \pi^- X$

¹ Calculated from $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.54 \pm 0.06$ MeV obtained by combining the data with $\chi_{c1}(3872)$ produced in B^+ decays from AAIJ 20S and inclusive b -hadron decays from AAIJ 20AD and using $m_{\psi(2S)} = 3686.097$ MeV. Breit-Wigner parametrization.

² The mass difference for the $\chi_{c1}(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.

³ Width consistent with detector resolution.

⁴ A possible equal mixture of two states with a mass difference greater than 3.6 MeV/ c^2 is excluded at 95% CL.

⁵ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.

⁶ From a fit of a relativistic S -wave Breit-Wigner convolved with the detector resolution. The width of $\chi_{c1}(3872)$ is constrained to the PDG 22 value. Systematic errors not evaluated.

⁷ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays and $m_{\psi(2S)} = 3686.097 \pm 0.010$ MeV. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.

⁸ Using Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.

⁹ Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.

¹⁰ Could be a different state.

¹¹ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.

¹² Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96$ MeV. Superseded by AUBERT 06.

¹³ Superseded by AALTONEN 09AU.

¹⁴ Superseded by CHOI 11.

¹⁵ A lower mass value can be due to an incorrect momentum scale for soft pions.

$\chi_{c1}(3872)$ MASS FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$3873.71^{+0.56}_{-0.50} \pm 0.13$		¹ HIRATA	23 BELL	$B^0 \rightarrow D^0 \bar{D}^{*0} K^0$, $B^+ \rightarrow D^0 \bar{D}^{*0} K^+$
$3872.9^{+0.6}_{-0.4} \pm 0.4$	50	^{2,3} AUSHEV	10 BELL	$B \rightarrow \bar{D}^{*0} D^0 K$
$3875.1^{+0.7}_{-0.5} \pm 0.5$	33 ± 6	³ AUBERT	08B BABR	$B \rightarrow \bar{D}^{*0} D^0 K$
3875.2 ± 0.7	24 ± 6	^{3,4} GOKHROO	06 BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$

¹ From a fit of a Breit-Wigner function with energy dependent width.² Calculated from the measured $m_{\chi_{c1}(3872)} - m_{D^{*0}} - m_{\bar{D}^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$ MeV.³ Experiments report $D^{*0} \bar{D}^0$ invariant mass above $D^{*0} \bar{D}^0$ threshold because D^{*0} decay products are kinematically constrained to the D^{*0} mass, even though the D^{*0} may decay off-shell.⁴ Superseded by AUSHEV 10. **$m_{\chi_{c1}(3872)} - m_{J/\psi}$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$774.9 \pm 3.1 \pm 3.0$	522	ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$

 $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$185.598 \pm 0.067 \pm 0.068$	15.6k	¹ AAIJ	20AD LHCB	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
185.54 ± 0.06	19.8k	² AAIJ	20S LHCB	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
187.4 ± 1.4	25	³ AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$

¹ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.² Combining $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.49 \pm 0.06 \pm 0.03$ MeV from AAIJ 20S and the measured mass difference from AAIJ 20AD. Breit-Wigner parametrization.³ Superseded by AUBERT 06. **$\chi_{c1}(3872)$ WIDTH**

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.21 OUR AVERAGE			Error includes scale factor of 1.1.		
$1.39 \pm 0.24 \pm 0.10$		15.6k	¹ AAIJ	20AD LHCB	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
$0.96^{+0.19}_{-0.18} \pm 0.21$		4.2k	² AAIJ	20S LHCB	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<2.4	90		ABLIKIM	14 BES3	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
<1.2	90		CHOI	11 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
<3.3	90		AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
<4.1	90	69	AUBERT	06 BABR	$B \rightarrow K \pi^+ \pi^- J/\psi$
<2.3	90	36	³ CHOI	03 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$

¹Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization.

²Using Breit-Wigner parametrization. Partially overlapping dataset with that of AAIJ 20AD.

³Superseded by CHOI 11.

$\chi_{c1}(3872)$ WIDTH FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5.2^{+2.2}_{-1.5} \pm 0.4$		¹ HIRATA	23	BELL $B^0 \rightarrow D^0 \bar{D}^{*0} K^0$, $B^+ \rightarrow D^0 \bar{D}^{*0} K^+$
$3.9^{+2.8+0.2}_{-1.4-1.1}$	50	² AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
$3.0^{+1.9}_{-1.4} \pm 0.9$	33 ± 6	AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$

¹From a fit of a Breit-Wigner function with energy dependent width.

²With a measured value of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow D^{*0} \bar{D}^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.

$\chi_{c1}(3872)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $e^+ e^-$	$< 2.7 \times 10^{-7}$	90%
Γ_2 $\pi^+ \pi^- \pi^0$	< 1.0 %	90%
Γ_3 $\pi^+ \pi^- J/\psi(1S)$	(4.3 ± 1.4) %	
Γ_4 $\pi^+ \pi^- \pi^0 J/\psi(1S)$	not seen	
Γ_5 $\omega \eta_c(1S)$	< 40 %	90%
Γ_6 $\rho(770)^0 J/\psi(1S)$	(3.4 ± 1.1) %	
Γ_7 $\omega J/\psi(1S)$	(5.0 ± 1.9) %	
Γ_8 $\phi \phi$	not seen	
Γ_9 $D^0 \bar{D}^0 \pi^0$	(55 ± 28) %	
Γ_{10} $\bar{D}^{*0} D^0$	(46 ± 16) %	
Γ_{11} $\gamma \gamma$	< 13 %	90%
Γ_{12} $D^0 \bar{D}^0$	< 32 %	90%
Γ_{13} $D^+ D^-$	< 22 %	90%
Γ_{14} $\pi^0 \chi_{c2}$	< 5 %	90%
Γ_{15} $\pi^0 \chi_{c1}$	$(3.8^{+1.9}_{-1.7})$ %	
Γ_{16} $\pi^0 \chi_{c0}$	< 16 %	90%
Γ_{17} $\pi^+ \pi^- \eta_c(1S)$	< 16 %	90%
Γ_{18} $\pi^0 \pi^0 \chi_{c0}$	< 7 %	90%
Γ_{19} $\pi^0 \pi^0 \chi_{c1}$	< 5 %	90%
Γ_{20} $\pi^0 \pi^0 \chi_{c2}$	< 2.2 %	90%
Γ_{21} $\pi^+ \pi^- \chi_{c0}$	< 2.4 %	90%
Γ_{22} $\pi^+ \pi^- \chi_{c1}$	$< 8 \times 10^{-3}$	90%
Γ_{23} $\rho \bar{\rho}$	$< 2.7 \times 10^{-5}$	95%
Γ_{24} $\pi^+ \pi^- \eta$	$< 5 \times 10^{-3}$	90%

Radiative decays

Γ_{25}	$\gamma D^+ D^-$	< 4	%	90%
Γ_{26}	$\gamma \bar{D}^0 D^0$	< 7	%	90%
Γ_{27}	$\gamma J/\psi$	$(10 \pm 4) \times 10^{-3}$		
Γ_{28}	$\gamma \chi_{c1}$	< 1.0	%	90%
Γ_{29}	$\gamma \chi_{c2}$	< 4	%	90%
Γ_{30}	$\gamma \psi(2S)$	possibly seen		
Γ_{31}	$\gamma \psi_2(3823)$	< 3.3	$\times 10^{-3}$	90%

C-violating decays

Γ_{32}	$\eta J/\psi$	< 2.1	%	90%
---------------	---------------	---------	---	-----

 $\chi_{c1}(3872)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$					Γ_1
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.32	90	¹ ABLIKIM	230 BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 4.3	90	² ABLIKIM	15v BES3	$4.0\text{--}4.4 e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
< 280	90	³ YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
¹ Fit to cross section using a total width value of 1.19 ± 0.21 MeV and $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = (3.8 \pm 1.2)\%$ from PDG 20.					
² ABLIKIM 15v reports this limit from the measurement of $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-)/\Gamma < 0.13$ eV using $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma = 3\%$.					
³ Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^- \ell^+ \ell^-$. Assuming that $\Gamma(\pi^+\pi^- J/\psi)$ of $\chi_{c1}(3872)$ is the same as that of $\psi(2S)$ (85.4 keV).					

 $\chi_{c1}(3872) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_3 \Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
$< 7.5 \times 10^{-3}$	90	¹ ABLIKIM	230 BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.13	90	ABLIKIM	15v BES3	$4.0\text{--}4.4 e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
< 6.2	90	^{2,3} AUBERT	05D BABR	$10.6 e^+e^- \rightarrow K^+K^- \pi^+\pi^- \gamma$	
< 8.3	90	³ DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
< 10	90	⁴ YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
¹ Fit to cross section using a total width value of 1.19 ± 0.21 MeV from PDG 20.					
² Using $B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+\pi^-) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.					
³ Assuming $\chi_{c1}(3872)$ has $J^{PC} = 1^{--}$.					
⁴ Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^- \ell^+ \ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10)\%$.					

$\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ $\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_3\Gamma_{11}/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.5^{+4.1}_{-3.8} \pm 0.7$	3	¹ TERAMOTO	21	BELL	$e^+e^- \rightarrow \gamma^*\gamma$ at $\Upsilon(\text{nS})$
<12.9	90	² DOBBS	05	CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi\gamma$

¹ Measured in single-tag two-photon production assuming Q^2 dependence of a $c\bar{c}$ meson model. Here, $\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)$ is the reduced two-photon decay width, $\tilde{\Gamma}_{\gamma\gamma}$.

² Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.

 $\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_7\Gamma_{11}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7	90	¹ LEES	12AD BABR	$e^+e^- \rightarrow e^+e^- \omega J/\psi$
--------	----	-------------------	-----------	---

¹ Assuming $\chi_{c1}(3872)$ has spin 2.

 $\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_{11}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<11.1	90	LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$
---------	----	------	-----------	---

 $\chi_{c1}(3872)$ BRANCHING RATIOS $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

<1.0	90	^{1,2} YIN	23	BELL $B^+ \rightarrow \chi_{c1}(3872) K^+$
--------	----	--------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3	90	^{2,3} YIN	23	BELL $B^0 \rightarrow \chi_{c1}(3872) K^0$
--------	----	--------------------	----	--

¹ YIN 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

² Assuming the decay products, $\pi^+\pi^-\pi^0$, are uniformly distributed in phase space. The limit is the 90% "credible" upper limit (i.e. Bayesian).

³ YIN 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \chi_{c1}(3872) K^0)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^0 \rightarrow \chi_{c1}(3872) K^0) = 1.1 \times 10^{-4}$.

 $\Gamma(\pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.043±0.014 OUR AVERAGE

$0.043 \pm 0.002 \pm 0.013$		¹ AAIJ	20S	LHCB $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
-----------------------------	--	-------------------	-----	---

$0.047 \pm 0.005 \pm 0.015$		² CHOI	11	BELL $B^+ \rightarrow \pi^+ \pi^- J/\psi K^+$
-----------------------------	--	-------------------	----	---

$0.045 \pm 0.009 \pm 0.014$	93	^{3,4} AUBERT	08Y	BABR $B \rightarrow \chi_{c1}(3872) K$
-----------------------------	----	-----------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	151	⁵ BALA	15	BELL $B \rightarrow \chi_{c1}(3872) K \pi$
------	-----	-------------------	----	--

$0.069 \pm 0.022 \pm 0.022$	30	⁶ AUBERT	05R	BABR $B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$
-----------------------------	----	---------------------	-----	---

$0.074 \pm 0.016 \pm 0.023$	36	⁷ CHOI	03	BELL $B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$
-----------------------------	----	-------------------	----	---

¹AAIJ 20S reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (7.95 \pm 0.15 \pm 0.33) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²CHOI 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.63 \pm 0.82 \pm 0.52) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³AUBERT 08Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴superseded by LEES 20C

⁵BALA 15 reports $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872)K^+\pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872)K^0\pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.

⁶Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁷CHOI 03 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] / [B(B^+ \rightarrow \psi(2S)K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S)K^+) = (6.24 \pm 0.21) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\pi^+\pi^-\pi^0 J/\psi(1S))/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
not seen	² SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

¹WANG 11B reports $B(\Upsilon(2S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi) < 2.4 \times 10^{-6}$ at 95% CL.

²SHEN 10A reports $B(\Upsilon(1S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi) < 2.8 \times 10^{-6}$ at 95% CL.

$\Gamma(\omega\eta_c(1S))/\Gamma_{\text{total}}$

Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	¹ VINOKUROVA 15	BELL	$B^+ \rightarrow \omega\eta_c K^+$

¹VINOKUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega\eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6.9 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.

$\Gamma(\rho(770)^0 J/\psi(1S))/\Gamma(\pi^+\pi^- J/\psi(1S))$

Γ_6/Γ_3

VALUE (%)	DOCUMENT ID	TECN	COMMENT
78.6$\pm$2.3$\pm$2.0	¹ AAIJ 23S	LHCB	$B^+ \rightarrow K^+ J/\psi \pi^+\pi^-$

¹Assuming pure ρ contribution only, i.e. excluding the contribution from ρ - ω interference. Using $B(\rho^0 \rightarrow \pi^+\pi^-) = 100\%$.

$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.032 \pm 0.012 \pm 0.010$ 21 ± 7 ¹ DEL-AMO-SA..10B BABR $B^+ \rightarrow \omega J/\psi K^+$

¹ DEL-AMO-SANCHEZ 10B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow \chi_{c1}(3872) K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi \omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.

 $\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_7/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.16 $\pm$ 0.24 OUR AVERAGE Error includes scale factor of 1.2.

$1.24 \pm 0.33 \pm 0.10$ ^{1,2} AAIJ 23S LHCB $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$

$1.6^{+0.4}_{-0.3} \pm 0.2$ ³ ABLIKIM 19v BES $e^+ e^- \rightarrow \gamma \omega J/\psi$

0.8 ± 0.3 ⁴ DEL-AMO-SA..10B BABR $B \rightarrow \omega J/\psi K$

¹ AAIJ 23S reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] \times [B(\omega(782) \rightarrow \pi^+ \pi^-)] = (1.9 \pm 0.4 \pm 0.3) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^-) = (1.53 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Excluding ρ - ω interference effects.

³ Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.

⁴ Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen ¹ AAIJ 17BB LHCB pp at 7, 8 TeV

¹ AAIJ 17BB reports $B(b \rightarrow \chi_{c1}(3872) \text{ anything}) \times B(\chi_{c1}(3872) \rightarrow \phi\phi) < 4.5 \times 10^{-7}$ at 95% CL.

 $\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

$0.55^{+0.20}_{-0.23} \pm 0.17$ 17 ¹ GOKHROO 06 BELL $B^+ \rightarrow D^0 \bar{D}^0 \pi^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.32 90 ² CHISTOV 04 BELL Sup. by GOKHROO 06

¹ GOKHROO 06 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 0.6 \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

$\Gamma(D^0 \bar{D}^0 \pi^0) / \Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_9 / Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.16	90	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 $\Gamma(\bar{D}^{*0} D^0) / \Gamma_{\text{total}}$ Γ_{10} / Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.46 ± 0.16 OUR AVERAGE				

$0.52^{+0.13}_{-0.11} \pm 0.16$	$1,2$	HIRATA	23	BELL	$e^+ e^- \rightarrow \gamma(4S)$
$0.42 \pm 0.10 \pm 0.13$	41^{+9}_{-8}	3 AUSHEV	10	BELL	$B^+ \rightarrow D^{*0} \bar{D}^0 K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.90 \pm 0.32 \pm 0.28$	27 ± 6	4 AUBERT	08B	BABR	$B^+ \rightarrow \bar{D}^{*0} D^0 K^+$

1 HIRATA 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.97^{+0.21}_{-0.18} \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 Assumes equal production of B^+ and B^0 at the $\gamma(4S)$.

3 AUSHEV 10 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

4 AUBERT 08B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\bar{D}^{*0} D^0) / \Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{10} / Γ_3

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
11.77 ± 3.09	50	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

 $\Gamma(\gamma\gamma) / \Gamma_{\text{total}}$ Γ_{11} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.13	90	¹ WICHT	08	BELL	$e^+ e^- \rightarrow \gamma(4S)$
¹ WICHT 08 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 2.4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$					

 $\Gamma(D^0 \bar{D}^0) / \Gamma_{\text{total}}$ Γ_{12} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.32	90	¹ CHISTOV	04	BELL	$B \rightarrow K D^0 \bar{D}^0$
¹ CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)]$ $< 6 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$					

 $\Gamma(D^+ D^-) / \Gamma_{\text{total}}$ Γ_{13} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	¹ CHISTOV	04	BELL
$B \rightarrow K D^+ D^-$				
¹ CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^+ D^-)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)]$ $< 4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$				

$$\Gamma(\pi^0 \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S)) \quad \Gamma_{14}/\Gamma_3$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	ABLIKIM	19U	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$$\Gamma(\pi^0 \chi_{c1})/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04 90 ¹ BHARDWAJ 19 BELL $B^\pm \rightarrow \pi^0 \chi_{c1} K^\pm$
¹ BHARDWAJ 19 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^0 \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 8.1 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

$$\Gamma(\pi^0 \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S)) \quad \Gamma_{15}/\Gamma_3$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$88^{+33}_{-27} \pm 10$	10.8	ABLIKIM	19U	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$$\Gamma(\pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S)) \quad \Gamma_{16}/\Gamma_3$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3.6	90	ABLIKIM	22D	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19 90 ABLIKIM 19U BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	¹ VINOKUROVA 15	BELL	$B^+ \rightarrow \pi^+ \pi^- \eta_c K^+$

¹ VINOKUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 3.0 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

$$\Gamma(\pi^0 \pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S)) \quad \Gamma_{18}/\Gamma_3$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	ABLIKIM	22D	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$$\Gamma(\pi^0 \pi^0 \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S)) \quad \Gamma_{19}/\Gamma_3$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	ABLIKIM	24BU	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$$\Gamma(\pi^0 \pi^0 \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S)) \quad \Gamma_{20}/\Gamma_3$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	ABLIKIM	24BU	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$$\Gamma(\pi^+ \pi^- \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S)) \quad \Gamma_{21}/\Gamma_3$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.56	90	ABLIKIM	22D	BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$\Gamma(\pi^+\pi^-\chi_{c1})/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-3}$	90	¹ BHARDWAJ 16	BELL	$B^+ \rightarrow \pi^+\pi^-\chi_{c1}K^+$
¹ BHARDWAJ 16 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.				

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7 \times 10^{-5}$	95	¹ AAIJ 17AD	LHCB	$B^+ \rightarrow p\bar{p}K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<9 \times 10^{-5}$	95	² AAIJ 13S	LHCB	$B^+ \rightarrow p\bar{p}K^+$
¹ AAIJ 17AD reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 0.5 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.				
² AAIJ 13S reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.				

 $\Gamma(\pi^+\pi^-\eta)/\Gamma(\pi^+\pi^-J/\psi(1S))$ Γ_{24}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.12	90	ABLIKIM 24K	BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

———— Radiative decays ————

 $\Gamma(\gamma D^+D^-)/\Gamma(\pi^+\pi^-J/\psi(1S))$ Γ_{25}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.99	90	ABLIKIM 20W	BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

 $\Gamma(\gamma \bar{D}^0D^0)/\Gamma(\pi^+\pi^-J/\psi(1S))$ Γ_{26}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.58	90	ABLIKIM 20W	BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

 $\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.6^{+2.7}_{-2.5} \pm 3.0$		¹ BHARDWAJ 11	BELL	$B^\pm \rightarrow \gamma J/\psi K^\pm$

• • • We do not use the following data for averages, fits, limits, etc. • • •

15 $\pm 4 \pm 5$	20	² AUBERT 09B	BABR	$B^+ \rightarrow \gamma J/\psi K^+$
18 $\pm 6 \pm 6$	19	³ AUBERT,BE 06M	BABR	$B^+ \rightarrow \gamma J/\psi K^+$
¹ BHARDWAJ 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

³ Superseded by AUBERT 09B. AUBERT, BE 06M reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma J/\psi)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

Γ_{27}/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.52±0.19 OUR AVERAGE	Error includes scale factor of 1.2.			
0.38±0.20±0.01	8 ± 4	ABLIKIM	24X BES3	$e^+ e^- \rightarrow \omega \chi_{c1}(3872)$
0.79±0.28		ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$

Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	¹ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1} \gamma K^\pm$

¹ BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

$\Gamma(\gamma \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

Γ_{28}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.89	90	CHOI	03 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$

$\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$

Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	¹ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2} \gamma K^\pm$

¹ BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \chi_{c2})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 6.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 1.9 \times 10^{-4}$.

$\Gamma(\gamma \psi(2S))/\Gamma_{\text{total}}$

Γ_{30}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
possibly seen	36 ± 9	¹ AAIJ	14AH LHCb	$B^+ \rightarrow \gamma \psi(2S) K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen		² BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma \psi(2S) K^+$
0.051±0.015±0.016	25 ± 7	³ AUBERT	09B BABR	$B^+ \rightarrow \gamma \psi(2S) K^+$

¹ From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi \gamma$ decays with a statistical significance of 4.4σ .

² BHARDWAJ 11 reports $B(B^+ \rightarrow K^+ \chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma \psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.

³ AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (1.9 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma \psi(2S))/\Gamma(\pi^+ \pi^- J/\psi(1S))$

Γ_{30}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42	90	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$\Gamma(\pi^+ \pi^- \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

Γ_{22}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.18	90	ABLIKIM	24S BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$			Γ_{30}/Γ_{27}		
VALUE	CL%	EVS	DOCUMENT ID	TECN	COMMENT
$1.67 \pm 0.21 \pm 0.13$			¹ AAIJ	24AD LHCb	$B^+ \rightarrow \gamma\psi(2S)K^+$ and $\gamma J/\psi K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.59	90	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$
$2.46 \pm 0.64 \pm 0.29$	36 ± 9	² AAIJ	14AH LHCb	$B^+ \rightarrow \gamma\psi(2S)K^+$
<2.1	90	BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$
3.4 ± 1.4		AUBERT	09B BABR	$B^+ \rightarrow \gamma c\bar{c}K'$

¹ AAIJ 24AD reports this ratio as $1.67 \pm 0.21 \pm 0.12 \pm 0.04$ where the last uncertainty is due to the uncertainties of the branching fractions of $\psi(2S)$ and J/ψ mesons. We have added the last two uncertainties in quadrature.

² From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ . Superseded by AAIJ 24AD.

$\Gamma(\gamma\psi_2(3823))/\Gamma(\pi^+\pi^- J/\psi(1S))$					Γ_{31}/Γ_3
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.075	90	¹ ABLIKIM	24Z	BES3	$e^+e^- \rightarrow \gamma\psi_2(3823)$

¹ Using $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = 0.343 \pm 0.010$. We have assumed that $B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = 1$.

C-violating decays

$\Gamma(\eta J/\psi)/\Gamma_{\text{total}}$					Γ_{32}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.021	90	^{1,2} IWASHITA	14	BELL	$B \rightarrow K \eta J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	90	³ AUBERT	04Y BABR	$B \rightarrow K\eta J/\psi$
-------	----	---------------------	----------	------------------------------

¹ IWASHITA 14 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.

² IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals.

³ AUBERT 04Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 7.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 1.9 \times 10^{-4}$.

$\chi_{c1}(3872)$ REFERENCES

AAIJ	24AD	JHEP 2411 121	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	24BU	PR D110 072015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24C	PRL 132 151903	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24K	PR D109 L011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24S	PR D109 L071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24X	PR D110 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24Z	PR D110 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	23AP	JHEP 2307 084	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	23S	PR D108 L011103	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23O	PR D107 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23W	PRL 130 151904	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HIRATA	23	PR D107 112011	H. Hirata <i>et al.</i>	(BELLE Collab.)
YIN	23	PR D107 052004	J.H. Yin <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22D	PR D105 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
TERAMOTO	21	PRL 126 122001	Y. Teramoto <i>et al.</i>	(BELLE Collab.)

AAIJ	20AD	PR D102 092005	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	20W	PRL 124 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	20C	PRL 124 152001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19U	PRL 122 202001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AGHASYAN	18A	PL B783 334	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also		JHEP 1702 088 (errat.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)
AAIJ	13Q	PRL 110 222001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AALTONEN	09AU	PRL 103 152001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>	
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)