

$\Omega(2012)^-$ $I(J^P) = 0(?^-)$ Status: ***

Seen in $\Xi^0 K^-$ and $\Xi^- K_S^0$ decays with a combined significance of 8.3 standard deviations.

 $\Omega(2012)^-$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2012.5±0.6 OUR AVERAGE				
2012.5±0.7±0.5	844	JIA	25	BELL In $\gamma(1S, 2S, 3S)$
2012.4±0.7±0.6	520	YELTON	18A	BELL In $\gamma(1S, 2S, 3S)$

 $\Omega(2012)^-$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.4^{+2.5}_{-2.0}±1.6	520	YELTON	18A	BELL In $\gamma(1S), \gamma(2S), \gamma(3S)$

 $\Omega(2012)^-$ DECAY MODES

Branching fractions are given relative to the one **DEFINED AS 1**.

Mode	Fraction (Γ_i/Γ)
Γ_1 ΞK	
Γ_2 $(\Xi\pi)K$	
Γ_3 $\Xi^0 K^-$	DEFINED AS 1
Γ_4 $\Xi^- \bar{K}^0$	0.83 ± 0.21
Γ_5 $\Xi^0 \pi^0 K^-$	
Γ_6 $\Xi^0 \pi^- \bar{K}^0$	
Γ_7 $\Xi^- \pi^0 \bar{K}^0$	
Γ_8 $\Xi^- \pi^+ K^-$	

 $\Omega(2012)^-$ BRANCHING RATIOS

$\Gamma((\Xi\pi)K)/\Gamma(\Xi K)$	Γ_2/Γ_1
0.99 ±0.26±0.06	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<0.119	90 JIA 19 BELL In $\gamma(1S, 2S, 3S)$

$\Gamma(\Xi^0 K^-)/\Gamma(\Xi^- \bar{K}^0)$	Γ_3/Γ_4
1.2±0.3	

$\Gamma(\Xi^0 \pi^0 K^-)/\Gamma(\Xi^0 K^-)$				Γ_5/Γ_3
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.304	90	JIA	19	BELL In $\gamma(1S, 2S, 3S)$
$\Gamma(\Xi^0 \pi^- \bar{K}^0)/\Gamma(\Xi^0 K^-)$				Γ_6/Γ_3
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.213	90	JIA	19	BELL In $\gamma(1S, 2S, 3S)$
$\Gamma(\Xi^0 \pi^- \bar{K}^0)/\Gamma(\Xi^- \bar{K}^0)$				Γ_6/Γ_4
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.256	90	JIA	19	BELL In $\gamma(1S, 2S, 3S)$
$\Gamma(\Xi^- \pi^0 \bar{K}^0)/\Gamma(\Xi^- \bar{K}^0)$				Γ_7/Γ_4
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.811	90	JIA	19	BELL In $\gamma(1S, 2S, 3S)$
$\Gamma(\Xi^- \pi^+ K^-)/\Gamma(\Xi^0 K^-)$				Γ_8/Γ_3
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.078	90	JIA	19	BELL In $\gamma(1S, 2S, 3S)$
$\Gamma(\Xi^- \pi^+ K^-)/\Gamma(\Xi^- \bar{K}^0)$				Γ_8/Γ_4
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.093	90	JIA	19	BELL In $\gamma(1S, 2S, 3S)$

$\Omega(2012)^-$ REFERENCES

JIA	25	PL B860 139224	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	19	PR D100 032006	S. Jia <i>et al.</i>	(BELLE Collab.)
YELTON	18A	PRL 121 052003	J. Yelton <i>et al.</i>	(BELLE Collab.)