

$I^G(J^{PC}) = 0^-(1^{--})$

$\phi(1020)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1019.460 ±0.016 OUR AVERAGE				
1019.443 ±0.010 ±0.060 1.28M	1	ACHASOV 24	SND	$e^+e^- \rightarrow K_L^0 K_S^0$
1019.463 ±0.061 2.3M	2	KOZYREV 18	CMD3	$e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
1019.462 ±0.042 ±0.056 28k	3	LEES 14H	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1019.51 ±0.02 ±0.05	4	LEES 13Q	BABR	$e^+e^- \rightarrow K^+ K^- \gamma$
1019.30 ±0.02 ±0.10 105k		AKHMETSHIN 06	CMD2	$0.98-1.06 e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.52 ±0.05 ±0.05 17.4k		AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta \gamma$
1019.483 ±0.011 ±0.025 272k	5	AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
1019.42 ±0.05 1900k	6	ACHASOV 01E	SND	$e^+e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
1019.40 ±0.04 ±0.05 23k		AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta \gamma$
1019.36 ±0.12	7	ACHASOV 00B	SND	$e^+e^- \rightarrow \eta \gamma$
1019.38 ±0.07 ±0.08 2200	8	AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \pi^+ \pi^- \geq 2\gamma$
1019.51 ±0.07 ±0.10 11169		AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.5 ±0.4		BARBERIS 98	OMEG 450	$pp \rightarrow pp2K^+2K^-$
1019.42 ±0.06 55600		AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \text{hadrons}$
1019.7 ±0.3 2012		DAVENPORT 86	MPSF 400	$pA \rightarrow 4K X$
1019.7 ±0.1 ±0.1 5079		ALBRECHT 85D	ARG 10	$e^+e^- \rightarrow K^+ K^- X$
1019.3 ±0.1 1500		ARENTON 82	AEMS 11.8	polar. $pp \rightarrow KK$
1019.67 ±0.17 25080	9	PELLINEN 82	RVUE	
1019.52 ±0.13 3681		BUKIN 78C	OLYA	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1019.0330±0.0084	10	DUBNICKA 24	RVUE	$e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
1019.761 ±0.130 34M	11	IGNATOV 24	CMD3	$e^+e^- \rightarrow \pi^+ \pi^-$
1018.4 ±0.5 ±0.1	12	ALBRECHT 20	CBAR 0.9	$\bar{p}p \rightarrow K^+ K^- \pi^0$
1019.21 ±0.04 ±0.03	13	HOID 20	RVUE	$e^+e^- \rightarrow \pi^0 \gamma$
1019.54 ±0.10 ±0.51	14	AAIJ 19H	LHCB	$pp \rightarrow D^\pm X$
1019.20 ±0.02 ±0.01	15	HOFERICHT... 19	RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
1019.469 ±0.061 1.7M		KOZYREV 18	CMD3	$e^+e^- \rightarrow K^+ K^-$
1019.457 ±0.061 610k		KOZYREV 16	CMD3	$e^+e^- \rightarrow K_S^0 K_L^0$
1019.48 ±0.01		LEES 13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$
1019.441 ±0.008 ±0.080 542k	16	AKHMETSHIN 08	CMD2	$1.02 e^+e^- \rightarrow K^+ K^-$

1019.63	± 0.07		12540	¹⁷ AUBERT,B	05J	BABR	$D^0 \rightarrow \overline{K}^0 K^+ K^-$
1019.8	± 0.7			ARMSTRONG	86	OMEG	$85 \pi^+ / p p \rightarrow$ $\pi^+ / p 4 K p$
1020.1	± 0.11		5526	¹⁷ ATKINSON	86	OMEG	$20-70 \gamma p$
1019.7	± 1.0			BEBEK	86	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
1019.411	± 0.008		642k	¹⁸ DIJKSTRA	86	SPEC	$100-200 \pi^\pm, \overline{p}, p,$ $K^\pm, \text{ on Be}$
1020.9	± 0.2			¹⁷ FRAME	86	OMEG	$13 K^+ p \rightarrow \phi K^+ p$
1021.0	± 0.2			¹⁷ ARMSTRONG	83B	OMEG	$18.5 K^- p \rightarrow$ $K^- K^+ \Lambda$
1020.0	± 0.5			¹⁷ ARMSTRONG	83B	OMEG	$18.5 K^- p \rightarrow$ $K^- K^+ \Lambda$
1019.7	± 0.3			¹⁷ BARATE	83	GOLI	$190 \pi^- \text{Be} \rightarrow 2\mu X$
1019.8	± 0.2	± 0.5	766	IVANOV	81	OLYA	$1-1.4 e^+ e^- \rightarrow$ $K^+ K^-$
1019.4	± 0.5		337	COOPER	78B	HBC	$0.7-0.8 \overline{p} p \rightarrow$ $K_S^0 K_L^0 \pi^+ \pi^-$
1020	± 1		383	¹⁷ BALDI	77	CNTR	$10 \pi^- p \rightarrow \pi^- \phi p$
1018.9	± 0.6		800	COHEN	77	ASPK	$6 \pi^\pm N \rightarrow$ $K^+ K^- N$
1019.7	± 0.5		454	KALBFLEISCH	76	HBC	$2.18 K^- p \rightarrow$ $\Lambda K \overline{K}$
1019.4	± 0.8		984	BESCH	74	CNTR	$2 \gamma p \rightarrow p K^+ K^-$
1020.3	± 0.4		100	BALLAM	73	HBC	$2.8-9.3 \gamma p$
1019.4	± 0.7			BINNIE	73B	CNTR	$\pi^- p \rightarrow \phi n$
1019.6	± 0.5		120	¹⁹ AGUILAR-...	72B	HBC	$3.9, 4.6 K^- p \rightarrow$ $\Lambda K^+ K^-$
1019.9	± 0.5		100	¹⁹ AGUILAR-...	72B	HBC	$3.9, 4.6 K^- p \rightarrow$ $K^- p K^+ K^-$
1020.4	± 0.5		131	COLLEY	72	HBC	$10 K^+ p \rightarrow K^+ p \phi$
1019.9	± 0.3		410	STOTTLE...	71	HBC	$2.9 K^- p \rightarrow$ $\Sigma / \Lambda K \overline{K}$

¹ From a fit of the cross section in the energy region $1.000 < \sqrt{s} < 1.100$ GeV using a vector meson dominance model with contribution from $\phi(1020)$, $\rho(770)$, $\omega(782)$ and higher vector resonances. The the first three points near the threshold are removed in the fit.

² Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.

³ Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

⁴ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.

⁵ Update of AKHMETSHIN 99D

⁶ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.

⁷ Using a total width of 4.43 ± 0.05 MeV. Systematic uncertainty included.

⁸ Using a total width of 4.43 ± 0.05 MeV.

⁹ PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DE-GROOT 74.

¹⁰ From the fit to data of CMD3, BABAR and BES3.

- ¹¹ From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit. Systematic errors not estimated.
- ¹² Width fixed at 4.2 MeV.
- ¹³ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 1019.457 ± 0.020 MeV.
- ¹⁴ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUE 18.
- ¹⁵ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.
- ¹⁶ Strongly correlated with AKHMETSHIN 04.
- ¹⁷ Systematic errors not evaluated.
- ¹⁸ Weighted and scaled average of 12 measurements of DIJKSTRA 86.
- ¹⁹ Mass errors enlarged by us to $\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

$\phi(1020)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.249 ± 0.013 OUR AVERAGE		Error includes scale factor of 1.1.		
$4.212 \pm 0.20 \pm 0.13$	1.28M	¹ ACHASOV	24	SND $e^+e^- \rightarrow K_L^0 K_S^0$
4.245 ± 0.013	2.3M	² KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
$4.205 \pm 0.103 \pm 0.067$	28k	³ LEES	14H	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
$4.29 \pm 0.04 \pm 0.07$		⁴ LEES	13Q	BABR $e^+e^- \rightarrow K^+ K^- \gamma$
$4.30 \pm 0.06 \pm 0.17$	105k	AKHMETSHIN 06	CMD2	$0.98-1.06 e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
$4.280 \pm 0.033 \pm 0.025$	272k	⁵ AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
4.21 ± 0.04	1900k	⁶ ACHASOV	01E	SND $e^+e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
4.44 ± 0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow$ hadrons
4.5 ± 0.7	1500	ARENTON	82	AEMS 11.8 polar. $pp \rightarrow KK$
4.2 ± 0.6	766	⁷ IVANOV	81	OLYA $1-1.4 e^+e^- \rightarrow K^+ K^-$
4.3 ± 0.6		⁷ CORDIER	80	DM1 $e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
4.36 ± 0.29	3681	⁷ BUKIN	78C	OLYA $e^+e^- \rightarrow$ hadrons
4.4 ± 0.6	984	⁷ BESCH	74	CNTR $2 \gamma p \rightarrow p K^+ K^-$
4.67 ± 0.72	681	⁷ BALAKIN	71	OSPK $e^+e^- \rightarrow$ hadrons
4.09 ± 0.29		BIZOT	70	OSPK $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.151 ± 0.018		⁸ DUBNICKA	24	RVUE $e^+e^- \rightarrow K^+ K^-, K_S^0 K_L^0$
4.681 ± 0.277	34M	⁹ IGNATOV	24	CMD3 $e^+e^- \rightarrow \pi^+ \pi^-$
$4.07 \pm 0.13 \pm 0.01$		¹⁰ HOID	20	RVUE $e^+e^- \rightarrow \pi^0 \gamma$
$4.23 \pm 0.04 \pm 0.02$		¹¹ HOFERICH...	19	RVUE $e^+e^- \rightarrow \pi^+ \pi^- \pi^0$
4.249 ± 0.015	1.7M	KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+ K^-$
4.240 ± 0.017	610k	KOZYREV	16	CMD3 $e^+e^- \rightarrow K_S^0 K_L^0$
4.37 ± 0.02		LEES	13F	BABR $D^+ \rightarrow K^+ K^- \pi^+$
$4.24 \pm 0.02 \pm 0.03$	542k	¹² AKHMETSHIN 08	CMD2	$1.02 e^+e^- \rightarrow K^+ K^-$

4.28 ±0.13	12540	¹³ AUBERT,B	05J	BABR	$D^0 \rightarrow \bar{K}^0 K^+ K^-$
4.45 ±0.06	271k	DIJKSTRA	86	SPEC	100 $\pi^- \text{Be}$
3.6 ±0.8	337	⁷ COOPER	78B	HBC	0.7–0.8 $\bar{p}p \rightarrow$ $K_S^0 K_L^0 \pi^+ \pi^-$
4.5 ±0.50	1300	^{7,13} AKERLOF	77	SPEC	400 $pA \rightarrow K^+ K^- X$
4.5 ±0.8	500	^{7,13} AYRES	74	ASPK	3–6 $\pi^- p \rightarrow$ $K^+ K^- n, K^- p \rightarrow$ $K^+ K^- \Lambda/\Sigma^0$
3.81 ±0.37		COSME	74B	OSPK	$e^+ e^- \rightarrow K_L^0 K_S^0$
3.8 ±0.7	454	⁷ BORENSTEIN	72	HBC	2.18 $K^- p \rightarrow K \bar{K} n$

¹ From a fit of the cross section in the energy region $1.000 < \sqrt{s} < 1.100$ GeV using a vector meson dominance model with contribution from $\phi(1020)$, $\rho(770)$, $\omega(782)$ and higher vector resonances. The the first three points near the threshold are removed in the fit.

² Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.

³ Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

⁴ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.

⁵ Update of AKHMETSHIN 99D

⁶ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

⁷ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁸ From the fit to data of CMD3, BABAR and BES3.

⁹ From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit. Systematic errors not estimated.

¹⁰ The values were extracted from a dispersively improved Breit-Wigner parameterization.

¹¹ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.

¹² Strongly correlated with AKHMETSHIN 04.

¹³ Systematic errors not evaluated.

$\phi(1020)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $K^+ K^-$	(49.9 ±0.5) %	S=1.5
Γ_2 $K_L^0 K_S^0$	(33.6 ±0.4) %	S=1.3
Γ_3 $\rho\pi + \pi^+ \pi^- \pi^0$	(14.9 ±0.4) %	S=1.3
Γ_4 $\pi^+ \pi^- \pi^0$		
Γ_5 $\eta\gamma$	(1.306±0.024) %	S=1.2
Γ_6 $\pi^0\gamma$	(1.33 ±0.05) × 10 ⁻³	
Γ_7 $\ell^+ \ell^-$	—	
Γ_8 $e^+ e^-$	(2.964±0.033) × 10 ⁻⁴	S=1.4
Γ_9 $\mu^+ \mu^-$	(2.86 ±0.22) × 10 ⁻⁴	S=1.2
Γ_{10} $\eta e^+ e^-$	(1.08 ±0.04) × 10 ⁻⁴	

Γ_{11}	$\pi^+\pi^-$	$(9.5 \pm 1.9) \times 10^{-5}$	S=2.0
Γ_{12}	$\omega\pi^0$	$(4.7 \pm 0.5) \times 10^{-5}$	
Γ_{13}	$\omega\gamma$	< 5	CL=84%
Γ_{14}	$\rho\gamma$	$< 1.2 \times 10^{-5}$	CL=90%
Γ_{15}	$\pi^+\pi^-\gamma$	$(4.1 \pm 1.3) \times 10^{-5}$	
Γ_{16}	$f_0(980)\gamma$	$(3.22 \pm 0.19) \times 10^{-4}$	S=1.1
Γ_{17}	$\pi^0\pi^0\gamma$	$(1.13 \pm 0.06) \times 10^{-4}$	
Γ_{18}	$\pi^+\pi^-\pi^+\pi^-$	$(3.9^{+2.8}_{-2.2}) \times 10^{-6}$	
Γ_{19}	$\pi^+\pi^+\pi^-\pi^-\pi^0$	$< 4.6 \times 10^{-6}$	CL=90%
Γ_{20}	$\pi^0e^+e^-$	$(1.33^{+0.07}_{-0.10}) \times 10^{-5}$	
Γ_{21}	$\pi^0\eta\gamma$	$(7.27 \pm 0.30) \times 10^{-5}$	S=1.5
Γ_{22}	$a_0(980)\gamma$	$(7.6 \pm 0.6) \times 10^{-5}$	
Γ_{23}	$K^0\bar{K}^0\gamma$	$< 1.9 \times 10^{-8}$	CL=90%
Γ_{24}	$\eta'(958)\gamma$	$(6.23 \pm 0.21) \times 10^{-5}$	
Γ_{25}	$\eta\pi^0\pi^0\gamma$	$< 2 \times 10^{-5}$	CL=90%
Γ_{26}	$\mu^+\mu^-\gamma$	$(1.4 \pm 0.5) \times 10^{-5}$	
Γ_{27}	$\rho\gamma\gamma$	$< 1.2 \times 10^{-4}$	CL=90%
Γ_{28}	$\eta\pi^+\pi^-$	$< 1.8 \times 10^{-5}$	CL=90%
Γ_{29}	$\eta\mu^+\mu^-$	$< 9.4 \times 10^{-6}$	CL=90%
Γ_{30}	$\eta U \rightarrow \eta e^+e^-$	$< 1 \times 10^{-6}$	CL=90%
Γ_{31}	invisible	$< 1.7 \times 10^{-4}$	CL=90%

Lepton Family number (LF) violating modes

Γ_{32}	$e^\pm\mu^\mp$	$LF < 2 \times 10^{-6}$	CL=90%
---------------	----------------	-------------------------	--------

CONSTRAINED FIT INFORMATION

An overall fit to 30 branching ratios uses 83 measurements and one constraint to determine 14 parameters. The overall fit has a $\chi^2 = 89.3$ for 70 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	−74										
x_3	−67	0									
x_5	−26	27	3								
x_6	−15	17	2	8							
x_8	64	−77	−9	−36	−22						
x_9	−6	7	1	3	2	−9					
x_{11}	−4	4	0	2	1	−6	1				
x_{12}	−5	6	1	3	2	−8	1	0			
x_{16}	0	0	0	0	0	0	0	0	0		
x_{17}	−13	15	2	18	4	−19	2	1	2	0	
x_{18}	−1	1	0	1	0	−2	0	0	0	0	
x_{22}	0	0	0	0	0	0	0	0	0	0	
x_{24}	−8	9	1	32	3	−12	1	1	1	1	0
	x_1	x_2	x_3	x_5	x_6	x_8	x_9	x_{11}	x_{12}	x_{16}	

x_{18}	0		
x_{22}	0	0	
x_{24}	6	0	0
	x_{17}	x_{18}	x_{22}

$\phi(1020)$ PARTIAL WIDTHS

$\Gamma(\eta\gamma)$					Γ_5
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$58.9 \pm 0.5 \pm 2.4$	ACHASOV	00	SND $e^+ e^- \rightarrow \eta\gamma$		

$\Gamma(\pi^0\gamma)$					Γ_6
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$5.40 \pm 0.16^{+0.43}_{-0.40}$	ACHASOV	00	SND $e^+ e^- \rightarrow \pi^0\gamma$		

$\Gamma(\ell^+ \ell^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.320 \pm 0.017 \pm 0.015$	¹ AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow \mu^+ \mu^-$
-----------------------------	---------------------------	------	--

¹ Weighted average of Γ_{ee} and $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ from AMBROSINO 05 assuming lepton universality.

$\Gamma(e^+ e^-)$ Γ_8

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1.27 ± 0.04 OUR EVALUATION

1.251 ± 0.021 OUR AVERAGE Error includes scale factor of 1.1.

$1.235 \pm 0.006 \pm 0.022$	¹ AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow \phi$
$1.32 \pm 0.05 \pm 0.03$	² AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow e^+ e^-$
1.28 ± 0.05	AKHMETSHIN 95	CMD2	$1.02 e^+ e^- \rightarrow \phi$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² From forward-backward asymmetry and using $\Gamma_{\text{total}} = 4.26 \pm 0.05$ MeV from the 2004 edition of this Review.

$(\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-))^{1/2}$ $(\Gamma_8 \Gamma_9)^{1/2}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1.320 ± 0.018 ± 0.017	AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow \mu^+ \mu^-$
------------------------------	--------------	------	--

$\phi(1020) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_1 \Gamma_8 / \Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.6340 ± 0.0070 ± 0.0039		¹ LEES	13Q BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
---------------------------------	--	-------------------	----------	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.669 \pm 0.001 \pm 0.023$	1.7M	KOZYREV	18 CMD3	$e^+ e^- \rightarrow K^+ K^-$
-----------------------------	------	---------	---------	-------------------------------

¹ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations. The first error combines statistical and systematic uncertainties. The second one is due to the parametrization of the charged kaon form factor and mass calibration.

$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_2 \Gamma_8 / \Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.4200 ± 0.0033 ± 0.0123	28k	¹ LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$
---------------------------------	-----	-------------------	----------	--

¹ Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)] \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_3 \Gamma_8 / \Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

184.1 ± 2.1 ± 8.0	¹ LEES	21B BABR	$10.5 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
--------------------------	-------------------	----------	---


¹ From the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$ with contributions from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

$\phi(1020) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$

$\Gamma(K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_1 / \Gamma \times \Gamma_8 / \Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
14.79 ± 0.30 OUR FIT		Error includes scale factor of 1.5.		
14.6 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		
15.789 ± 0.541	1.7M	KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-$
14.27 ± 0.05 ± 0.31	542k	AKHMETSHIN	08	CMD2 $1.02 e^+ e^- \rightarrow K^+ K^-$
13.93 ± 0.14 ± 0.99	1000k	¹ ACHASOV	01E	SND $e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.



$\Gamma(K_L^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_2 / \Gamma \times \Gamma_8 / \Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
9.97 ± 0.08 OUR FIT				
9.94 ± 0.08 OUR AVERAGE				
9.85 ± 0.03 ± 0.10	1.28M	¹ ACHASOV	24	SND $e^+ e^- \rightarrow K_L^0 K_S^0$
10.078 ± 0.223	610k	² KOZYREV	16	CMD3 $e^+ e^- \rightarrow K_S^0 K_L^0$
10.01 ± 0.04 ± 0.17	272k	³ AKHMETSHIN	04	CMD2 $e^+ e^- \rightarrow K_L^0 K_S^0$
10.27 ± 0.07 ± 0.34	500k	⁴ ACHASOV	01E	SND $e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$

¹ From a fit of the cross section in the energy region $1.000 < \sqrt{s} < 1.100$ GeV using a vector meson dominance model with contribution from $\phi(1020)$, $\rho(770)$, $\omega(782)$ and higher vector resonances. The the first three points near the threshold are removed in the fit.

² KOZYREV 16 also reports $\Gamma(e^+e^-) B(\phi \rightarrow K_S^0 K_L^0) = (0.428 \pm 0.001 \pm 0.009)$ keV.

³ Update of AKHMETSHIN 99D

⁴ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.42 ± 0.11 OUR FIT	Error includes scale factor of 1.2.			
4.51 ± 0.14 OUR AVERAGE				
4.51 $\pm 0.16 \pm 0.11$	105k	AKHMETSHIN 06	CMD2	0.98–1.06 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.665 $\pm 0.042 \pm 0.261$	400k	¹ ACHASOV	01E SND	$e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$
4.35 $\pm 0.27 \pm 0.08$	11169	² AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.38 ± 0.12		BENAYOUN	10 RVUE	0.4–1.05 e^+e^-
4.30 $\pm 0.08 \pm 0.21$		³ AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

² Recalculated by us from the cross section in the peak.

³ Superseded by LEES 21B.

$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma \times \Gamma_8/\Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.87 ± 0.07 OUR FIT	Error includes scale factor of 1.2.			
3.93 ± 0.09 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
4.050 $\pm 0.067 \pm 0.118$	33k	¹ ACHASOV	07B SND	0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
4.093 $^{+0.040}_{-0.043} \pm 0.247$	17.4k	² AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
3.850 $\pm 0.041 \pm 0.159$	23k	^{3,4} AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
4.00 $\pm 0.04 \pm 0.11$		⁵ ACHASOV	00 SND	$e^+e^- \rightarrow \eta\gamma$
3.53 $\pm 0.08 \pm 0.17$	2200	^{6,7} AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \eta\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.19 ± 0.06		⁸ BENAYOUN	10 RVUE	0.4–1.05 e^+e^-

¹ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.

² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).


⁵ From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow 2\gamma) = (39.21 \pm 0.34) \times 10^{-2}$.

⁶ Recalculated by the authors from the cross section in the peak.

⁷ From the $\eta \rightarrow \pi^+\pi^-\pi^0$ decay and using $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.1 \pm 0.5) \times 10^{-2}$.

⁸ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

WEIGHTED AVERAGE
3.93±0.09 (Error scaled by 1.3)



$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
3.94±0.16 OUR FIT				
3.95±0.17 OUR AVERAGE				
4.04±0.09±0.19		¹ ACHASOV 16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
3.75±0.11±0.29	18k	AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.29±0.11		² BENAYOUN 10	RVUE	0.4–1.05 e^+e^-
3.67±0.10 ^{+0.27} _{-0.25}		³ ACHASOV 00	SND	$e^+e^- \rightarrow \pi^0\gamma$


¹ From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$ resonances, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 00.

² A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

³ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\pi^0 \rightarrow 2\gamma) = (98.798 \pm 0.032) \times 10^{-2}$.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
8.5 ±0.6 OUR FIT	Error includes scale factor of 1.2.		
8.8 ±0.9 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
8.36±0.59±0.37	ACHASOV 01G	SND	$e^+e^- \rightarrow \mu^+\mu^-$
9.9 ±1.4 ±0.9	¹ ACHASOV 99c	SND	$e^+e^- \rightarrow \mu^+\mu^-$
14.4 ±3.0	² VASSERMAN 81	OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
8.6 ±5.9	² AUGUSTIN 73	OSPK	$e^+e^- \rightarrow \mu^+\mu^-$


¹ Recalculated by the authors from the cross section in the peak.² Recalculated by us from the cross section in the peak.WEIGHTED AVERAGE
8.8±0.9 (Error scaled by 1.5) $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{11}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8 ±0.6 OUR FIT	Error includes scale factor of 2.0.			
2.8 ±0.5 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
3.51±0.33±0.24	34M	¹ IGNATOV 24	CMD3	$e^+e^- \rightarrow \pi^+\pi^-$
2.1 ±0.3 ±0.3		² ACHASOV 00C	SND	$e^+e^- \rightarrow \pi^+\pi^-$
1.95 ^{+1.15} _{-0.87}		³ GOLUBEV 86	ND	$e^+e^- \rightarrow \pi^+\pi^-$
6.01 ^{+3.19} _{-2.51}		³ VASSERMAN 81	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.31±0.99		⁴ BENAYOUN 13	RVUE	0.4–1.05 e^+e^-

¹ From a fit of the pion form factor in the energy range $0.32 < \sqrt{s} < 1.2$ GeV using the GOUNARIS 68 parametrization with the complex phase of the $\rho - \omega$ interference with ω and ϕ masses and widths constrained by the values and their errors from PDG 22, and leaving $\rho(1450)$, $\rho(1700)$ resonances as free parameters of the fit. A fit without the constraints from PDG 22 gives a value $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}^2 = (3.65 \pm 0.24) \times 10^{-8}$.

² Recalculated by the authors from the cross section in the peak.³ Recalculated by us from the cross section in the peak.⁴ A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$, $K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data.

WEIGHTED AVERAGE
 2.8 ± 0.5 (Error scaled by 1.8)



$\Gamma(\omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
1.40±0.15 OUR FIT			
1.37±0.17±0.01	^{1,2} AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0, 2\pi^0\gamma$
¹ Recalculated by the authors from the cross section at the peak.			
² AMBROSINO 08G reports $[\Gamma(\phi(1020) \rightarrow \omega\pi^0)/\Gamma_{\text{total}} \times \Gamma(\phi(1020) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (1.22 \pm 0.13 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
3.34±0.17 OUR FIT			
3.33^{+0.04+0.19}_{-0.09-0.20}	¹ AMBROSINO 07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
¹ Calculated by the authors from the cross section at the peak.			

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{18}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2^{+0.8}_{-0.7} OUR FIT				
1.17±0.52±0.64	3285	¹ AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
¹ Recalculated by the authors from the cross section in the peak.				

$\phi(1020)$ BRANCHING RATIOS $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.499±0.005 OUR FIT Error includes scale factor of 1.5.**0.493±0.010 OUR AVERAGE**

0.492±0.012	2913	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow K^+ K^-$
0.44 ±0.05	321	KALBFLEISCH 76	HBC	$2.18 K^- p \rightarrow \Lambda K^+ K^-$
0.49 ±0.06	270	DEGROOT 74	HBC	$4.2 K^- p \rightarrow \Lambda \phi$
0.540±0.034	565	BALAKIN 71	OSPK	$e^+ e^- \rightarrow K^+ K^-$
0.48 ±0.04	252	LINDSEY 66	HBC	$2.1\text{--}2.7 K^- p \rightarrow \Lambda K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.493±0.003±0.007		¹ AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow K^+ K^-$
0.476±0.017	1000k	² ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .² Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$. $\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.336±0.004 OUR FIT Error includes scale factor of 1.3.**0.331±0.009 OUR AVERAGE**

0.335±0.010	40644	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$
0.326±0.035		DOLINSKY 91	ND	$e^+ e^- \rightarrow K_L^0 K_S^0$
0.310±0.024		DRUZHININ 84	ND	$e^+ e^- \rightarrow K_L^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.336±0.002±0.006		¹ AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow K_S^0 K_L^0$
0.351±0.013	500k	² ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
0.27 ±0.03	133	KALBFLEISCH 76	HBC	$2.18 K^- p \rightarrow \Lambda K_L^0 K_S^0$
0.257±0.030	95	³ BALAKIN 71	OSPK	$e^+ e^- \rightarrow K_L^0 K_S^0$
0.40 ±0.04	167	LINDSEY 66	HBC	$2.1\text{--}2.7 K^- p \rightarrow \Lambda K_L^0 K_S^0$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .² Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$.³ Balakin error increased by Paul. $\Gamma(K_L^0 K_S^0)/\Gamma(K^+ K^-)$ Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.674±0.014 OUR FIT Error includes scale factor of 1.4.**0.740±0.031 OUR AVERAGE**

0.70 ±0.06	2732	BUKIN 78C	OLYA	$e^+ e^- \rightarrow K_L^0 K_S^0$
0.82 ±0.08		LOSTY 78	HBC	$4.2 K^- p \rightarrow \phi \text{ hyperon}$
0.71 ±0.05		LAVEN 77	HBC	$10 K^- p \rightarrow K^+ K^- \Lambda$
0.71 ±0.08		LYONS 77	HBC	$3\text{--}4 K^- p \rightarrow \Lambda \phi$
0.89 ±0.10	144	AGUILAR-... 72B	HBC	$3.9, 4.6 K^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.644 ± 0.017		¹ DUBNICKA	24	RVUE	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0$
0.638 ± 0.022	2.3M	² KOZYREV	18	CMD3	$e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$
0.68 ± 0.03		³ AKHMETSHIN	95	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$

¹ From a fit to CMD3, BABAR and BES3 data.

² The prediction taking into account phase-space difference, radiative corrections, isospin breaking, and the Sommerfeld-Gamow-Sakharov factor gives 0.630.

³ Theoretical analysis of BRAMON 00 taking into account phase-space difference, electromagnetic radiative corrections, as well as isospin breaking, predicts 0.62. FLOREZ-BAEZ 08 predicts 0.63 considering also structure-dependent radiative corrections. FIS-CHBACH 02 calculates additional corrections caused by the close threshold and predicts 0.68. See also BENAYOUN 01 and DUBYNSKIY 07. BENAYOUN 12 obtains 0.71 ± 0.01 in the HLS model.

$\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$ $\Gamma_2/(\Gamma_1+\Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.403 ± 0.005 OUR FIT Error includes scale factor of 1.4.

0.45 ± 0.04 OUR AVERAGE

0.44 ± 0.07		¹ LONDON	66	HBC	$2.24 K^-p \rightarrow \Lambda K\bar{K}$
0.48 ± 0.07	52	BADIER	65B	HBC	$3 K^-p$
0.40 ± 0.10	34	SCHLEIN	63	HBC	$1.95 K^-p \rightarrow \Lambda K\bar{K}$

¹ This is probably not affected by their controversial background subtraction; the value is from their numbers of $K_1 K_2$ vs $K^+ K^-$ events.

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.149 ± 0.004 OUR FIT Error includes scale factor of 1.3.

0.151 ± 0.009 OUR AVERAGE Error includes scale factor of 1.7.

0.161 ± 0.008	11761	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.143 ± 0.007		DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.155 \pm 0.002 \pm 0.005$		¹ AKHMETSHIN 11	CMD2	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.159 ± 0.008	400k	² ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$
$0.145 \pm 0.009 \pm 0.003$	11169	³ AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.139 ± 0.007		⁴ PARROUR	76B	OSPK e^+e^-

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

³ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁴ Using $\Gamma(\phi) = 4.1$ MeV. If interference between the $\rho\pi$ and 3π modes is neglected, the fraction of the $\rho\pi$ is more than 80% at the 90% confidence level.

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K^+K^-)$ Γ_3/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.299 ± 0.010 OUR FIT Error includes scale factor of 1.4.

0.232 ± 0.017 OUR AVERAGE

$0.230 \pm 0.014 \pm 0.010$		ABLIKIM	25A	BES3	$D_s^+ \rightarrow \phi\pi^+$
0.28 ± 0.09	34	AGUILAR-...	72B	HBC	$3.9, 4.6 K^-p$

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K\bar{K}) \quad \Gamma_3/(\Gamma_1+\Gamma_2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.178±0.005 OUR FIT	Error includes scale factor of 1.3.		
0.24 ±0.04 OUR AVERAGE			
0.237±0.039	CERRADA	77B HBC	4.2 $K^- p \rightarrow \Lambda 3\pi$
0.30 ±0.15	LONDON	66 HBC	2.24 $K^- p \rightarrow \Lambda \pi^+\pi^-\pi^0$

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K_L^0 K_S^0) \quad \Gamma_3/\Gamma_2$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.443±0.012 OUR FIT	Error includes scale factor of 1.2.			
0.51 ±0.05 OUR AVERAGE				
0.56 ±0.07	3681	BUKIN	78c OLYA	$e^+e^- \rightarrow K_L^0 K_S^0, \pi^+\pi^-\pi^0$
0.47 ±0.06	516	COSME	74 OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

$$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$\simeq 0.0087$		1.98M	^{1,2} ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.0006	90		³ ACHASOV	02 SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.23	90		³ CORDIER	80 DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<0.20	90		³ PARROUR	76B OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ From a fit without limitations on charged and neutral ρ masses and widths.

² Adding the direct and $\omega\pi$ contributions and considering the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

³ Neglecting the interference between the $\rho\pi$ and $\pi^+\pi^-\pi^0$.

$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.306±0.024 OUR FIT	Error includes scale factor of 1.2.			
1.26 ±0.04 OUR AVERAGE				
1.246±0.025±0.057	10k	¹ ACHASOV	98F SND	$e^+e^- \rightarrow 7\gamma$
1.18 ±0.11	279	² AKHMETSHIN	95 CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0 3\gamma$
1.30 ±0.06		³ DRUZHININ	84 ND	$e^+e^- \rightarrow 3\gamma$
1.4 ±0.2		⁴ DRUZHININ	84 ND	$e^+e^- \rightarrow 6\gamma$
0.88 ±0.20	290	KURDADZE	83C OLYA	$e^+e^- \rightarrow 3\gamma$
1.35 ±0.29		ANDREWS	77 CNTR	6.7–10 γ Cu
1.5 ±0.4	54	³ COSME	76 OSPK	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.38 ±0.02 ±0.02		⁵ AKHMETSHIN	11 CMD2	1.02 $e^+e^- \rightarrow \eta\gamma$
1.37 ±0.05 ±0.02	33k	⁶ ACHASOV	07B SND	0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
1.373±0.014±0.085	17.4k	^{7,8} AKHMETSHIN	05 CMD2	0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
1.287±0.013±0.063		^{9,10} AKHMETSHIN	01B CMD2	$e^+e^- \rightarrow \eta\gamma$
1.338±0.012±0.052		¹¹ ACHASOV	00 SND	$e^+e^- \rightarrow \eta\gamma$
1.18 ±0.03 ±0.06	2200	¹² AKHMETSHIN	99F CMD2	$e^+e^- \rightarrow \eta\gamma$
1.21 ±0.07		¹³ BENAYOUN	96 RVUE	0.54–1.04 $e^+e^- \rightarrow \eta\gamma$

¹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.2 \pm 0.4) \times 10^{-2}$.

² From $\pi^+\pi^-\pi^0$ decay mode of η .

³ From 2γ decay mode of η .

⁴ From $3\pi^0$ decay mode of η .

⁵ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

⁶ ACHASOV 07B reports $[\Gamma(\phi(1020) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow e^+e^-)] = (4.050 \pm 0.067 \pm 0.118) \times 10^{-6}$ which we divide by our best value $B(\phi(1020) \rightarrow e^+e^-) = (2.964 \pm 0.033) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

⁷ Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

⁸ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

¹⁰ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

¹¹ From the $\eta \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹² From $\pi^+\pi^-\pi^0$ decay mode of η and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹³ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.33 $\pm$ 0.05 OUR FIT

1.31 $\pm$ 0.13 OUR AVERAGE

1.30 $\pm$ 0.13		DRUZHININ	84	ND	$e^+e^- \rightarrow 3\gamma$
-----------------	--	-----------	----	----	------------------------------

1.4 $\pm$ 0.5	32	COSME	76	OSPK	e^+e^-
---------------	----	-------	----	------	----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.367 $\pm$ 0.072		¹ ACHASOV	16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
-------------------	--	----------------------	-----	-----	--

1.258 $\pm$ 0.037 $\pm$ 0.077	18k	^{2,3} AKHMETSHIN	05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
-------------------------------	-----	---------------------------	----	------	--

1.226 $\pm$ 0.036 $^{+0.096}_{-0.089}$		⁴ ACHASOV	00	SND	$e^+e^- \rightarrow \pi^0\gamma$
--	--	----------------------	----	-----	----------------------------------

1.26 $\pm$ 0.17		⁵ BENAYOUN	96	RVUE	0.54–1.04 $e^+e^- \rightarrow \pi^0\gamma$
-----------------	--	-----------------------	----	------	--

¹ Using $B(\phi \rightarrow e^+e^-)$ from PDG 15. Supersedes ACHASOV 00.

² Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$.

³ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

⁴ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁵ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

$\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$ Γ_5/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.9 $\pm$ 0.3 $^{+0.7}_{-0.8}$	ACHASOV	00	SND $e^+e^- \rightarrow \eta\gamma, \pi^0\gamma$
---------------------------------	---------	----	--

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.964±0.033 OUR FIT	Error includes scale factor of 1.4.			
2.98 ±0.07 OUR AVERAGE	Error includes scale factor of 1.1.			
2.93 ±0.14	1900k	¹ ACHASOV	01E SND	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$
2.88 ±0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \text{hadrons}$
3.00 ±0.21	3681	BUKIN	78C OLYA	$e^+e^- \rightarrow \text{hadrons}$
3.10 ±0.14		² PARROUR	76 OSPK	e^+e^-
3.3 ±0.3		COSME	74 OSPK	$e^+e^- \rightarrow \text{hadrons}$
2.81 ±0.25	681	BALAKIN	71 OSPK	$e^+e^- \rightarrow \text{hadrons}$
3.50 ±0.27		CHATELUS	71 OSPK	e^+e^-

¹From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S^0 K_L^0$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

²Using total width 4.2 MeV. They detect 3π mode and observe significant interference with ω tail. This is accounted for in the result quoted above.

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.86 ±0.22 OUR FIT	Error includes scale factor of 1.2.		
2.96 ±0.16 OUR AVERAGE	Error includes scale factor of 1.1.		
2.69 ±0.46	¹ HAYES	71 CNTR	8.3,9.8 $\gamma C \rightarrow \mu^+\mu^- X$
2.17 ±0.60	¹ EARLES	70 CNTR	6.0 $\gamma C \rightarrow \mu^+\mu^- X$
• • • We use the following data for averages but not for fits. • • •			
3.045±0.049±0.148	² AAIJ	24B LHCB	$D^+, D_s^+ \rightarrow \pi^+\phi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.87 ±0.20 ±0.14	³ ACHASOV	01G SND	$e^+e^- \rightarrow \mu^+\mu^-$
3.30 ±0.45 ±0.32	⁴ ACHASOV	99C SND	$e^+e^- \rightarrow \mu^+\mu^-$
4.83 ±1.02	⁵ VASSERMAN	81 OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
2.87 ±1.98	⁵ AUGUSTIN	73 OSPK	$e^+e^- \rightarrow \mu^+\mu^-$

¹Neglecting interference between resonance and continuum.

²Value not independent from the measurement of $B(\phi \rightarrow \mu^+\mu^-)/B(\phi \rightarrow e^+e^-)$.

³Using $B(\phi \rightarrow e^+e^-) = (2.91 \pm 0.07) \times 10^{-4}$.

⁴Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁵Recalculated by us using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

 $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_9/Γ_8

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.022±0.012±0.048	24k	AAIJ	24B LHCB	$D^+, D_s^+ \rightarrow \pi^+\phi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.026±0.020±0.056	7.5k	AAIJ	24B LHCB	$D^+ \rightarrow \pi^+\phi$
1.017±0.013±0.051	16.7k	AAIJ	24B LHCB	$D_s^+ \rightarrow \pi^+\phi$

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.08 ± 0.04 OUR AVERAGE				
1.075 ± 0.007 ± 0.038	30k	¹ BABUSCI 15	KLOE	1.02 $e^+ e^- \rightarrow \eta e^+ e^-$
1.19 ± 0.19 ± 0.12	213	² ACHASOV 01B	SND	$e^+ e^- \rightarrow \eta e^+ e^-$
1.14 ± 0.10 ± 0.06	355	³ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.13 ± 0.14 ± 0.07	183	⁴ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.21 ± 0.14 ± 0.09	130	⁵ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.04 ± 0.20 ± 0.08	42	⁶ AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$
1.3 $^{+0.8}_{-0.6}$	7	GOLUBEV 85	ND	$e^+ e^- \rightarrow \eta e^+ e^-$

¹ Using $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.23)\%$ from PDG 12.² Using $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.32)\%$, $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06)\%$, and $B(\phi \rightarrow e^+ e^-) = (3.00 \pm 0.06) \times 10^{-4}$.³ The average of the branching ratios separately obtained from the $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+ \pi^- \pi^0$ decays.⁴ From $\eta \rightarrow \gamma\gamma$ decays and using $B(\eta \rightarrow \gamma\gamma) = (39.33 \pm 0.25) \times 10^{-2}$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = (4.75 \pm 11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.⁵ From $\eta \rightarrow 3\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$, $B(\eta \rightarrow \pi^+ \pi^- \gamma) = (4.75 \pm 0.11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.⁶ From $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\pi^0 \rightarrow e^+ e^- \gamma) = (1.198 \pm 0.032) \times 10^{-2}$, $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (23.0 \pm 0.4) \times 10^{-2}$, $B(\phi \rightarrow \pi^+ \pi^- \pi^0) = (15.5 \pm 0.6) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$. $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.71 ± 0.11 ± 0.09		¹ ACHASOV 00C	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
0.65 $^{+0.38}_{-0.29}$		¹ GOLUBEV 86	ND	$e^+ e^- \rightarrow \pi^+ \pi^-$
2.01 $^{+1.07}_{-0.84}$		¹ VASSERMAN 81	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$
<6.6	95	BUKIN 78B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$
<2.7	95	ALVENSLEB... 72	CNTR	6.7 $\gamma C \rightarrow C \pi^+ \pi^-$

¹ Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$. $\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.7 ± 0.5 OUR FIT			
5.2 $^{+1.3}_{-1.1}$	^{1,2} AULCHENKO 00A	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.4 ± 0.6	³ AMBROSINO 08G	KLOE	$e^+ e^- \rightarrow \pi^+ \pi^- 2\pi^0, 2\pi^0 \gamma$
~ 5.4	⁴ ACHASOV 00E	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
5.5 $^{+1.6}_{-1.4} \pm 0.3$	^{2,5} AULCHENKO 00A	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
4.8 $^{+1.9}_{-1.7} \pm 0.8$	⁴ ACHASOV 99	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

¹Using the 1996 and 1998 data.²(2.3 ± 0.3)% correction for other decay modes of the $\omega(782)$ applied.³Not independent of the corresponding $\Gamma(\omega\pi^0) \times \Gamma(e^+e^-) / \Gamma^2(\text{total})$.⁴Using the 1996 data.⁵Using the 1998 data.

$\Gamma(\omega\gamma)/\Gamma_{\text{total}}$						Γ_{13}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.05	84	LINDSEY	66	HBC	2.1–2.7 $K^- p \rightarrow \Lambda \pi^+ \pi^-$ neutrals	

$\Gamma(\rho\gamma)/\Gamma_{\text{total}}$						Γ_{14}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
< 0.12	90	¹ AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 7	90	AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$		
<200	84	LINDSEY	66	HBC	2.1–2.7 $K^- p \rightarrow \Lambda \pi^+ \pi^-$ neutrals	

¹Supersedes AKHMETSHIN 97C.

$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$						Γ_{15}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.41 \pm 0.12 \pm 0.04$		30175	¹ AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 0.3	90		² AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
<600	90		KALBFLEISCH 75	HBC	2.18 $K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$	
< 70	90		COSME	74	OSPK $e^+e^- \rightarrow \pi^+\pi^-\gamma$	
<400	90		LINDSEY	65	HBC 2.1–2.7 $K^- p \rightarrow \Lambda \pi^+ \pi^-$ neutrals	

¹For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible. Supersedes AKHMETSHIN 97C.²For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible.

$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}}$						Γ_{16}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
3.22 ± 0.19 OUR FIT	Error includes scale factor of 1.1.					
3.21 ± 0.19 OUR AVERAGE						
$3.21^{+0.03}_{-0.09} \pm 0.18$			¹ AMBROSINO 07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$2.90 \pm 0.21 \pm 1.54$			² AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma, \pi^0\pi^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
4.47 ± 0.21	2438		³ ALOISIO	02D	KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$3.5 \pm 0.3^{+1.3}_{-0.5}$	419		^{4,5} ACHASOV	00H	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$1.93 \pm 0.46 \pm 0.50$	27188		⁶ AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
$3.05 \pm 0.25 \pm 0.72$	268		⁷ AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
1.5 ± 0.5	268		⁸ AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	

$3.42 \pm 0.30 \pm 0.36$	164	⁴ ACHASOV	98I	SND	$e^+e^- \rightarrow 5\gamma$
< 1	90	⁹ AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 7	90	¹⁰ AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 20	90	DRUZHININ	87	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ Obtained by the authors taking into account the $\pi^+\pi^-$ decay mode. Includes a component due to $\pi\pi$ production via the $f_0(500)$ meson. Supersedes ALOISIO 02D.

² From the combined fit of the photon spectra in the reactions $e^+e^- \rightarrow \pi^+\pi^-\gamma$, $\pi^0\pi^0\gamma$.

³ From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution. Superseded by AMBROSINO 07.

⁴ Assuming that the $\pi^0\pi^0\gamma$ final state is completely determined by the $f_0\gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$.

⁵ Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

⁶ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.

⁷ Neglecting other intermediate mechanisms ($\rho\pi$, $\sigma\gamma$).

⁸ A narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.

⁹ For destructive interference with the Bremsstrahlung process

¹⁰ For constructive interference with the Bremsstrahlung process

$\Gamma(f_0(980)\gamma)/\Gamma(\eta\gamma)$

Γ_{16}/Γ_5

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.47 ± 0.15 OUR FIT	Error includes scale factor of 1.1.			
2.6 ± 0.2 ^{+0.8}_{-0.3}	419	¹ ACHASOV	00H	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ Assuming that the $\pi^0\pi^0\gamma$ final state is completely determined by the $f_0\gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$

Γ_{17}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.06 OUR AVERAGE					
1.07 ^{+0.01} _{-0.03} ^{+0.06} _{-0.06}			¹ AMBROSINO	07	KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$
$1.08 \pm 0.17 \pm 0.09$		268	AKHMETSHIN	99C	CMD2 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.09 \pm 0.03 \pm 0.05$		2438	ALOISIO	02D	KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$
$1.158 \pm 0.093 \pm 0.052$		419	^{2,3} ACHASOV	00H	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
< 10		90	DRUZHININ	87	ND $e^+e^- \rightarrow 5\gamma$

¹ Supersedes ALOISIO 02D.

² Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

³ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\eta\gamma)$

Γ_{17}/Γ_5

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.04 OUR FIT				
$0.865 \pm 0.070 \pm 0.017$	419	¹ ACHASOV	00H	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.90 \pm 0.08 \pm 0.07$	164	ACHASOV	98I	SND $e^+e^- \rightarrow 5\gamma$

¹ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.5 ± 2.7 ± 1.6		6.8k	¹ AKHMETSHIN 17	CMD3	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.93 ± 1.74 ± 2.14		3.3k	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
< 870	90		CORDIER 79	WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

¹ Using the cross section at the ϕ meson peak $\sigma(\phi) = 4172 \pm 42$ nb, the nonresonant cross section $\sigma(0) = 1.263 \pm 0.027$ nb and $\text{Re}(Z) = 0.146 \pm 0.030$, $\text{Im}(Z) = -0.002 \pm 0.024$ for the complex amplitude of the $\phi \rightarrow \pi^+\pi^-\pi^+\pi^-$ transition.

 $\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 150	95	BARKOV 88	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.33^{+0.07}_{-0.10} OUR AVERAGE					
1.35 ± 0.05 ^{+0.05} _{-0.10}		9.5k	¹ ANASTASI 16B	KLOE	$e^+e^- \rightarrow \pi^0 e^+ e^-$
1.01 ± 0.28 ± 0.29		52	² ACHASOV 02D	SND	$e^+e^- \rightarrow \pi^0 e^+ e^-$
1.22 ± 0.34 ± 0.21		46	³ AKHMETSHIN 01C	CMD2	$e^+e^- \rightarrow \pi^0 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 12	90		DOLINSKY 88	ND	$e^+e^- \rightarrow \pi^0 e^+ e^-$

¹ Using $B(\pi^0 \rightarrow \gamma\gamma)$ from the 2014 Edition of this Review (PDG 14).


² Using various branching ratios from the 2000 Edition of this Review (PDG 00).

³ Using $B(\pi^0 \rightarrow \gamma\gamma) = 0.98798 \pm 0.00032$, $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$, and $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$.

 $\Gamma(\pi^0 \eta \gamma)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.27 ± 0.30 OUR AVERAGE					Error includes scale factor of 1.5. See the ideogram below.
7.06 ± 0.22		16.9k	¹ AMBROSINO 09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
8.51 ± 0.51 ± 0.57		607	² ALOISIO 02C	KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
7.96 ± 0.60 ± 0.40		197	³ ALOISIO 02C	KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
8.8 ± 1.4 ± 0.9		36	⁴ ACHASOV 00F	SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
9.0 ± 2.4 ± 1.0		80	AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.01 ± 0.10 ± 0.20		13.3k	^{2,5} AMBROSINO 09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
7.12 ± 0.13 ± 0.22		3.6k	^{3,6} AMBROSINO 09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
8.3 ± 2.3 ± 1.2		20	ACHASOV 98B	SND	$e^+e^- \rightarrow 5\gamma$
< 250	90		DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^0\eta\gamma$

- ¹ Combined results of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decay modes measurements.
- ² From the decay mode $\eta \rightarrow \gamma\gamma$.
- ³ From the decay mode $\eta \rightarrow \pi^+\pi^-\pi^0$.
- ⁴ Supersedes ACHASOV 98B.
- ⁵ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.31 \pm 0.20)\%$.
- ⁶ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.73 \pm 0.28)\%$.



$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$			Γ_{22}/Γ		
VALUE (units 10^{-5})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
7.6±0.6 OUR FIT					
7.6±0.6 OUR AVERAGE					
7.4±0.7			¹ ALOISIO	02C KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
8.8±1.7		36	² ACHASOV	00F SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
11 ±2			³ GOKALP	02 RVUE	$e^+e^- \rightarrow \eta\pi^0\gamma$
<500	90		DOLINSKY	91 ND	$e^+e^- \rightarrow \pi^0\eta\gamma$

¹ Using $M_{a_0(980)}=984.8$ MeV and assuming $a_0(980)\gamma$ dominance.

² Assuming $a_0(980)\gamma$ dominance in the $\eta\pi^0\gamma$ final state.

³ Using data of ACHASOV 00F.

$\Gamma(f_0(980)\gamma)/\Gamma(a_0(980)\gamma)$ Γ_{16}/Γ_{22}

VALUE	DOCUMENT ID	TECN	COMMENT
6.1 ± 0.6	¹ ALOISIO	02C	KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$

¹ Using results of ALOISIO 02D and assuming that $f_0(980)$ decays into $\pi\pi$ only and $a_0(980)$ into $\eta\pi$ only.

 $\Gamma(K^0\bar{K}^0\gamma)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.9 \times 10^{-8}$	90	AMBROSINO	09C	KLOE $e^+e^- \rightarrow K_S^0 K_S^0 \gamma$

 $\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.23 ± 0.21 OUR FIT					
6.24 ± 0.30 OUR AVERAGE					
$6.23 \pm 0.27 \pm 0.12$	3407		¹ AMBROSINO	07A	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-7\gamma$
$6.7^{+2.8}_{-2.4} \pm 0.8$	12		² AULCHENKO	03B	SND $e^+e^- \rightarrow \eta'\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$6.7^{+5.0}_{-4.2} \pm 1.5$	7		AULCHENKO	03B	SND $e^+e^- \rightarrow 7\gamma$
$6.10 \pm 0.61 \pm 0.43$	120		³ ALOISIO	02E	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-3\gamma$
$8.2^{+2.1}_{-1.9} \pm 1.1$	21		⁴ AKHMETSHIN	00B	CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$
$4.9^{+2.2}_{-1.8} \pm 0.6$	9		⁵ AKHMETSHIN	00F	CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^- \geq 2\gamma$
6.4 ± 1.6	30		⁶ AKHMETSHIN	00F	CMD2 $e^+e^- \rightarrow \eta'(958)\gamma$
$6.7^{+3.4}_{-2.9} \pm 1.0$	5		⁷ AULCHENKO	99	SND $e^+e^- \rightarrow \pi^+\pi^-3\gamma$
<11	90		AULCHENKO	98	SND $e^+e^- \rightarrow 7\gamma$
$12^{+7}_{-5} \pm 2$	6		⁴ AKHMETSHIN	97B	CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$
<41	90		DRUZHININ	87	ND $e^+e^- \rightarrow \gamma\eta\pi^+\pi^-$

¹ AMBROSINO 07A reports $[\Gamma(\phi(1020) \rightarrow \eta'(958)\gamma)/\Gamma_{\text{total}}] / [B(\phi(1020) \rightarrow \eta\gamma)] = (4.77 \pm 0.09 \pm 0.19) \times 10^{-3}$ which we multiply by our best value $B(\phi(1020) \rightarrow \eta\gamma) = (1.306 \pm 0.024) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Averaging AULCHENKO 03B with AULCHENKO 99.

³ Using $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033)\%$.

⁴ Using the value $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06) \times 10^{-2}$.

⁵ Using $B(\phi \rightarrow K_L^0 K_S^0) = (33.8 \pm 0.6)\%$.

⁶ Averaging AKHMETSHIN 00B with AKHMETSHIN 00F.

⁷ Using the value $B(\eta' \rightarrow \eta\pi^+\pi^-) = (43.7 \pm 1.5) \times 10^{-2}$ and $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.31) \times 10^{-2}$.

$\Gamma(\eta'(958)\gamma)/\Gamma(K_L^0 K_S^0)$ Γ_{24}/Γ_2

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.85$\pm$0.06 OUR FIT				
1.46$^{+0.64}_{-0.54}$ $\pm$0.18	9	¹ AKHMETSHIN 00F	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^- \geq 2\gamma$

¹ Using various branching ratios of K_S^0 , K_L^0 , η , η' from the 2000 edition (The European Physical Journal **C15** 1 (2000)) of this Review.

 $\Gamma(\eta'(958)\gamma)/\Gamma(\eta\gamma)$ Γ_{24}/Γ_5

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.77$\pm$0.15 OUR FIT				
4.78$\pm$0.20 OUR AVERAGE				
4.77 $\pm$ 0.09 $\pm$ 0.19	3407	AMBROSINO 07A	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-7\gamma$
4.70 $\pm$ 0.47 $\pm$ 0.31	120	¹ ALOISIO 02E	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$
6.5 $^{+1.7}_{-1.5}$ $\pm$ 0.8	21	AKHMETSHIN 00B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.5 $^{+5.2}_{-4.0}$ $\pm$ 1.4	6	² AKHMETSHIN 97B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-3\gamma$

¹ From the decay mode $\eta' \rightarrow \eta\pi^+\pi^-$, $\eta \rightarrow \gamma\gamma$.

² Superseded by AKHMETSHIN 00B.

 $\Gamma(\eta\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	AULCHENKO 98	SND	$e^+e^- \rightarrow 7\gamma$

 $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.43$\pm$0.45$\pm$0.14	27188	¹ AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.3 $\pm$ 1.0	824 $\pm$ 33	² AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

¹ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.

² For $E_\gamma > 20$ MeV.

 $\Gamma(\rho\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	AULCHENKO 08	CMD2	$\phi \rightarrow \pi^+\pi^-\gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5	90	AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.8	90	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 6.1	90	AULCHENKO 08	CMD2	$\phi \rightarrow \eta\pi^+\pi^-$
<30	90	AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$

$\Gamma(\eta\mu^+\mu^-)/\Gamma_{\text{total}}$				Γ_{29}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9.4	90	AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$

$\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$				Γ_{30}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1 $\times 10^{-6}$	90	¹ BABUSCI	13B KLOE	1.02 $e^+e^- \rightarrow \eta e^+e^-$

¹ For a narrow vector U with mass between 5 and 470 MeV, from the combined analysis of $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow \pi^0\pi^0\pi^0$ from ARCHILLI 12. Measured 90% CL limits as a function of m_U range from 2.2×10^{-8} to 10^{-6} .

$\Gamma(\text{invisible})/\Gamma(K^+K^-)$				Γ_{31}/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.4 $\times 10^{-4}$	90	ABLIKIM	18S BES3	$J/\psi \rightarrow \phi\eta \rightarrow \phi\pi^+\pi^-\pi^0$

———— Lepton Family number (LF) violating modes ————

$\Gamma(e^\pm\mu^\mp)/\Gamma_{\text{total}}$				Γ_{32}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2 $\times 10^{-6}$	90	ACHASOV	10A SND	$e^+e^- \rightarrow e^\pm\mu^\mp$

$\pi^+\pi^-\pi^0 / \rho\pi$ AMPLITUDE RATIO a_1 IN DECAY OF $\phi \rightarrow \pi^+\pi^-\pi^0$

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1$\pm$1.2 OUR AVERAGE					
10.1 $\pm$ 4.4 $\pm$ 1.7	80k		¹ AKHMETSHIN 06	CMD2	1.017–1.021 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0 $\pm$ 1.1 $\pm$ 0.6	1.98M		^{2,3} ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

–6 < a_1 < 6	500k	³ ACHASOV	02	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
–16 < a_1 < 11	90	9.8k	^{1,4} AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$

¹ Dalitz plot analysis taking into account interference between the contact and $\rho\pi$ amplitudes.

² From a fit without limitations on charged and neutral ρ masses and widths.

³ Recalculated by us to match the notations of AKHMETSHIN 98.

⁴ Assuming zero phase for the contact term.

PARAMETER β IN $\phi \rightarrow P e^+ e^-$ DECAYS

In the one-pole approximation the electromagnetic transition form factor for $\phi \rightarrow P e^+ e^-$ ($P = \pi, \eta$) is given as a function of the $e^+ e^-$ invariant mass squared, q^2 , by the expression:

$$|F(q^2)|^2 = (1 - q^2/\Lambda^2)^{-2},$$

where vector meson dominance predicts parameter $\Lambda \approx 0.770$ GeV ($\Lambda^{-2} \approx 1.687$ GeV $^{-2}$). The slope of this form factor, $\beta = dF/dq^2(q^2=0)$, equals Λ^{-2} in this approximation.

The measurements below obtain β in the one-pole approximation.

PARAMETER β IN $\phi \rightarrow \pi^0 e^+ e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
2.02$\pm$0.11	9.5k	¹ ANASTASI	16B	KLOE 1.02 $e^+ e^- \rightarrow \pi^0 e^+ e^-$

¹ The error combines statistical and systematic uncertainties.

PARAMETER β IN $\phi \rightarrow \eta e^+ e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29$\pm$0.13 OUR AVERAGE				
1.28 $\pm$ 0.10 $^{+0.09}_{-0.08}$	30k	BABUSCI	15	KLOE 1.02 $e^+ e^- \rightarrow \eta e^+ e^-$
3.8 $\pm$ 1.8	213	¹ ACHASOV	01B	SND 1.02 $e^+ e^- \rightarrow \eta e^+ e^-$

¹ The uncertainty is statistical only. The systematic one is negligible, in comparison.

$\phi(1020)$ REFERENCES

ABLIKIM	25A	PRL 134 011904	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	24B	JHEP 2405 293	R. Aaij <i>et al.</i>	(LHCb Collab.)
ACHASOV	24	PR D110 072001	M.N. Achasov <i>et al.</i>	(SND Collab.)
DUBNICKA	24	PR D110 054019	S. Dubnicka <i>et al.</i>	(SLOV, CMNS)
IGNATOV	24	PR D109 112002	F.V. Ignatov <i>et al.</i>	(CMD-3 Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)
HOERICHT...	19	JHEP 1908 137	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>	
KOZYREV	18	PL B779 64	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
AKHMETSHIN	17	PL B768 345	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)
ANASTASI	16B	PL B757 362	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)
KOZYREV	16	PL B760 314	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
BABUSCI	15	PL B742 1	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
BABUSCI	13B	PL B720 111	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ARCHILLI	12	PL B706 251	F. Archilli <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	12	EPJ C72 1848	M. Benayoun <i>et al.</i>	
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
AKHMETSHIN	11	PL B695 412	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
ACHASOV	10A	PR D81 057102	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)

BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>	
AMBROSINO	09C	PL B679 10	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AKHMETSHIN	08	PL B669 217	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AULCHENKO	08	JETPL 88 85	V. Aulchenko <i>et al.</i>	(CMD-2 Collab.)
		Translated from ZETFP 88 93.		
FLOREZ-BAEZ	08	PR D78 077301	F.V. Florez-Baez, G. Lopez Castro	
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	07A	PL B648 267	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
DUBYSKIY	07	PR D75 113001	S. Dubynskiy <i>et al.</i>	
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	06	PL B642 203	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AMBROSINO	05	PL B608 199	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)
AULCHENKO	03B	JETP 97 24	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 124 28.		
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02D	JETPL 75 449	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 75 539.		
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)
ALOISIO	02E	PL B541 45	A. Aloisio <i>et al.</i>	(KLOE Collab.)
FISCHBACH	02	PL B526 355	E. Fischbach, A.W. Overhauser, B. Woodahl	
GOKALP	02	JP G28 2783	A. Gokalp <i>et al.</i>	
ACHASOV	01B	PL B504 275	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)
ACHASOV	01G	PRL 86 1698	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
AKHMETSHIN	01	PL B501 191	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01C	PL B503 237	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
BENAYOUN	01	EPJ C22 503	M. Benayoun, H.B. O'Connell	
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00B	JETP 90 17	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 117 22.		
ACHASOV	00C	PL B474 188	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 72 411.		
ACHASOV	00E	NP B569 158	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00B	PL B473 337	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	00E	PL B491 81	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	00F	PL B494 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 117 1067.		
BRAMON	00	PL B486 406	A. Bramon <i>et al.</i>	
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)
ACHASOV	99	PL B449 122	M.N. Achasov <i>et al.</i>	
ACHASOV	99C	PL B456 304	M.N. Achasov <i>et al.</i>	
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99D	PL B466 385	R.R. Akhmetshin <i>et al.</i>	
Also		PL B508 217 (errat.)	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	99F	PL B460 242	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AULCHENKO	99	JETPL 69 97	V.M. Aulchenko <i>et al.</i>	
		Translated from ZETFP 69 87.		
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98F	JETPL 68 573	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>	
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AULCHENKO	98	PL B436 199	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)
BARBERIS	98	PL B432 436	D. Barberis <i>et al.</i>	(Omega Expt.)
AKHMETSHIN	97B	PL B415 445	R.R. Akhmetshin <i>et al.</i>	(NOVO, BOST, PITT+)
AKHMETSHIN	97C	PL B415 452	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)

BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)
AKHMETSHIN	95	PL B364 199	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko	
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)
BARKOV	88	SJNP 47 248	L.M. Barkov <i>et al.</i>	(NOVO)
DOLINSKY	88	Translated from YAF 47	393.	
		SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)
DRUZHININ	87	Translated from YAF 48	442.	
		ZPHY C37 1	V.P. Druzhinin <i>et al.</i>	(NOVO)
ARMSTRONG	86	PL 166B 245	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BEBEK	86	PRL 56 1893	C. Bebek <i>et al.</i>	(CLEO Collab.)
DAVENPORT	86	PR D33 2519	T.F. Davenport	(TUFTS, ARIZ, FNAL, FSU, NDAM+)
DIJKSTRA	86	ZPHY C31 375	H. Dijkstra <i>et al.</i>	(ANIK, BRIS, CERN+)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
GOLUBEV	86	SJNP 44 409	V.B. Golubev <i>et al.</i>	(NOVO)
ALBRECHT	85D	Translated from YAF 44	633.	
		PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GOLUBEV	85	SJNP 41 756	V.B. Golubev <i>et al.</i>	(NOVO)
DRUZHININ	84	Translated from YAF 41	1183.	
		PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)
KURDADZE	83C	JETPL 38 366	L.M. Kurdadze <i>et al.</i>	(NOVO)
ARENTO	82	Translated from ZETFP 38	306.	
		PR D25 2241	M.W. Arenton <i>et al.</i>	(ANL, ILL)
PELLINEN	82	PS 25 599	A. Pellinen, M. Roos	(HELS)
DAUM	81	PL 100B 439	C. Daum <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
Also		Private Comm.	S.I. Eidelman	(NOVO)
VASSERMAN	81	PL 99B 62	I.B. Vasserman <i>et al.</i>	(NOVO)
Also		SJNP 35 240	L.M. Kurdadze <i>et al.</i>	
CORDIER	80	Translated from YAF 35	352.	
		NP B172 13	A. Cordier <i>et al.</i>	(LALO)
CORDIER	79	PL 81B 389	A. Cordier <i>et al.</i>	(LALO)
BUKIN	78B	SJNP 27 521	A.D. Bukin <i>et al.</i>	(NOVO)
BUKIN	78C	Translated from YAF 27	985.	
		SJNP 27 516	A.D. Bukin <i>et al.</i>	(NOVO)
COOPER	78B	Translated from YAF 27	976.	
		NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)
LOSTY	78	NP B133 38	M.J. Losty <i>et al.</i>	(CERN, AMST, NIJM+)
AKERLOF	77	PRL 39 861	C.W. Akerlof <i>et al.</i>	(FNAL, MICH, PURD)
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)
BALDI	77	PL 68B 381	R. Baldi <i>et al.</i>	(GEVA)
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+)
COHEN	77	PRL 38 269	D. Cohen <i>et al.</i>	(ANL)
LAVEN	77	NP B127 43	H. Laven <i>et al.</i>	(AACH3, BERL, CERN, LOIC+)
LYONS	77	NP B125 207	L. Lyons, A.M. Cooper, A.G. Clark	(OXF)
COSME	76	PL 63B 352	G. Cosme <i>et al.</i>	(ORSAY)
KALBFLEISCH	76	PR D13 22	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
PARROUR	76	PL 63B 357	G. Parrou <i>et al.</i>	(ORSAY)
PARROUR	76B	PL 63B 362	G. Parrou <i>et al.</i>	(ORSAY)
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)
AYRES	74	PRL 32 1463	D.S. Ayres <i>et al.</i>	(ANL)
BESCH	74	NP B70 257	H.J. Besch <i>et al.</i>	(BONN)
COSME	74	PL 48B 155	G. Cosme <i>et al.</i>	(ORSAY)
COSME	74B	PL 48B 159	G. Cosme <i>et al.</i>	(ORSAY)
DEGROOT	74	NP B74 77	A.J. de Groot <i>et al.</i>	(AMST, NIJM)
AUGUSTIN	73	PRL 30 462	J.E. Augustin <i>et al.</i>	(ORSAY)
BALLAM	73B	PR D7 3150	J. Ballam <i>et al.</i>	(SLAC, LBL)
BINNIE	73B	PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)
ALVENSLEB...	72	PRL 28 66	H. Alvensleben <i>et al.</i>	(MIT, DESY)
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)
COLLEY	72	NP B50 1	D.C. Colley <i>et al.</i>	(BIRM, GLAS)
BALAKIN	71	PL 34B 328	V.E. Balakin <i>et al.</i>	(NOVO)
CHATELUS	71	Thesis LAL 1247	Y. Chatelus	(STRB)
Also		PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)
HAYES	71	PR D4 899	S. Hayes <i>et al.</i>	(CORN)
STOTTLE...	71	Thesis ORO 2504 170	A.R. Stottlemeyer	(UMD)

BIZOT	70	PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)
Also		Liverpool Sym. 69	J.P. Perez-y-Jorba	
EARLES	70	PRL 25 1312	D.R. Earles <i>et al.</i>	(NEAS)
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	
LINDSEY	66	PR 147 913	J.S. Lindsey, G. Smith	(LRL)
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IGJPC
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)
LINDSEY	65	PRL 15 221	J.S. Lindsey, G.A. Smith	(LRL)
LINDSEY	65	data included in LINDSEY 66.		
SCHLEIN	63	PRL 10 368	P.E. Schlein <i>et al.</i>	(UCLA) IGJP
