

$\phi_3(1850)$

$$I^G(J^{PC}) = 0^-(3^--)$$

$\phi_3(1850)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1854± 7 OUR AVERAGE				
1855±10		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
1870 ⁺³⁰ ₋₂₀	430	ARMSTRONG	82 OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
1850±10	123	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \Lambda$

$\phi_3(1850)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
87⁺²⁸₋₂₃ OUR AVERAGE Error includes scale factor of 1.2.				
64±31		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
160 ⁺⁹⁰ ₋₅₀	430	ARMSTRONG	82 OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
80 ⁺⁴⁰ ₋₃₀	123	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \Lambda$

$\phi_3(1850)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K}$	seen
Γ_2 $K \bar{K}^*(892) + \text{c.c.}$	seen

$\phi_3(1850)$ BRANCHING RATIOS

$\Gamma(K \bar{K}^*(892) + \text{c.c.})/\Gamma(K \bar{K})$				Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	COMMENT	
0.55^{+0.85}_{-0.45}	ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.8 ±0.4	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \pi \Lambda$	

$\phi_3(1850)$ REFERENCES

ASTON	88E	PL B208 324	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) IGJPC
ARMSTRONG	82	PL 110B 77	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+) JP
ALHARRAN	81B	PL 101B 357	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)