

$\Sigma(1580) \ 3/2^-$

$I(J^P) = 1(\frac{3}{2}^-)$ Status: *

OMITTED FROM SUMMARY TABLE

Seen in the isospin-1 $\overline{K}N$ cross section at BNL (LI 73, CARROLL 76) and in a partial-wave analysis of $K^-p \rightarrow \Lambda\pi^0$ for c.m. energies 1560–1600 MeV by LITCHFIELD 74. LITCHFIELD 74 finds $J^P = 3/2^-$. Not seen by ENGLER 78, CAMERON 78C, OLMSTED 04, nor by PRAKHOV 04.

Neither ZHANG 13A nor SARANTSEV 19 see any evidence for this state.

$\Sigma(1580)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1607^{+13}_{-11}	¹ KAMANO	15	DPWA Multichannel
--------------------	---------------------	----	-------------------

¹From the preferred solution A in KAMANO 15. Solution B reports $M = 1492^{+4}_{-7}$ MeV.

–2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

253^{+30}_{-18}	² KAMANO	15	DPWA Multichannel
-------------------	---------------------	----	-------------------

²From the preferred solution A in KAMANO 15. Solution B reports $M = 138^{+8}_{-14}$ MeV.

$\Sigma(1580)$ POLE RESIDUES

The “normalized residue” is the residue divided by $\Gamma_{pole}/2$.

Normalized residue in $N\overline{K} \rightarrow \Sigma(1580) \rightarrow N\overline{K}$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
---------	-----------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.00778	51	³ KAMANO	15	DPWA Multichannel
---------	----	---------------------	----	-------------------

³From the preferred solution A in KAMANO 15.

Normalized residue in $N\overline{K} \rightarrow \Sigma(1580) \rightarrow \Sigma\pi$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
---------	-----------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0625	–6	⁴ KAMANO	15	DPWA Multichannel
--------	----	---------------------	----	-------------------

⁴From the preferred solution A in KAMANO 15.

Normalized residue in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Lambda\pi$

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.059	156	⁵ KAMANO	15	DPWA Multichannel
⁵ From the preferred solution A in KAMANO 15.				

Normalized residue in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma(1385)\pi$, S-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0368	−18	⁶ KAMANO	15	DPWA Multichannel
⁶ From the preferred solution A in KAMANO 15.				

Normalized residue in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma(1385)\pi$, D-wave

MODULUS	PHASE (°)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0103	123	⁷ KAMANO	15	DPWA Multichannel
⁷ From the preferred solution A in KAMANO 15.				

$\Sigma(1580)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1580 OUR ESTIMATE			
1583±4	⁸ CARROLL	76	DPWA Isospin-1 total σ
1582±4	⁹ LITCHFIELD	74	DPWA $K^- p \rightarrow \Lambda\pi^0$
⁸ CARROLL 76 sees a total-cross-section bump with $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}} = 0.06$.			
⁹ The main effect observed by LITCHFIELD 74 is in the $\Lambda\pi$ final state; the $\bar{K}N$ and $\Sigma\pi$ couplings are estimated from a multichannel fit including total-cross-section data of LI 73.			

$\Sigma(1580)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
15	¹⁰ CARROLL	76	DPWA Isospin-1 total σ
11±4	¹¹ LITCHFIELD	74	DPWA $K^- p \rightarrow \Lambda\pi^0$
¹⁰ CARROLL 76 sees a total-cross-section bump with $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}} = 0.06$.			
¹¹ The main effect observed by LITCHFIELD 74 is in the $\Lambda\pi$ final state; the $\bar{K}N$ and $\Sigma\pi$ couplings are estimated from a multichannel fit including total-cross-section data of LI 73.			

$\Sigma(1580)$ DECAY MODES

Mode	
Γ_1	$N\bar{K}$
Γ_2	$\Lambda\pi$
Γ_3	$\Sigma\pi$
Γ_4	$\Sigma(1385)\pi$, S -wave
Γ_5	$\Sigma(1385)\pi$, D -wave
Γ_6	$N\bar{K}^*(892)$, $S=1/2$, D -wave
Γ_7	$N\bar{K}^*(892)$, $S=3/2$, S -wave
Γ_8	$N\bar{K}^*(892)$, $S=3/2$, D -wave

$\Sigma(1580)$ BRANCHING RATIOS

See “Sign conventions for resonance couplings” in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$+0.03 \pm 0.01$	¹² LITCHFIELD	74	DPWA $\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.003	¹³ KAMANO	15	DPWA Multichannel	
¹² The main effect observed by LITCHFIELD 74 is in the $\Lambda\pi$ final state; the $\bar{K}N$ and $\Sigma\pi$ couplings are estimated from a multichannel fit including total-cross-section data of LI 73.				
¹³ From the preferred solution A in KAMANO 15.				

$\Gamma(\Lambda\pi)/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.490	¹⁴ KAMANO	15	DPWA Multichannel	
¹⁴ From the preferred solution A in KAMANO 15.				

$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$				Γ_3/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.387	¹⁵ KAMANO	15	DPWA Multichannel	
¹⁵ From the preferred solution A in KAMANO 15.				

$\Gamma(\Sigma(1385)\pi, S\text{-wave})/\Gamma_{\text{total}}$				Γ_4/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.12	¹⁶ KAMANO	15	DPWA Multichannel	
¹⁶ From the preferred solution A in KAMANO 15.				

$\Gamma(\Sigma(1385)\pi, D\text{-wave})/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.001¹⁷ KAMANO 15 DPWA Multichannel

¹⁷From the preferred solution A in KAMANO 15.

 $\Gamma(N\bar{K}^*(892), S=1/2, D\text{-wave})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen¹⁸ KAMANO 15 DPWA Multichannel

¹⁸From the preferred solution A in KAMANO 15.

 $\Gamma(N\bar{K}^*(892), S=3/2, S\text{-wave})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen¹⁹ KAMANO 15 DPWA Multichannel

¹⁹From the preferred solution A in KAMANO 15.

 $\Gamma(N\bar{K}^*(892), S=3/2, D\text{-wave})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen²⁰ KAMANO 15 DPWA Multichannel

²⁰From the preferred solution A in KAMANO 15.

 $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Lambda\pi$ $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen CAMERON 78c HBC $K^0 p \rightarrow \Lambda\pi^+$

not seen ENGLER 78 HBC $K_L^0 p \rightarrow \Lambda\pi^+$

$+0.10 \pm 0.02$ ²¹ LITCHFIELD 74 DPWA $K^- p \rightarrow \Lambda\pi^0$

²¹The main effect observed by LITCHFIELD 74 is in the $\Lambda\pi$ final state; the $\bar{K}N$ and $\Sigma\pi$ couplings are estimated from a multichannel fit including total-cross-section data of LI 73.

 $(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma\pi$ $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen CAMERON 78c HBC $K^0 p \rightarrow \Sigma^0\pi^+$

not seen ENGLER 78 HBC $K_L^0 p \rightarrow \Sigma^0\pi^+$

$+0.03 \pm 0.04$ ²² LITCHFIELD 74 DPWA $\bar{K}N$ multichannel

²²The main effect observed by LITCHFIELD 74 is in the $\Lambda\pi$ final state; the $\bar{K}N$ and $\Sigma\pi$ couplings are estimated from a multichannel fit including total-cross-section data of LI 73.

$\Sigma(1580)$ REFERENCES

SARANTSEV	19	EPJ A55 180	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
KAMANO	15	PR C92 025205	H. Kamano <i>et al.</i>	(ANL, OSAK)
ZHANG	13A	PR C88 035205	H. Zhang <i>et al.</i>	(KSU)
OLMSTED	04	PL B588 29	J. Olmsted <i>et al.</i>	(BNL Crystal Ball Collab.)
PRAKHOV	04	PR C69 042202	S. Prakhov <i>et al.</i>	(BNL Crystal Ball Collab.)
CAMERON	78C	NP B132 189	W. Cameron <i>et al.</i>	(BGNA, EDIN, GLAS+) I
ENGLER	78	PR D18 3061	A. Engler <i>et al.</i>	(CMU, ANL)
CARROLL	76	PRL 37 806	A.S. Carroll <i>et al.</i>	(BNL) I
LITCHFIELD	74	PL 51B 509	P.J. Litchfield	(CERN) IJP
LI	73	Purdue Conf. 283	K.K. Li	(BNL) I
