

$$J^{PC} = 0^-(1^--)$$

$\tau(2S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10023.4±0.5	¹ SHAMOV 23	RVUE	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10022.7±0.4	² SHAMOV 23	RVUE	$e^+e^- \rightarrow$ hadrons
10023.5±0.5	^{3,4} ARTAMONOV 00	MD1	$e^+e^- \rightarrow$ hadrons
10023.6±0.5	^{5,6} BARU 86B	MD1	$e^+e^- \rightarrow$ hadrons
10023.1±0.4	⁷ BARBER 84	ARG	$e^+e^- \rightarrow$ hadrons
¹ Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.			
² Obtained by reanalysing ARGUS and Crystal Ball data (BARBER 84), but not authored by the ARGUS and Crystal Ball collaboration.			
³ Reanalysis of BARU 86B using new electron mass (COHEN 87).			
⁴ Superseded by SHAMOV 23.			
⁵ Reanalysis of ARTAMONOV 84.			
⁶ Superseded by ARTAMONOV 00.			
⁷ Reanalysed by SHAMOV 23.			

$m\tau(2S) - m\tau(1S)$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
562.81±0.11 OUR AVERAGE				
562.71±0.04±0.20	3.7M	¹ AAIJ 24AC	LHCB	$\tau \rightarrow \mu^+\mu^-$
562.85±0.03±0.12	89k	¹ AAIJ 24AC	LHCB	$\tau(2S) \rightarrow \tau(1S)\pi^+\pi^-$
¹ Observed in prompt $p\bar{p}$ production.				

$\tau(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID
31.98±2.63 OUR EVALUATION	See the Note on “Width Determinations of the τ States”

$\tau(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \quad \tau(1S)\pi^+\pi^-$	(17.85± 0.26) %	
$\Gamma_2 \quad \tau(1S)\pi^0\pi^0$	(8.6 ± 0.4) %	
$\Gamma_3 \quad \tau^+\tau^-$	(2.00± 0.21) %	
$\Gamma_4 \quad \mu^+\mu^-$	(1.93± 0.17) %	S=2.2
$\Gamma_5 \quad e^+e^-$	(1.91± 0.16) %	
$\Gamma_6 \quad \tau(1S)\pi^0$	< 4 × 10 ^{−5}	CL=90%
$\Gamma_7 \quad \tau(1S)\eta$	(2.9 ± 0.4) × 10 ^{−4}	S=2.0
$\Gamma_8 \quad J/\psi(1S) \text{ anything}$	< 6 × 10 ^{−3}	CL=90%

Γ_9	$J/\psi(1S)\eta_c$	< 5.4	$\times 10^{-6}$	CL=90%
Γ_{10}	$J/\psi(1S)\chi_{c0}$	< 3.4	$\times 10^{-6}$	CL=90%
Γ_{11}	$J/\psi(1S)\chi_{c1}$	< 1.2	$\times 10^{-6}$	CL=90%
Γ_{12}	$J/\psi(1S)\chi_{c2}$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{13}	$J/\psi(1S)\eta_c(2S)$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{14}	$J/\psi(1S)X(3940)$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{15}	$J/\psi(1S)X(4160)$	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{16}	χ_{c1} anything	$(2.2 \pm 0.5) \times 10^{-4}$		
Γ_{17}	$\chi_{c1}(1P)^0 X_{tetra}$	< 3.67	$\times 10^{-5}$	CL=90%
Γ_{18}	χ_{c2} anything	$(2.3 \pm 0.8) \times 10^{-4}$		
Γ_{19}	$\psi(2S)\eta_c$	< 5.1	$\times 10^{-6}$	CL=90%
Γ_{20}	$\psi(2S)\chi_{c0}$	< 4.7	$\times 10^{-6}$	CL=90%
Γ_{21}	$\psi(2S)\chi_{c1}$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{22}	$\psi(2S)\chi_{c2}$	< 1.9	$\times 10^{-6}$	CL=90%
Γ_{23}	$\psi(2S)\eta_c(2S)$	< 3.3	$\times 10^{-6}$	CL=90%
Γ_{24}	$\psi(2S)X(3940)$	< 3.9	$\times 10^{-6}$	CL=90%
Γ_{25}	$\psi(2S)X(4160)$	< 3.9	$\times 10^{-6}$	CL=90%
Γ_{26}	$T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-$	< 1.0	$\times 10^{-6}$	CL=90%
Γ_{27}	$T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-$	< 1.67	$\times 10^{-5}$	CL=90%
Γ_{28}	$T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp$	< 7.3	$\times 10^{-6}$	CL=90%
Γ_{29}	$T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-$	< 1.35	$\times 10^{-5}$	CL=90%
Γ_{30}	$T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-$	< 2.67	$\times 10^{-5}$	CL=90%
Γ_{31}	$T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp$	< 2.72	$\times 10^{-5}$	CL=90%
Γ_{32}	$T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%
Γ_{33}	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp$	< 1.11	$\times 10^{-5}$	CL=90%
Γ_{34}	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp$	< 2.11	$\times 10^{-5}$	CL=90%
Γ_{35}	2H anything	$(2.78^{+0.30}_{-0.26}) \times 10^{-5}$		S=1.2
Γ_{36}	hadrons	$(94 \pm 11) \%$		
Γ_{37}	ggg	$(58.8 \pm 1.2) \%$		
Γ_{38}	$\gamma g g$	$(1.87 \pm 0.28) \%$		
Γ_{39}	$\phi K^+ K^-$	$(1.6 \pm 0.4) \times 10^{-6}$		
Γ_{40}	$\omega \pi^+ \pi^-$	< 2.58	$\times 10^{-6}$	CL=90%
Γ_{41}	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(2.3 \pm 0.7) \times 10^{-6}$		
Γ_{42}	$\phi f_2'(1525)$	< 1.33	$\times 10^{-6}$	CL=90%
Γ_{43}	$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$	CL=90%
Γ_{44}	$\rho(770) a_2(1320)$	< 8.8	$\times 10^{-7}$	CL=90%
Γ_{45}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(1.5 \pm 0.6) \times 10^{-6}$		
Γ_{46}	$K_1(1270)^\pm K^\mp$	< 3.22	$\times 10^{-6}$	CL=90%
Γ_{47}	$K_1(1400)^\pm K^\mp$	< 8.3	$\times 10^{-7}$	CL=90%
Γ_{48}	$b_1(1235)^\pm \pi^\mp$	< 4.0	$\times 10^{-7}$	CL=90%
Γ_{49}	$\rho \pi$	< 1.16	$\times 10^{-6}$	CL=90%
Γ_{50}	$\pi^+ \pi^- \pi^0$	< 8.0	$\times 10^{-7}$	CL=90%
Γ_{51}	$\omega \pi^0$	< 1.63	$\times 10^{-6}$	CL=90%

Γ_{52}	$\pi^+\pi^-\pi^0\pi^0$	$(1.30 \pm 0.28) \times 10^{-5}$	
Γ_{53}	$K_S^0 K^+ \pi^- + \text{c.c.}$	$(1.14 \pm 0.33) \times 10^{-6}$	
Γ_{54}	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$< 4.22 \times 10^{-6}$	CL=90%
Γ_{55}	$K^*(892)^- K^+ + \text{c.c.}$	$< 1.45 \times 10^{-6}$	CL=90%
Γ_{56}	$f_1(1285) \text{ anything}$	$(2.2 \pm 1.6) \times 10^{-3}$	
Γ_{57}	$f_1(1285) X_{tetra}$	$< 6.47 \times 10^{-5}$	CL=90%
Γ_{58}	$D_s^+ D_{s1}(2536)^-$		
Γ_{59}	$D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0$	$(1.6 \pm 0.4) \times 10^{-5}$	
Γ_{60}	$D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-$	$(8.4 \pm 2.3) \times 10^{-6}$	
Γ_{61}	$D_s^{*+} D_{s1}(2536)^-$		
Γ_{62}	$D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0$	$(1.4 \pm 0.4) \times 10^{-5}$	
Γ_{63}	$D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-$	$(8.2 \pm 3.1) \times 10^{-6}$	
Γ_{64}	$D_s^+ D_{s2}^*(2573)^-$		
Γ_{65}	$D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0$	$(1.4 \pm 0.4) \times 10^{-5}$	
Γ_{66}	$D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-$	$(6.9 \pm 3.0) \times 10^{-6}$	
Γ_{67}	$D_s^{*+} D_{s2}^*(2573)^-$		
Γ_{68}	$D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0$	$(9 \pm 5) \times 10^{-6}$	
Γ_{69}	$D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-$	$(5 \pm 6) \times 10^{-6}$	
Γ_{70}	Sum of 100 exclusive modes	$(2.90 \pm 0.30) \times 10^{-3}$	

Radiative decays

Γ_{71}	$\gamma \chi_{b1}(1P)$	$(6.9 \pm 0.4) \%$	
Γ_{72}	$\gamma \chi_{b2}(1P)$	$(7.15 \pm 0.35) \%$	
Γ_{73}	$\gamma \chi_{b0}(1P)$	$(3.8 \pm 0.4) \%$	
Γ_{74}	$\gamma f_0(1710)$	$< 5.9 \times 10^{-4}$	CL=90%
Γ_{75}	$\gamma f_2'(1525)$	$< 5.3 \times 10^{-4}$	CL=90%
Γ_{76}	$\gamma f_2(1270)$	$< 2.41 \times 10^{-4}$	CL=90%
Γ_{77}	$\gamma f_J(2220)$		
Γ_{78}	$\gamma \eta_c(1S)$	$< 2.7 \times 10^{-5}$	CL=90%
Γ_{79}	$\gamma \chi_{c0}$	$< 1.0 \times 10^{-4}$	CL=90%
Γ_{80}	$\gamma \chi_{c1}$	$< 3.6 \times 10^{-6}$	CL=90%
Γ_{81}	$\gamma \chi_{c2}$	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{82}	$\gamma \chi_{c1}(3872)$	$< 1.8 \times 10^{-5}$	CL=90%
Γ_{83}	$\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi$	$< 2.4 \times 10^{-6}$	CL=90%

Γ_{84}	$\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%
Γ_{85}	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%
Γ_{86}	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{87}	$\gamma\eta_b(1S)$	$(5.5 \pm 1.1 - 0.9) \times 10^{-4}$		S=1.2
Γ_{88}	$\gamma\eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	< 3.7	$\times 10^{-6}$	CL=90%
Γ_{89}	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	< 4.9	$\times 10^{-6}$	CL=90%
Γ_{90}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95	$\times 10^{-4}$	CL=95%
Γ_{91}	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$	CL=90%
Γ_{92}	$\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	< 8.3	$\times 10^{-6}$	CL=90%

Lepton Family number (LF) violating modes

Γ_{93}	$e^\pm \tau^\mp$	LF	< 1.12	$\times 10^{-6}$	CL=90%
Γ_{94}	$\mu^\pm \tau^\mp$	LF	< 2.3	$\times 10^{-7}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$

FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements to determine 2 parameters. The overall fit has a $\chi^2 = 11.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

$$x_7 \begin{vmatrix} & 2 \\ & x_1 \end{vmatrix}$$

$\Upsilon(2S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_4 \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
$6.5 \pm 1.5 \pm 1.0$	KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$

$\Gamma(\Upsilon(1S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_1 \Gamma_5 / \Gamma$
VALUE (eV)	EVT5	DOCUMENT ID	TECN	COMMENT
$105.4 \pm 1.0 \pm 4.2$	11.8k	¹ AUBERT	08BP BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$

¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{36}\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.577±0.009 OUR AVERAGE			
0.581±0.004±0.009	¹ ROSNER	06 CLEO	10.0 $e^+e^- \rightarrow \text{hadrons}$
0.552±0.031±0.017	¹ BARU	96 MD1	$e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.04 ±0.02	¹ JAKUBOWSKI	88 CBAL	$e^+e^- \rightarrow \text{hadrons}$
0.58 ±0.03 ±0.04	² GILES	84B CLEO	$e^+e^- \rightarrow \text{hadrons}$
0.60 ±0.12 ±0.07	² ALBRECHT	82 DASP	$e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.07 ^{+0.09} _{-0.05}	² NICZYPORUK	81C LENA	$e^+e^- \rightarrow \text{hadrons}$
0.41 ±0.18	² BOCK	80 CNTR	$e^+e^- \rightarrow \text{hadrons}$

¹ Radiative corrections evaluated following KURAEV 85.² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85. $\Upsilon(2S)$ PARTIAL WIDTHS $\Gamma(e^+e^-)$ Γ_5

VALUE (keV)	DOCUMENT ID
0.612±0.011 OUR EVALUATION	

 $\Upsilon(2S)$ BRANCHING RATIOS $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_1/Γ

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
17.85±0.26 OUR FIT				
17.92±0.26 OUR AVERAGE				
16.8 ±1.1 ±1.3	906k	¹ LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
17.80±0.05±0.37	170k	² LEES	11L BABR	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02±0.02±0.61	851k	³ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
17.22±0.17±0.75	11.8k	⁴ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
19.2 ±0.2 ±1.0	52.6k	⁵ ALEXANDER	98 CLE2	$\pi^+\pi^-\ell^+\ell^-$, $\pi^+\pi^- \text{MM}$
18.1 ±0.5 ±1.0	11.6k	ALBRECHT	87 ARG	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
16.9 ±4.0		GELPHMAN	85 CBAL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
19.1 ±1.2 ±0.6		BESSON	84 CLEO	$\pi^+\pi^- \text{MM}$
18.9 ±2.6		FONSECA	84 CUSB	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$
21 ±7	7	NICZYPORUK	81B LENA	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$

¹ LEES 11C reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.² Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.³ A weighted average of the inclusive and exclusive results.⁴ Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.⁵ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 $\pm$ 0.4 OUR AVERAGE				
8.43 $\pm$ 0.16 $\pm$ 0.42	38k	¹ BHARI 09	CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.2 $\pm$ 0.6 $\pm$ 0.8	275	² ALEXANDER 98	CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.5 $\pm$ 1.9 $\pm$ 1.9	25	ALBRECHT 87	ARG	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 $\pm$ 1.5		GELPHMAN 85	CBAL	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 $\pm$ 2.3		FONSECA 84	CUSB	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$. $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.462 $\pm$ 0.037	¹ BHARI 09	CLEO	$e^+e^- \rightarrow \Upsilon(2S)$

¹ Not independent of other values reported by BHARI 09. $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_3/Γ


VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 $\pm$ 0.21 OUR AVERAGE				
2.00 $\pm$ 0.12 $\pm$ 0.18	22k	¹ BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$
1.7 $\pm$ 1.5 $\pm$ 0.6		HAAS 84B	CLEO	$e^+e^- \rightarrow \tau^+\tau^-$

¹ BESSON 07 reports $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0193 $\pm$ 0.0017 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.				
0.0203 $\pm$ 0.0003 $\pm$ 0.0008	120k	ADAMS	05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0122 $\pm$ 0.0028 $\pm$ 0.0019		¹ KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0138 $\pm$ 0.0025 $\pm$ 0.0015		KAARSBERG	89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.009 $\pm$ 0.006 $\pm$ 0.006		² ALBRECHT	85	ARG	$e^+e^- \rightarrow \mu^+\mu^-$
0.018 $\pm$ 0.008 $\pm$ 0.005		HAAS	84B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.038 90 NICZYPORUK 81C LENA $e^+e^- \rightarrow \mu^+\mu^-$ ¹ Taking into account interference between the resonance and continuum.² Re-evaluated using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$.



$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$					Γ_3/Γ_4
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
$1.04 \pm 0.04 \pm 0.05$	22k	BESSON	07	CLEO	$e^+e^- \rightarrow \Upsilon(2S)$

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 4	90	¹ TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$
< 18	90	² HE	08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<110	90	ALEXANDER	98	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<800	90	LURZ	87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

¹TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)]$
 $< 2.3 \times 10^{-4}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$.
²Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$					Γ_6/Γ_1
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<2.3	90	TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$

Γ_7/Γ

VALUE (units 10^{-4})

CL%

EVTs

DOCUMENT ID

TECN


COMMENT

2.9 ±0.4	OUR FIT	Error includes scale factor of 2.0.			
2.9 ±0.4	OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.			
2.39±0.31±0.14	112	¹ LEES	11L	BABR	$\Upsilon(2S) \rightarrow \ell^+ \ell^- \eta$
2.1 ^{+0.7} _{-0.6} ±0.3	14	² HE	08A	CLEO	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$

• • • We use the following data for averages but not for fits. • • •

3.55±0.32±0.05	241	³ TAMPONI	13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\eta$
< 9	90	^{1,4} AUBERT	08BP	BABR	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$
< 28	90	ALEXANDER	98	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \eta$
< 50	90	ALBRECHT	87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ MM
< 70	90	LURZ	87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ MM
< 20	90	FONSECA	84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma\gamma, \pi^+ \pi^- \pi^0)$

¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.
² Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.
³ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+ \pi^-) = (17.85 \pm 0.26) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
⁴ Using $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.



$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_1

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.64±0.25 OUR FIT Error includes scale factor of 2.0.**1.99±0.14±0.11** 241 TAMPONI 13 BELL $e^+e^- \rightarrow \Upsilon(1S)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35±0.17±0.08 ¹ LEES 11L BABR $\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$ < 5.2 90 ² AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$ ¹ Not independent of other values reported by LEES 11L.² Not independent of other values reported by AUBERT 08BP. $\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$ Γ_6/Γ_7

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13 90 TAMPONI 13 BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$ $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.006 90 MASCHMANN 90 CBAL $e^+e^- \rightarrow \text{hadrons}$ $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<5.4 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$ $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<3.4 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$ $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.2 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$ $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$ $\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.5 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$ $\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$ $\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10⁻⁶ 90 YANG 14 BELL $e^+e^- \rightarrow J/\psi X$

$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.24 \pm 0.44 \pm 0.20$	376	JIA	17	BELL $\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\chi_{c1}(1P)^0 X_{tetra})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 36.7 \times 10^{-6}$	90	¹ JIA	17A	BELL $e^+e^- \rightarrow \text{hadrons}$

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 36.7×10^{-6} .

 $\Gamma(\chi_{c2} \text{ anything})/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.28 \pm 0.73 \pm 0.34$	JIA	17	BELL $\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$

 $\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.1 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.7 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.5 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.9 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.3 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$

 $\Gamma(T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.0 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$.

$$\Gamma(T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-)/\Gamma_{\text{total}} \quad \Gamma_{27}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<16.7 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$				

$$\Gamma(T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp)/\Gamma_{\text{total}} \quad \Gamma_{28}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.3 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm)$.				

$$\Gamma(T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-)/\Gamma_{\text{total}} \quad \Gamma_{29}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<13.5 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$				

$$\Gamma(T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-)/\Gamma_{\text{total}} \quad \Gamma_{30}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<26.7 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$				

$$\Gamma(T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp)/\Gamma_{\text{total}} \quad \Gamma_{31}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<27.2 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$				

$$\Gamma(T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-)/\Gamma_{\text{total}} \quad \Gamma_{32}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$				

$$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp)/\Gamma_{\text{total}} \quad \Gamma_{33}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<11.1 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$				

$$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp)/\Gamma_{\text{total}} \quad \Gamma_{34}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<21.1 \times 10^{-6}$	90	¹ JIA 18	BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$
¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$				

$$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}} \quad \Gamma_{35}/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.78^{+0.30}_{-0.26}$ OUR AVERAGE				Error includes scale factor of 1.2.

$2.64 \pm 0.11^{+0.26}_{-0.21}$		LEES	14G	BABR	$e^+e^- \rightarrow \overline{2H} X$
$3.37 \pm 0.50 \pm 0.25$	58	ASNER	07	CLEO	$e^+e^- \rightarrow \overline{2H} X$

$\Gamma(ggg)/\Gamma_{\text{total}}$
 Γ_{37}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
58.8 ± 1.2	6M	¹ BESSON	06A CLEO	$\Upsilon(2S) \rightarrow \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+\pi^-\Upsilon(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0\pi^0\Upsilon(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

 $\Gamma(\gamma gg)/\Gamma(ggg)$
 Γ_{38}/Γ_{37}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.18 \pm 0.04 \pm 0.47$	6M	BESSON	06A CLEO	$\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$
 Γ_{39}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.58 \pm 0.33 \pm 0.18$	58	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$
 Γ_{40}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.58	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$
 Γ_{41}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.32 \pm 0.40 \pm 0.54$	135	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$
 Γ_{42}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.33	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$
 Γ_{43}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.57	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$
 Γ_{44}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.88	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$
 Γ_{45}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.53 \pm 0.52 \pm 0.19$	32	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$
 Γ_{46}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.22	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$					Γ_{47}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.83	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$
$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$					Γ_{48}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.40	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					Γ_{49}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.16	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{50}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.80	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					Γ_{51}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.63	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{52}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$13.0 \pm 1.9 \pm 2.1$	261 ± 37	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{53}/Γ
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.14 \pm 0.30 \pm 0.13$	40 ± 10	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.2	90	¹ DOBBS	12A		$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$
¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.					
$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{54}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<4.22	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$
$\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{55}/Γ
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.45	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$
$\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$					Γ_{56}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.20 \pm 1.50 \pm 0.63$	2.9k	JIA	17A	BELL	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(f_1(1285)X_{tetra})/\Gamma_{total}$ Γ_{57}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<64.7 \times 10^{-6}$	90	¹ JIA	17A	BELL $e^+e^- \rightarrow \text{hadrons}$

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 7.8×10^{-6} to 64.7×10^{-6} .

 $\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{total}$ Γ_{59}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.6 \pm 0.3 \pm 0.2$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{total}$ Γ_{60}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.84 \pm 0.18 \pm 0.15$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{total}$ Γ_{62}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.4 \pm 0.4 \pm 0.2$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{total}$ Γ_{63}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.82 \pm 0.25 \pm 0.19$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{total}$ Γ_{65}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.4 \pm 0.4 \pm 0.2$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{total}$ Γ_{66}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.69 \pm 0.20 \pm 0.22$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{total}$ Γ_{68}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.9 \pm 0.5 \pm 0.2$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{total}$ Γ_{69}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.54 \pm 0.31 \pm 0.47$	GAO	23	BELL e^+e^- at 10.52 GeV

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{total}$ Γ_{70}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	COMMENT
0.29 ± 0.03	^{1,2} DOBBS	12A $\Upsilon(2S) \rightarrow \text{hadrons}$

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$						Γ_{71}/Γ
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT		
0.069 ± 0.004 OUR AVERAGE						
0.0693 ± 0.0012 ± 0.0041	407k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$	
0.069 ± 0.005 ± 0.009		EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$	
0.091 ± 0.018 ± 0.022		ALBRECHT	85E	ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$	
0.065 ± 0.007 ± 0.012		NERNST	85	CBAL	$e^+e^- \rightarrow \gamma X$	
0.080 ± 0.017 ± 0.016		HAAS	84	CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$	
0.059 ± 0.014		KLOPFEN...	83	CUSB	$e^+e^- \rightarrow \gamma X$	

$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$						Γ_{72}/Γ
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT		
0.0715 ± 0.0035 OUR AVERAGE						
0.0724 ± 0.0011 ± 0.0040	410k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$	
0.074 ± 0.005 ± 0.008		EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$	
0.098 ± 0.021 ± 0.024		ALBRECHT	85E	ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$	
0.058 ± 0.007 ± 0.010		NERNST	85	CBAL	$e^+e^- \rightarrow \gamma X$	
0.102 ± 0.018 ± 0.021		HAAS	84	CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$	
0.061 ± 0.014		KLOPFEN...	83	CUSB	$e^+e^- \rightarrow \gamma X$	

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$						Γ_{73}/Γ
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT		
0.038 ± 0.004 OUR AVERAGE						
0.0375 ± 0.0012 ± 0.0047	198k	ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$	
0.034 ± 0.005 ± 0.006		EDWARDS	99	CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$	
0.064 ± 0.014 ± 0.016		ALBRECHT	85E	ARG	$e^+e^- \rightarrow \gamma \text{conv. } X$	
0.036 ± 0.008 ± 0.009		NERNST	85	CBAL	$e^+e^- \rightarrow \gamma X$	
0.044 ± 0.023 ± 0.009		HAAS	84	CLEO	$e^+e^- \rightarrow \gamma \text{conv. } X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.035 ± 0.014		KLOPFEN...	83	CUSB	$e^+e^- \rightarrow \gamma X$	

$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$						Γ_{74}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<59	90	¹ ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 5.9	90	² ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$	
¹ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.						
² Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+ \pi^-$.						

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$						Γ_{75}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<53	90	¹ ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$	
¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K \bar{K}) = 0.71$.						

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$						Γ_{76}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<24.1	90	¹ ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$	
¹ Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$.						

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	¹ ALBRECHT	89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
--------	----	-----------------------	----	-----	---

¹Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.

 $\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.7 \times 10^{-5}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

 $\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.0 \times 10^{-4}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<3.6 \times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.5 \times 10^{-5}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

 $\Gamma(\gamma \chi_{c1}(3872))/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.8 \times 10^{-5}$	90	¹ WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	-------------------	-----	------	-------------------------------------

¹WANG 11B reports $[\Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 0.8 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 4.3 \times 10^{-2}$.

 $\Gamma(\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.4 \times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

 $\Gamma(\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.8 \times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

 $\Gamma(\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.2 \times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

 $\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.3 \times 10^{-6}$	90	WANG	11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------	----	------	-----	------	-------------------------------------

$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$5.5^{+1.1}_{-0.9}$ OUR AVERAGE Error includes scale factor of 1.2.

$6.1^{+0.6+0.9}_{-0.7-0.6}$	29k	FULSOM	18	BELL	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------------	-----	--------	----	------	-------------------------------------

$3.9 \pm 1.1^{+1.1}_{-0.9}$	13 $\pm$ 5k	¹ AUBERT	09AQ	BABR	$\Upsilon(2S) \rightarrow \gamma X$
-----------------------------	-------------	---------------------	------	------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 21	90	LEES	11J	BABR	$\Upsilon(2S) \rightarrow X\gamma$
--------	----	------	-----	------	------------------------------------

< 8.4	90	¹ BONVICINI	10	CLEO	$\Upsilon(2S) \rightarrow \gamma X$
---------	----	------------------------	----	------	-------------------------------------

< 5.1	90	² ARTUSO	05	CLEO	$e^+e^- \rightarrow \gamma X$
---------	----	---------------------	----	------	-------------------------------

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

² Superseded by BONVICINI 10.

 $\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 3.7 \times 10^{-6}$	90	SANDILYA	13	BELL	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$
------------------------	----	----------	----	------	---

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

< 4.9	90	SANDILYA	13	BELL	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$
---------	----	----------	----	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$46.2^{+29.7}_{-14.2} \pm 10.6$	10	¹ DOBBS	12		$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$
---------------------------------	----	--------------------	----	--	---

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{90}/Γ
 (1.5 GeV $< m_X < 5.0$ GeV)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 1.95	95	ROSNER	07A	CLEO	$e^+e^- \rightarrow \gamma X$
----------	----	--------	-----	------	-------------------------------

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{ hadrons})/\Gamma_{\text{total}}$ Γ_{91}/Γ
 (0.3 GeV $< m_{A^0} < 7$ GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 8 \times 10^{-5}$	90	¹ LEES	11H	BABR	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$
----------------------	----	-------------------	-----	------	---

¹ For a narrow scalar or pseudoscalar, A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 8.3	90	¹ AUBERT	09Z	BABR	$e^+e^- \rightarrow A^0 \rightarrow \gamma \mu^+ \mu^-$
---------	----	---------------------	-----	------	---

¹ For a narrow scalar or pseudoscalar, A^0 , with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} range from 0.26 – 8.3×10^{-6} .

LEPTON FAMILY NUMBER (LF) VIOLATING MODES $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.12 90 DHAMIJA 24 BELL $e^+ e^- \rightarrow e^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2 90 LEES 10B BABR $e^+ e^- \rightarrow e^\pm \tau^\mp$

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 0.23 90 DHAMIJA 24 BELL $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.3 90 LEES 10B BABR $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

<14.4 95 LOVE 08A CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

 $\tau(2S)$ Cross-Particle Branching Ratios $B(\tau(2S) \rightarrow \pi^+ \pi^-) \times B(\tau(3S) \rightarrow \tau(2S) X)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.78 \pm 0.02 \pm 0.11$ 906k LEES 11C BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$

 $\tau(2S)$ REFERENCES

AAIJ	24AC	JHEP 2410 122	R. Aaij <i>et al.</i>	(LHCb Collab.)
DHAMIJA	24	JHEP 2402 187	R. Dhamija <i>et al.</i>	(BELLE Collab.)
GAO	23	PR D108 112015	B.S. Gao <i>et al.</i>	(BELLE Collab.)
SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)
FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BESSION	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSION	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)

ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)
BARU	86B	ZPHY C32 622 (errat.)	S.E. Baru <i>et al.</i>	(NOVO)
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
GELPHMAN	85	PR D32 2893	D. Gelphman <i>et al.</i>	(Crystal Ball Collab.)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
Translated from YAF 41 733.				
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>	
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)