

# N BARYONS

## ( $S = 0, I = 1/2$ )

$p, N^+ = uud; \quad n, N^0 = udd$

**p**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass  $m = 1.007276466621 \pm 0.000000000053$  u

Mass  $m = 938.27208816 \pm 0.00000029$  MeV [a]

$|m_p - m_{\bar{p}}|/m_p < 7 \times 10^{-10}$ , CL = 90% [b]

$|\frac{q_{\bar{p}}}{m_{\bar{p}}}|/(\frac{q_p}{m_p}) = 1.000000000003 \pm 0.000000000016$

$|q_p + q_{\bar{p}}|/e < 7 \times 10^{-10}$ , CL = 90% [b]

$|q_p + q_e|/e < 1 \times 10^{-21}$  [c]

Magnetic moment  $\mu = 2.7928473446 \pm 0.0000000008 \mu_N$

$(\mu_p + \mu_{\bar{p}}) / \mu_p = (0.002 \pm 0.004) \times 10^{-6}$

Electric dipole moment  $d < 0.021 \times 10^{-23}$  e cm

Electric polarizability  $\alpha = (11.5 \pm 0.4) \times 10^{-4}$  fm<sup>3</sup> (S = 1.1)

Magnetic polarizability  $\beta = (2.31 \pm 0.29) \times 10^{-4}$  fm<sup>3</sup> (S = 1.1)

$p$  SPIN POLARIZABILITY  $\gamma_{E1E1} = (-3.0 \pm 0.7) \times 10^{-4}$  fm<sup>4</sup>

$p$  SPIN POLARIZABILITY  $\gamma_{M1M1} = (3.7 \pm 0.5) \times 10^{-4}$  fm<sup>4</sup>

$p$  SPIN POLARIZABILITY  $\gamma_{E1M2} = (-1.2 \pm 1.0) \times 10^{-4}$  fm<sup>4</sup>

$p$  SPIN POLARIZABILITY  $\gamma_{M1E2} = (2.0 \pm 0.8) \times 10^{-4}$  fm<sup>4</sup>

Charge radius,  $\mu p$  Lamb shift =  $0.84087 \pm 0.00039$  fm [d]

Charge radius =  $0.8409 \pm 0.0004$  fm [d]

Magnetic radius =  $0.851 \pm 0.026$  fm [e]

Mean life  $\tau > 9 \times 10^{29}$  years, CL = 90% ( $p \rightarrow$  invisible mode)

See the “Note on Nucleon Decay” in our 1994 edition (Phys. Rev. **D50**, 1173) for a short review.

The “partial mean life” limits tabulated here are the limits on  $\tau/B_i$ , where  $\tau$  is the total mean life and  $B_i$  is the branching fraction for the mode in question. For  $N$  decays,  $p$  and  $n$  indicate proton and neutron partial lifetimes.

| $p$ DECAY MODES                   | Partial mean life<br>( $10^{30}$ years) | Confidence level | $p$<br>(MeV/c) |
|-----------------------------------|-----------------------------------------|------------------|----------------|
| <b>Antilepton + meson</b>         |                                         |                  |                |
| $N \rightarrow e^+ \pi$           | $> 5300 (n), > 24000 (p)$               | 90%              | 459            |
| $N \rightarrow \mu^+ \pi$         | $> 3500 (n), > 16000 (p)$               | 90%              | 453            |
| $N \rightarrow \nu \pi$           | $> 1100 (n), > 390 (p)$                 | 90%              | 459            |
| $p \rightarrow e^+ \eta$          | $> 10000$                               | 90%              | 309            |
| $p \rightarrow \mu^+ \eta$        | $> 4700$                                | 90%              | 297            |
| $n \rightarrow \nu \eta$          | $> 158$                                 | 90%              | 310            |
| $N \rightarrow e^+ \rho$          | $> 217 (n), > 720 (p)$                  | 90%              | 149            |
| $N \rightarrow \mu^+ \rho$        | $> 228 (n), > 570 (p)$                  | 90%              | 113            |
| $N \rightarrow \nu \rho$          | $> 19 (n), > 162 (p)$                   | 90%              | 149            |
| $p \rightarrow e^+ \omega$        | $> 1600$                                | 90%              | 143            |
| $p \rightarrow \mu^+ \omega$      | $> 2800$                                | 90%              | 105            |
| $n \rightarrow \nu \omega$        | $> 108$                                 | 90%              | 144            |
| $N \rightarrow e^+ K$             | $> 17 (n), > 1000 (p)$                  | 90%              | 339            |
| $N \rightarrow \mu^+ K$           | $> 26 (n), > 4500 (p)$                  | 90%              | 329            |
| $N \rightarrow \nu K$             | $> 86 (n), > 5900 (p)$                  | 90%              | 339            |
| $n \rightarrow \nu K_S^0$         | $> 260$                                 | 90%              | 338            |
| $p \rightarrow e^+ K^*(892)^0$    | $> 84$                                  | 90%              | 45             |
| $N \rightarrow \nu K^*(892)$      | $> 78 (n), > 51 (p)$                    | 90%              | 45             |
| <b>Antilepton + mesons</b>        |                                         |                  |                |
| $p \rightarrow e^+ \pi^+ \pi^-$   | $> 82$                                  | 90%              | 448            |
| $p \rightarrow e^+ \pi^0 \pi^0$   | $> 147$                                 | 90%              | 449            |
| $n \rightarrow e^+ \pi^- \pi^0$   | $> 52$                                  | 90%              | 449            |
| $p \rightarrow \mu^+ \pi^+ \pi^-$ | $> 133$                                 | 90%              | 425            |
| $p \rightarrow \mu^+ \pi^0 \pi^0$ | $> 101$                                 | 90%              | 427            |
| $n \rightarrow \mu^+ \pi^- \pi^0$ | $> 74$                                  | 90%              | 427            |
| $n \rightarrow e^+ K^0 \pi^-$     | $> 18$                                  | 90%              | 319            |
| <b>Lepton + meson</b>             |                                         |                  |                |
| $n \rightarrow e^- \pi^+$         | $> 65$                                  | 90%              | 459            |
| $n \rightarrow \mu^- \pi^+$       | $> 49$                                  | 90%              | 453            |
| $n \rightarrow e^- \rho^+$        | $> 62$                                  | 90%              | 150            |
| $n \rightarrow \mu^- \rho^+$      | $> 7$                                   | 90%              | 115            |
| $n \rightarrow e^- K^+$           | $> 32$                                  | 90%              | 340            |
| $n \rightarrow \mu^- K^+$         | $> 57$                                  | 90%              | 330            |

**Lepton + mesons**

|                                   |         |     |     |
|-----------------------------------|---------|-----|-----|
| $p \rightarrow e^- \pi^+ \pi^+$   | $> 30$  | 90% | 448 |
| $n \rightarrow e^- \pi^+ \pi^0$   | $> 29$  | 90% | 449 |
| $p \rightarrow \mu^- \pi^+ \pi^+$ | $> 17$  | 90% | 425 |
| $n \rightarrow \mu^- \pi^+ \pi^0$ | $> 34$  | 90% | 427 |
| $p \rightarrow e^- \pi^+ K^+$     | $> 75$  | 90% | 320 |
| $p \rightarrow \mu^- \pi^+ K^+$   | $> 245$ | 90% | 279 |

**Antilepton + photon(s)**

|                                   |         |     |     |
|-----------------------------------|---------|-----|-----|
| $p \rightarrow e^+ \gamma$        | $> 670$ | 90% | 469 |
| $p \rightarrow \mu^+ \gamma$      | $> 478$ | 90% | 463 |
| $n \rightarrow \nu \gamma$        | $> 550$ | 90% | 470 |
| $p \rightarrow e^+ \gamma \gamma$ | $> 100$ | 90% | 469 |
| $n \rightarrow \nu \gamma \gamma$ | $> 219$ | 90% | 470 |

**Antilepton + single massless**

|                         |         |     |   |
|-------------------------|---------|-----|---|
| $p \rightarrow e^+ X$   | $> 790$ | 90% | — |
| $p \rightarrow \mu^+ X$ | $> 410$ | 90% | — |

**Three (or more) leptons**

|                                   |                      |     |     |
|-----------------------------------|----------------------|-----|-----|
| $p \rightarrow e^+ e^+ e^-$       | $> 34000$            | 90% | 469 |
| $p \rightarrow e^+ \mu^+ \mu^-$   | $> 9200$             | 90% | 457 |
| $p \rightarrow e^+ \nu \nu$       | $> 170$              | 90% | 469 |
| $n \rightarrow e^+ e^- \nu$       | $> 257$              | 90% | 470 |
| $n \rightarrow \mu^+ e^- \nu$     | $> 83$               | 90% | 464 |
| $n \rightarrow \mu^+ \mu^- \nu$   | $> 79$               | 90% | 458 |
| $p \rightarrow \mu^+ e^+ e^-$     | $> 23000$            | 90% | 463 |
| $p \rightarrow \mu^- e^+ e^+$     | $> 19000$            | 90% | 463 |
| $p \rightarrow \mu^+ \mu^+ \mu^-$ | $> 10000$            | 90% | 439 |
| $p \rightarrow \mu^+ \nu \nu$     | $> 220$              | 90% | 463 |
| $p \rightarrow e^- \mu^+ \mu^+$   | $> 11000$            | 90% | 457 |
| $n \rightarrow 3\nu$              | $> 5 \times 10^{-4}$ | 90% | 470 |

**Inclusive modes**

|                                           |                |     |   |
|-------------------------------------------|----------------|-----|---|
| $N \rightarrow e^+ \text{anything}$       | $> 0.6 (n, p)$ | 90% | — |
| $N \rightarrow \mu^+ \text{anything}$     | $> 12 (n, p)$  | 90% | — |
| $N \rightarrow e^+ \pi^0 \text{anything}$ | $> 0.6 (n, p)$ | 90% | — |

 **$\Delta B = 2$  dinucleon modes**

The following are lifetime limits per iron nucleus.

|                              |          |     |   |
|------------------------------|----------|-----|---|
| $pp \rightarrow \pi^+ \pi^+$ | $> 72.2$ | 90% | — |
| $pn \rightarrow \pi^+ \pi^0$ | $> 170$  | 90% | — |
| $nn \rightarrow \pi^+ \pi^-$ | $> 0.7$  | 90% | — |
| $nn \rightarrow \pi^0 \pi^0$ | $> 404$  | 90% | — |
| $pp \rightarrow K^+ K^+$     | $> 170$  | 90% | — |
| $pp \rightarrow e^+ e^+$     | $> 5.8$  | 90% | — |

|                                        |          |     |   |
|----------------------------------------|----------|-----|---|
| $pp \rightarrow e^+ \mu^+$             | $> 3.6$  | 90% | — |
| $pp \rightarrow \mu^+ \mu^+$           | $> 1.7$  | 90% | — |
| $pn \rightarrow e^+ \bar{\nu}$         | $> 260$  | 90% | — |
| $pn \rightarrow \mu^+ \bar{\nu}$       | $> 200$  | 90% | — |
| $pn \rightarrow \tau^+ \bar{\nu}_\tau$ | $> 29$   | 90% | — |
| $nn \rightarrow \text{invisible}$      | $> 1.4$  | 90% | — |
| $nn \rightarrow \nu_e \bar{\nu}_e$     | $> 1.4$  | 90% | — |
| $nn \rightarrow \nu_\mu \bar{\nu}_\mu$ | $> 1.4$  | 90% | — |
| $pn \rightarrow \text{invisible}$      | $> 0.06$ | 90% | — |
| $pp \rightarrow \text{invisible}$      | $> 0.11$ | 90% | — |

**$\bar{p}$  DECAY MODES**

| $\bar{p}$ DECAY MODES                     | Partial mean life<br>(years) | Confidence level | $p$<br>(MeV/c) |
|-------------------------------------------|------------------------------|------------------|----------------|
| $\bar{p} \rightarrow e^- \gamma$          | $> 7 \times 10^5$            | 90%              | 469            |
| $\bar{p} \rightarrow \mu^- \gamma$        | $> 5 \times 10^4$            | 90%              | 463            |
| $\bar{p} \rightarrow e^- \pi^0$           | $> 4 \times 10^5$            | 90%              | 459            |
| $\bar{p} \rightarrow \mu^- \pi^0$         | $> 5 \times 10^4$            | 90%              | 453            |
| $\bar{p} \rightarrow e^- \eta$            | $> 2 \times 10^4$            | 90%              | 309            |
| $\bar{p} \rightarrow \mu^- \eta$          | $> 8 \times 10^3$            | 90%              | 297            |
| $\bar{p} \rightarrow e^- K_S^0$           | $> 900$                      | 90%              | 337            |
| $\bar{p} \rightarrow \mu^- K_S^0$         | $> 4 \times 10^3$            | 90%              | 326            |
| $\bar{p} \rightarrow e^- K_L^0$           | $> 9 \times 10^3$            | 90%              | 337            |
| $\bar{p} \rightarrow \mu^- K_L^0$         | $> 7 \times 10^3$            | 90%              | 326            |
| $\bar{p} \rightarrow e^- \gamma \gamma$   | $> 2 \times 10^4$            | 90%              | 469            |
| $\bar{p} \rightarrow \mu^- \gamma \gamma$ | $> 2 \times 10^4$            | 90%              | 463            |
| $\bar{p} \rightarrow e^- \omega$          | $> 200$                      | 90%              | 143            |

**$n$**

$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$

Mass  $m = 1.0086649160 \pm 0.0000000005$  u  
Mass  $m = 939.5654205 \pm 0.0000005$  MeV [a]  
 $(m_n - m_{\bar{n}}) / m_n = (9 \pm 5) \times 10^{-5}$   
 $m_n - m_p = 1.2933324 \pm 0.0000005$  MeV  
                   $= 0.00138844919(45)$  u  
Mean life  $\tau = 878.4 \pm 0.5$  s (S = 1.8)  
           $c\tau = 2.6335 \times 10^8$  km  
Magnetic moment  $\mu = -1.9130427 \pm 0.0000005 \mu_N$   
Electric dipole moment  $d < 0.18 \times 10^{-25}$  ecm, CL = 90%  
Mean-square charge radius  $\langle r_n^2 \rangle = -0.1155 \pm 0.0017$  fm<sup>2</sup>  
Magnetic radius  $\sqrt{\langle r_M^2 \rangle} = 0.864^{+0.009}_{-0.008}$  fm  
Electric polarizability  $\alpha = (11.8 \pm 1.1) \times 10^{-4}$  fm<sup>3</sup>

Magnetic polarizability  $\beta = (3.7 \pm 1.2) \times 10^{-4} \text{ fm}^3$   
Charge  $q = (-0.2 \pm 0.8) \times 10^{-21} \text{ e}$   
Mean  $n\bar{n}$ -oscillation time  $> 8.6 \times 10^7 \text{ s}$ , CL = 90% (free  $n$ )  
Mean  $n\bar{n}$ -oscillation time  $> 4.7 \times 10^8 \text{ s}$ , CL = 90% <sup>[f]</sup> (bound  $n$ )  
Mean  $nn'$ -oscillation time  $> 448 \text{ s}$ , CL = 90% <sup>[g]</sup>

**$pe^- \nu_e$  decay parameters <sup>[h]</sup>**

$\lambda \equiv g_A / g_V = -1.2753 \pm 0.0013 \quad (S = 2.7)$   
 $A = -0.11958 \pm 0.00021 \quad (S = 1.2)$   
 $B = 0.9807 \pm 0.0030$   
 $C = -0.2377 \pm 0.0026$   
 $a = -0.1044 \pm 0.0007$   
 $\phi_{AV} = (180.017 \pm 0.026)^\circ$  <sup>[i]</sup>  
 $D = (-1.2 \pm 2.0) \times 10^{-4}$  <sup>[j]</sup>  
 $R = 0.004 \pm 0.013$  <sup>[j]</sup>  
Fierz interference term  $b = 0.017 \pm 0.020$

See the proton listings for many other neutron decay modes.

| <b><math>n</math> DECAY MODES</b>                          | Fraction ( $\Gamma_i/\Gamma$ )     |                  | Confidence level  | $\frac{p}{p}$<br>(MeV/c) |
|------------------------------------------------------------|------------------------------------|------------------|-------------------|--------------------------|
| $pe^- \bar{\nu}_e$                                         | 100                                | %                |                   | 1                        |
| $pe^- \bar{\nu}_e \gamma$                                  | [k] $(9.2 \pm 0.7) \times 10^{-3}$ |                  |                   | 1                        |
| hydrogen-atom $\bar{\nu}_e$                                | $< 2.7$                            | $\times 10^{-3}$ | 95%               | 1.19                     |
| <b>Charge conservation (<math>Q</math>) violating mode</b> |                                    |                  |                   |                          |
| $p\nu_e \bar{\nu}_e$                                       | $Q$                                | $< 8$            | $\times 10^{-27}$ | 68%<br>1                 |

$N(1440) \ 1/2^+$

$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$

Re(pole position) = 1360 to 1380 ( $\approx 1370$ ) MeV  
 $-2\text{Im}(\text{pole position}) = 180$  to  $205$  ( $\approx 190$ ) MeV  
Breit-Wigner mass = 1410 to 1470 ( $\approx 1440$ ) MeV  
Breit-Wigner full width = 250 to 450 ( $\approx 350$ ) MeV

| <b><math>N(1440)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 55–75 %                        | 398         |
| $N\eta$                                 | <1 %                           | †           |
| $N\pi\pi$                               | 17–50 %                        | 347         |
| $\Delta(1232)\pi$ , $P$ -wave           | 6–27 %                         | 147         |
| $N\sigma$                               | 11–23 %                        | –           |
| $p\gamma$ , helicity=1/2                | 0.035–0.048 %                  | 414         |
| $n\gamma$ , helicity=1/2                | 0.02–0.04 %                    | 413         |

$$N(1520) \ 3/2^-$$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Re(pole position) = 1505 to 1515 ( $\approx 1510$ ) MeV  
–2Im(pole position) = 105 to 120 ( $\approx 110$ ) MeV  
Breit-Wigner mass = 1510 to 1520 ( $\approx 1515$ ) MeV  
Breit-Wigner full width = 100 to 120 ( $\approx 110$ ) MeV

| <b><math>N(1520)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 55–65 %                        | 453         |
| $N\eta$                                 | 0.07–0.09 %                    | 142         |
| $N\pi\pi$                               | 25–35 %                        | 410         |
| $\Delta(1232)\pi$                       | 22–34 %                        | 225         |
| $\Delta(1232)\pi$ , $S$ -wave           | 15–23 %                        | 225         |
| $\Delta(1232)\pi$ , $D$ -wave           | 7–11 %                         | 225         |
| $N\rho$                                 | 10–16 %                        | †           |
| $N\rho$ , $S=3/2$ , $S$ -wave           | 10–16 %                        | †           |
| $N\rho$ , $S=1/2$ , $D$ -wave           | 0.2–0.4 %                      | †           |
| $N\sigma$                               | <10 %                          | –           |
| $p\gamma$                               | 0.31–0.52 %                    | 467         |
| $p\gamma$ , helicity=1/2                | 0.01–0.02 %                    | 467         |
| $p\gamma$ , helicity=3/2                | 0.30–0.50 %                    | 467         |
| $n\gamma$                               | 0.30–0.53 %                    | 466         |
| $n\gamma$ , helicity=1/2                | 0.04–0.10 %                    | 466         |
| $n\gamma$ , helicity=3/2                | 0.25–0.45 %                    | 466         |

$$N(1535) \ 1/2^-$$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

Re(pole position) = 1500 to 1520 ( $\approx 1510$ ) MeV  
–2Im(pole position) = 80 to 130 ( $\approx 110$ ) MeV  
Breit-Wigner mass = 1515 to 1545 ( $\approx 1530$ ) MeV  
Breit-Wigner full width = 125 to 175 ( $\approx 150$ ) MeV

| <b><i>N</i>(1535) DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------|--------------------------------|-------------|
| $N\pi$                            | 32–52 %                        | 464         |
| $N\eta$                           | 30–55 %                        | 176         |
| $N\pi\pi$                         | 4–31 %                         | 422         |
| $\Delta(1232)\pi$ , $D$ -wave     | 1–4 %                          | 240         |
| $N\rho$                           | 2–17 %                         | †           |
| $N\rho$ , $S=1/2$ , $S$ -wave     | 2–16 %                         | †           |
| $N\rho$ , $S=3/2$ , $D$ -wave     | <1 %                           | †           |
| $N\sigma$                         | 2–10 %                         | –           |
| $N(1440)\pi$                      | 5–12 %                         | †           |
| $p\gamma$ , helicity=1/2          | 0.15–0.30 %                    | 477         |
| $n\gamma$ , helicity=1/2          | 0.01–0.25 %                    | 477         |

***N*(1650) 1/2<sup>–</sup>**

$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$

Re(pole position) = 1650 to 1680 ( $\approx$  1665) MeV  
– 2Im(pole position) = 100 to 170 ( $\approx$  135) MeV  
Breit-Wigner mass = 1635 to 1665 ( $\approx$  1650) MeV  
Breit-Wigner full width = 100 to 150 ( $\approx$  125) MeV

| <b><i>N</i>(1650) DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------|--------------------------------|-------------|
| $N\pi$                            | 50–70 %                        | 547         |
| $N\eta$                           | 15–35 %                        | 348         |
| $\Lambda K$                       | 5–15 %                         | 169         |
| $N\pi\pi$                         | 20–58 %                        | 514         |
| $\Delta(1232)\pi$ , $D$ -wave     | 6–18 %                         | 345         |
| $N\rho$                           | 12–22 %                        | †           |
| $N\rho$ , $S=1/2$ , $S$ -wave     | <4 %                           | †           |
| $N\rho$ , $S=3/2$ , $D$ -wave     | 12–18 %                        | †           |
| $N\sigma$                         | 2–18 %                         | –           |
| $N(1440)\pi$                      | 6–26 %                         | 150         |
| $p\gamma$ , helicity=1/2          | 0.04–0.20 %                    | 558         |
| $n\gamma$ , helicity=1/2          | 0.003–0.17 %                   | 557         |

***N*(1675) 5/2<sup>–</sup>**

$I(J^P) = \frac{1}{2}(\frac{5}{2}^-)$

Re(pole position) = 1650 to 1660 ( $\approx$  1655) MeV  
– 2Im(pole position) = 120 to 150 ( $\approx$  135) MeV  
Breit-Wigner mass = 1665 to 1680 ( $\approx$  1675) MeV  
Breit-Wigner full width = 130 to 160 ( $\approx$  145) MeV

| <b><math>N(1675)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 38–42 %                        | 564         |
| $N\eta$                                 | < 1 %                          | 376         |
| $\Lambda K$                             | <0.04 %                        | 216         |
| $N\pi\pi$                               | 25–45 %                        | 532         |
| $\Delta(1232)\pi$ , $D$ -wave           | 23–37 %                        | 366         |
| $N\rho$                                 | 0.1–0.9 %                      | †           |
| $N\rho$ , $S=1/2$                       | <0.2 %                         | †           |
| $N\rho$ , $S=3/2$ , $D$ -wave           | 0.1–0.7 %                      | †           |
| $N\sigma$                               | 3–7 %                          | –           |
| $p\gamma$                               | 0–0.02 %                       | 575         |
| $p\gamma$ , helicity=1/2                | 0–0.01 %                       | 575         |
| $p\gamma$ , helicity=3/2                | 0–0.01 %                       | 575         |
| $n\gamma$                               | 0–0.15 %                       | 574         |
| $n\gamma$ , helicity=1/2                | 0–0.05 %                       | 574         |
| $n\gamma$ , helicity=3/2                | 0–0.10 %                       | 574         |

$$N(1680) \ 5/2^+$$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^+)$$

Re(pole position) = 1660 to 1680 ( $\approx 1670$ ) MeV  
– 2Im(pole position) = 110 to 135 ( $\approx 120$ ) MeV  
Breit-Wigner mass = 1680 to 1690 ( $\approx 1685$ ) MeV  
Breit-Wigner full width = 115 to 130 ( $\approx 120$ ) MeV

| <b><math>N(1680)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 60–70 %                        | 571         |
| $N\eta$                                 | <1 %                           | 386         |
| $N\pi\pi$                               | 28–53 %                        | 539         |
| $\Delta(1232)\pi$                       | 11–23 %                        | 374         |
| $\Delta(1232)\pi$ , $P$ -wave           | 4–10 %                         | 374         |
| $\Delta(1232)\pi$ , $F$ -wave           | 1–13 %                         | 374         |
| $N\rho$                                 | 8–11 %                         | †           |
| $N\rho$ , $S=3/2$ , $P$ -wave           | 6–8 %                          | †           |
| $N\rho$ , $S=3/2$ , $F$ -wave           | 2–3 %                          | †           |
| $N\sigma$                               | 9–19 %                         | –           |
| $p\gamma$                               | 0.21–0.32 %                    | 581         |
| $p\gamma$ , helicity=1/2                | 0.001–0.011 %                  | 581         |
| $p\gamma$ , helicity=3/2                | 0.20–0.32 %                    | 581         |
| $n\gamma$                               | 0.021–0.046 %                  | 581         |
| $n\gamma$ , helicity=1/2                | 0.004–0.029 %                  | 581         |
| $n\gamma$ , helicity=3/2                | 0.01–0.024 %                   | 581         |

**$N(1700) \ 3/2^-$**

$$I(J^P) = \tfrac{1}{2}(\tfrac{3}{2}^-)$$

Re(pole position) = 1650 to 1750 ( $\approx 1700$ ) MeV  
 $-2\text{Im}(\text{pole position}) = 100$  to  $300$  ( $\approx 200$ ) MeV  
Breit-Wigner mass = 1650 to 1800 ( $\approx 1720$ ) MeV  
Breit-Wigner full width = 100 to 300 ( $\approx 200$ ) MeV

| <b><math>N(1700)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 7–17 %                         | 594         |
| $N\eta$                                 | 1–2 %                          | 422         |
| $N\omega$                               | 10–34 %                        | †           |
| $\Lambda K$                             | 1–2 %                          | 283         |
| $N\pi\pi$                               | >89 %                          | 564         |
| $\Delta(1232)\pi$                       | 55–85 %                        | 402         |
| $\Delta(1232)\pi$ , $S$ -wave           | 50–80 %                        | 402         |
| $\Delta(1232)\pi$ , $D$ -wave           | 4–14 %                         | 402         |
| $N\rho$ , $S=3/2$ , $S$ -wave           | 32–44 %                        | 75          |
| $N\sigma$                               | 2–14 %                         | –           |
| $N(1440)\pi$                            | 3–11 %                         | 225         |
| $N(1520)\pi$                            | <4 %                           | 145         |
| $p\gamma$                               | 0.01–0.05 %                    | 604         |
| $p\gamma$ , helicity=1/2                | 0.0–0.024 %                    | 604         |
| $p\gamma$ , helicity=3/2                | 0.002–0.026 %                  | 604         |
| $n\gamma$                               | 0.01–0.13 %                    | 603         |
| $n\gamma$ , helicity=1/2                | 0.0–0.09 %                     | 603         |
| $n\gamma$ , helicity=3/2                | 0.01–0.05 %                    | 603         |

**$N(1710) \ 1/2^+$**

$$I(J^P) = \tfrac{1}{2}(\tfrac{1}{2}^+)$$

Re(pole position) = 1650 to 1750 ( $\approx 1700$ ) MeV  
 $-2\text{Im}(\text{pole position}) = 80$  to  $160$  ( $\approx 120$ ) MeV  
Breit-Wigner mass = 1680 to 1740 ( $\approx 1710$ ) MeV  
Breit-Wigner full width = 80 to 200 ( $\approx 140$ ) MeV

| <b><i>N</i>(1710) DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------|--------------------------------|-------------|
| $N\pi$                            | 5–20 %                         | 588         |
| $N\eta$                           | 10–50 %                        | 412         |
| $N\omega$                         | 1–5 %                          | †           |
| $\Lambda K$                       | 5–25 %                         | 269         |
| $\Sigma K$                        | seen                           | 138         |
| $N\pi\pi$                         | 14–48 %                        | 557         |
| $\Delta(1232)\pi$ , $P$ -wave     | 3–9 %                          | 394         |
| $N\rho$ , $S=1/2$ , $P$ -wave     | 11–23 %                        | †           |
| $N\sigma$                         | <16 %                          | –           |
| $N(1535)\pi$                      | 9–21 %                         | 113         |
| $p\gamma$ , helicity=1/2          | 0.002–0.08 %                   | 598         |
| $n\gamma$ , helicity=1/2          | 0.0–0.02%                      | 597         |

***N*(1720) 3/2<sup>+</sup>**

$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$

Re(pole position) = 1660 to 1710 ( $\approx$  1680) MeV  
–2Im(pole position) = 150 to 300 ( $\approx$  200) MeV  
Breit-Wigner mass = 1680 to 1750 ( $\approx$  1720) MeV  
Breit-Wigner full width = 150 to 400 ( $\approx$  250) MeV

| <b><i>N</i>(1720) DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------|--------------------------------|-------------|
| $N\pi$                            | 8–14 %                         | 594         |
| $N\eta$                           | 1–5 %                          | 422         |
| $N\omega$                         | 12–40 %                        | †           |
| $\Lambda K$                       | 4–19 %                         | 283         |
| $N\pi\pi$                         | >50 %                          | 564         |
| $\Delta(1232)\pi$                 | 47–89 %                        | 402         |
| $\Delta(1232)\pi$ , $P$ -wave     | 47–77 %                        | 402         |
| $\Delta(1232)\pi$ , $F$ -wave     | <12 %                          | 402         |
| $N\rho$ , $S=1/2$ , $P$ -wave     | 1–2 %                          | 75          |
| $N\sigma$                         | 2–14 %                         | –           |
| $N(1440)\pi$                      | <2 %                           | 225         |
| $N(1520)\pi$ , $S$ -wave          | 1–5 %                          | 145         |
| $p\gamma$                         | 0.05–0.25 %                    | 604         |
| $p\gamma$ , helicity=1/2          | 0.05–0.15 %                    | 604         |
| $p\gamma$ , helicity=3/2          | 0.002–0.16 %                   | 604         |
| $n\gamma$                         | 0.0–0.016 %                    | 603         |
| $n\gamma$ , helicity=1/2          | 0.0–0.01 %                     | 603         |
| $n\gamma$ , helicity=3/2          | 0.0–0.015 %                    | 603         |

**$N(1875) \ 3/2^-$**

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

was  $N(2080)$

Re(pole position) = 1850 to 1950 ( $\approx 1900$ ) MeV  
 $-2\text{Im}(\text{pole position}) = 100$  to  $220$  ( $\approx 160$ ) MeV  
Breit-Wigner mass = 1850 to 1920 ( $\approx 1875$ ) MeV  
Breit-Wigner full width = 120 to 250 ( $\approx 200$ ) MeV

| <b><math>N(1875)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 3–11 %                         | 695         |
| $N\eta$                                 | 3–16 %                         | 559         |
| $N\omega$                               | 15–25 %                        | 371         |
| $\Lambda K$                             | 1–2 %                          | 454         |
| $\Sigma K$                              | 0.3–1.1 %                      | 384         |
| $N\pi\pi$                               | >56 %                          | 670         |
| $\Delta(1232)\pi$                       | 4–44 %                         | 520         |
| $\Delta(1232)\pi$ , $S$ -wave           | 2–21 %                         | 520         |
| $\Delta(1232)\pi$ , $D$ -wave           | 2–23 %                         | 520         |
| $N\rho$ , $S=3/2$ , $S$ -wave           | 36–56 %                        | 379         |
| $N\sigma$                               | 16–60 %                        | –           |
| $N(1440)\pi$                            | 2–8 %                          | 365         |
| $N(1520)\pi$                            | <2 %                           | 301         |
| $\Lambda K^*(892)$                      | <0.2 %                         | †           |
| $p\gamma$                               | 0.001–0.025 %                  | 703         |
| $p\gamma$ , helicity=1/2                | 0.001–0.021 %                  | 703         |
| $p\gamma$ , helicity=3/2                | <0.003 %                       | 703         |
| $n\gamma$                               | <0.040 %                       | 702         |
| $n\gamma$ , helicity=1/2                | <0.007 %                       | 702         |
| $n\gamma$ , helicity=3/2                | <0.033 %                       | 702         |

**$N(1880) \ 1/2^+$**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Re(pole position) = 1820 to 1900 ( $\approx 1860$ ) MeV  
 $-2\text{Im}(\text{pole position}) = 180$  to  $280$  ( $\approx 230$ ) MeV  
Breit-Wigner mass = 1830 to 1930 ( $\approx 1880$ ) MeV  
Breit-Wigner full width = 200 to 400 ( $\approx 300$ ) MeV

| <b><math>N(1880)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 3–31 %                         | 698         |
| $N\eta$                                 | 1–55 %                         | 563         |
| $N\omega$                               | 12–28 %                        | 377         |
| $\Lambda K$                             | 1–3 %                          | 459         |
| $\Sigma K$                              | 10–24 %                        | 389         |
| $N\pi\pi$                               | >32 %                          | 673         |
| $\Delta(1232)\pi$                       | 5–42 %                         | 524         |
| $N\rho$ , $S=1/2$ , $P$ -wave           | 19–45 %                        | 385         |
| $N\sigma$                               | 8–40 %                         | 539         |
| $N(1535)\pi$                            | 4–12 %                         | 293         |
| $Na_0(980)$                             | 1–5 %                          | †           |
| $\Lambda K^*(892)$                      | 0.5–1.1 %                      | †           |
| $p\gamma$ , helicity=1/2                | seen                           | 706         |
| $n\gamma$ , helicity=1/2                | 0.002–0.63 %                   | 705         |

$N(1895) \ 1/2^-$

$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$

was  $N(2090)$

Re(pole position) = 1890 to 1930 ( $\approx 1910$ ) MeV  
–2Im(pole position) = 80 to 140 ( $\approx 110$ ) MeV  
Breit-Wigner mass = 1870 to 1920 ( $\approx 1895$ ) MeV  
Breit-Wigner full width = 80 to 200 ( $\approx 120$ ) MeV

| <b><math>N(1895)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 2–18 %                         | 707         |
| $N\eta$                                 | 15–45 %                        | 575         |
| $N\eta'$                                | 10–40 %                        | †           |
| $N\omega$                               | 16–40 %                        | 395         |
| $\Lambda K$                             | 3–23 %                         | 473         |
| $\Sigma K$                              | 6–20 %                         | 405         |
| $N\pi\pi$                               | 17–74 %                        | 683         |
| $\Delta(1232)\pi$ , $D$ -wave           | 3–11 %                         | 535         |
| $N\rho$                                 | 14–50 %                        | 403         |
| $N\rho$ , $S=1/2$ , $S$ -wave           | <18 %                          | 403         |
| $N\rho$ , $S=3/2$ , $D$ -wave           | 14–32 %                        | 403         |
| $N\sigma$                               | <13 %                          | –           |
| $N(1440)\pi$                            | 2–12 %                         | 382         |

|                          |              |     |
|--------------------------|--------------|-----|
| $\Lambda K^*(892)$       | 4–9 %        | †   |
| $p\gamma$ , helicity=1/2 | 0.01–0.06 %  | 715 |
| $n\gamma$ , helicity=1/2 | 0.003–0.05 % | 715 |

$N(1900) \ 3/2^+$

$$I(J^P) = \tfrac{1}{2}(\tfrac{3}{2}^+)$$

Re(pole position) = 1900 to 1940 ( $\approx$  1920) MeV  
–2Im(pole position) = 90 to 160 ( $\approx$  130) MeV  
Breit-Wigner mass = 1890 to 1950 ( $\approx$  1920) MeV  
Breit-Wigner full width = 100 to 320 ( $\approx$  200) MeV

| <b><math>N(1900)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 1–20 %                         | 723         |
| $N\eta$                                 | 2–14 %                         | 595         |
| $N\eta'$                                | 4–8 %                          | 151         |
| $N\omega$                               | 7–13 %                         | 424         |
| $\Lambda K$                             | 2–20 %                         | 495         |
| $\Sigma K$                              | 3–7 %                          | 431         |
| $N\pi\pi$                               | >56 %                          | 699         |
| $\Delta(1232)\pi$                       | 30–70 %                        | 553         |
| $\Delta(1232)\pi$ , $P$ -wave           | 9–25 %                         | 553         |
| $\Delta(1232)\pi$ , $F$ -wave           | 21–45 %                        | 553         |
| $N\rho$ , $S=1/2$                       | 25–40 %                        | 432         |
| $N\sigma$                               | 1–7 %                          | –           |
| $N(1520)\pi$                            | 7–23 %                         | 341         |
| $N(1535)\pi$                            | 4–10 %                         | 328         |
| $\Lambda K^*(892)$                      | < 0.2 %                        | †           |
| $p\gamma$                               | 0.001–0.025 %                  | 731         |
| $p\gamma$ , helicity=1/2                | 0.001–0.021 %                  | 731         |
| $p\gamma$ , helicity=3/2                | <0.003 %                       | 731         |
| $n\gamma$                               | <0.040 %                       | 730         |
| $n\gamma$ , helicity=1/2                | <0.007 %                       | 730         |
| $n\gamma$ , helicity=3/2                | <0.033 %                       | 730         |

$N(2060) \ 5/2^-$

$$I(J^P) = \tfrac{1}{2}(\tfrac{5}{2}^-)$$

was  $N(2200)$

Re(pole position) = 2020 to 2130 ( $\approx$  2070) MeV  
–2Im(pole position) = 350 to 430 ( $\approx$  400) MeV  
Breit-Wigner mass = 2030 to 2200 ( $\approx$  2100) MeV  
Breit-Wigner full width = 300 to 450 ( $\approx$  400) MeV

| <b><math>N(2060)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 7–12 %                         | 834         |
| $N\eta$                                 | 2–38 %                         | 729         |
| $N\omega$                               | 1–7 %                          | 600         |
| $\Lambda K$                             | 10–20 %                        | 644         |
| $\Sigma K$                              | 1–5 %                          | 593         |
| $N\pi\pi$                               | 12–52 %                        | 814         |
| $\Delta(1232)\pi$ , $D$ -wave           | 4–10 %                         | 680         |
| $N\rho$                                 | 5–33 %                         | 605         |
| $N\rho$ , $S=1/2$ , $P$ -wave           | <10 %                          | 605         |
| $N\rho$ , $S=3/2$ , $D$ -wave           | 5–23 %                         | 605         |
| $N\sigma$                               | 3–9 %                          | —           |
| $N(1440)\pi$                            | 4–14 %                         | 544         |
| $N(1520)\pi$ , $P$ -wave                | 9–21 %                         | 490         |
| $N(1680)\pi$ , $S$ -wave                | 8–22 %                         | 353         |
| $\Lambda K^*(892)$                      | 0.3–1.3 %                      | 307         |
| $p\gamma$                               | 0.03–0.19 %                    | 840         |
| $p\gamma$ , helicity=1/2                | 0.02–0.08 %                    | 840         |
| $p\gamma$ , helicity=3/2                | 0.01–0.10 %                    | 840         |
| $n\gamma$                               | 0.003–0.07 %                   | 840         |
| $n\gamma$ , helicity=1/2                | 0.001–0.02 %                   | 840         |
| $n\gamma$ , helicity=3/2                | 0.002–0.05 %                   | 840         |

**$N(2100) 1/2^+$**

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Re(pole position) = 2050 to 2150 ( $\approx 2100$ ) MeV  
–2Im(pole position) = 240 to 340 ( $\approx 300$ ) MeV  
Breit-Wigner mass = 2050 to 2150 ( $\approx 2100$ ) MeV  
Breit-Wigner full width = 200 to 320 ( $\approx 260$ ) MeV

| <b><math>N(2100)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 8–32 %                         | 834         |
| $N\eta$                                 | 5–45 %                         | 729         |
| $N\eta'$                                | 5–11 %                         | 451         |
| $N\omega$                               | 10–25 %                        | 600         |
| $\Lambda K$                             | <1.0 %                         | 644         |
| $N\pi\pi$                               | >55 %                          | 814         |
| $\Delta(1232)\pi$ , $P$ -wave           | 6–14 %                         | 680         |
| $N\rho$ , $S=1/2$ , $P$ -wave           | 35–70                          | 605         |
| $N\sigma$                               | 14–35 %                        | –           |
| $N(1535)\pi$                            | 26–34 %                        | 478         |
| $\Lambda K^*(892)$                      | 3–11 %                         | 307         |
| $p\gamma$ , helicity=1/2                | 0.001–0.13 %                   | 840         |
| $n\gamma$ , helicity=1/2                | 0.004–0.09 %                   | 840         |

$$N(2120) \ 3/2^-$$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Re(pole position) = 2050 to 2150 ( $\approx 2100$ ) MeV  
– 2Im(pole position) = 200 to 360 ( $\approx 280$ ) MeV  
Breit-Wigner mass = 2060 to 2160 ( $\approx 2120$ ) MeV  
Breit-Wigner full width = 260 to 360 ( $\approx 300$ ) MeV

| <b><math>N(2120)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 5–15 %                         | 846         |
| $N\eta$                                 | 1–5 %                          | 743         |
| $N\eta'$                                | 2–6 %                          | 474         |
| $N\omega$                               | 4–20 %                         | 617         |
| $\Lambda K$                             | 6–11 %                         | 660         |
| $N\pi\pi$                               | >27 %                          | 827         |
| $\Delta(1232)\pi$                       | >23 %                          | 693         |
| $\Delta(1232)\pi$ , $S$ -wave           | 15–70 %                        | 693         |
| $\Delta(1232)\pi$ , $D$ -wave           | 8–45 %                         | 693         |
| $N\rho$ , $S=3/2$ , $S$ -wave           | < 3 %                          | 622         |
| $N\sigma$                               | 4–15 %                         | –           |
| $N(1535)\pi$                            | 7–23 %                         | 494         |
| $\Lambda K^*(892)$                      | < 0.2 %                        | 339         |
| $p\gamma$                               | 0.16–2.1 %                     | 852         |
| $p\gamma$ , helicity=1/2                | 0.07–0.80 %                    | 852         |
| $p\gamma$ , helicity=3/2                | 0.09–1.3 %                     | 852         |

|                          |              |     |
|--------------------------|--------------|-----|
| $n\gamma$                | 0.04–0.72 %  | 852 |
| $n\gamma$ , helicity=1/2 | 0.04–0.60 %  | 852 |
| $n\gamma$ , helicity=3/2 | 0.001–0.12 % | 852 |

**$N(2190) \ 7/2^-$**

$$I(J^P) = \frac{1}{2}(\frac{7}{2}^-)$$

Re(pole position) = 1950 to 2150 ( $\approx$  2050) MeV  
–2Im(pole position) = 300 to 500 ( $\approx$  400) MeV  
Breit-Wigner mass = 2140 to 2220 ( $\approx$  2180) MeV  
Breit-Wigner full width = 300 to 500 ( $\approx$  400) MeV

| <b><math>N(2190)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 10–20 %                        | 882         |
| $N\eta$                                 | 1–5 %                          | 785         |
| $N\omega$                               | 8–20 %                         | 667         |
| $\Lambda K$                             | 0.2–0.8 %                      | 705         |
| $N\pi\pi$                               | 22–51 %                        | 864         |
| $\Delta(1232)\pi$ , $D$ -wave           | 19–31 %                        | 734         |
| $N\rho$ , $S=3/2$ , $D$ -wave           | <11 %                          | 672         |
| $N\sigma$                               | 3–9 %                          | –           |
| $\Lambda K^*(892)$                      | 0.2–0.8 %                      | 423         |
| $p\gamma$                               | <0.08 %                        | 888         |
| $p\gamma$ , helicity=1/2                | <0.06 %                        | 888         |
| $p\gamma$ , helicity=3/2                | <0.02 %                        | 888         |
| $n\gamma$                               | <0.04 %                        | 888         |
| $n\gamma$ , helicity=1/2                | <0.01 %                        | 888         |
| $n\gamma$ , helicity=3/2                | <0.03 %                        | 888         |

**$N(2220) \ 9/2^+$**

$$I(J^P) = \frac{1}{2}(\frac{9}{2}^+)$$

Re(pole position) = 2130 to 2200 ( $\approx$  2150) MeV  
–2Im(pole position) = 360 to 480 ( $\approx$  400) MeV  
Breit-Wigner mass = 2200 to 2300 ( $\approx$  2250) MeV  
Breit-Wigner full width = 350 to 500 ( $\approx$  400) MeV

| <b><math>N(2220)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 15–30 %                        | 924         |

$N(2250) \ 9/2^-$

$I(J^P) = \frac{1}{2}(\frac{9}{2}^-)$

Re(pole position) = 2100 to 2200 ( $\approx 2150$ ) MeV  
– 2Im(pole position) = 350 to 500 ( $\approx 420$ ) MeV  
Breit-Wigner mass = 2250 to 2320 ( $\approx 2280$ ) MeV  
Breit-Wigner full width = 300 to 600 ( $\approx 500$ ) MeV

| <b><math>N(2250)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 5–15 %                         | 941         |
| $N\eta$                                 | <5 %                           | 852         |
| $\Lambda K$                             | 1–3 %                          | 777         |

$N(2600) \ 11/2^-$

$I(J^P) = \frac{1}{2}(\frac{11}{2}^-)$

Breit-Wigner mass = 2550 to 2750 ( $\approx 2600$ ) MeV  
Breit-Wigner full width = 500 to 800 ( $\approx 650$ ) MeV

| <b><math>N(2600)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\pi$                                  | 3–8 %                          | 1126        |

$\Delta$  BARYONS

$(S=0, I=3/2)$

$\Delta^{++} = uuu, \ \Delta^+ = uud, \ \Delta^0 = udd, \ \Delta^- = ddd$

$\Delta(1232) \ 3/2^+$

$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$

Re(pole position) = 1209 to 1211 ( $\approx 1210$ ) MeV  
– 2Im(pole position) = 98 to 102 ( $\approx 100$ ) MeV  
Breit-Wigner mass (mixed charges) = 1230 to 1234 ( $\approx 1232$ ) MeV  
Breit-Wigner full width (mixed charges) = 114 to 120 ( $\approx 117$ ) MeV

| $\Delta(1232)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$                     | 99.4 %                         | 229         |
| $N\gamma$                  | 0.55–0.65 %                    | 259         |
| $N\gamma$ , helicity=1/2   | 0.11–0.13 %                    | 259         |
| $N\gamma$ , helicity=3/2   | 0.44–0.52 %                    | 259         |
| $p e^+ e^-$                | $(4.2 \pm 0.7) \times 10^{-5}$ | 259         |

$\Delta(1600) \ 3/2^+$

$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$

Re(pole position) = 1470 to 1590 ( $\approx 1520$ ) MeV  
–2Im(pole position) = 150 to 320 ( $\approx 280$ ) MeV  
Breit-Wigner mass = 1500 to 1640 ( $\approx 1570$ ) MeV  
Breit-Wigner full width = 200 to 300 ( $\approx 250$ ) MeV

| $\Delta(1600)$ DECAY MODES    | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------|--------------------------------|-------------|
| $N\pi$                        | 8–24%                          | 492         |
| $N\pi\pi$                     | 58–84 %                        | 454         |
| $\Delta(1232)\pi$             | 58–82 %                        | 276         |
| $\Delta(1232)\pi$ , $P$ -wave | 72–82%                         | 276         |
| $\Delta(1232)\pi$ , $F$ -wave | <2%                            | 276         |
| $N(1440)\pi$                  | 17–27%                         | †           |
| $N\gamma$                     | 0.001–0.035 %                  | 505         |
| $N\gamma$ , helicity=1/2      | 0.0–0.02 %                     | 505         |
| $N\gamma$ , helicity=3/2      | 0.001–0.015 %                  | 505         |

$\Delta(1620) \ 1/2^-$

$I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$

Re(pole position) = 1590 to 1610 ( $\approx 1600$ ) MeV  
–2Im(pole position) = 80 to 140 ( $\approx 110$ ) MeV  
Breit-Wigner mass = 1590 to 1630 ( $\approx 1610$ ) MeV  
Breit-Wigner full width = 110 to 150 ( $\approx 130$ ) MeV

| <b><math>\Delta(1620)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------------------|--------------------------------|-------------|
| $N\pi$                                       | 25–35 %                        | 520         |
| $N\pi\pi$                                    | >67 %                          | 484         |
| $\Delta(1232)\pi$ , $D$ -wave                | 44–72 %                        | 311         |
| $N\rho$                                      | 23–32%                         | †           |
| $N\rho$ , $S=1/2$ , $S$ -wave                | 23–32%                         | †           |
| $N\rho$ , $S=3/2$ , $D$ -wave                | <0.04%                         | †           |
| $N(1440)\pi$                                 | <9 %                           | 98          |
| $N\gamma$ , helicity=1/2                     | 0.03–0.10 %                    | 532         |

$$\Delta(1700) \ 3/2^-$$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^-)$$

Re(pole position) = 1640 to 1690 ( $\approx 1665$ ) MeV  
– 2Im(pole position) = 200 to 300 ( $\approx 250$ ) MeV  
Breit-Wigner mass = 1690 to 1730 ( $\approx 1710$ ) MeV  
Breit-Wigner full width = 220 to 380 ( $\approx 300$ ) MeV

| <b><math>\Delta(1700)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------------------|--------------------------------|-------------|
| $N\pi$                                       | 10–20 %                        | 588         |
| $N\pi\pi$                                    | >31 %                          | 557         |
| $\Delta(1232)\pi$                            | 9–70 %                         | 394         |
| $\Delta(1232)\pi$ , $S$ -wave                | 5–54 %                         | 394         |
| $\Delta(1232)\pi$ , $D$ -wave                | 4–16 %                         | 394         |
| $N\rho$ , $S=3/2$ , $S$ -wave                | 22–32%                         | †           |
| $N(1520)\pi$ , $P$ -wave                     | 1–5 %                          | 133         |
| $N(1535)\pi$                                 | 0.5–1.5 %                      | 113         |
| $\Delta(1232)\eta$                           | 3–7 %                          | †           |
| $N\gamma$                                    | 0.22–0.60 %                    | 598         |
| $N\gamma$ , helicity=1/2                     | 0.12–0.30 %                    | 598         |
| $N\gamma$ , helicity=3/2                     | 0.10–0.30 %                    | 598         |

$$\Delta(1900) \ 1/2^-$$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$$

Re(pole position) = 1830 to 1900 ( $\approx 1865$ ) MeV  
– 2Im(pole position) = 180 to 300 ( $\approx 240$ ) MeV  
Breit-Wigner mass = 1840 to 1920 ( $\approx 1860$ ) MeV  
Breit-Wigner full width = 180 to 320 ( $\approx 250$ ) MeV

| $\Delta(1900)$ DECAY MODES    | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------|--------------------------------|-------------|
| $N\pi$                        | 4–12%                          | 685         |
| $\Sigma K$                    | seen                           | 367         |
| $N\pi\pi$                     | > 52%                          | 660         |
| $\Delta(1232)\pi$ , $D$ -wave | 30–70%                         | 509         |
| $N\rho$                       | 22–60 %                        | 361         |
| $N\rho$ , $S=1/2$ , $S$ -wave | 11–35%                         | 361         |
| $N\rho$ , $S=3/2$ , $D$ -wave | 11–25%                         | 361         |
| $N(1440)\pi$                  | 3–32%                          | 353         |
| $N(1520)\pi$                  | 2–10%                          | 288         |
| $\Delta(1232)\eta$            | < 2%                           | 251         |
| $N\gamma$ , helicity=1/2      | 0.06–0.43 %                    | 693         |

$\Delta(1905) \ 5/2^+$

$$I(J^P) = \frac{3}{2}(\frac{5}{2}^+)$$

Re(pole position) = 1750 to 1800 ( $\approx 1770$ ) MeV  
– 2Im(pole position) = 260 to 340 ( $\approx 300$ ) MeV  
Breit-Wigner mass = 1855 to 1910 ( $\approx 1880$ ) MeV  
Breit-Wigner full width = 270 to 400 ( $\approx 330$ ) MeV

| $\Delta(1905)$ DECAY MODES    | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------|--------------------------------|-------------|
| $N\pi$                        | 9–15%                          | 698         |
| $N\pi\pi$                     | > 65%                          | 673         |
| $\Delta(1232)\pi$             | > 48%                          | 524         |
| $\Delta(1232)\pi$ , $P$ -wave | 8–43%                          | 524         |
| $\Delta(1232)\pi$ , $F$ -wave | 40–58%                         | 524         |
| $N\rho$ , $S=3/2$ , $P$ -wave | 17–35%                         | 385         |
| $N(1535)\pi$                  | < 1 %                          | 293         |
| $N(1680)\pi$ , $P$ -wave      | 5–15%                          | 133         |
| $\Delta(1232)\eta$            | 2–6%                           | 282         |
| $N\gamma$                     | 0.012–0.036 %                  | 706         |
| $N\gamma$ , helicity=1/2      | 0.002–0.006 %                  | 706         |
| $N\gamma$ , helicity=3/2      | 0.01–0.03 %                    | 706         |

$\Delta(1910) \ 1/2^+$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^+)$$

Re(pole position) = 1800 to 1900 ( $\approx 1850$ ) MeV  
– 2Im(pole position) = 200 to 500 ( $\approx 350$ ) MeV  
Breit-Wigner mass = 1850 to 1950 ( $\approx 1900$ ) MeV  
Breit-Wigner full width = 200 to 400 ( $\approx 300$ ) MeV

| <b><math>\Delta(1910)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------------------|--------------------------------|-------------|
| $N\pi$                                       | 10–30%                         | 710         |
| $\Sigma K$                                   | 4–14%                          | 410         |
| $\Delta(1232)\pi$                            | 34–66%                         | 539         |
| $N(1440)\pi$                                 | 3–45%                          | 386         |
| $\Delta(1232)\eta$                           | 5–13%                          | 310         |
| $N\gamma$ , helicity=1/2                     | 0.0–0.02 %                     | 718         |

**$\Delta(1920)$   $3/2^+$**

$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$

Re(pole position) = 1850 to 1950 ( $\approx 1900$ ) MeV  
– 2Im(pole position) = 200 to 400 ( $\approx 300$ ) MeV  
Breit-Wigner mass = 1870 to 1970 ( $\approx 1920$ ) MeV  
Breit-Wigner full width = 240 to 360 ( $\approx 300$ ) MeV

| <b><math>\Delta(1920)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------------------|--------------------------------|-------------|
| $N\pi$                                       | 5–20 %                         | 723         |
| $\Sigma K$                                   | 2–6 %                          | 431         |
| $N\pi\pi$                                    | >46 %                          | 699         |
| $\Delta(1232)\pi$                            | >46 %                          | 553         |
| $\Delta(1232)\pi$ , $P$ -wave                | 2–28 %                         | 553         |
| $\Delta(1232)\pi$ , $F$ -wave                | 44–72 %                        | 553         |
| $N(1440)\pi$ , $P$ -wave                     | 4–86 %                         | 403         |
| $N(1520)\pi$ , $S$ -wave                     | <5 %                           | 341         |
| $N(1535)\pi$                                 | <2 %                           | 328         |
| $N a_0(980)$                                 | seen                           | 41          |
| $\Delta(1232)\eta$                           | 5–17 %                         | 336         |
| $N\gamma$                                    | 0.01–0.84 %                    | 731         |
| $N\gamma$ , helicity=1/2                     | 0.0–0.42 %                     | 731         |
| $N\gamma$ , helicity=3/2                     | 0.01–0.42 %                    | 731         |

**$\Delta(1930)$   $5/2^-$**

$I(J^P) = \frac{3}{2}(\frac{5}{2}^-)$

Re(pole position) = 1820 to 1880 ( $\approx 1850$ ) MeV  
– 2Im(pole position) = 300 to 450 ( $\approx 320$ ) MeV  
Breit-Wigner mass = 1900 to 2000 ( $\approx 1950$ ) MeV  
Breit-Wigner full width = 200 to 400 ( $\approx 300$ ) MeV

| $\Delta(1930)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$                     | 5–15 %                         | 742         |
| $N\gamma$                  | 0.0–0.01 %                     | 749         |
| $N\gamma$ , helicity=1/2   | 0.0–0.005 %                    | 749         |
| $N\gamma$ , helicity=3/2   | 0.0–0.004 %                    | 749         |

$\Delta(1950) \ 7/2^+$

$I(J^P) = \frac{3}{2}(\frac{7}{2}^+)$

Re(pole position) = 1870 to 1890 ( $\approx 1880$ ) MeV  
– 2Im(pole position) = 220 to 260 ( $\approx 240$ ) MeV  
Breit-Wigner mass = 1915 to 1950 ( $\approx 1930$ ) MeV  
Breit-Wigner full width = 235 to 335 ( $\approx 285$ ) MeV

| $\Delta(1950)$ DECAY MODES    | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------|--------------------------------|-------------|
| $N\pi$                        | 35–45 %                        | 729         |
| $\Sigma K$                    | 0.3–0.5 %                      | 441         |
| $N\pi\pi$                     | 37–77 %                        | 706         |
| $\Delta(1232)\pi$ , $F$ -wave | 1–9 %                          | 560         |
| $N(1680)\pi$ , $P$ -wave      | 3–9 %                          | 191         |
| $\Delta(1232)\eta$            | < 0.6 %                        | 349         |
| $N\gamma$                     | 0.06–0.14 %                    | 737         |
| $N\gamma$ , helicity=1/2      | 0.03–0.05 %                    | 737         |
| $N\gamma$ , helicity=3/2      | 0.04–0.09 %                    | 737         |

$\Delta(2200) \ 7/2^-$

$I(J^P) = \frac{3}{2}(\frac{7}{2}^-)$

Re(pole position) = 2050 to 2150 ( $\approx 2100$ ) MeV  
– 2Im(pole position) = 260 to 420 ( $\approx 340$ ) MeV  
Breit-Wigner mass = 2150 to 2250 ( $\approx 2200$ ) MeV  
Breit-Wigner full width = 200 to 500 ( $\approx 350$ ) MeV

| $\Delta(2200)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$                     | 2–8 %                          | 894         |
| $\Sigma K$                 | 1–7 %                          | 672         |
| $N\pi\pi$                  | >45 %                          | 876         |
| $\Delta\pi$                | >45 %                          | 747         |
| $\Delta\pi$ , $D$ -wave    | >40 %                          | 747         |
| $\Delta\pi$ , $G$ -wave    | 5–25 %                         | 747         |
| $\Delta\eta$ , $D$ -wave   | seen                           | 614         |

$\Delta(2420)$   $11/2^+$

$I(J^P) = \frac{3}{2}(\frac{11}{2}^+)$

Re(pole position) = 2300 to 2500 ( $\approx 2400$ ) MeV  
–2Im(pole position) = 350 to 550 ( $\approx 450$ ) MeV  
Breit-Wigner mass = 2300 to 2600 ( $\approx 2450$ ) MeV  
Breit-Wigner full width = 300 to 700 ( $\approx 500$ ) MeV

| $\Delta(2420)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\pi$                     | 5–10 %                         | 1040        |

$\Lambda$  BARYONS

$(S = -1, I = 0)$

$\Lambda^0 = uds$

$\Lambda$

$I(J^P) = 0(\frac{1}{2}^+)$

Mass  $m = 1115.683 \pm 0.006$  MeV  
 $(m_\Lambda - m_{\bar\Lambda}) / m_\Lambda = (-0.1 \pm 1.1) \times 10^{-5}$  ( $S = 1.6$ )  
Mean life  $\tau = (2.617 \pm 0.010) \times 10^{-10}$  s ( $S = 1.5$ )  
 $(\tau_\Lambda - \tau_{\bar\Lambda}) / \tau_\Lambda = (0.9 \pm 3.2) \times 10^{-3}$   
 $c\tau = 7.845$  cm  
Magnetic moment  $\mu = -0.613 \pm 0.004 \mu_N$   
Electric dipole moment  $d < 1.5 \times 10^{-16}$  ecm, CL = 95%

**Decay parameters**

$p\pi^-$        $\alpha_- = 0.749 \pm 0.008$  (S = 2.4)  
 $\bar{p}\pi^+$        $\alpha_+ = -0.758 \pm 0.005$  (S = 1.2)  
 $\bar{n}\pi^0$        $\bar{\alpha}_0 = -0.675 \pm 0.011$  (S = 1.2)  
 $n\pi^0$        $\alpha_0 = 0.670 \pm 0.012$   
 $n\gamma$        $\alpha_\gamma = -0.16 \pm 0.11$   
 $p\pi^-$        $\phi_- = (-6.5 \pm 3.5)^\circ$   
"       $\gamma_- = 0.76$  [l]  
"       $\Delta_- = (8 \pm 4)^\circ$  [l]  
 $\bar{\alpha}_0 / \alpha_+$  in  $\bar{\Lambda} \rightarrow \bar{n}\pi^0, \bar{\Lambda} \rightarrow \bar{p}\pi^+ = 0.876_{-0.020}^{+0.022}$  (S = 1.4)  
 $(\alpha_0 + \bar{\alpha}_0)/(\alpha_0 - \bar{\alpha}_0)$  in  $\Lambda \rightarrow n\pi^0, \bar{\Lambda} \rightarrow \bar{n}\pi^0 = (0.1_{-1.1}^{+1.0}) \times 10^{-2}$   
 $R = |G_E/G_M|$  in  $\Lambda \rightarrow p\pi^-, \bar{\Lambda} \rightarrow \bar{p}\pi^+ = 0.96 \pm 0.14$   
 $\Delta\Phi = \Phi_E - \Phi_M$  in  $\Lambda \rightarrow p\pi^-, \bar{\Lambda} \rightarrow \bar{p}\pi^+ = 37 \pm 13$  degrees  
 $p e^- \bar{\nu}_e$        $g_A/g_V = -0.718 \pm 0.015$  [h]

| <b><math>\Lambda</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ )         | Confidence level | $P$<br>(MeV/c) |
|-----------------------------------------|----------------------------------------|------------------|----------------|
| $p\pi^-$                                | (64.1 $\pm$ 0.5 ) %                    |                  | 101            |
| $n\pi^0$                                | (35.9 $\pm$ 0.5 ) %                    |                  | 104            |
| $n\gamma$                               | ( 8.3 $\pm$ 0.7 ) $\times 10^{-4}$     |                  | 162            |
| $p\pi^-\gamma$                          | [n] ( 8.5 $\pm$ 1.4 ) $\times 10^{-4}$ |                  | 101            |
| $p e^- \bar{\nu}_e$                     | ( 8.34 $\pm$ 0.14) $\times 10^{-4}$    |                  | 163            |
| $p\mu^-\bar{\nu}_\mu$                   | ( 1.51 $\pm$ 0.19) $\times 10^{-4}$    |                  | 131            |

**Lepton (L) and/or Baryon (B) number violating decay modes**

|                |     |       |                  |     |     |
|----------------|-----|-------|------------------|-----|-----|
| $\pi^+ e^-$    | L,B | < 6   | $\times 10^{-7}$ | 90% | 549 |
| $\pi^+ \mu^-$  | L,B | < 6   | $\times 10^{-7}$ | 90% | 544 |
| $\pi^- e^+$    | L,B | < 4   | $\times 10^{-7}$ | 90% | 549 |
| $\pi^- \mu^+$  | L,B | < 6   | $\times 10^{-7}$ | 90% | 544 |
| $K^+ e^-$      | L,B | < 2   | $\times 10^{-6}$ | 90% | 449 |
| $K^+ \mu^-$    | L,B | < 3   | $\times 10^{-6}$ | 90% | 441 |
| $K^- e^+$      | L,B | < 2   | $\times 10^{-6}$ | 90% | 449 |
| $K^- \mu^+$    | L,B | < 3   | $\times 10^{-6}$ | 90% | 441 |
| $K_S^0 \nu$    | L,B | < 2   | $\times 10^{-5}$ | 90% | 447 |
| $\bar{p}\pi^+$ | B   | < 9   | $\times 10^{-7}$ | 90% | 101 |
| invisible      |     | < 7.4 | $\times 10^{-5}$ | 90% | —   |

**$\Lambda(1405) 1/2^-$**

$I(J^P) = 0(\frac{1}{2}^-)$

Mass  $m = 1405.1_{-1.0}^{+1.3}$  MeV  
Full width  $\Gamma = 50.5 \pm 2.0$  MeV  
Below  $\bar{K}N$  threshold

| <b><math>\Lambda(1405)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $\Sigma \pi$                                  | 100 %                          | 155         |

$\Lambda(1520) \ 3/2^-$

$I(J^P) = 0(\frac{3}{2}^-)$

Mass  $m = 1518$  to  $1520$  ( $\approx 1519$ ) MeV [o]  
Full width  $\Gamma = 15$  to  $17$  ( $\approx 16$ ) MeV [o]

| <b><math>\Lambda(1520)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $N \overline{K}$                              | (45 $\pm$ 1 ) %                | 242         |
| $\Sigma \pi$                                  | (42 $\pm$ 1 ) %                | 268         |
| $\Lambda \pi \pi$                             | (10 $\pm$ 1 ) %                | 259         |
| $\Sigma \pi \pi$                              | ( 0.9 $\pm$ 0.1 ) %            | 168         |
| $\Lambda \gamma$                              | ( 0.85 $\pm$ 0.15 ) %          | 350         |

$\Lambda(1600) \ 1/2^+$

$I(J^P) = 0(\frac{1}{2}^+)$

Mass  $m = 1570$  to  $1630$  ( $\approx 1600$ ) MeV  
Full width  $\Gamma = 150$  to  $250$  ( $\approx 200$ ) MeV

| <b><math>\Lambda(1600)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $N \overline{K}$                              | 15–30 %                        | 343         |
| $\Sigma \pi$                                  | 10–60 %                        | 338         |
| $\Lambda \sigma$                              | (19 $\pm$ 4) %                 | —           |
| $\Sigma(1385) \pi$                            | ( 9 $\pm$ 4 ) %                | 158         |

$\Lambda(1670) \ 1/2^-$

$I(J^P) = 0(\frac{1}{2}^-)$

Mass  $m = 1670$  to  $1678$  ( $\approx 1674$ ) MeV  
Full width  $\Gamma = 25$  to  $35$  ( $\approx 30$ ) MeV

| <b><math>\Lambda(1670)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $N\overline{K}$                               | 20–30 %                        | 418         |
| $\Sigma\pi$                                   | 25–55 %                        | 398         |
| $\Lambda\eta$                                 | 10–25 %                        | 88          |
| $\Sigma(1385)\pi$ , $D$ -wave                 | ( 6.0 $\pm$ 2.0 ) %            | 235         |
| $N\overline{K}^*(892)$ , $S=3/2$ , $D$ -wave  | ( 5 $\pm$ 4 ) %                | †           |
| $\Lambda\sigma$                               | (20 $\pm$ 8 ) %                | –           |

$\Lambda(1690) \ 3/2^-$

$I(J^P) = 0(\frac{3}{2}^-)$

Mass  $m = 1685$  to  $1695$  ( $\approx 1690$ ) MeV  
Full width  $\Gamma = 60$  to  $80$  ( $\approx 70$ ) MeV

| <b><math>\Lambda(1690)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $N\overline{K}$                               | 20–30 %                        | 433         |
| $\Sigma\pi$                                   | 20–40 %                        | 410         |
| $\Lambda\sigma$                               | (5.0 $\pm$ 2.0) %              | –           |
| $\Lambda\pi\pi$                               | $\sim 25$ %                    | 419         |
| $\Sigma\pi\pi$                                | $\sim 20$ %                    | 358         |
| $\Sigma(1385)\pi$ , $S$ -wave                 | (9 $\pm 5$ ) %                 | 251         |
| $\Sigma(1385)\pi$ , $D$ -wave                 | (3.0 $\pm$ 2.0) %              | 251         |

$\Lambda(1800) \ 1/2^-$

$I(J^P) = 0(\frac{1}{2}^-)$

Mass  $m = 1750$  to  $1850$  ( $\approx 1800$ ) MeV  
Full width  $\Gamma = 150$  to  $250$  ( $\approx 200$ ) MeV

| <b><math>\Lambda(1800)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $N\overline{K}$                               | 25–40 %                        | 528         |
| $\Sigma\pi$                                   | (27 $\pm$ 6) %                 | 494         |
| $\Lambda\sigma$                               | (15 $\pm$ 4) %                 | –           |
| $\Sigma(1385)\pi$                             | ( 9 $\pm$ 4 ) %                | 349         |
| $\Lambda\eta$                                 | 0.01 to 0.10                   | 326         |
| $N\overline{K}^*(892)$                        | seen                           | †           |

$\Lambda(1810) \ 1/2^+$

$I(J^P) = 0(\frac{1}{2}^+)$

Mass  $m = 1740$  to  $1840$  ( $\approx 1790$ ) MeV  
Full width  $\Gamma = 50$  to  $170$  ( $\approx 110$ ) MeV

| <b><math>\Lambda(1810)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $N\overline{K}$                               | 0.05 to 0.35                   | 520         |
| $\Sigma\pi$                                   | (16 $\pm$ 5) %                 | 487         |
| $\Sigma(1385)\pi$                             | (40 $\pm$ 15) %                | 340         |
| $N\overline{K}^*(892)$                        | 30–60 %                        | †           |

$\Lambda(1820) \ 5/2^+$

$I(J^P) = 0(\frac{5}{2}^+)$

Mass  $m = 1815$  to  $1825$  ( $\approx 1820$ ) MeV  
Full width  $\Gamma = 70$  to  $90$  ( $\approx 80$ ) MeV

| <b><math>\Lambda(1820)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------------|--------------------------------|-------------|
| $N\overline{K}$                               | 55–65 %                        | 545         |
| $\Sigma\pi$                                   | 8–14 %                         | 509         |
| $\Sigma(1385)\pi$                             | 5–10 %                         | 366         |
| $N\overline{K}^*(892)$ , $S=3/2$ , $P$ -wave  | (3.0 $\pm$ 1.0) %              | †           |

$\Lambda(1830) \ 5/2^-$

$I(J^P) = 0(\frac{5}{2}^-)$

Mass  $m = 1820$  to  $1830$  ( $\approx 1825$ ) MeV  
Full width  $\Gamma = 60$  to  $120$  ( $\approx 90$ ) MeV

| <b><math>\Lambda(1830)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | Scale factor | $p$<br>(MeV/c) |
|-----------------------------------------------|--------------------------------|--------------|----------------|
| $N\overline{K}$                               | 0.04 to 0.08                   |              | 549            |
| $\Sigma\pi$                                   | 35–75 %                        |              | 512            |
| $\Sigma(1385)\pi$                             | >15 %                          |              | 370            |
| $\Sigma(1385)\pi$ , $D$ -wave                 | (40 $\pm$ 15) %                | 3.2          | 370            |

$\Lambda(1890) \ 3/2^+$

$I(J^P) = 0(\frac{3}{2}^+)$

Mass  $m = 1870$  to  $1910$  ( $\approx 1890$ ) MeV  
Full width  $\Gamma = 80$  to  $160$  ( $\approx 120$ ) MeV

| $\Lambda(1890)$ DECAY MODES   | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------|--------------------------------|-------------|
| $N\overline{K}$               | 0.24 to 0.36                   | 599         |
| $\Sigma\pi$                   | 3–10 %                         | 560         |
| $\Sigma(1385)\pi$             | seen                           | 423         |
| $\Sigma(1385)\pi$ , $P$ -wave | $(6.0 \pm 3.0) \%$             | 423         |
| $\Sigma(1385)\pi$ , $F$ -wave | $(4.0 \pm 2.0) \%$             | 423         |
| $N\overline{K}^*(892)$        | seen                           | 236         |

$\Lambda(2100) \ 7/2^-$

$I(J^P) = 0(\frac{7}{2}^-)$

Mass  $m = 2090$  to  $2110$  ( $\approx 2100$ ) MeV  
Full width  $\Gamma = 100$  to  $250$  ( $\approx 200$ ) MeV

| $\Lambda(2100)$ DECAY MODES                  | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------------------|--------------------------------|-------------|
| $N\overline{K}$                              | 25–35 %                        | 751         |
| $\Sigma\pi$                                  | $\sim 5 \%$                    | 705         |
| $\Lambda\eta$                                | $< 3 \%$                       | 617         |
| $\Xi K$                                      | $< 3 \%$                       | 491         |
| $\Lambda\omega$                              | $< 8 \%$                       | 443         |
| $\Sigma(1385)\pi$ , $G$ -wave                | $(1.0 \pm 1.0) \%$             | 584         |
| $N\overline{K}^*(892)$                       | 10–20 %                        | 515         |
| $N\overline{K}^*(892)$ , $S=3/2$ , $D$ -wave | $(4.0 \pm 2.0) \%$             | 515         |

$\Lambda(2110) \ 5/2^+$

$I(J^P) = 0(\frac{5}{2}^+)$

Mass  $m = 2050$  to  $2130$  ( $\approx 2090$ ) MeV  
Full width  $\Gamma = 200$  to  $300$  ( $\approx 250$ ) MeV

| $\Lambda(2110)$ DECAY MODES           | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------------|--------------------------------|-------------|
| $N\overline{K}$                       | 5–25 %                         | 744         |
| $\Sigma\pi$                           | 10–40 %                        | 698         |
| $\Lambda\omega$                       | seen                           | 432         |
| $\Lambda\omega$ , $S=3/2$ , $P$ -wave | $(5.0\pm 2.0)$ %               | 432         |
| $\Sigma(1385)\pi$                     | seen                           | 576         |
| $N\overline{K}^*(892)$                | 10–60 %                        | 505         |

$\Lambda(2350) \ 9/2^+$

$I(J^P) = 0(\frac{9}{2}^+)$

Mass  $m = 2340$  to  $2370$  ( $\approx 2350$ ) MeV  
Full width  $\Gamma = 100$  to  $250$  ( $\approx 150$ ) MeV

| $\Lambda(2350)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------|--------------------------------|-------------|
| $N\overline{K}$             | $\sim 12$ %                    | 915         |
| $\Sigma\pi$                 | $\sim 10$ %                    | 867         |

$\Sigma$  BARYONS

$(S = -1, I = 1)$

$\Sigma^+ = uus, \quad \Sigma^0 = uds, \quad \Sigma^- = dds$

$\Sigma^+$

$I(J^P) = 1(\frac{1}{2}^+)$

Mass  $m = 1189.37 \pm 0.07$  MeV ( $S = 2.2$ )  
Mean life  $\tau = (0.8018 \pm 0.0026) \times 10^{-10}$  s  
 $c\tau = 2.404$  cm  
 $(\tau_{\Sigma^+} - \tau_{\Sigma^-}) / \tau_{\Sigma^+} = -0.0006 \pm 0.0012$   
Magnetic moment  $\mu = 2.458 \pm 0.010 \mu_N$  ( $S = 2.1$ )  
 $(\mu_{\Sigma^+} + \mu_{\Sigma^-}) / \mu_{\Sigma^+} = 0.014 \pm 0.015$   
 $\Gamma(\Sigma^+ \rightarrow n\ell^+\nu) / \Gamma(\Sigma^- \rightarrow n\ell^-\overline{\nu}_\ell) < 0.043$

**Decay parameters**

$p\pi^0 \quad \alpha_0 = -0.982 \pm 0.014$   
 $\bar{p}\pi^0 \quad \bar{\alpha}_0 = 0.99 \pm 0.04$   
 $(\alpha_0 + \bar{\alpha}_0) / (\alpha_0 - \bar{\alpha}_0) = 0.00 \pm 0.04$   
 $p\pi^0 \quad \phi_0 = (36 \pm 34)^\circ$   
"  $\gamma_0 = 0.16 [I]$   
"  $\Delta_0 = (187 \pm 6)^\circ [I]$   
 $n\pi^+ \quad \alpha_+ = (4.89 \pm 0.26) \times 10^{-2}$   
"  $\phi_+ = (167 \pm 20)^\circ \quad (S = 1.1)$   
 $\bar{\alpha}_- \text{ FOR } \bar{\Sigma}^- \rightarrow \bar{n}\pi^- = (-5.7 \pm 0.5) \times 10^{-2}$   
 $\bar{\alpha}_- / \bar{\alpha}_0 = (-5.7 \pm 0.6) \times 10^{-2}$   
 $(\alpha_+ + \bar{\alpha}_-) / (\alpha_+ - \bar{\alpha}_-) = (-8 \pm 6) \times 10^{-2}$   
"  $\gamma_+ = -0.97 [I]$   
"  $\Delta_+ = (-73^{+133}_{-10})^\circ [I]$   
 $p\gamma \quad \alpha_\gamma = -0.69 \pm 0.05$

| $\Sigma^+$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ )           | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|------------------------|------------------------------------------|-----------------------------------|----------------|
| $p\pi^0$               | $(51.47 \pm 0.30) \%$                    |                                   | 189            |
| $n\pi^+$               | $(48.43 \pm 0.30) \%$                    |                                   | 185            |
| $p\gamma$              | $(1.04 \pm 0.06) \times 10^{-3}$         | $S=2.4$                           | 225            |
| $n\pi^+\gamma$         | $[n] \quad (4.5 \pm 0.5) \times 10^{-4}$ |                                   | 185            |
| $\Lambda e^+ \nu_e$    | $(2.3 \pm 0.4) \times 10^{-5}$           |                                   | 71             |

**$\Delta S = \Delta Q$  (SQ) violating modes or  
 $\Delta S = 1$  weak neutral current (S1) modes**


|                  |    |                                      |        |     |
|------------------|----|--------------------------------------|--------|-----|
| $ne^+ \nu_e$     | SQ | $< 5 \times 10^{-6}$                 | CL=90% | 224 |
| $n\mu^+ \nu_\mu$ | SQ | $< 3.0 \times 10^{-5}$               | CL=90% | 202 |
| $pe^+ e^-$       | S1 | $< 7 \times 10^{-6}$                 |        | 225 |
| $p\mu^+ \mu^-$   | S1 | $(2.4^{+1.7}_{-1.3}) \times 10^{-8}$ |        | 121 |
| $p\gamma_{dark}$ |    | $< 3.2 \times 10^{-5}$               | CL=90% | —   |



$I(J^P) = 1(\frac{1}{2}^+)$

Mass  $m = 1192.642 \pm 0.024$  MeV  
 $m_{\Sigma^-} - m_{\Sigma^0} = 4.807 \pm 0.035$  MeV  $(S = 1.1)$   
 $m_{\Sigma^0} - m_\Lambda = 76.959 \pm 0.023$  MeV  
Mean life  $\tau = (7.4 \pm 0.7) \times 10^{-20}$  s  
 $c\tau = 2.22 \times 10^{-11}$  m  
Transition magnetic moment  $|\mu_{\Sigma\Lambda}| = 1.61 \pm 0.08 \mu_N$   
Magnetic moment  $\mu = (-1.7 \pm 2.8) \times 10^{-3}$   
Magnetic moment  $\mu = (2.1 \pm 3.0) \times 10^{-3}$   
Magnetic moment  $\mu = (0.4 \pm 3.2) \times 10^{-3}$

| $\Sigma^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $\rho$<br>(MeV/c) |
|------------------------|--------------------------------|------------------|-------------------|
| $\Lambda\gamma$        | 100 %                          |                  | 74                |
| $\Lambda\gamma\gamma$  | < 3 %                          | 90%              | 74                |
| $\Lambda e^+ e^-$      | [p] $5 \times 10^{-3}$         |                  | 74                |



$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass  $m = 1197.449 \pm 0.029$  MeV (S = 1.1)  
 $m_{\Sigma^-} - m_{\Sigma^+} = 8.08 \pm 0.08$  MeV (S = 1.9)  
 $m_{\Sigma^-} - m_{\Lambda} = 81.766 \pm 0.029$  MeV (S = 1.1)  
Mean life  $\tau = (1.479 \pm 0.011) \times 10^{-10}$  s (S = 1.3)  
 $c\tau = 4.434$  cm  
Magnetic moment  $\mu = -1.160 \pm 0.025 \mu_N$  (S = 1.7)  
 $\Sigma^-$  charge radius =  $0.78 \pm 0.10$  fm

Decay parameters

$n\pi^-$   $\alpha_- = -0.068 \pm 0.008$   
"  $\phi_- = (10 \pm 15)^\circ$   
"  $\gamma_- = 0.98$  [l]  
"  $\Delta_- = (249^{+12}_{-120})^\circ$  [l]  
 $ne^- \bar{\nu}_e$   $g_A/g_V = 0.340 \pm 0.017$  [h]  
"  $f_2(0)/f_1(0) = 0.97 \pm 0.14$   
"  $D = 0.11 \pm 0.10$   
 $\Lambda e^- \bar{\nu}_e$   $g_V/g_A = 0.01 \pm 0.10$  [h] (S = 1.5)  
"  $g_{WM}/g_A = 2.4 \pm 1.7$  [h]

| $\Sigma^-$ DECAY MODES                | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$<br>(MeV/c) |
|---------------------------------------|--------------------------------|------------------|----------------|
| $n\pi^-$                              | $(99.848\pm0.005)\%$           |                  | 193            |
| $n\pi^-\gamma$                        | [n] $(4.6\pm0.6)\times10^{-4}$ |                  | 193            |
| $ne^-\bar{\nu}_e$                     | $(1.017\pm0.034)\times10^{-3}$ |                  | 230            |
| $n\mu^-\bar{\nu}_\mu$                 | $(4.5\pm0.4)\times10^{-4}$     |                  | 210            |
| $\Lambda e^-\bar{\nu}_e$              | $(5.73\pm0.27)\times10^{-5}$   |                  | 79             |
| $\Sigma^+\chi$                        | $<1.2\times10^{-4}$            | 90%              | —              |
| Lepton number ( $L$ ) violating modes |                                |                  |                |
| $pe^-e^-$                             | $L < 6.7\times10^{-5}$         | 90%              | 231            |

$\Sigma(1385) \ 3/2^+$

$I(J^P) = 1(\frac{3}{2}^+)$

$\Sigma(1385)^+$  mass  $m = 1382.83 \pm 0.34$  MeV ( $S = 1.9$ )  
 $\Sigma(1385)^0$  mass  $m = 1383.7 \pm 1.0$  MeV ( $S = 1.4$ )  
 $\Sigma(1385)^-$  mass  $m = 1387.2 \pm 0.5$  MeV ( $S = 2.2$ )  
 $\Sigma(1385)^+$  full width  $\Gamma = 36.2 \pm 0.7$  MeV  
 $\Sigma(1385)^0$  full width  $\Gamma = 36 \pm 5$  MeV  
 $\Sigma(1385)^-$  full width  $\Gamma = 39.4 \pm 2.1$  MeV ( $S = 1.7$ )  
Below  $\bar{K}N$  threshold

| $\Sigma(1385)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$<br>(MeV/c) |
|----------------------------|--------------------------------|------------------|----------------|
| $\Lambda\pi$               | $(87.0\pm1.5)\%$               |                  | 208            |
| $\Sigma\pi$                | $(11.7\pm1.5)\%$               |                  | 129            |
| $\Lambda\gamma$            | $(1.25^{+0.13}_{-0.12})\%$     |                  | 241            |
| $\Sigma^+\gamma$           | $(7.0\pm1.7)\times10^{-3}$     |                  | 180            |
| $\Sigma^-\gamma$           | $<2.4\times10^{-4}$            | 90%              | 173            |

$\Sigma(1660) \ 1/2^+$

$I(J^P) = 1(\frac{1}{2}^+)$

Re(pole position) =  $1585 \pm 20$  MeV  
 $-2\text{Im}(\text{pole position}) = 290^{+140}_{-40}$  MeV  
Mass  $m = 1640$  to  $1680$  ( $\approx 1660$ ) MeV  
Full width  $\Gamma = 100$  to  $300$  ( $\approx 200$ ) MeV

| $\Sigma(1660)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ )  | $p$ (MeV/c) |
|----------------------------|---------------------------------|-------------|
| $N\bar{K}$                 | 0.05 to 0.15 ( $\approx 0.10$ ) | 405         |
| $\Lambda\pi$               | (35 $\pm 12$ ) %                | 440         |
| $\Sigma\pi$                | (37 $\pm 10$ ) %                | 387         |
| $\Sigma\sigma$             | (20 $\pm 8$ ) %                 | —           |
| $\Lambda(1405)\pi$         | ( 4.0 $\pm$ 2.0) %              | 199         |

$\Sigma(1670) 3/2^-$

$I(J^P) = 1(\frac{3}{2}^-)$

Mass  $m = 1665$  to  $1685$  ( $\approx 1675$ ) MeV  
Full width  $\Gamma = 40$  to  $100$  ( $\approx 70$ ) MeV

| $\Sigma(1670)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------|--------------------------------|-------------|
| $N\bar{K}$                 | 0.06 to 0.12                   | 419         |
| $\Lambda\pi$               | 5–15 %                         | 452         |
| $\Sigma\pi$                | 30–60 %                        | 398         |
| $\Sigma\sigma$             | (7.0 $\pm$ 3.0) %              | —           |

$\Sigma(1750) 1/2^-$

$I(J^P) = 1(\frac{1}{2}^-)$

Mass  $m = 1700$  to  $1800$  ( $\approx 1750$ ) MeV  
Full width  $\Gamma = 100$  to  $200$  ( $\approx 150$ ) MeV

| $\Sigma(1750)$ DECAY MODES    | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------|--------------------------------|-------------|
| $N\bar{K}$                    | 0.06 to 0.12                   | 486         |
| $\Lambda\pi$                  | (14 $\pm 5$ ) %                | 507         |
| $\Sigma\pi$                   | (16 $\pm 4$ ) %                | 456         |
| $\Sigma\eta$                  | 15–55 %                        | 98          |
| $\Sigma(1385)\pi$ , $D$ -wave | < 1 %                          | 305         |
| $\Lambda(1520)\pi$            | ( 2.0 $\pm$ 1.0) %             | 175         |
| $N\bar{K}^*(892)$ , $S=1/2$   | ( 8 $\pm 4$ ) %                | †           |

$\Sigma(1775) 5/2^-$

$I(J^P) = 1(\frac{5}{2}^-)$

Mass  $m = 1770$  to  $1780$  ( $\approx 1775$ ) MeV  
Full width  $\Gamma = 105$  to  $135$  ( $\approx 120$ ) MeV

| <b><math>\Sigma(1775)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------------------|--------------------------------|-------------|
| $N\bar{K}$                                   | 37–43%                         | 508         |
| $\Lambda\pi$                                 | 14–20%                         | 525         |
| $\Sigma\pi$                                  | 2–5%                           | 475         |
| $\Sigma(1385)\pi$                            | 8–12%                          | 327         |
| $\Lambda(1520)\pi$ , $P$ -wave               | 17–23%                         | 202         |

$\Sigma(1910) \ 3/2^-$

$I(J^P) = 1(\frac{3}{2}^-)$

was  $\Sigma(1940)$

Mass  $m = 1870$  to  $1950$  ( $\approx 1910$ ) MeV  
Full width  $\Gamma = 150$  to  $300$  ( $\approx 220$ ) MeV

| <b><math>\Sigma(1910)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ )  | $p$ (MeV/c) |
|----------------------------------------------|---------------------------------|-------------|
| $N\bar{K}$                                   | 0.01 to 0.05 ( $\approx 0.02$ ) | 615         |
| $\Lambda\pi$                                 | ( 6 $\pm$ 4 ) %                 | 619         |
| $\Sigma\pi$                                  | (86 $\pm$ 21 ) %                | 574         |
| $\Sigma(1385)\pi$                            | seen                            | 439         |
| $\Lambda(1520)\pi$                           | seen                            | 329         |
| $\Delta(1232)\bar{K}$                        | ( 3.0 $\pm$ 1.0 ) %             | 377         |
| $N\bar{K}^*(892)$                            | seen                            | 274         |
| $N\bar{K}^*(892)$ , $S=1/2$ , $D$ -wave      | ( 1.0 $\pm$ 1.0 ) %             | 274         |

$\Sigma(1915) \ 5/2^+$

$I(J^P) = 1(\frac{5}{2}^+)$

Mass  $m = 1900$  to  $1935$  ( $\approx 1915$ ) MeV  
Full width  $\Gamma = 80$  to  $160$  ( $\approx 120$ ) MeV

| $\Sigma(1915)$ DECAY MODES              | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\bar{K}$                              | 0.05 to 0.15                   | 618         |
| $\Lambda\pi$                            | ( 6.0 $\pm$ 2.0 ) %            | 623         |
| $\Sigma\pi$                             | (10.0 $\pm$ 2.0) %             | 577         |
| $\Sigma(1385)\pi$ , $P$ -wave           | ( 2.0 $\pm$ 2.0 ) %            | 443         |
| $\Sigma(1385)\pi$ , $F$ -wave           | ( 4.0 $\pm$ 2.0 ) %            | 443         |
| $\Lambda(1520)\pi$ , $D$ -wave          | ( 8.0 $\pm$ 2.0 ) %            | 334         |
| $N\bar{K}^*(892)$ , $S=1/2$ , $F$ -wave | ( 5.0 $\pm$ 3.0 ) %            | 282         |
| $N\bar{K}^*(892)$ , $S=3/2$ , $F$ -wave | ( 5.0 $\pm$ 2.0 ) %            | 282         |
| $\Delta\bar{K}$ , $P$ -wave             | (16 $\pm$ 5 ) %                | 383         |
| $\Delta\bar{K}$ , $F$ -wave             | ( 5.0 $\pm$ 3.0 ) %            | 383         |

$\Sigma(2030) \ 7/2^+$

 $I(J^P) = 1(\frac{7}{2}^+)$

Mass  $m$  = 2025 to 2040 ( $\approx$  2030) MeV  
Full width  $\Gamma$  = 150 to 200 ( $\approx$  180) MeV

| $\Sigma(2030)$ DECAY MODES              | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------------------|--------------------------------|-------------|
| $N\bar{K}$                              | 17–23 %                        | 702         |
| $\Lambda\pi$                            | 17–23 %                        | 700         |
| $\Sigma\pi$                             | 5–10 %                         | 657         |
| $\Xi K$                                 | <2 %                           | 422         |
| $\Sigma(1385)\pi$                       | 5–15 %                         | 532         |
| $\Sigma(1385)\pi$ , $F$ -wave           | ( 1.0 $\pm$ 1.0 ) %            | 532         |
| $\Lambda(1520)\pi$                      | 10–20 %                        | 431         |
| $\Delta(1232)\bar{K}$                   | 10–20 %                        | 498         |
| $\Delta(1232)\bar{K}$ , $F$ -wave       | (15 $\pm$ 5 ) %                | 498         |
| $\Delta(1232)\bar{K}$ , $H$ -wave       | ( 1.0 $\pm$ 1.0 ) %            | 498         |
| $N\bar{K}^*(892)$ , $S=3/2$ , $F$ -wave | (14 $\pm$ 8 ) %                | 439         |

# $\Xi$ BARYONS

## ( $S = -2$ , $I = 1/2$ )

$$\Xi^0 = uss, \quad \Xi^- = dss$$

 $\Xi^0$ 

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$P$  is not yet measured; + is the quark model prediction.

$$\text{Mass } m = 1314.86 \pm 0.20 \text{ MeV}$$

$$m_{\Xi^-} - m_{\Xi^0} = 6.85 \pm 0.21 \text{ MeV}$$

$$\text{Mean life } \tau = (2.90 \pm 0.09) \times 10^{-10} \text{ s}$$

$$c\tau = 8.71 \text{ cm}$$

$$\text{Magnetic moment } \mu = -1.250 \pm 0.014 \mu_N$$

### Decay parameters

$$\Lambda\pi^0 \quad \alpha = -0.348 \pm 0.009$$

$$\alpha \text{ FOR } \Xi^0 \rightarrow \bar{\Lambda}\pi^0 = 0.379 \pm 0.004$$

$$" \quad \phi = (0.3 \pm 0.6)^\circ$$

$$\phi \text{ ANGLE FOR } \Xi^0 \rightarrow \bar{\Lambda}\pi^0 \text{ with } \tan\phi = \beta/\gamma = -0.3 \pm 0.6$$

degrees

$$\Delta\phi_{CP}(\Xi^0) = (\phi_{\Xi^0} + \phi_{\Xi^0})/2 = 0.0 \pm 0.4 \text{ degrees}$$

$$A_{CP} \text{ FOR } \Xi^0 \rightarrow \Lambda\pi^0, \Xi^0 \rightarrow \bar{\Lambda}\pi^0 = (-5 \pm 7) \times 10^{-3}$$

$$" \quad \gamma = 0.85 [I]$$

$$" \quad \Delta = (218_{-19}^{+12})^\circ [I]$$

$$\Lambda\gamma \quad \alpha = -0.70 \pm 0.07$$

$$\Lambda e^+ e^- \quad \alpha = -0.8 \pm 0.2$$

$$\Sigma^0\gamma \quad \alpha = -0.69 \pm 0.06$$

$$\Sigma^+ e^- \bar{\nu}_e \quad g_1(0)/f_1(0) = 1.22 \pm 0.05$$

$$\Sigma^+ e^- \bar{\nu}_e \quad f_2(0)/f_1(0) = 2.0 \pm 0.9$$

| $\Xi^0$ DECAY MODES            | Fraction ( $\Gamma_i/\Gamma$ )   | Confidence level | $p$<br>(MeV/c) |
|--------------------------------|----------------------------------|------------------|----------------|
| $\Lambda\pi^0$                 | $(99.524 \pm 0.012) \%$          |                  | 135            |
| $\Lambda\gamma$                | $(1.17 \pm 0.07) \times 10^{-3}$ |                  | 184            |
| $\Lambda e^+ e^-$              | $(7.6 \pm 0.6) \times 10^{-6}$   |                  | 184            |
| $\Sigma^0 \gamma$              | $(3.33 \pm 0.10) \times 10^{-3}$ |                  | 117            |
| $\Sigma^+ e^- \bar{\nu}_e$     | $(2.52 \pm 0.08) \times 10^{-4}$ |                  | 120            |
| $\Sigma^+ \mu^- \bar{\nu}_\mu$ | $(2.33 \pm 0.35) \times 10^{-6}$ |                  | 64             |

**$\Delta S = \Delta Q$  (SQ) violating modes or  
 $\Delta S = 2$  forbidden (S2) modes**

|                          |    |         |                  |     |     |
|--------------------------|----|---------|------------------|-----|-----|
| $\Sigma^- e^+ \nu_e$     | SQ | $< 1.6$ | $\times 10^{-4}$ | 90% | 112 |
| $\Sigma^- \mu^+ \nu_\mu$ | SQ | $< 9$   | $\times 10^{-4}$ | 90% | 49  |
| $p\pi^-$                 | S2 | $< 8$   | $\times 10^{-6}$ | 90% | 299 |
| $p e^- \bar{\nu}_e$      | S2 | $< 1.3$ | $\times 10^{-3}$ |     | 323 |
| $p \mu^- \bar{\nu}_\mu$  | S2 | $< 1.3$ | $\times 10^{-3}$ |     | 309 |



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$P$  is not yet measured; + is the quark model prediction.

Mass  $m = 1321.71 \pm 0.07$  MeV

$$(m_{\Xi^-} - m_{\Xi^+}) / m_{\Xi^-} = (-3 \pm 9) \times 10^{-5}$$

$$\text{Mean life } \tau = (1.639 \pm 0.015) \times 10^{-10} \text{ s}$$

$$c\tau = 4.91 \text{ cm}$$

$$(\tau_{\Xi^-} - \tau_{\Xi^+}) / \tau_{\Xi^-} = -0.01 \pm 0.07$$

$$\text{Magnetic moment } \mu = -0.6507 \pm 0.0025 \mu_N$$

$$(\mu_{\Xi^-} + \mu_{\Xi^+}) / |\mu_{\Xi^-}| = +0.01 \pm 0.05$$

**Decay parameters**

$$\Lambda\pi^- \quad \alpha = -0.385 \pm 0.008 \quad (S = 2.4)$$

$$\bar{\Lambda}\pi^+ \quad \bar{\alpha} = 0.374^{+0.005}_{-0.006}$$

$$(\alpha + \bar{\alpha}) / (\alpha - \bar{\alpha}) \text{ for } \Xi^- \rightarrow \Lambda\pi^-, \Xi^+ \rightarrow \bar{\Lambda}\pi^+ = (-9^{+11}_{-8}) \times 10^{-3}$$

$$[\alpha(\Xi^-)\alpha_-(\Lambda) - \bar{\alpha}\Xi^+\alpha_+(\bar{\Lambda})] / [\text{sum}] = (0 \pm 7) \times 10^{-4}$$

$$\Lambda\pi^- \quad \phi_- = (-1.5 \pm 0.6)^\circ$$

$$" \quad \gamma = 0.89 [l]$$

$$" \quad \Delta = (175.9 \pm 1.5)^\circ [l]$$

$$\bar{\Lambda}\pi^+ \quad \phi_+ = (0.6^{+0.7}_{-1.0})^\circ$$

$$\Delta\phi_{CP} = (\phi_- + \phi_+)/2 = (-0.2^{+0.5}_{-0.6})^\circ$$

$$\Lambda e^- \bar{\nu}_e \quad g_A/g_V = -0.25 \pm 0.05 [h]$$

| $\Xi^-$ DECAY MODES          | Fraction ( $\Gamma_i/\Gamma$ )     | Confidence level    | $\rho$<br>(MeV/c) |
|------------------------------|------------------------------------|---------------------|-------------------|
| $\Lambda\pi^-$               | $(99.887\pm0.035)\%$               |                     | 140               |
| $\Sigma^-\gamma$             | $(1.27\pm0.23)\times10^{-4}$       |                     | 118               |
| $\Lambda e^-\bar{\nu}_e$     | $(5.63\pm0.31)\times10^{-4}$       |                     | 190               |
| $\Lambda\mu^-\bar{\nu}_\mu$  | $(3.5^{+3.5}_{-2.2})\times10^{-4}$ |                     | 163               |
| $\Sigma^0 e^-\bar{\nu}_e$    | $(8.7\pm1.7)\times10^{-5}$         |                     | 123               |
| $\Sigma^0\mu^-\bar{\nu}_\mu$ | $<8$                               | $\times10^{-4}$ 90% | 70                |
| $\Xi^0 e^-\bar{\nu}_e$       | $<2.59$                            | $\times10^{-4}$ 90% | 7                 |

$\Delta S = 2$  forbidden ( $S2$ ) modes

|                            |      |        |                 |     |     |
|----------------------------|------|--------|-----------------|-----|-----|
| $n\pi^-$                   | $S2$ | $<1.9$ | $\times10^{-5}$ | 90% | 304 |
| $ne^-\bar{\nu}_e$          | $S2$ | $<3.2$ | $\times10^{-3}$ | 90% | 327 |
| $n\mu^-\bar{\nu}_\mu$      | $S2$ | $<1.5$ | %               | 90% | 314 |
| $p\pi^-\pi^-$              | $S2$ | $<4$   | $\times10^{-4}$ | 90% | 223 |
| $p\pi^-e^-\bar{\nu}_e$     | $S2$ | $<4$   | $\times10^{-4}$ | 90% | 305 |
| $p\pi^-\mu^-\bar{\nu}_\mu$ | $S2$ | $<4$   | $\times10^{-4}$ | 90% | 251 |
| $p\mu^-\mu^-$              | $L$  | $<4$   | $\times10^{-8}$ | 90% | 272 |

$\Xi(1530) 3/2^+$

$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$

$\Xi(1530)^0$  mass  $m = 1531.80 \pm 0.32$  MeV ( $S = 1.3$ )  
 $\Xi(1530)^-$  mass  $m = 1535.0 \pm 0.6$  MeV  
 $\Xi(1530)^0$  full width  $\Gamma = 9.1 \pm 0.5$  MeV  
 $\Xi(1530)^-$  full width  $\Gamma = 9.9^{+1.7}_{-1.9}$  MeV

| $\Xi(1530)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $\rho$<br>(MeV/c) |
|-------------------------|--------------------------------|------------------|-------------------|
| $\Xi\pi$                | 100 %                          |                  | 158               |
| $\Xi\gamma$             | $<3.7\%$                       | 90%              | 202               |

$\Xi(1690)$

$I(J^P) = \frac{1}{2}(??)$

Mass  $m = 1690 \pm 10$  MeV [o]  
Full width  $\Gamma = 20 \pm 15$  MeV

| $\Xi(1690)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\Lambda \bar{K}$       | seen                           | 240         |
| $\Sigma \bar{K}$        | seen                           | 70          |
| $\Xi \pi$               | seen                           | 311         |
| $\Xi^- \pi^+ \pi^-$     | possibly seen                  | 213         |

$\Xi(1820) \ 3/2^-$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Mass  $m = 1823 \pm 5$  MeV [o]  
Full width  $\Gamma = 24^{+15}_{-10}$  MeV [o]

| $\Xi(1820)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\Lambda \bar{K}$       | large                          | 402         |
| $\Sigma \bar{K}$        | small                          | 324         |
| $\Xi \pi$               | small                          | 421         |
| $\Xi(1530) \pi$         | small                          | 237         |

$\Xi(1950)$

$$I(J^P) = \frac{1}{2}(?^?)$$

Mass  $m = 1950 \pm 15$  MeV [o]  
Full width  $\Gamma = 60 \pm 20$  MeV [o]

| $\Xi(1950)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\Lambda \bar{K}$       | seen                           | 522         |
| $\Sigma \bar{K}$        | possibly seen                  | 460         |
| $\Xi \pi$               | seen                           | 519         |

$\Xi(2030)$

$$I(J^P) = \frac{1}{2}(\geq \frac{5}{2}^?)$$

Mass  $m = 2025 \pm 5$  MeV [o]  
Full width  $\Gamma = 20^{+15}_{-5}$  MeV [o]

| $\Xi(2030)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------|--------------------------------|-------------|
| $\Lambda \bar{K}$       | $\sim 20\%$                    | 585         |
| $\Sigma \bar{K}$        | $\sim 80\%$                    | 529         |
| $\Xi \pi$               | small                          | 574         |
| $\Xi(1530) \pi$         | small                          | 416         |
| $\Lambda \bar{K} \pi$   | small                          | 499         |
| $\Sigma \bar{K} \pi$    | small                          | 428         |

$$\Omega \text{ BARYONS}$$
$$(S = -3, I = 0)$$
$$\Omega^- = sss$$

$\Omega^-$

$$I(J^P) = 0(\frac{3}{2}^+)$$

$J^P = \frac{3}{2}^+$  is the quark-model prediction; and  $J = 3/2$  is fairly well established.

Mass  $m = 1672.45 \pm 0.29$  MeV  
 $(m_{\Omega^-} - m_{\bar{\Omega}^+}) / m_{\Omega^-} = (-1 \pm 8) \times 10^{-5}$   
Mean life  $\tau = (0.821 \pm 0.011) \times 10^{-10}$  s  
 $c\tau = 2.461$  cm  
 $(\tau_{\Omega^-} - \tau_{\bar{\Omega}^+}) / \tau_{\Omega^-} = 0.00 \pm 0.05$   
Magnetic moment  $\mu = -2.02 \pm 0.05 \mu_N$

Decay parameters

$\alpha(\Omega^-) \alpha_-(\Lambda)$  FOR  $\Omega^- \rightarrow \Lambda K^- = 0.0115 \pm 0.0015$   
 $\Lambda K^- \quad \alpha = 0.0154 \pm 0.0020$   
 $\Lambda K^-, \bar{\Lambda} K^+ \quad (\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) = -0.02 \pm 0.13$   
 $\Xi^0 \pi^- \quad \alpha = 0.09 \pm 0.14$   
 $\Xi^- \pi^0 \quad \alpha = 0.05 \pm 0.21$

| $\Omega^-$ DECAY MODES                                             | Fraction ( $\Gamma_i/\Gamma$ )       | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|--------------------------------------------------------------------|--------------------------------------|-----------------------------------|----------------|
| $\Lambda K^-$                                                      | $(67.7 \pm 0.7) \%$                  |                                   | 211            |
| $\Xi^0 \pi^-$                                                      | $(24.3 \pm 0.7) \%$                  | S=1.5                             | 294            |
| $\Xi^- \pi^0$                                                      | $(8.55 \pm 0.33) \%$                 |                                   | 289            |
| $\Xi^- \pi^+ \pi^-$                                                | $(3.7^{+0.7}_{-0.6}) \times 10^{-4}$ |                                   | 189            |
| $\Xi(1530)^0 \pi^-$                                                | $< 7 \times 10^{-5}$                 | CL=90%                            | 17             |
| $\Xi^0 e^- \bar{\nu}_e$                                            | $(5.6 \pm 2.8) \times 10^{-3}$       |                                   | 319            |
| $\Xi^- \gamma$                                                     | $< 4.6 \times 10^{-4}$               | CL=90%                            | 314            |
| <b><math>\Delta S = 2</math> forbidden (<math>S2</math>) modes</b> |                                      |                                   |                |
| $\Lambda \pi^-$                                                    | $S2 \quad < 2.9 \times 10^{-6}$      | CL=90%                            | 449            |
| $\Sigma^0 \pi^-$                                                   | $< 5.4 \times 10^{-4}$               | CL=90%                            | 393            |
| $n K^-$                                                            | $< 2.4 \times 10^{-4}$               | CL=90%                            | 415            |

$\Omega(2012)^-$

$I(J^P) = 0(?^-)$

Mass  $m = 2012.5 \pm 0.6$  MeV  
Full width  $\Gamma = 6.4^{+3.0}_{-2.6}$  MeV

Branching fractions are given relative to the one **DEFINED AS 1**.

| $\Omega(2012)^-$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\Xi^0 K^-$                  | <b>DEFINED AS 1</b>            | 403         |
| $\Xi^- \bar{K}^0$            | $0.83 \pm 0.21$                | 392         |

$\Omega(2250)^-$

$I(J^P) = 0(?^?)$

Mass  $m = 2252 \pm 9$  MeV  
Full width  $\Gamma = 55 \pm 18$  MeV

| $\Omega(2250)^-$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\Xi^- \pi^+ K^-$            | seen                           | 532         |
| $\Xi(1530)^0 K^-$            | seen                           | 437         |

# CHARMED BARYONS

## ( $C = +1$ )

$$\Lambda_c^+ = u d c, \quad \Sigma_c^{++} = u u c, \quad \Sigma_c^+ = u d c, \quad \Sigma_c^0 = d d c, \\ \Xi_c^+ = u s c, \quad \Xi_c^0 = d s c, \quad \Omega_c^0 = s s c$$

$\Lambda_c^+$

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass  $m = 2286.46 \pm 0.14$  MeV

Mean life  $\tau = (202.6 \pm 1.0) \times 10^{-15}$  s

$c\tau = 60.75$   $\mu\text{m}$

### Decay asymmetry parameters

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda \pi^+ = -0.768 \pm 0.015 \quad (S = 3.4)$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda \rho^+ = -0.76 \pm 0.07$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^+ \pi^0 = -0.484 \pm 0.027$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^+ \eta = -0.99 \pm 0.06$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^+ \eta' = -0.46 \pm 0.07$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^0 \pi^+ = -0.466 \pm 0.018$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma(1385)^+ \pi^0 = -0.92 \pm 0.09$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma(1385)^0 \pi^+ = -0.79 \pm 0.11$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda \ell^+ \nu_\ell = -0.875 \pm 0.033$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow p K_S^0 = -0.754 \pm 0.010$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda K^+ = -0.546 \pm 0.035$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Sigma^0 K^+ = -0.54 \pm 0.20$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda(1405) \pi^+ = 0.58 \pm 0.28$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda(1520) \pi^+ = 0.93 \pm 0.09$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda(1600) \pi^+ = 0.2 \pm 0.5$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda(1670) \pi^+ = 0.82 \pm 0.08$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda(1690) \pi^+ = 0.958 \pm 0.034$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Lambda(2000) \pi^+ = -0.57 \pm 0.19$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Delta(1232)^{++} K^- = 0.55 \pm 0.04$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Delta(1600)^{++} K^- = -0.50 \pm 0.18$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Delta(1700)^{++} K^- = 0.22 \pm 0.08$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \bar{K}_0^*(700)^0 p = -0.1 \pm 0.7$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \bar{K}_0^*(1430)^0 p = 0.34 \pm 0.14$$

$$\alpha \text{ FOR } \Lambda_c^+ \rightarrow \Xi^0 K^+ = 0.01 \pm 0.16$$

$$\begin{aligned}
&(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda \pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} \pi^- = 0.020 \pm 0.016 \\
&(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Sigma^0 \pi^+, \bar{\Lambda}_c^- \rightarrow \bar{\Sigma}^0 \pi^- = -0.02 \pm 0.05 \\
&(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda e^+ \nu_e, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} e^- \bar{\nu}_e = 0.00 \pm 0.04 \\
&(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Lambda K^+, \bar{\Lambda}_c^- \rightarrow \bar{\Lambda} K^- = -0.02 \pm 0.11 \\
&(\alpha + \bar{\alpha})/(\alpha - \bar{\alpha}) \text{ in } \Lambda_c^+ \rightarrow \Sigma^0 K^+, \bar{\Lambda}_c^- \rightarrow \bar{\Sigma}^0 K^- = 0.1 \pm 0.4 \\
&A_{CP}(\Lambda X) \text{ in } \Lambda_c \rightarrow \Lambda X, \bar{\Lambda}_c \rightarrow \bar{\Lambda} X = (2 \pm 7)\% \\
&A_{CP}(\Lambda K^+) \text{ in } \Lambda_c \rightarrow \Lambda K^+, \bar{\Lambda}_c \rightarrow \bar{\Lambda} K^- = 0.021 \pm 0.026 \\
&A_{CP}(\Sigma^0 K^+) \text{ in } \Lambda_c \rightarrow \Sigma^0 K^+, \bar{\Lambda}_c \rightarrow \bar{\Sigma}^0 K^- = 0.03 \pm 0.05 \\
&\Delta A_{CP} = A_{CP}(\Lambda_c^+ \rightarrow p K^+ K^-) - A_{CP}(\Lambda_c^+ \rightarrow p \pi^+ \pi^-) = \\
&\quad (0.3 \pm 1.1)\%
\end{aligned}$$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction  $\Lambda_c^+ \rightarrow p \bar{K}^*(892)^0$  seen in  $\Lambda_c^+ \rightarrow p K^- \pi^+$  has been multiplied up to include  $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$  decays.

| $\Lambda_c^+$ DECAY MODES                                                                       | Fraction ( $\Gamma_i/\Gamma$ )           | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|-------------------------------------------------------------------------------------------------|------------------------------------------|-----------------------------------|----------------|
| <b>Hadronic modes with a <math>p</math> or <math>n</math>: <math>S = -1</math> final states</b> |                                          |                                   |                |
| $p K_S^0$                                                                                       | ( 1.61 $\pm$ 0.07 ) %                    | S=1.1                             | 873            |
| $p K_L^0$                                                                                       | ( 1.67 $\pm$ 0.07 ) %                    |                                   | 873            |
| $p K^- \pi^+$                                                                                   | ( 6.35 $\pm$ 0.25 ) %                    | S=1.3                             | 823            |
| $p \bar{K}_0^*(700)^0$                                                                          | ( 1.9 $\pm$ 0.6 ) $\times 10^{-3}$       |                                   | 719            |
| $p \bar{K}^*(892)^0$                                                                            | [q] ( 1.41 $\pm$ 0.07 ) %                |                                   | 685            |
| $p \bar{K}_0^*(1430)$                                                                           | ( 9.3 $\pm$ 1.8 ) $\times 10^{-3}$       |                                   | †              |
| $\Delta(1232)^{++} K^-$                                                                         | ( 1.79 $\pm$ 0.09 ) %                    |                                   | 710            |
| $\Delta(1600)^{++} K^-$                                                                         | ( 2.9 $\pm$ 1.0 ) $\times 10^{-3}$       |                                   | —              |
| $\Delta(1700)^{++} K^-$                                                                         | ( 2.5 $\pm$ 0.6 ) $\times 10^{-3}$       |                                   | —              |
| $\Lambda(1405)^0 \pi^+$                                                                         | ( 4.9 $\pm$ 1.9 ) $\times 10^{-3}$       |                                   | —              |
| $\Lambda(1520) \pi^+$                                                                           | [q] ( 1.18 $\pm$ 0.16 ) $\times 10^{-3}$ |                                   | 628            |
| $\Lambda(1600) \pi^+$                                                                           | ( 3.3 $\pm$ 1.2 ) $\times 10^{-3}$       |                                   | 571            |
| $\Lambda(1670) \pi^+$                                                                           | ( 7.5 $\pm$ 2.1 ) $\times 10^{-4}$       |                                   | 516            |
| $\Lambda(1690) \pi^+$                                                                           | ( 7.6 $\pm$ 2.3 ) $\times 10^{-4}$       |                                   | 504            |
| $\Lambda(2000) \pi^+$                                                                           | ( 6.1 $\pm$ 0.7 ) $\times 10^{-3}$       |                                   | 234            |
| $p K^- \pi^+$ nonresonant                                                                       | ( 3.5 $\pm$ 0.4 ) %                      |                                   | 823            |
| $p K_S^0 \pi^0$                                                                                 | ( 1.99 $\pm$ 0.12 ) %                    |                                   | 823            |
| $p K_L^0 \pi^0$                                                                                 | ( 2.02 $\pm$ 0.14 ) %                    |                                   | 823            |
| $n K_S^0 \pi^+$                                                                                 | ( 1.86 $\pm$ 0.09 ) %                    |                                   | 821            |
| $n K_S^0 K^+$                                                                                   | ( 3.9 $\pm$ 1.7 ) $\times 10^{-4}$       |                                   | 612            |
| $n K_S^0 \pi^+ \pi^0$                                                                           | ( 8.5 $\pm$ 1.3 ) $\times 10^{-3}$       |                                   | 756            |

|                                         |                                  |       |     |
|-----------------------------------------|----------------------------------|-------|-----|
| $nK^-\pi^+\pi^+$                        | ( 1.90 ± 0.12 ) %                |       | 756 |
| $p\bar{K}^0\eta$                        | ( 8.9 ± 0.6 ) × 10 <sup>-3</sup> | S=1.1 | 568 |
| $pK_S^0\pi^+\pi^-$                      | ( 1.62 ± 0.11 ) %                | S=1.1 | 754 |
| $pK_L^0\pi^+\pi^-$                      | ( 1.69 ± 0.11 ) %                |       | 754 |
| $pK^-\pi^+\pi^0$                        | ( 4.52 ± 0.28 ) %                | S=1.5 | 759 |
| $pK^*(892)^-\pi^+$                      | [q] ( 1.4 ± 0.5 ) %              |       | 580 |
| $p(K^-\pi^+)_{\text{nonresonant}}\pi^0$ | ( 4.6 ± 0.8 ) %                  |       | 759 |
| $\Delta(1232)\bar{K}^*(892)$            | seen                             |       | 419 |
| $pK^-2\pi^+\pi^-$                       | ( 1.4 ± 1.0 ) × 10 <sup>-3</sup> |       | 671 |
| $pK^-\pi^+2\pi^0$                       | ( 1.0 ± 0.5 ) %                  |       | 678 |

**Hadronic modes with a  $p$  or  $n$ :  $S = 0$  final states**

|                                    |                                                                                     |        |     |
|------------------------------------|-------------------------------------------------------------------------------------|--------|-----|
| $p\pi^0$                           | ( 1.6 $\begin{smallmatrix} + \\ - \end{smallmatrix}$ 0.7 / 0.6 ) × 10 <sup>-4</sup> |        | 945 |
| $n\pi^+$                           | ( 6.6 ± 1.3 ) × 10 <sup>-4</sup>                                                    |        | 944 |
| $p\eta$                            | ( 1.49 ± 0.08 ) × 10 <sup>-3</sup>                                                  | S=1.1  | 856 |
| $p\eta'$                           | ( 4.9 ± 0.9 ) × 10 <sup>-4</sup>                                                    |        | 639 |
| $p\omega(782)^0$                   | ( 8.9 ± 1.1 ) × 10 <sup>-4</sup>                                                    | S=1.2  | 751 |
| $p\pi^+\pi^-$                      | ( 4.67 ± 0.24 ) × 10 <sup>-3</sup>                                                  |        | 927 |
| $pf_0(980)$                        | [q] ( 3.5 ± 2.3 ) × 10 <sup>-3</sup>                                                |        | 614 |
| $p\rho(770)^0$                     | ( 1.5 ± 0.4 ) × 10 <sup>-3</sup>                                                    |        | —   |
| $n\pi^+\pi^0$                      | ( 6.4 ± 0.9 ) × 10 <sup>-3</sup>                                                    |        | 927 |
| $nK^+\pi^0$                        | < 7.1 × 10 <sup>-4</sup>                                                            | CL=90% | 824 |
| $n\pi^+\pi^-\pi^+$                 | ( 4.5 ± 0.8 ) × 10 <sup>-3</sup>                                                    |        | 895 |
| $p2\pi^+2\pi^-$                    | ( 2.3 ± 1.5 ) × 10 <sup>-3</sup>                                                    |        | 852 |
| $pK^+K^-$                          | ( 1.08 ± 0.05 ) × 10 <sup>-3</sup>                                                  |        | 616 |
| $p\phi$                            | [q] ( 1.05 ± 0.14 ) × 10 <sup>-3</sup>                                              | S=1.1  | 590 |
| $pK^+K^- \text{ non-}\phi$         | ( 5.3 ± 1.2 ) × 10 <sup>-4</sup>                                                    |        | 616 |
| $pK_S^0K_S^0$                      | ( 2.38 ± 0.18 ) × 10 <sup>-4</sup>                                                  |        | 610 |
| $p\phi\pi^0$                       | ( 10 ± 4 ) × 10 <sup>-5</sup>                                                       |        | 460 |
| $pK^+K^-\pi^0 \text{ nonresonant}$ | < 6.3 × 10 <sup>-5</sup>                                                            | CL=90% | 494 |

**Hadronic modes with a hyperon:  $S = -1$  final states**

|                                                                  |                                  |       |     |
|------------------------------------------------------------------|----------------------------------|-------|-----|
| $\Lambda\pi^+$                                                   | ( 1.31 ± 0.05 ) %                | S=1.1 | 864 |
| $\Lambda(1670)\pi^+, \Lambda(1670) \rightarrow \eta\Lambda$      | ( 3.5 ± 0.5 ) × 10 <sup>-3</sup> |       | —   |
| $\Lambda\pi^+\pi^0$                                              | ( 7.10 ± 0.34 ) %                | S=1.1 | 844 |
| $\Lambda\rho^+$                                                  | ( 4.1 ± 0.5 ) %                  |       | 636 |
| $\Sigma(1385)^+\pi^0, \Sigma^+ \rightarrow \Lambda\pi^+$         | ( 5.1 ± 0.7 ) × 10 <sup>-3</sup> |       | —   |
| $\Sigma(1385)^0\pi^+, \Sigma^0 \rightarrow \Lambda\pi^0$         | ( 5.6 ± 0.8 ) × 10 <sup>-3</sup> |       | —   |
| $\Lambda\pi^-2\pi^+$                                             | ( 3.67 ± 0.26 ) %                | S=1.4 | 807 |
| $\Sigma(1385)^+\pi^+\pi^-, \Sigma^{*+} \rightarrow \Lambda\pi^+$ | ( 1.0 ± 0.5 ) %                  |       | 688 |
| $\Sigma(1385)^-2\pi^+, \Sigma^{*-} \rightarrow \Lambda\pi^-$     | ( 7.7 ± 1.4 ) × 10 <sup>-3</sup> |       | 688 |
| $\Lambda\pi^+\rho^0$                                             | ( 1.5 ± 0.6 ) %                  |       | 524 |
| $\Sigma(1385)^+\rho^0, \Sigma^{*+} \rightarrow \Lambda\pi^+$     | ( 5 ± 4 ) × 10 <sup>-3</sup>     |       | 363 |

|                                                         |                         |                  |        |     |
|---------------------------------------------------------|-------------------------|------------------|--------|-----|
| $\Lambda\pi^-2\pi^+$ nonresonant                        | < 1.1                   | %                | CL=90% | 807 |
| $\Lambda\pi^-\pi^02\pi^+$ total                         | ( 2.3 $\pm$ 0.8 )       | %                |        | 757 |
| $\Lambda\pi^+\eta$                                      | [q] ( 1.87 $\pm$ 0.11 ) | %                | S=1.1  | 691 |
| $\Sigma(1385)^+\eta$                                    | [q] ( 9.1 $\pm$ 2.0 )   | $\times 10^{-3}$ |        | 570 |
| $\Lambda\pi^+\omega$                                    | [q] ( 1.5 $\pm$ 0.5 )   | %                |        | 517 |
| $\Lambda\pi^-\pi^02\pi^+$ , no $\eta$ or $\omega$       | < 8                     | $\times 10^{-3}$ | CL=90% | 757 |
| $\Lambda K^+\bar{K}^0$                                  | ( 5.7 $\pm$ 1.1 )       | $\times 10^{-3}$ | S=2.0  | 443 |
| $\Xi(1690)^0K^+, \Xi^{*0} \rightarrow \Lambda\bar{K}^0$ | ( 1.6 $\pm$ 0.5 )       | $\times 10^{-3}$ |        | 286 |
| $\Sigma^0\pi^+$                                         | ( 1.29 $\pm$ 0.05 )     | %                |        | 825 |
| $\Sigma^0\pi^+\eta$                                     | ( 7.6 $\pm$ 0.8 )       | $\times 10^{-3}$ |        | 635 |
| $\Sigma^+\pi^0$                                         | ( 1.26 $\pm$ 0.10 )     | %                | S=1.1  | 827 |
| $\Sigma^+\eta$                                          | ( 3.2 $\pm$ 0.5 )       | $\times 10^{-3}$ |        | 713 |
| $\Sigma^+\eta'$                                         | ( 4.2 $\pm$ 0.9 )       | $\times 10^{-3}$ |        | 391 |
| $\Sigma^+\pi^+\pi^-$                                    | ( 4.54 $\pm$ 0.20 )     | %                | S=1.2  | 804 |
| $\Sigma^+\rho^0$                                        | < 1.7                   | %                | CL=95% | 575 |
| $\Sigma^-2\pi^+$                                        | ( 1.87 $\pm$ 0.18 )     | %                |        | 799 |
| $\Sigma^0\pi^+\pi^0$                                    | ( 3.6 $\pm$ 0.4 )       | %                |        | 803 |
| $\Sigma^+\pi^0\pi^0$                                    | ( 1.57 $\pm$ 0.14 )     | %                |        | 806 |
| $\Sigma^0\pi^-2\pi^+$                                   | ( 1.12 $\pm$ 0.31 )     | %                |        | 763 |
| $\Sigma^+\omega$                                        | ( 1.72 $\pm$ 0.20 )     | %                |        | 569 |
| $\Sigma^-\pi^02\pi^+$                                   | ( 2.1 $\pm$ 0.4 )       | %                |        | 762 |
| $\Sigma^+K^+K^-$                                        | ( 3.6 $\pm$ 0.4 )       | $\times 10^{-3}$ | S=1.1  | 349 |
| $\Sigma^+\phi$                                          | [q] ( 4.0 $\pm$ 0.5 )   | $\times 10^{-3}$ | S=1.1  | 295 |
| $\Xi(1690)^0K^+, \Xi^{*0} \rightarrow \Sigma^+K^-$      | ( 1.03 $\pm$ 0.25 )     | $\times 10^{-3}$ |        | 286 |
| $\Sigma^+K^+K^-$ nonresonant                            | < 8                     | $\times 10^{-4}$ | CL=90% | 349 |
| $\Xi^0K^+$                                              | ( 5.5 $\pm$ 0.7 )       | $\times 10^{-3}$ |        | 653 |
| $\Xi^-K^+\pi^+$                                         | ( 6.3 $\pm$ 0.5 )       | $\times 10^{-3}$ | S=1.1  | 565 |
| $\Xi^0K^+\pi^0$                                         | ( 7.8 $\pm$ 1.7 )       | $\times 10^{-3}$ |        | 574 |
| $\Xi(1530)^0K^+$                                        | ( 4.9 $\pm$ 0.6 )       | $\times 10^{-3}$ | S=1.1  | 473 |

**Hadronic modes with a hyperon:  $S = 0$  final states**

|                         |                       |                  |        |     |
|-------------------------|-----------------------|------------------|--------|-----|
| $\Lambda K^+$           | ( 6.48 $\pm$ 0.31 )   | $\times 10^{-4}$ |        | 781 |
| $\Lambda K^+\pi^0$      | ( 1.48 $\pm$ 0.29 )   | $\times 10^{-3}$ |        | 722 |
| $\Lambda K^+\pi^+\pi^-$ | ( 4.2 $\pm$ 1.6 )     | $\times 10^{-4}$ |        | 637 |
| $\Sigma^0K^+$           | ( 3.73 $\pm$ 0.31 )   | $\times 10^{-4}$ |        | 735 |
| $\Sigma^+K_S^0$         | ( 4.8 $\pm$ 1.4 )     | $\times 10^{-4}$ |        | 736 |
| $\Sigma^0K^+\pi^+\pi^-$ | < 2.6                 | $\times 10^{-4}$ | CL=90% | 574 |
| $\Sigma^0K^+\pi^0$      | < 1.8                 | $\times 10^{-3}$ | CL=90% | 670 |
| $\Sigma^+K^+\pi^-$      | ( 2.04 $\pm$ 0.26 )   | $\times 10^{-3}$ |        | 670 |
| $\Sigma^+K^*(892)^0$    | [q] ( 3.5 $\pm$ 1.0 ) | $\times 10^{-3}$ |        | 470 |
| $\Sigma^+K^+\pi^-\pi^0$ | < 1.1                 | $\times 10^{-3}$ | CL=90% | 581 |
| $\Sigma^-K^+\pi^+$      | ( 3.8 $\pm$ 1.2 )     | $\times 10^{-4}$ |        | 664 |

**Doubly Cabibbo-suppressed modes**

|             |                                  |     |
|-------------|----------------------------------|-----|
| $pK^+\pi^-$ | $(1.13 \pm 0.17) \times 10^{-4}$ | 823 |
|-------------|----------------------------------|-----|

**Semileptonic modes**

|                                                         |                                |     |
|---------------------------------------------------------|--------------------------------|-----|
| $\Lambda e^+ \nu_e$                                     | $(3.56 \pm 0.13) \%$           | 871 |
| $\Lambda \pi^+ \pi^- e^+ \nu_e$                         | $< 3.9 \times 10^{-4}$ CL=90%  | 843 |
| $pK^- e^+ \nu_e$                                        | $(8.8 \pm 1.8) \times 10^{-4}$ | 874 |
| $pK_S^0 \pi^- e^+ \nu_e$                                | $< 3.3 \times 10^{-4}$ CL=90%  | 821 |
| $\Lambda(1520) e^+ \nu_e$                               | $(1.0 \pm 0.5) \times 10^{-3}$ | 639 |
| $\Lambda(1405)^0 e^+ \nu_e, \Lambda^0 \rightarrow pK^-$ | $(4.2 \pm 1.9) \times 10^{-4}$ | —   |
| $\Lambda \mu^+ \nu_\mu$                                 | $(3.48 \pm 0.17) \%$           | 867 |

**Inclusive modes**

|                    |                      |   |
|--------------------|----------------------|---|
| $e^+$ anything     | $(4.06 \pm 0.13) \%$ | — |
| $p$ anything       | $(50 \pm 16) \%$     | — |
| $n$ anything       | $(32.6 \pm 1.6) \%$  | — |
| $\Lambda$ anything | $(38.2 \pm 2.9) \%$  | — |
| $K_S^0$ anything   | $(9.9 \pm 0.7) \%$   | — |
| 3prongs            | $(24 \pm 8) \%$      | — |

**$\Delta C = 1$  weak neutral current ( $C1$ ) modes, or  
Lepton Family number ( $LF$ ), or Lepton number ( $L$ ), or  
Baryon number ( $B$ ) violating modes**

|                            |       |                        |        |     |
|----------------------------|-------|------------------------|--------|-----|
| $pe^+e^-$                  | $C1$  | $< 5.5 \times 10^{-6}$ | CL=90% | 951 |
| $p\mu^+\mu^-$ non-resonant | $C1$  | $< 2.9 \times 10^{-8}$ | CL=90% | 937 |
| $pe^+\mu^-$                | $LF$  | $< 9.9 \times 10^{-6}$ | CL=90% | 947 |
| $pe^-\mu^+$                | $LF$  | $< 1.9 \times 10^{-5}$ | CL=90% | 947 |
| $\bar{p}2e^+$              | $L,B$ | $< 2.7 \times 10^{-6}$ | CL=90% | 951 |
| $\bar{p}2\mu^+$            | $L,B$ | $< 9.4 \times 10^{-6}$ | CL=90% | 937 |
| $\bar{p}e^+\mu^+$          | $L,B$ | $< 1.6 \times 10^{-5}$ | CL=90% | 947 |
| $\Sigma^-\mu^+\mu^+$       | $L$   | $< 7.0 \times 10^{-4}$ | CL=90% | 812 |

**Radiative modes**

|                   |                        |        |     |
|-------------------|------------------------|--------|-----|
| $\Sigma^+ \gamma$ | $< 2.5 \times 10^{-4}$ | CL=90% | 834 |
|-------------------|------------------------|--------|-----|

**Exotic modes**

|             |                            |        |   |
|-------------|----------------------------|--------|---|
| $p\gamma D$ | $[r] < 8.0 \times 10^{-5}$ | CL=90% | — |
|-------------|----------------------------|--------|---|

 **$\Lambda_c(2595)^+$** 

$$I(J^P) = 0(\frac{1}{2}^-)$$

The spin-parity follows from the fact that  $\Sigma_c(2455)\pi$  decays, with little available phase space, are dominant. This assumes that  $J^P = 1/2^+$  for the  $\Sigma_c(2455)$ .

Mass  $m = 2592.25 \pm 0.28 \text{ MeV}$   
 $m - m_{\Lambda_c^+} = 305.79 \pm 0.24 \text{ MeV}$   
Full width  $\Gamma = 2.6 \pm 0.6 \text{ MeV}$

$\Lambda_c^+ \pi \pi$  and its submode  $\Sigma_c(2455) \pi$  — the latter just barely — are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass; and the submode seems to dominate.

| $\Lambda_c(2595)^+$ DECAY MODES  | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \pi^+ \pi^-$        | [s] —                          | 117         |
| $\Sigma_c(2455)^{++} \pi^-$      | $24 \pm 7 \%$                  | 3           |
| $\Sigma_c(2455)^0 \pi^+$         | $24 \pm 7 \%$                  | 3           |
| $\Lambda_c^+ \pi^+ \pi^-$ 3-body | $18 \pm 10 \%$                 | 117         |
| $\Lambda_c^+ \pi^0$              | [t] not seen                   | 258         |
| $\Lambda_c^+ \gamma$             | not seen                       | 288         |

$\Lambda_c(2625)^+$

$I(J^P) = 0(\frac{3}{2}^-)$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

Mass  $m = 2628.00 \pm 0.15 \text{ MeV}$   
 $m - m_{\Lambda_c^+} = 341.54 \pm 0.05 \text{ MeV}$   
Full width  $\Gamma < 0.52 \text{ MeV}$ , CL = 90%

$\Lambda_c^+ \pi \pi$  and its submode  $\Sigma(2455) \pi$  are the only strong decays allowed to an excited  $\Lambda_c^+$  having this mass.

| $\Lambda_c(2625)^+$ DECAY MODES  | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$ (MeV/c) |
|----------------------------------|--------------------------------|------------------|-------------|
| $\Lambda_c^+ \pi^+ \pi^-$        | [u] (50 $\pm$ 7 ) %            |                  | 184         |
| $\Sigma_c(2455)^{++} \pi^-$      | ( 2.6 $\pm$ 0.4 ) %            |                  | 103         |
| $\Sigma_c(2455)^0 \pi^+$         | ( 2.6 $\pm$ 0.4 ) %            |                  | 103         |
| $\Lambda_c^+ \pi^+ \pi^-$ 3-body | large                          |                  | 184         |
| $\Lambda_c^+ \pi^0$              | [t] < 50 %                     | 90%              | 293         |
| $\Lambda_c^+ \gamma$             | < 26 %                         | 90%              | 319         |

$\Lambda_c(2860)^+$

$I(J^P) = 0(\frac{3}{2}^+)$

Mass  $m = 2856.1^{+2.3}_{-6.0} \text{ MeV}$   
Full width  $\Gamma = 68^{+12}_{-22} \text{ MeV}$

| $\Lambda_c(2860)^+$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------|--------------------------------|-------------|
| $D^0 p$                         | seen                           | 259         |

 **$\Lambda_c(2880)^+$** 

$$I(J^P) = 0(\frac{5}{2}^+)$$

Mass  $m = 2881.63 \pm 0.24$  MeV $m - m_{\Lambda_c^+} = 595.17 \pm 0.28$  MeVFull width  $\Gamma = 5.6_{-0.6}^{+0.8}$  MeV

| $\Lambda_c(2880)^+$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \pi^+ \pi^-$       | seen                           | 471         |
| $\Sigma_c(2455)^{0,++} \pi^\pm$ | seen                           | 376         |
| $\Sigma_c(2520)^{0,++} \pi^\pm$ | seen                           | 317         |
| $p D^0$                         | seen                           | 316         |

 **$\Lambda_c(2940)^+$** 

$$I(J^P) = 0(\frac{3}{2}^-)$$

 $J^P = 3/2^-$  is favored, but is not certainMass  $m = 2939.6_{-1.5}^{+1.3}$  MeVFull width  $\Gamma = 20_{-5}^{+6}$  MeV

| $\Lambda_c(2940)^+$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------|--------------------------------|-------------|
| $p D^0$                         | seen                           | 420         |
| $\Sigma_c(2455)^{0,++} \pi^\pm$ | seen                           | —           |

 **$\Sigma_c(2455)$** 

$$I(J^P) = 1(\frac{1}{2}^+)$$

 $\Sigma_c(2455)^{++}$  mass  $m = 2453.97 \pm 0.14$  MeV $\Sigma_c(2455)^+$  mass  $m = 2452.65_{-0.16}^{+0.22}$  MeV $\Sigma_c(2455)^0$  mass  $m = 2453.75 \pm 0.14$  MeV $m_{\Sigma_c(2455)^{++}} - m_{\Lambda_c^+} = 167.510 \pm 0.017$  MeV $m_{\Sigma_c(2455)^+} - m_{\Lambda_c^+} = 166.19_{-0.08}^{+0.16}$  MeV $m_{\Sigma_c(2455)^0} - m_{\Lambda_c^+} = 167.290 \pm 0.017$  MeV $m_{\Sigma_c(2455)^{++}} - m_{\Sigma_c(2455)^0} = 0.220 \pm 0.013$  MeV $m_{\Sigma_c(2455)^+} - m_{\Sigma_c(2455)^0} = -1.10_{-0.08}^{+0.16}$  MeV

$$\Sigma_c(2455)^{++} \text{ full width } \Gamma = 1.89^{+0.09}_{-0.18} \text{ MeV} \quad (S = 1.1)$$

$$\Sigma_c(2455)^+ \text{ full width } \Gamma = 2.3 \pm 0.4 \text{ MeV}$$

$$\Sigma_c(2455)^0 \text{ full width } \Gamma = 1.83^{+0.11}_{-0.19} \text{ MeV} \quad (S = 1.2)$$

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

| <b><math>\Sigma_c(2455)</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \pi$                              | $\approx 100 \%$               | 94          |

## $\Sigma_c(2520)$

$$I(J^P) = 1(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$$\Sigma_c(2520)^{++} \text{ mass } m = 2518.41 \pm 0.22 \text{ MeV} \quad (S = 1.3)$$

$$\Sigma_c(2520)^+ \text{ mass } m = 2517.4^{+0.7}_{-0.5} \text{ MeV}$$

$$\Sigma_c(2520)^0 \text{ mass } m = 2518.48 \pm 0.21 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Lambda_c^+} = 231.95 \pm 0.18 \text{ MeV} \quad (S = 1.8)$$

$$m_{\Sigma_c(2520)^+} - m_{\Lambda_c^+} = 230.9^{+0.7}_{-0.5} \text{ MeV}$$

$$m_{\Sigma_c(2520)^0} - m_{\Lambda_c^+} = 232.02 \pm 0.15 \text{ MeV} \quad (S = 1.4)$$

$$m_{\Sigma_c(2520)^{++}} - m_{\Sigma_c(2520)^0} = 0.01 \pm 0.15 \text{ MeV}$$

$$\Sigma_c(2520)^{++} \text{ full width } \Gamma = 14.78^{+0.30}_{-0.40} \text{ MeV}$$

$$\Sigma_c(2520)^+ \text{ full width } \Gamma = 17.2^{+4.0}_{-2.2} \text{ MeV}$$

$$\Sigma_c(2520)^0 \text{ full width } \Gamma = 15.3^{+0.4}_{-0.5} \text{ MeV}$$

$\Lambda_c^+ \pi$  is the only strong decay allowed to a  $\Sigma_c$  having this mass.

| $\Sigma_c(2520)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \pi$            | $\approx 100\%$                | 179         |

|                                                                        |                   |
|------------------------------------------------------------------------|-------------------|
| $\Sigma_c(2800)$                                                       | $I(J^P) = 1(?^?)$ |
| $\Sigma_c(2800)^{++}$ mass $m = 2801^{+4}_{-6}$ MeV                    |                   |
| $\Sigma_c(2800)^+$ mass $m = 2792^{+14}_{-5}$ MeV                      |                   |
| $\Sigma_c(2800)^0$ mass $m = 2806^{+5}_{-7}$ MeV (S = 1.3)             |                   |
| $m_{\Sigma_c(2800)^{++}} - m_{\Lambda_c^+} = 514^{+4}_{-6}$ MeV        |                   |
| $m_{\Sigma_c(2800)^+} - m_{\Lambda_c^+} = 505^{+14}_{-5}$ MeV          |                   |
| $m_{\Sigma_c(2800)^0} - m_{\Lambda_c^+} = 519^{+5}_{-7}$ MeV (S = 1.3) |                   |
| $\Sigma_c(2800)^{++}$ full width $\Gamma = 75^{+22}_{-17}$ MeV         |                   |
| $\Sigma_c(2800)^+$ full width $\Gamma = 60^{+60}_{-40}$ MeV            |                   |
| $\Sigma_c(2800)^0$ full width $\Gamma = 72^{+22}_{-15}$ MeV            |                   |

| $\Sigma_c(2800)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \pi$            | seen                           | 443         |


|           |                                       |
|-----------|---------------------------------------|
| $\Xi_c^+$ | $I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ |
|-----------|---------------------------------------|

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2467.71 \pm 0.23$  MeV (S = 1.3)  
Mean life  $\tau = (453 \pm 5) \times 10^{-15}$  s  
 $c\tau = 135.8 \mu\text{m}$

Branching fractions marked with a footnote, e.g. [a], have been corrected for decay modes not observed in the experiments. For example, the sub-mode fraction  $\Xi_c^+ \rightarrow \Sigma^+ \bar{K}^*(892)^0$  seen in  $\Xi_c^+ \rightarrow \Sigma^+ K^- \pi^+$  has been multiplied up to include  $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$  decays.

| $\Xi_c^+$ DECAY MODES                               | Fraction ( $\Gamma_i/\Gamma$ )     | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|-----------------------------------------------------|------------------------------------|-----------------------------------|----------------|
| <b>Cabibbo-favored (<math>S = -2</math>) decays</b> |                                    |                                   |                |
| $p 2K_S^0$                                          | $(2.5 \pm 1.3) \times 10^{-3}$     |                                   | 766            |
| $\Lambda \bar{K}^0 \pi^+$                           | —                                  |                                   | 852            |
| $\Sigma(1385)^+ \bar{K}^0$                          | [q] $(2.9 \pm 2.0) \%$             |                                   | 746            |
| $\Lambda K^- 2\pi^+$                                | $(9 \pm 4) \times 10^{-3}$         |                                   | 787            |
| $\Lambda \bar{K}^*(892)^0 \pi^+$                    | [q] $< 5 \times 10^{-3}$           | CL=90%                            | 608            |
| $\Sigma(1385)^+ K^- \pi^+$                          | [q] $< 6 \times 10^{-3}$           | CL=90%                            | 678            |
| $\Sigma^+ K^- \pi^+$                                | $(2.7 \pm 1.2) \%$                 |                                   | 810            |
| $\Sigma^+ \bar{K}^*(892)^0$                         | [q] $(2.3 \pm 1.1) \%$             |                                   | 658            |
| $\Sigma^0 K^- 2\pi^+$                               | $(8 \pm 5) \times 10^{-3}$         |                                   | 735            |
| $\Xi^0 \pi^+$                                       | $(1.6 \pm 0.8) \%$                 |                                   | 876            |
| $\Xi^- 2\pi^+$                                      | $(2.9 \pm 1.3) \%$                 |                                   | 851            |
| $\Xi(1530)^0 \pi^+$                                 | [q] $< 2.9 \times 10^{-3}$         | CL=90%                            | 749            |
| $\Xi(1620)^0 \pi^+$                                 | seen                               |                                   | —              |
| $\Xi(1690)^0 \pi^+$                                 | seen                               |                                   | 644            |
| $\Xi^0 \pi^+ \pi^0$                                 | $(6.7 \pm 3.5) \%$                 |                                   | 856            |
| $\Xi^0 \pi^- 2\pi^+$                                | $(5.0 \pm 2.6) \%$                 |                                   | 818            |
| $\Xi^0 e^+ \nu_e$                                   | $(7 \pm 4) \%$                     |                                   | 884            |
| $\Omega^- K^+ \pi^+$                                | $(2.0 \pm 1.5) \times 10^{-3}$     |                                   | 399            |
| <b>Cabibbo-suppressed decays</b>                    |                                    |                                   |                |
| $p K^- \pi^+$                                       | $(6.2 \pm 3.0) \times 10^{-3}$     | S=1.5                             | 944            |
| $p \bar{K}^*(892)^0$                                | [q] $(3.3 \pm 1.7) \times 10^{-3}$ |                                   | 828            |
| $\Sigma^+ \pi^+ \pi^-$                              | $(1.4 \pm 0.8) \%$                 |                                   | 922            |
| $\Sigma^- 2\pi^+$                                   | $(5.1 \pm 3.4) \times 10^{-3}$     |                                   | 918            |
| $\Sigma^+ K^+ K^-$                                  | $(4.3 \pm 2.5) \times 10^{-3}$     |                                   | 579            |
| $\Sigma^+ \phi$                                     | [q] $< 3.2 \times 10^{-3}$         | CL=90%                            | 549            |
| $\Xi(1690)^0 K^+, \Xi^0 \rightarrow \Sigma^+ K^-$   | $< 1.3 \times 10^{-3}$             | CL=90%                            | 501            |
| $p \phi(1020)$                                      | $(1.2 \pm 0.6) \times 10^{-4}$     |                                   | 751            |



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

$$\text{Mass } m = 2470.44 \pm 0.28 \text{ MeV} \quad (S = 1.2)$$

$$m_{\Xi_c^0} - m_{\Xi_c^+} = 2.72 \pm 0.23 \text{ MeV} \quad (S = 1.1)$$

$$\text{Mean life } \tau = (150.4 \pm 2.8) \times 10^{-15} \text{ s} \quad (S = 1.4)$$

$$c\tau = 45.1 \text{ } \mu\text{m}$$

### Decay asymmetry parameters

$$\Xi^- \pi^+ \quad \alpha = -0.64 \pm 0.05$$

$$\alpha \text{ FOR } \Xi_c^0 \rightarrow \Xi^+ \pi^- = 0.61 \pm 0.05$$

$$\alpha \text{ FOR } \Xi_c^0 \rightarrow \Lambda \bar{K}^*(892)^0 = 0.15 \pm 0.22$$

$$\alpha \text{ FOR } \Xi_c^0 \rightarrow \Sigma^+ K^*(892)^- = -0.52 \pm 0.30$$

$$\alpha \text{ FOR } \Xi_c^0 \rightarrow \Xi^0 \pi^0 = -0.90 \pm 0.27$$

$$\tau_{mix}, \Xi_c^0 - \Xi_c^+ \text{ oscillation period} > 1.3 \times 10^{-12} \text{ s}$$

| $\Xi_c^0$ DECAY MODES                                        | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $\frac{P}{\text{MeV}/c}$ |
|--------------------------------------------------------------|--------------------------------|------------------|--------------------------|
| <b>Cabibbo-favored decays</b>                                |                                |                  |                          |
| $p K^- K^- \pi^+$                                            | $(4.9 \pm 1.0) \times 10^{-3}$ |                  | 676                      |
| $p K^- \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow K^- \pi^+$ | $(2.0 \pm 0.6) \times 10^{-3}$ |                  | 413                      |
| $p K^- K^- \pi^+ (\text{no } \bar{K}^{*0})$                  | $(3.0 \pm 0.8) \times 10^{-3}$ |                  | 676                      |
| $\Lambda K_S^0$                                              | $(3.2 \pm 0.6) \times 10^{-3}$ |                  | 906                      |
| $\Lambda K^- \pi^+$                                          | $(1.45 \pm 0.28) \%$           |                  | 856                      |
| $\Lambda \bar{K}^*(892)^0$                                   | $(2.6 \pm 0.6) \times 10^{-3}$ |                  | 717                      |
| $\Lambda \bar{K}^0 \pi^+ \pi^-$                              | seen                           |                  | 786                      |
| $\Lambda K^- \pi^+ \pi^+ \pi^-$                              | seen                           |                  | 703                      |
| $\Sigma^0 K_S^0$                                             | $(5.4 \pm 1.4) \times 10^{-4}$ |                  | 864                      |
| $\Sigma^+ K^-$                                               | $(1.8 \pm 0.4) \times 10^{-3}$ |                  | 868                      |
| $\Sigma^0 \bar{K}^*(892)^0$                                  | $(9.9 \pm 1.9) \times 10^{-3}$ |                  | 658                      |
| $\Sigma^+ K^*(892)^-$                                        | $(4.9 \pm 1.3) \times 10^{-3}$ |                  | 661                      |
| $\Xi^- \pi^+$                                                | $(1.43 \pm 0.27) \%$           |                  | 875                      |
| $\Xi^- \pi^+ \pi^+ \pi^-$                                    | $(4.8 \pm 2.3) \%$             |                  | 816                      |
| $\Xi^0 \pi^0$                                                | $(6.9 \pm 1.4) \times 10^{-3}$ |                  | 879                      |
| $\Xi^0 \eta$                                                 | $(1.6 \pm 0.4) \times 10^{-3}$ |                  | 771                      |
| $\Xi^0 \eta'$                                                | $(1.1 \pm 0.4) \times 10^{-3}$ |                  | 479                      |
| $\Xi^0 \phi, \phi \rightarrow K^+ K^-$                       | $(5.2 \pm 1.2) \times 10^{-4}$ |                  | —                        |
| $\Xi^0 K^+ K^- \text{ nonresonant}$                          | $(5.6 \pm 1.2) \times 10^{-4}$ |                  | 444                      |
| $\Omega^- K^+$                                               | $(4.2 \pm 0.9) \times 10^{-3}$ |                  | 522                      |
| $\Xi^- e^+ \nu_e$                                            | $(1.05 \pm 0.20) \%$           |                  | 882                      |
| $\Xi^- \mu^+ \nu_\mu$                                        | $(1.01 \pm 0.21) \%$           |                  | 878                      |
| $\Xi^0 \gamma$                                               | $< 1.7 \times 10^{-4}$         | 90%              | 885                      |
| $\Xi^0 \mu^+ \mu^-$                                          | $< 6 \times 10^{-5}$           | 90%              | 869                      |
| $\Xi^0 e^+ e^-$                                              | $< 1.0 \times 10^{-4}$         | 90%              | 885                      |

Cabibbo-suppressed decays

|                                |                                |     |
|--------------------------------|--------------------------------|-----|
| $\Lambda_c^+ \pi^-$            | $(5.5 \pm 1.1) \times 10^{-3}$ | 115 |
| $\Xi^- K^+$                    | $(3.9 \pm 1.1) \times 10^{-4}$ | 789 |
| $\Lambda K^+ K^-$ (no $\phi$ ) | $(4.1 \pm 1.3) \times 10^{-4}$ | 648 |
| $\Lambda \phi$                 | $(4.9 \pm 1.3) \times 10^{-4}$ | 621 |



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.


Mass  $m = 2578.2 \pm 0.5$  MeV (S = 1.1)

$$m_{\Xi_c'^+} - m_{\Xi_c^+} = 110.5 \pm 0.4 \text{ MeV}$$

$$m_{\Xi_c'^+} - m_{\Xi_c'^0} = -0.5 \pm 0.6 \text{ MeV}$$

The  $\Xi_c'^+ - \Xi_c^+$  mass difference is too small for any strong decay to occur.

| $\Xi_c'^+$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------|--------------------------------|-------------|
| $\Xi_c^+ \gamma$       | seen                           | 108         |



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2578.7 \pm 0.5$  MeV

$$m_{\Xi_c'^0} - m_{\Xi_c^0} = 108.3 \pm 0.4 \text{ MeV}$$

The  $\Xi_c^{\prime 0} - \Xi_c^0$  mass difference is too small for any strong decay to occur.

| $\Xi_c^{\prime 0}$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^{\prime 0} \gamma$      | seen                           | 106         |

### $\Xi_c(2645)$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

$$\Xi_c(2645)^+ \text{ mass } m = 2645.10 \pm 0.30 \text{ MeV} \quad (S = 1.2)$$

$$\Xi_c(2645)^0 \text{ mass } m = 2646.16 \pm 0.25 \text{ MeV} \quad (S = 1.3)$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c^0} = 174.67 \pm 0.09 \text{ MeV}$$

$$m_{\Xi_c(2645)^0} - m_{\Xi_c^+} = 178.45 \pm 0.10 \text{ MeV}$$

$$m_{\Xi_c(2645)^+} - m_{\Xi_c(2645)^0} = -1.06 \pm 0.27 \text{ MeV} \quad (S = 1.1)$$

$$\Xi_c(2645)^+ \text{ full width } \Gamma = 2.14 \pm 0.19 \text{ MeV} \quad (S = 1.1)$$

$$\Xi_c(2645)^0 \text{ full width } \Gamma = 2.35 \pm 0.22 \text{ MeV}$$

$\Xi_c \pi$  is the only strong decay allowed to a  $\Xi_c$  resonance having this mass.

| $\Xi_c(2645)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Xi_c^0 \pi^+$           | seen                           | 102         |
| $\Xi_c^+ \pi^-$           | seen                           | 106         |

### $\Xi_c(2790)$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

$J^P$  has not been measured;  $\frac{1}{2}^-$  is the quark-model prediction.

$$\Xi_c(2790)^+ \text{ mass } = 2791.9 \pm 0.5 \text{ MeV}$$

$$\Xi_c(2790)^0 \text{ mass } = 2793.9 \pm 0.5 \text{ MeV}$$

$$m_{\Xi_c(2790)^+} - m_{\Xi_c^{\prime 0}} = 213.20 \pm 0.22 \text{ MeV}$$

$$m_{\Xi_c(2790)^0} - m_{\Xi_c^{\prime +}} = 215.70 \pm 0.22 \text{ MeV}$$

$$m_{\Xi_c(2790)^+} - m_{\Xi_c(2790)^0} = -2.0 \pm 0.7 \text{ MeV}$$

$$\Xi_c(2790)^+ \text{ width } = 8.9 \pm 1.0 \text{ MeV}$$

$$\Xi_c(2790)^0 \text{ width } = 10.0 \pm 1.1 \text{ MeV}$$

| $\Xi_c(2790)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Xi_c' \pi$              | seen                           | 159         |
| $\Lambda_c^+ K^-$         | seen                           | 98          |

 **$\Xi_c(2815)$** 

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

$J^P$  has not been measured;  $\frac{3}{2}^-$  is the quark-model prediction.

$$\Xi_c(2815)^+ \text{ mass } m = 2816.51 \pm 0.25 \text{ MeV} \quad (S = 1.2)$$

$$\Xi_c(2815)^0 \text{ mass } m = 2819.79 \pm 0.30 \text{ MeV} \quad (S = 1.1)$$

$$m_{\Xi_c(2815)^+} - m_{\Xi_c^+} = 348.80 \pm 0.10 \text{ MeV}$$

$$m_{\Xi_c(2815)^0} - m_{\Xi_c^0} = 349.35 \pm 0.11 \text{ MeV}$$

$$m_{\Xi_c(2815)^+} - m_{\Xi_c(2815)^0} = -3.27 \pm 0.27 \text{ MeV}$$

$$\Xi_c(2815)^+ \text{ full width } \Gamma = 2.43 \pm 0.26 \text{ MeV}$$

$$\Xi_c(2815)^0 \text{ full width } \Gamma = 2.54 \pm 0.25 \text{ MeV}$$

The  $\Xi_c \pi \pi$  modes are consistent with being entirely via  $\Xi_c(2645) \pi$ .

| $\Xi_c(2815)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Xi_c' \pi$              | seen                           | 188         |
| $\Xi_c(2645) \pi$         | seen                           | 102         |
| $\Xi_c^0 \gamma$          | seen                           | 325         |

 **$\Xi_c(2970)$** 

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

was  $\Xi_c(2980)$

$$\Xi_c(2970)^+ m = 2964.3 \pm 1.5 \text{ MeV} \quad (S = 3.9)$$

$$\Xi_c(2970)^0 m = 2967.1 \pm 1.7 \text{ MeV} \quad (S = 6.7)$$

$$m_{\Xi_c(2970)^+} - m_{\Xi_c^+} = 496.6 \pm 1.5 \text{ MeV} \quad (S = 3.7)$$

$$m_{\Xi_c(2970)^0} - m_{\Xi_c^0} = 496.7 \pm 1.8 \text{ MeV} \quad (S = 5.3)$$

$$m_{\Xi_c(2970)^+} - m_{\Xi_c(2970)^0} = -2.8 \pm 1.9 \text{ MeV} \quad (S = 4.8)$$

$$\Xi_c(2970)^+ \text{ width } \Gamma = 20.9_{-3.5}^{+2.4} \text{ MeV} \quad (S = 1.2)$$

| $\Xi_c(2970)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \bar{K} \pi$ | seen                           | 223         |
| $\Sigma_c(2455) \bar{K}$  | seen                           | 122         |
| $\Lambda_c^+ \bar{K}$     | not seen                       | 410         |
| $\Lambda_c^+ K^-$         | seen                           | 410         |
| $\Xi_c 2\pi$              | seen                           | 381         |
| $\Xi_c' \pi$              | seen                           | —           |
| $\Xi_c(2645) \pi$         | seen                           | 274         |

$\Xi_c(3055)$

$I(J^P) = ?(??)$

Mass  $m = 3055.9 \pm 0.4$  MeV  
Full width  $\Gamma = 7.8 \pm 1.9$  MeV

| $\Xi_c(3055)$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------|--------------------------------|-------------|
| $\Sigma^{++} K^-$         | seen                           | —           |
| $\Lambda D^+$             | seen                           | 316         |

$\Xi_c(3080)$

$I(J^P) = \frac{1}{2}(??)$

$\Xi_c(3080)^+ m = 3077.2 \pm 0.4$  MeV  
 $\Xi_c(3080)^0 m = 3079.9 \pm 1.4$  MeV ( $S = 1.3$ )  
 $\Xi_c(3080)^+$  width  $\Gamma = 3.6 \pm 1.1$  MeV ( $S = 1.5$ )  
 $\Xi_c(3080)^0$  width  $\Gamma = 5.6 \pm 2.2$  MeV

| $\Xi_c(3080)$ DECAY MODES                         | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------------------------|--------------------------------|-------------|
| $\Lambda_c^+ \bar{K} \pi$                         | seen                           | 415         |
| $\Sigma_c(2455) \bar{K}$                          | seen                           | 342         |
| $\Sigma_c(2455)^{++} K^-$                         | seen                           | 342         |
| $\Sigma_c(2520)^{++} K^-$                         | seen                           | 239         |
| $\Sigma_c(2455) \bar{K} + \Sigma_c(2520) \bar{K}$ | seen                           | —           |
| $\Lambda_c^+ \bar{K}$                             | not seen                       | 536         |
| $\Lambda_c^+ \bar{K} \pi^+ \pi^-$                 | not seen                       | 144         |
| $\Lambda D^+$                                     | seen                           | 362         |

$\Omega_c^0$

$I(J^P) = 0(\frac{1}{2}^+)$

$J^P$  has not been measured;  $\frac{1}{2}^+$  is the quark-model prediction.

Mass  $m = 2695.3 \pm 0.4$  MeV

Mean life  $\tau = (273 \pm 12) \times 10^{-15}$  s

$c\tau = 82$   $\mu\text{m}$

No absolute branching fractions have been measured. The following are branching *ratios* relative to  $\Omega^- \pi^+$ .

| $\Omega_c^0$ DECAY MODES                                                                      | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$ (MeV/c) |
|-----------------------------------------------------------------------------------------------|--------------------------------|------------------|-------------|
| <b>Cabibbo-favored (<math>S = -3</math>) decays — relative to <math>\Omega^- \pi^+</math></b> |                                |                  |             |
| $\Omega^- \pi^+$                                                                              | <b>DEFINED AS 1</b>            |                  | 821         |
| $\Omega^- \pi^+ \pi^0$                                                                        | $1.80 \pm 0.33$                |                  | 797         |
| $\Omega^- \rho^+$                                                                             | $>1.3$                         | 90%              | 532         |
| $\Omega^- \pi^- 2\pi^+$                                                                       | $0.31 \pm 0.05$                |                  | 753         |
| $\Omega^- e^+ \nu_e$                                                                          | $1.98 \pm 0.29$                |                  | 829         |
| $\Omega^- \mu^+ \nu_\mu$                                                                      | $1.94 \pm 0.21$                |                  | 824         |
| $\Xi^0 \bar{K}^0$                                                                             | $1.64 \pm 0.29$                |                  | 950         |
| $\Xi^0 K^- \pi^+$                                                                             | $1.20 \pm 0.18$                |                  | 901         |
| $\Xi^0 \bar{K}^{*0}, \bar{K}^{*0} \rightarrow K^- \pi^+$                                      | $0.68 \pm 0.16$                |                  | 764         |
| $\Omega(2012)^- \pi^+, \Omega(2012)^- \rightarrow \Xi^0 K^-$                                  | $0.12 \pm 0.05$                |                  | —           |
| $\Xi^- \bar{K}^0 \pi^+$                                                                       | $2.12 \pm 0.28$                |                  | 895         |
| $\Omega(2012)^- \pi^+, \Omega(2012)^- \rightarrow \Xi^- \bar{K}^0$                            | $0.12 \pm 0.06$                |                  | —           |
| $\Xi^- K^- 2\pi^+$                                                                            | $0.63 \pm 0.09$                |                  | 830         |
| $\Xi(1530)^0 K^- \pi^+, \Xi^{*0} \rightarrow \Xi^- \pi^+$                                     | $0.21 \pm 0.06$                |                  | 757         |

|                               |                 |     |     |
|-------------------------------|-----------------|-----|-----|
| $\Xi^- \bar{K}^{*0} \pi^+$    | $0.34 \pm 0.11$ |     | 653 |
| $p K^- K^- \pi^+$             | seen            |     | 864 |
| $\Sigma^+ K^- K^- \pi^+$      | $<0.32$         | 90% | 689 |
| $\Lambda \bar{K}^0 \bar{K}^0$ | $1.72 \pm 0.35$ |     | 837 |

**Singly Cabibbo-suppressed modes — relative to  $\Omega^- \pi^+$**

|                |                   |   |
|----------------|-------------------|---|
| $\Xi^- \pi^+$  | $0.161 \pm 0.010$ | — |
| $\Omega^- K^+$ | $0.061 \pm 0.006$ | — |

**Doubly Cabibbo-suppressed modes — relative to  $\Omega^- \pi^+$**

|             |         |     |   |
|-------------|---------|-----|---|
| $\Xi^- K^+$ | $<0.07$ | 90% | — |
|-------------|---------|-----|---|

$\Omega_c(2770)^0$

$I(J^P) = 0(\frac{3}{2}^+)$

$J^P$  has not been measured;  $\frac{3}{2}^+$  is the quark-model prediction.

Mass  $m = 2766.0^{+0.9}_{-1.0}$  MeV

$m_{\Omega_c(2770)^0} - m_{\Omega_c^0} = 70.7^{+0.8}_{-0.9}$  MeV

The  $\Omega_c(2770)^0$ – $\Omega_c^0$  mass difference is too small for any strong decay to occur.

| $\Omega_c(2770)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Omega_c^0 \gamma$            | presumably 100%                | 70          |

$\Omega_c(3000)^0$

$I(J^P) = ?(?^?)$

Mass  $m = 3000.46 \pm 0.25$  MeV

Full width  $\Gamma = 3.8^{+1.6}_{-0.4}$  MeV

| $\Omega_c(3000)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^+ K^-$                  | seen                           | 182         |

$\Omega_c(3050)^0$

$I(J^P) = ?(?^?)$

Mass  $m = 3050.17 \pm 0.19$  MeV

Full width  $\Gamma < 1.8$  MeV, CL = 95%

| $\Omega_c(3050)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^+ K^-$                  | seen                           | 278         |

$\Omega_c(3065)^0$

$I(J^P) = ?(??)$

Mass  $m = 3065.58 \pm 0.21$  MeV  
Full width  $\Gamma = 3.4^{+0.7}_{-0.8}$  MeV ( $S = 1.7$ )

| $\Omega_c(3065)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^+ K^-$                  | seen                           | 303         |

$\Omega_c(3090)^0$

$I(J^P) = ?(??)$

Mass  $m = 3090.15 \pm 0.26$  MeV  
Full width  $\Gamma = 8.5^{+0.8}_{-1.7}$  MeV

| $\Omega_c(3090)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^+ K^-$                  | seen                           | 340         |

$\Omega_c(3120)^0$

$I(J^P) = ?(??)$

Mass  $m = 3118.98^{+0.27}_{-0.35}$  MeV  
Full width  $\Gamma < 2.5$  MeV, CL = 95%

| $\Omega_c(3120)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^+ K^-$                  | seen                           | 379         |

$\Omega_c(3185)^0$

$I(J^P) = ?(??)$

Mass  $m = 3185^{+7.6}_{-1.9}$  MeV  
Full width  $\Gamma = 50^{+12}_{-21}$  MeV

| $\Omega_c(3185)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^+ K^-$                  | seen                           | 460         |

$\Omega_c(3327)^0$

$I(J^P) = ?(?^?)$

Mass  $m = 3327.1^{+1.2}_{-1.8}$  MeV

Full width  $\Gamma = 20^{+14}_{-5}$  MeV

| $\Omega_c(3327)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_c^+ K^-$                  | seen                           | 610         |

DOUBLY CHARMED BARYONS

( $C = +2$ )

$\Xi_{cc}^{++} = ucc, \Xi_{cc}^+ = dcc, \Omega_{cc}^+ = scc$

$\Xi_{cc}^{++}$

$I(J^P) = ?(?^?)$

Mass  $m = 3621.6 \pm 0.4$  MeV

Mean life  $\tau = (256 \pm 27) \times 10^{-15}$  s

| $\Xi_{cc}^{++}$ DECAY MODES                                                                          | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$ (MeV/c) |
|------------------------------------------------------------------------------------------------------|--------------------------------|------------------|-------------|
| $\Lambda_c^+ K^- \pi^+ \pi^+$                                                                        | <b>DEFINED AS 1</b>            |                  | 880         |
| $\Xi_c^+ \pi^+, \Xi_c^+ \rightarrow p K^- \pi^+$                                                     | $0.0022 \pm 0.0006$            |                  | —           |
| $\Xi_c^{\prime+} \pi^+, \Xi_c^{\prime+} \rightarrow \Xi_c^+ \gamma, \Xi_c^+ \rightarrow p K^- \pi^+$ | $0.0031 \pm 0.0010$            |                  | —           |
| $D^+ p K^- \pi^+$                                                                                    | $<0.017$                       | 90%              | 562         |

# BOTTOM BARYONS

## ( $B = -1$ )

$$\Lambda_b^0 = udb, \Sigma_b^0 = udb, \Sigma_b^+ = uub, \Sigma_b^- = ddb$$

$$\Xi_b^0 = usb, \Xi_b^- = dsb, \Omega_b^- = ssb$$

$\Lambda_b^0$

$$I(J^P) = 0(\frac{1}{2}^+)$$

$I(J^P)$  not yet measured;  $0(\frac{1}{2}^+)$  is the quark model prediction.

$$\text{Mass } m = 5619.57 \pm 0.16 \text{ MeV}$$

$$m_{\Lambda_b^0} - m_{B^0} = 339.2 \pm 1.4 \text{ MeV}$$

$$m_{\Lambda_b^0} - m_{B^+} = 339.72 \pm 0.28 \text{ MeV}$$

$$\text{Mean life } \tau = (1.468 \pm 0.009) \times 10^{-12} \text{ s}$$

$$c\tau = 440.1 \text{ } \mu\text{m}$$

$$A_{CP}(\Lambda_b \rightarrow p\pi^-) = -0.025 \pm 0.029 \quad (S = 1.2)$$

$$A_{CP}(\Lambda_b \rightarrow pK^-) = -0.025 \pm 0.022$$

$$A_{CP}(\Lambda_b \rightarrow DpK^-) = 0.12 \pm 0.09$$

$$A_{CP}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = 0.007 \pm 0.009$$

$$A_{CP}(\Lambda_b^0 \rightarrow \Lambda_c^+ K^-) = -0.032 \pm 0.030$$

$$\Delta A_{CP}(pK^-/\pi^-) = 0.014 \pm 0.024$$

$$A_{CP}(\Lambda_b \rightarrow p\bar{K}^0\pi^-) = 0.22 \pm 0.13$$

$$\Delta A_{CP}(J/\psi p\pi^-/K^-) = (5.7 \pm 2.7) \times 10^{-2}$$

$$A_{CP}(\Lambda_b \rightarrow \Lambda K^+\pi^-) = -0.53 \pm 0.25$$

$$A_{CP}(\Lambda_b \rightarrow \Lambda K^+K^-) = -0.28 \pm 0.12$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow pK^- \mu^+ \mu^-) = (-4 \pm 5) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow p\pi^- \pi^+ \pi^-) = (1.1 \pm 2.6) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow (p\pi^- \pi^+ \pi^-)_{LBM}) = (4 \pm 4) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow p a_1(1260)^-) = (-1 \pm 4) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow N(1520)^0 \rho(770)^0) = (2 \pm 5) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow \Delta(1232)^{++} \pi^- \pi^-) = (0.1 \pm 3.3) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow pK^- \pi^+ \pi^-) = (3.2 \pm 1.3) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow (pK^- \pi^+ \pi^-)_{LBM}) = (3.5 \pm 1.6) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow N(1520)^0 K^*(892)^0) = (5.5 \pm 2.5) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow \Lambda(1520) \rho(770)^0) = (1 \pm 6) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow \Delta(1232)^{++} K^- \pi^-) = (4.4 \pm 2.7) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow pK_1(1410)^-) = (5 \pm 4) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow pK^- K^+ \pi^-) = (-7 \pm 5) \times 10^{-2}$$

$$\Delta A_{CP}(\Lambda_b^0 \rightarrow pK^- K^+ K^-) = (0.2 \pm 1.9) \times 10^{-2}$$

$$\begin{aligned}
\Delta A_{CP}(\Lambda_b^0 \rightarrow \Lambda(1520)\phi(1020)) &= (4 \pm 6) \times 10^{-2} \\
\Delta A_{CP}(\Lambda_b^0 \rightarrow (pK^-)_{highmass} \phi(1020)) &= (-0.7 \pm 3.4) \times 10^{-2} \\
\Delta A_{CP}(\Lambda_b^0 \rightarrow (pK^- K^+ K^-)_{LBM}) &= (2.7 \pm 2.4) \times 10^{-2} \\
A_{FB}^\ell(\mu\mu) \text{ in } \Lambda_b \rightarrow \Lambda\mu^+\mu^- &= -0.39 \pm 0.04 \\
\Delta(A_{FB}^\ell(\mu\mu)) \text{ in } \Lambda_b \rightarrow \Lambda\mu^+\mu^- &= -0.05 \pm 0.09 \\
A_{FB}^h(p\pi) \text{ in } \Lambda_b \rightarrow \Lambda(p\pi)\mu^+\mu^- &= -0.30 \pm 0.05 \\
A_{FB}^{\ell h} \text{ in } \Lambda_b \rightarrow \Lambda\mu^+\mu^- &= 0.25 \pm 0.04
\end{aligned}$$

The branching fractions  $B(b\text{-baryon} \rightarrow \Lambda\ell^-\bar{\nu}_\ell \text{ anything})$  and  $B(\Lambda_b^0 \rightarrow \Lambda_c^+\ell^-\bar{\nu}_\ell \text{ anything})$  are not pure measurements because the underlying measured products of these with  $B(b \rightarrow b\text{-baryon})$  were used to determine  $B(b \rightarrow b\text{-baryon})$ , as described in the note “Production and Decay of  $b$ -Flavored Hadrons.”

For inclusive branching fractions, e.g.,  $\Lambda_b \rightarrow \bar{\Lambda}_c \text{ anything}$ , the values usually are multiplicities, not branching fractions. They can be greater than one.

| $\Lambda_b^0$ DECAY MODES                                         | Fraction ( $\Gamma_i/\Gamma$ )           | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|-------------------------------------------------------------------|------------------------------------------|-----------------------------------|----------------|
| $J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)$           | $(5.8 \pm 0.8) \times 10^{-5}$           |                                   | 1740           |
| $pD^0\pi^-$                                                       | $(6.3 \pm 0.6) \times 10^{-4}$           |                                   | 2370           |
| $pD^+\pi^-\pi^-$                                                  | $(2.8 \pm 0.4) \times 10^{-4}$           |                                   | 2332           |
| $pD^*(2010)^+\pi^-\pi^-$                                          | $(5.3 \pm 1.0) \times 10^{-4}$           |                                   | 2277           |
| $pD^0K^-$                                                         | $(4.6 \pm 0.8) \times 10^{-5}$           |                                   | 2269           |
| $pJ/\psi\pi^-$                                                    | $(2.6 \pm_{-0.4}^{+0.5}) \times 10^{-5}$ |                                   | 1755           |
| $p\pi^- J/\psi, J/\psi \rightarrow \mu^+\mu^-$                    | $(1.6 \pm 0.8) \times 10^{-6}$           |                                   | —              |
| $pJ/\psi K^-$                                                     | $(3.2 \pm_{-0.5}^{+0.6}) \times 10^{-4}$ |                                   | 1589           |
| $p\eta_c(1S)K^-$                                                  | $(1.06 \pm 0.26) \times 10^{-4}$         |                                   | 1670           |
| $P_{c\bar{c}}(4312)^+K^-, P_{c\bar{c}}^+ \rightarrow p\eta_c(1S)$ | $< 2.5 \times 10^{-5}$                   | CL=95%                            | —              |
| $P_{c\bar{c}}(4380)^+K^-, P_{c\bar{c}}^+ \rightarrow pJ/\psi$     | [v] $(2.7 \pm 1.4) \times 10^{-5}$       |                                   | —              |
| $P_c(4450)^+K^-, P_c \rightarrow pJ/\psi$                         | [v] $(1.3 \pm 0.4) \times 10^{-5}$       |                                   | —              |
| $\chi_{c1}(1P)pK^-$                                               | $(7.6 \pm_{-1.3}^{+1.5}) \times 10^{-5}$ |                                   | 1242           |
| $\chi_{c1}(1P)p\pi^-$                                             | $(5.0 \pm_{-1.1}^{+1.3}) \times 10^{-6}$ |                                   | 1462           |
| $\chi_{c2}(1P)pK^-$                                               | $(7.7 \pm_{-1.4}^{+1.6}) \times 10^{-5}$ |                                   | 1198           |
| $\chi_{c2}(1P)p\pi^-$                                             | $(4.8 \pm 1.9) \times 10^{-6}$           |                                   | 1427           |
| $pJ/\psi(1S)\pi^+\pi^-K^-$                                        | $(6.6 \pm_{-1.1}^{+1.3}) \times 10^{-5}$ |                                   | 1410           |
| $p\psi(2S)K^-$                                                    | $(6.6 \pm_{-1.0}^{+1.2}) \times 10^{-5}$ |                                   | 1063           |

|                                                                                |                                      |        |      |
|--------------------------------------------------------------------------------|--------------------------------------|--------|------|
| $\chi_{c1}(3872)pK^-$                                                          | $(2.8 \pm 1.2) \times 10^{-5}$       |        | 837  |
| $\chi_{c1}(3872)\Lambda(1520)$                                                 | $(1.6 \pm 0.8) \times 10^{-5}$       |        | 721  |
| $\psi(2S)p\pi^-$                                                               | $(7.5^{+1.6}_{-1.4}) \times 10^{-6}$ |        | 1320 |
| $p\bar{K}^0\pi^-$                                                              | $(1.3 \pm 0.4) \times 10^{-5}$       |        | 2693 |
| $pK^0K^-$                                                                      | $< 3.5 \times 10^{-6}$               | CL=90% | 2639 |
| $\Lambda_c^+\pi^-$                                                             | $(4.9 \pm 0.4) \times 10^{-3}$       | S=1.2  | 2342 |
| $\Lambda_c^+K^-$                                                               | $(3.56 \pm 0.28) \times 10^{-4}$     | S=1.2  | 2314 |
| $\Lambda_c^+a_1(1260)^-$                                                       | seen                                 |        | 2153 |
| $\Lambda_c^+D^-$                                                               | $(4.6 \pm 0.6) \times 10^{-4}$       |        | 1886 |
| $\Lambda_c^+D_s^-$                                                             | $(1.10 \pm 0.10) \%$                 |        | 1833 |
| $\Lambda_c^+D_s^{*-}$                                                          | $(1.83 \pm 0.18) \%$                 |        | 1748 |
| $\Lambda_c^+\bar{D}^0K^-$                                                      | $(2.13 \pm 0.20) \times 10^{-3}$     |        | 1581 |
| $\Lambda_c^+\bar{D}^{*0}K^-$                                                   | $(6.6 \pm 0.7) \times 10^{-3}$       |        | 1471 |
| $\Lambda_c^+\pi^+\pi^-\pi^-$                                                   | $(7.6 \pm 1.1) \times 10^{-3}$       | S=1.1  | 2323 |
| $\Lambda_c(2595)^+\pi^-$                                                       | $(3.4 \pm 1.4) \times 10^{-4}$       |        | 2210 |
| $\Lambda_c(2595)^+ \rightarrow \Lambda_c^+\pi^+\pi^-$                          |                                      |        |      |
| $\Lambda_c(2625)^+\pi^-$                                                       | $(3.3 \pm 1.3) \times 10^{-4}$       |        | 2193 |
| $\Lambda_c(2625)^+ \rightarrow \Lambda_c^+\pi^+\pi^-$                          |                                      |        |      |
| $\Sigma_c(2455)^0\pi^+\pi^-$ , $\Sigma_c^0 \rightarrow \Lambda_c^+\pi^-$       | $(5.7 \pm 2.2) \times 10^{-4}$       |        | 2265 |
| $\Sigma_c(2455)^{++}\pi^-\pi^-$ , $\Sigma_c^{++} \rightarrow \Lambda_c^+\pi^+$ | $(3.2 \pm 1.5) \times 10^{-4}$       |        | 2265 |
| $\Sigma_c(2455)^{++}D^-K^-$                                                    | $(6.0 \pm 0.8) \times 10^{-4}$       |        | 1448 |
| $\Sigma_c(2455)^{++}D^{*-}K^-$                                                 | $(1.36 \pm 0.22) \times 10^{-3}$     |        | 1324 |
| $\Sigma_c(2520)^{++}D^-K^-$                                                    | $(2.8 \pm 0.5) \times 10^{-4}$       |        | 1392 |
| $\Sigma_c(2520)^{++}D^{*-}K^-$                                                 | $(5.4 \pm 1.1) \times 10^{-4}$       |        | 1262 |
| $\Lambda_c^+K^+K^-\pi^-$                                                       | $(1.02 \pm 0.11) \times 10^{-3}$     |        | 2184 |
| $\Lambda_c^+p\bar{p}\pi^-$                                                     | $(2.63 \pm 0.27) \times 10^{-4}$     |        | 1805 |
| $\Sigma_c(2455)^0p\bar{p}$ , $\Sigma_c^0 \rightarrow \Lambda_c^+\pi^-$         | $(2.3 \pm 0.5) \times 10^{-5}$       |        | —    |
| $\Sigma_c(2520)^0p\bar{p}$ , $\Sigma_c(2520)^0 \rightarrow \Lambda_c^+\pi^-$   | $(3.1 \pm 0.7) \times 10^{-5}$       |        | —    |
| $\Lambda_c^+\ell^-\bar{\nu}_\ell$ anything                                     | [x] $(10.9 \pm 2.2) \%$              |        | —    |
| $\Lambda_c^+\ell^-\bar{\nu}_\ell$                                              | $(6.2^{+1.4}_{-1.3}) \%$             |        | 2345 |
| $\Lambda_c^+\tau^-\bar{\nu}_\tau$                                              | $(1.9 \pm 0.5) \%$                   |        | 1933 |
| $\Lambda_c^+\pi^+\pi^-\ell^-\bar{\nu}_\ell$                                    | $(5.6 \pm 3.1) \%$                   |        | 2335 |
| $\Lambda_c(2595)^+\ell^-\bar{\nu}_\ell$                                        | $(7.9^{+4.0}_{-3.5}) \times 10^{-3}$ |        | 2212 |
| $\Lambda_c(2625)^+\ell^-\bar{\nu}_\ell$                                        | $(1.3^{+0.6}_{-0.5}) \%$             |        | 2195 |

|                         |                                      |        |      |
|-------------------------|--------------------------------------|--------|------|
| $p h^-$                 | $[y] < 2.3 \times 10^{-5}$           | CL=90% | 2730 |
| $p \pi^-$               | $(4.6 \pm 0.8) \times 10^{-6}$       |        | 2730 |
| $p K^-$                 | $(5.5 \pm 1.0) \times 10^{-6}$       |        | 2709 |
| $p D_s^-$               | $(1.25 \pm 0.13) \times 10^{-5}$     |        | 2364 |
| $p \mu^- \bar{\nu}_\mu$ | $(4.1 \pm 1.0) \times 10^{-4}$       |        | 2730 |
| $\Lambda \mu^+ \mu^-$   | $(1.08 \pm 0.28) \times 10^{-6}$     |        | 2695 |
| $p \pi^- \mu^+ \mu^-$   | $(6.9 \pm 2.5) \times 10^{-8}$       |        | 2720 |
| $p K^- e^+ e^-$         | $(3.1 \pm 0.6) \times 10^{-7}$       |        | 2708 |
| $p K^- \mu^+ \mu^-$     | $(2.6^{+0.5}_{-0.4}) \times 10^{-7}$ |        | 2685 |
| $\Lambda \gamma$        | $(7.1 \pm 1.7) \times 10^{-6}$       |        | 2699 |
| $\Lambda \eta$          | $(9^{+7}_{-5}) \times 10^{-6}$       |        | 2670 |
| $\Lambda \eta'(958)$    | $< 3.1 \times 10^{-6}$               | CL=90% | 2610 |
| $\Lambda \pi^+ \pi^-$   | $(4.6 \pm 1.9) \times 10^{-6}$       |        | 2692 |
| $\Lambda K^+ \pi^-$     | $(5.7 \pm 1.2) \times 10^{-6}$       |        | 2660 |
| $\Lambda K^+ K^-$       | $(1.61 \pm 0.22) \times 10^{-5}$     |        | 2605 |
| $\Lambda D^+ D^-$       | $(1.24 \pm 0.35) \times 10^{-4}$     |        | 1387 |
| $\Lambda \phi$          | $(9.8 \pm 2.6) \times 10^{-6}$       |        | 2599 |
| $p \pi^- \pi^+ \pi^-$   | $(2.12 \pm 0.21) \times 10^{-5}$     |        | 2715 |
| $p K^- K^+ \pi^-$       | $(4.1 \pm 0.6) \times 10^{-6}$       |        | 2612 |
| $p K^- \pi^+ \pi^-$     | $(5.1 \pm 0.5) \times 10^{-5}$       |        | 2675 |
| $p K^- K^+ K^-$         | $(1.27 \pm 0.13) \times 10^{-5}$     |        | 2524 |

|                                          |                                |             |
|------------------------------------------|--------------------------------|-------------|
| $\Lambda_b(5912)^0$                      | $J^P = \frac{1}{2}^-$          |             |
| Mass $m = 5912.16 \pm 0.16$ MeV          |                                |             |
| Full width $\Gamma < 0.25$ MeV, CL = 90% |                                |             |
| $\Lambda_b(5912)^0$ DECAY MODES          | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
| $\Lambda_b^0 \pi^+ \pi^-$                | seen                           | 86          |

|                                          |                       |  |
|------------------------------------------|-----------------------|--|
| $\Lambda_b(5920)^0$                      | $J^P = \frac{3}{2}^-$ |  |
| Mass $m = 5920.07 \pm 0.16$ MeV          |                       |  |
| Full width $\Gamma < 0.19$ MeV, CL = 90% |                       |  |

| $\Lambda_b(5920)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------|--------------------------------|-------------|
| $\Lambda_b^0 \pi^+ \pi^-$       | seen                           | 108         |

$\Lambda_b(6070)^0$

$$J^P = \frac{1}{2}^+$$

Quantum numbers based on quark model expectations.

$$\text{Mass } m = 6072.3 \pm 2.9 \text{ MeV}$$

$$\text{Full width } \Gamma = 72 \pm 11 \text{ MeV}$$

| $\Lambda_b(6070)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------|--------------------------------|-------------|
| $\Lambda_b^0 \pi^+ \pi^-$       | seen                           | 343         |

$\Lambda_b(6146)^0$

$$J^P = \frac{3}{2}^+$$

$$\text{Mass } m = 6146.2 \pm 0.4 \text{ MeV}$$

$$m_{\Lambda_b(6146)^0} - m_{\Lambda_b^0} = 526.55 \pm 0.34 \text{ MeV}$$

$$\text{Full width } \Gamma = 2.9 \pm 1.3 \text{ MeV}$$

| $\Lambda_b(6146)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------|--------------------------------|-------------|
| $\Lambda_b^0 \pi^+ \pi^-$       | seen                           | 427         |

$\Lambda_b(6152)^0$

$$J^P = \frac{5}{2}^+$$

$$\text{Mass } m = 6152.5 \pm 0.4 \text{ MeV}$$

$$m_{\Lambda_b(6152)^0} - m_{\Lambda_b^0} = 532.89 \pm 0.28 \text{ MeV}$$

$$m_{\Lambda_b(6152)^0} - m_{\Lambda_b(6146)^0} = 6.34 \pm 0.32 \text{ MeV}$$

$$\text{Full width } \Gamma = 2.1 \pm 0.9 \text{ MeV}$$

| $\Lambda_b(6152)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------|--------------------------------|-------------|
| $\Lambda_b^0 \pi^+ \pi^-$       | seen                           | 434         |

$\Sigma_b$

$I(J^P) = 1(\frac{1}{2}^+)$   
 $I, J, P$  need confirmation.

Mass  $m(\Sigma_b^+) = 5810.56 \pm 0.25$  MeV  
Mass  $m(\Sigma_b^-) = 5815.64 \pm 0.27$  MeV  
 $m_{\Sigma_b^+} - m_{\Sigma_b^-} = -5.06 \pm 0.18$  MeV  
 $\Gamma(\Sigma_b^+) = 5.0 \pm 0.5$  MeV  
 $\Gamma(\Sigma_b^-) = 5.3 \pm 0.5$  MeV

| $\Sigma_b$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------|--------------------------------|-------------|
| $\Lambda_b^0 \pi$      | dominant                       | 133         |

$\Sigma_b^*$

$I(J^P) = 1(\frac{3}{2}^+)$   
 $I, J, P$  need confirmation.

Mass  $m(\Sigma_b^{*+}) = 5830.32 \pm 0.27$  MeV  
Mass  $m(\Sigma_b^{*-}) = 5834.74 \pm 0.30$  MeV  
 $m_{\Sigma_b^{*+}} - m_{\Sigma_b^{*-}} = -4.37 \pm 0.33$  MeV ( $S = 1.6$ )  
 $m_{\Sigma_b^{*+}} - m_{\Sigma_b^+} = 19.73 \pm 0.18$   
 $m_{\Sigma_b^{*-}} - m_{\Sigma_b^-} = 19.09 \pm 0.22$   
 $\Gamma(\Sigma_b^{*+}) = 9.4 \pm 0.5$  MeV  
 $\Gamma(\Sigma_b^{*-}) = 10.4 \pm 0.8$  MeV ( $S = 1.3$ )  
 $m_{\Sigma_b^*} - m_{\Sigma_b} = 21.2 \pm 2.0$  MeV

| $\Sigma_b^*$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------|--------------------------------|-------------|
| $\Lambda_b^0 \pi$        | dominant                       | 159         |

$\Sigma_b(6097)^+$

$J^P = ?^?$

Mass  $m = 6095.8 \pm 1.7$  MeV  
Full width  $\Gamma = 31 \pm 6$  MeV

| $\Sigma_b(6097)^+$ DECAY MODES                             | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------------------------------------|--------------------------------|-------------|
| $\Lambda_b \pi^+ \times B(b \rightarrow \Sigma_b(6097)^+)$ | seen                           | —           |

$\Sigma_b(6097)^-$

$$J^P = ??$$

Mass  $m = 6098.0 \pm 1.8$  MeV  
Full width  $\Gamma = 29 \pm 4$  MeV

| $\Sigma_b(6097)^-$ DECAY MODES                             | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|------------------------------------------------------------|--------------------------------|-------------|
| $\Lambda_b \pi^- \times B(b \rightarrow \Sigma_b(6097)^-)$ | seen                           | —           |

$\Xi_b^-$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$I, J, P$  need confirmation.

$m(\Xi_b^-) = 5797.0 \pm 0.4$  MeV ( $S = 1.4$ )  
 $m_{\Xi_b^-} - m_{\Lambda_b^0} = 177.48 \pm 0.24$  MeV ( $S = 1.1$ )  
 $m_{\Xi_b^-} - m_{\Xi_b^0} = 5.9 \pm 0.5$  MeV  
Mean life  $\tau_{\Xi_b^-} = (1.578 \pm 0.021) \times 10^{-12}$  s

| $\Xi_b^-$ DECAY MODES                                                            | Fraction ( $\Gamma_i/\Gamma$ )          | Confidence level | $p$ (MeV/c) |
|----------------------------------------------------------------------------------|-----------------------------------------|------------------|-------------|
| $J/\psi \Xi^- \times B(b \rightarrow \Xi_b^-)$                                   | $(1.02^{+0.26}_{-0.21}) \times 10^{-5}$ |                  | 1782        |
| $J/\psi \Lambda K^- \times B(b \rightarrow \Xi_b^-)$                             | $(2.5 \pm 0.4) \times 10^{-6}$          |                  | 1631        |
| $p K^- K^- \times B(b \rightarrow \Xi_b^-)$                                      | $(3.7 \pm 0.8) \times 10^{-8}$          |                  | 2731        |
| $p K^- K^-$                                                                      | $(2.3 \pm 0.9) \times 10^{-6}$          |                  | 2731        |
| $p \pi^- \pi^-$                                                                  | $< 1.3 \times 10^{-6}$                  | 90%              | 2813        |
| $p K^- \pi^-$                                                                    | $(2.3 \pm 1.1) \times 10^{-6}$          |                  | 2783        |
| $\Lambda_b^0 \pi^- \times B(b \rightarrow \Xi_b^-)/B(b \rightarrow \Lambda_b^0)$ | $(7.0 \pm 0.9) \times 10^{-4}$          |                  | 99          |
| $\Xi_c^0 \pi^-$                                                                  | seen                                    |                  | 2367        |
| $\Xi_c^0 D_s^- \times B(b \rightarrow \Xi_b^-)/B(b \rightarrow \Lambda_b^0)$     | $(1.9 \pm 0.5) \times 10^{-3}$          |                  | —           |
| $\Sigma(1385) K^-$                                                               | $(2.6 \pm 2.3) \times 10^{-7}$          |                  | 2707        |
| $\Lambda(1405) K^-$                                                              | $(1.9 \pm 1.2) \times 10^{-7}$          |                  | 2702        |
| $\Lambda(1520) K^-$                                                              | $(7.6 \pm 3.2) \times 10^{-7}$          |                  | 2673        |
| $\Lambda(1670) K^-$                                                              | $(4.5 \pm 2.3) \times 10^{-7}$          |                  | 2629        |
| $\Sigma(1775) K^-$                                                               | $(2.2 \pm 1.5) \times 10^{-7}$          |                  | 2599        |
| $\Sigma(1915) K^-$                                                               | $(2.6 \pm 2.5) \times 10^{-7}$          |                  | 2553        |

|                  |         |                        |
|------------------|---------|------------------------|
| $J/\psi \Xi^-$   | seen    | —                      |
| $\psi(2S) \Xi^-$ | seen    | —                      |
| $\Xi^- \gamma$   | $< 1.3$ | $\times 10^{-4}$ 95% — |



$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$   
 $I, J, P$  need confirmation.

$m(\Xi_b^0) = 5791.7 \pm 0.4 \text{ MeV}$   
 $m_{\Xi_b^0} - m_{\Lambda_b^0} = 172.3 \pm 0.4 \text{ MeV}$   
Mean life  $\tau_{\Xi_b^0} = (1.477 \pm 0.032) \times 10^{-12} \text{ s}$

| $\Xi_b^0$ DECAY MODES                                                              | Fraction ( $\Gamma_i/\Gamma$ )   | Confidence level | $P$<br>(MeV/c) |
|------------------------------------------------------------------------------------|----------------------------------|------------------|----------------|
| $p D^0 K^- \times B(b \rightarrow \Xi_b^0)$                                        | $(1.7 \pm 0.6) \times 10^{-6}$   |                  | 2374           |
| $p \bar{K}^0 \pi^- \times B(b \rightarrow \Xi_b^0)/B(\bar{b} \rightarrow B^0)$     | $< 1.6 \times 10^{-6}$           | 90%              | 2783           |
| $p K^0 K^- \times B(b \rightarrow \Xi_b^0)/B(\bar{b} \rightarrow B^0)$             | $< 1.1 \times 10^{-6}$           | 90%              | 2730           |
| $\Lambda \pi^+ \pi^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$ | $< 1.7 \times 10^{-6}$           | 90%              | 2781           |
| $\Lambda K^- \pi^+ \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$   | $< 8 \times 10^{-7}$             | 90%              | 2751           |
| $\Lambda K^+ K^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$     | $< 3 \times 10^{-7}$             | 90%              | 2698           |
| $J/\psi \Lambda$                                                                   | seen                             |                  | 1868           |
| $J/\psi \Xi^0$                                                                     | seen                             |                  | 1785           |
| $\Xi_c^+ D_s^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$       | $(1.7 \pm 0.9) \times 10^{-3}$   |                  | —              |
| $\Lambda_c^+ K^- \times B(b \rightarrow \Xi_b^0)$                                  | $(6 \pm 4) \times 10^{-7}$       |                  | 2416           |
| $p K^- \pi^+ \pi^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$   | $(1.9 \pm 0.4) \times 10^{-6}$   |                  | 2766           |
| $p K^- K^- \pi^+ \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$     | $(1.73 \pm 0.31) \times 10^{-6}$ |                  | 2704           |
| $p K^- K^+ K^- \times B(b \rightarrow \Xi_b^0)/B(b \rightarrow \Lambda_b^0)$       | $(1.8 \pm 1.0) \times 10^{-7}$   |                  | 2620           |



$J^P = \frac{1}{2}^+$

Mass  $m = 5934.9 \pm 0.4 \text{ MeV}$   
Full width  $\Gamma = 0.03 \pm 0.032 \text{ MeV}$

| $\Xi'_b(5935)^-$ DECAY MODES                                                                | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|---------------------------------------------------------------------------------------------|--------------------------------|-------------|
| $\Xi_b^0 \pi^- \times B(\bar{b} \rightarrow \Xi'_b(5935)^-)/B(\bar{b} \rightarrow \Xi_b^0)$ | $(11.8 \pm 1.8) \%$            | 31          |

$\Xi_b(5945)^0$

$J^P = \frac{3}{2}^+$

Mass  $m = 5952.3 \pm 0.6$  MeV  
Full width  $\Gamma = 0.87 \pm 0.07$  MeV

| $\Xi_b(5945)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------|--------------------------------|-------------|
| $\Xi_b^- \pi^+$             | seen                           | 78          |

$\Xi_b(5955)^-$

$J^P = \frac{3}{2}^+$

Mass  $m = 5955.5 \pm 0.4$  MeV  
Full width  $\Gamma = 1.43 \pm 0.11$  MeV

| $\Xi_b(5955)^-$ DECAY MODES                                                                  | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|----------------------------------------------------------------------------------------------|--------------------------------|-------------|
| $\Xi_b^0 \pi^- \times B(\bar{b} \rightarrow \Xi_b^*(5955)^-)/B(\bar{b} \rightarrow \Xi_b^0)$ | $(20.7 \pm 3.5) \%$            | 84          |

$\Xi_b(6087)^0$

$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$   
 $J, P$  need confirmation.

Mass  $m = 6087.0 \pm 0.5$  MeV  
Full width  $\Gamma = 2.4 \pm 0.5$  MeV

| $\Xi_b(6087)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------|--------------------------------|-------------|
| $\Xi_b^0 \pi^+ \pi^-$       | seen                           | —           |

$\Xi_b(6095)^0$

$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$   
 $J, P$  need confirmation.

Mass  $m = 6095.1 \pm 0.4$  MeV  
Full width  $\Gamma = 0.50 \pm 0.35$  MeV

| $\Xi_b(6095)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------|--------------------------------|-------------|
| $\Xi_b^0 \pi^+ \pi^-$       | seen                           | —           |

$\Xi_b(6100)^-$

$J^P = \frac{3}{2}^-$   
 $J, P$  need confirmation.

Mass  $m = 6099.8 \pm 0.4$  MeV

Full width  $\Gamma = 0.94 \pm 0.31$  MeV

| $\Xi_b(6100)^-$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------|--------------------------------|-------------|
| $\Xi_b^- \pi^+ \pi^-$       | seen                           | 128         |

$\Xi_b(6227)^-$

$J^P = ??$

Mass  $m = 6227.9 \pm 0.9$  MeV

Full width  $\Gamma = 19.9 \pm 2.6$  MeV

| $\Xi_b(6227)^-$ DECAY MODES                                                        | Fraction ( $\Gamma_i/\Gamma$ )   | Scale factor | $p$ (MeV/c) |
|------------------------------------------------------------------------------------|----------------------------------|--------------|-------------|
| $\Lambda_b^0 K^- \times B(b \rightarrow \Xi_b(6227))/B(b \rightarrow \Lambda_b^0)$ | $(3.20 \pm 0.35) \times 10^{-3}$ |              | 336         |
| $\Xi_b^0 \pi^- \times B(b \rightarrow \Xi_b(6227))/B(b \rightarrow \Xi_b^0)$       | $(2.8 \pm 1.1) \%$               | 1.8          | 398         |

$\Xi_b(6227)^0$

$J^P = ??$

Mass  $m = 6226.8 \pm 1.6$  MeV

Full width  $\Gamma = 19^{+5}_{-4}$  MeV

| $\Xi_b(6227)^0$ DECAY MODES                                                    | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------------------------------------------------------|--------------------------------|-------------|
| $\Xi_b^- \pi^+ \times B(b \rightarrow \Xi_b(6227)^0)/B(b \rightarrow \Xi_b^-)$ | $(4.5 \pm 0.9) \%$             | 398         |

$\Xi_b(6327)^0$

$J^P = ??$

Mass  $m = 6327.28 \pm 0.35$  MeV

Full width  $\Gamma < 2.56$  MeV, CL = 95%

| $\Xi_b(6327)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------|--------------------------------|-------------|
| $\Lambda_b^0 K^- \pi^+$     | seen                           | 298         |

 **$\Xi_b(6333)^0$** 

$$J^P = ??$$

Mass  $m = 6332.69 \pm 0.28$  MeVFull width  $\Gamma < 1.92$  MeV, CL = 95%

| $\Xi_b(6333)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-----------------------------|--------------------------------|-------------|
| $\Lambda_b^0 K^- \pi^+$     | seen                           | 309         |

 **$\Omega_b^-$** 

$$I(J^P) = 0(\frac{1}{2}^+)$$

 $I, J, P$  need confirmation.Mass  $m = 6045.8 \pm 0.8$  MeV

$$m_{\Omega_b^-} - m_{\Lambda_b^0} = 426.4 \pm 2.2 \text{ MeV}$$

$$m_{\Omega_b^-} - m_{\Xi_b^-} = 248.5 \pm 0.6 \text{ MeV}$$

Mean life  $\tau = (1.64 \pm 0.16) \times 10^{-12}$  s

$$\tau(\Omega_b^-)/\tau(\Xi_b^-) \text{ mean life ratio} = 1.11 \pm 0.16$$

| $\Omega_b^-$ DECAY MODES                                   | Fraction ( $\Gamma_i/\Gamma$ )       | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|------------------------------------------------------------|--------------------------------------|-----------------------------------|----------------|
| $J/\psi \Omega^- \times B(b \rightarrow \Omega_b)$         | $(1.4^{+0.5}_{-0.4}) \times 10^{-6}$ | S=1.6                             | 1805           |
| $p K^- K^- \times B(\bar{b} \rightarrow \Omega_b)$         | $< 2.3 \times 10^{-9}$               | CL=90%                            | 2865           |
| $p \pi^- \pi^- \times B(\bar{b} \rightarrow \Omega_b)$     | $< 1.5 \times 10^{-8}$               | CL=90%                            | 2943           |
| $p K^- \pi^- \times B(\bar{b} \rightarrow \Omega_b)$       | $< 7 \times 10^{-9}$                 | CL=90%                            | 2915           |
| $\Omega_c^0 \pi^-$                                         | seen                                 |                                   | 2420           |
| $\Omega_c^0 \pi^-, \Omega_c^0 \rightarrow p K^- K^- \pi^+$ | seen                                 |                                   | —              |
| $\Xi_c^+ K^- \pi^-$                                        | seen                                 |                                   | 2473           |

 **$\Omega_b(6316)^-$** 

$$I(J^P) = ?(??)$$

 $I, J, P$  need confirmation.Mass  $m = 6315.6 \pm 0.6$  MeVFull width  $\Gamma < 4.2$  MeV, CL = 95%

| $\Omega_b(6316)^-$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_b^0 K^-$                  | seen                           | 168         |

$\Omega_b(6330)^-$

$I(J^P) = ?(?^?)$   
 $I, J, P$  need confirmation.

Mass  $m = 6330.3 \pm 0.6$  MeV  
Full width  $\Gamma < 4.7$  MeV, CL = 95%

| $\Omega_b(6330)^-$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_b^0 K^-$                  | seen                           | 206         |

$\Omega_b(6340)^-$

$I(J^P) = ?(?^?)$   
 $I, J, P$  need confirmation.

Mass  $m = 6339.7 \pm 0.6$  MeV  
Full width  $\Gamma < 1.8$  MeV, CL = 95%

| $\Omega_b(6340)^-$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_b^0 K^-$                  | seen                           | 227         |

$\Omega_b(6350)^-$

$I(J^P) = ?(?^?)$   
 $I, J, P$  need confirmation.

Mass  $m = 6349.8 \pm 0.6$  MeV  
Full width  $\Gamma < 3.2$  MeV, CL = 95%

| $\Omega_b(6350)^-$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|--------------------------------|--------------------------------|-------------|
| $\Xi_b^0 K^-$                  | seen                           | 248         |

$b$ -baryon ADMIXTURE ( $\Lambda_b, \Xi_b, \Omega_b$ )

These branching fractions are actually an average over weakly decaying  $b$ -baryons weighted by their production rates at the LHC, LEP, and Tevatron, branching ratios, and detection efficiencies. They scale with the  $b$ -baryon production fraction  $B(b \rightarrow b\text{-baryon})$ .

The branching fractions  $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{ anything})$  and  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{ anything})$  are not pure measurements because the underlying measured products of these with  $B(b \rightarrow b\text{-baryon})$  were used to determine  $B(b \rightarrow b\text{-baryon})$ , as described in the note “Production and Decay of  $b$ -Flavored Hadrons.”

For inclusive branching fractions, e.g.,  $B \rightarrow D^\pm \text{ anything}$ , the values usually are multiplicities, not branching fractions. They can be greater than one.

| $b\text{-baryon ADMIXTURE DECAY MODES}$<br>$(\Lambda_b, \Xi_b, \Omega_b)$ | Fraction ( $\Gamma_i/\Gamma$ ) | Scale factor | $p$<br>(MeV/c) |
|---------------------------------------------------------------------------|--------------------------------|--------------|----------------|
| $p\mu^- \bar{\nu}$ anything                                               | $(5.8^{+2.3}_{-2.0})\%$        |              | —              |
| $p\ell \bar{\nu}_\ell$ anything                                           | $(5.6 \pm 1.2)\%$              |              | —              |
| $p$ anything                                                              | $(70 \pm 22)\%$                |              | —              |
| $\Lambda \ell^- \bar{\nu}_\ell$ anything                                  | $(3.8 \pm 0.6)\%$              |              | —              |
| $\Lambda \ell^+ \nu_\ell$ anything                                        | $(3.2 \pm 0.8)\%$              |              | —              |
| $\Lambda$ anything                                                        | $(39 \pm 7)\%$                 |              | —              |
| $\Xi^- \ell^- \bar{\nu}_\ell$ anything                                    | $(4.6 \pm 1.4) \times 10^{-3}$ | 1.2          | —              |

EXOTIC BARYONS

$P_{c\bar{c}s}(4338)^0$

$$I(J^P) = 0(\frac{1}{2}^-)$$

Mass  $m = 4338.2 \pm 0.8$  MeV  
Full width  $\Gamma = 7.0 \pm 1.8$  MeV

| $P_{c\bar{c}s}(4338)^0$ DECAY MODES | Fraction ( $\Gamma_i/\Gamma$ ) | $p$ (MeV/c) |
|-------------------------------------|--------------------------------|-------------|
| $J/\psi \Lambda$                    | seen                           | —           |

## NOTES

- [a] The masses of the  $p$  and  $n$  are most precisely known in  $u$  (unified atomic mass units). The conversion factor to MeV,  $1\ u = 931.494061(21)\ \text{MeV}$ , is less well known than are the masses in  $u$ .
- [b] The  $|m_p - m_{\bar{p}}|/m_p$  and  $|q_p + q_{\bar{p}}|/e$  are not independent, and both use the more precise measurement of  $|q_{\bar{p}}/m_{\bar{p}}|/(q_p/m_p)$ .
- [c] The limit is from neutrality-of-matter experiments; it assumes  $q_n = q_p + q_e$ . See also the charge of the neutron.
- [d] The  $\mu p$  and  $e p$  values for the charge radius are much too different to average them. The disagreement is not yet understood.
- [e] There is a lot of disagreement about the value of the proton magnetic charge radius. See the Listings.
- [f] There is some controversy about whether nuclear physics and model dependence complicate the analysis for bound neutrons (from which the best limit comes). The first limit here is from reactor experiments with free neutrons.
- [g] Lee and Yang in 1956 proposed the existence of a mirror world in an attempt to restore global parity symmetry—thus a search for oscillations between the two worlds. Oscillations between the worlds would be maximal when the magnetic fields  $B$  and  $B'$  were equal. The limit for any  $B'$  in the range 0 to  $12.5\ \mu\text{T}$  is  $>12\ \text{s}$  (95% CL).
- [h] The parameters  $g_A$ ,  $g_V$ , and  $g_{WM}$  for semileptonic modes are defined by  $\bar{B}_f[\gamma_\lambda(g_V + g_A\gamma_5) + i(g_{WM}/m_{B_i})\sigma_{\lambda\nu}q^\nu]B_i$ , and  $\phi_{AV}$  is defined by  $g_A/g_V = |g_A/g_V|e^{i\phi_{AV}}$ . See the “Note on Baryon Decay Parameters” in the neutron Particle Listings.
- [i] Time-reversal invariance requires this to be  $0^\circ$  or  $180^\circ$ .
- [j] This coefficient is zero if time invariance is not violated.
- [k] This limit is for  $\gamma$  energies between 0.4 and 782 keV.
- [l] The decay parameters  $\gamma$  and  $\Delta$  are calculated from  $\alpha$  and  $\phi$  using
 
$$\gamma = \sqrt{1-\alpha^2} \cos\phi, \quad \tan\Delta = -\frac{1}{\alpha} \sqrt{1-\alpha^2} \sin\phi.$$
 See the “Note on Baryon Decay Parameters” in the neutron Particle Listings.
- [n] See the Listings for the pion momentum range used in this measurement.
- [o] Our estimate. See the Particle Listings for details.
- [p] A theoretical value using QED.
- [q] This branching fraction includes all the decay modes of the final-state resonance.
- [r] Here  $\gamma_D$  stands for a dark photon.

- [s] See AALTONEN 11H, Fig. 8, for the calculated ratio of  $\Lambda_c^+ \pi^0 \pi^0$  and  $\Lambda_c^+ \pi^+ \pi^-$  partial widths as a function of the  $\Lambda_c(2595)^+ - \Lambda_c^+$  mass difference. At our value of the mass difference, the ratio is about 4.
- [t] A test that the isospin is indeed 0, so that the particle is indeed a  $\Lambda_c^+$ .
- [u] In the isospin limit, this branching fraction would be 2/3, the other 1/3 being decays to  $\Lambda_c^+ \pi^0 \pi^0$ .
- [v]  $P_c^+$  is a pentaquark-charmonium state.
- [x] Not a pure measurement. See note at head of  $\Lambda_b^0$  Decay Modes.
- [y] Here  $h^-$  means  $\pi^-$  or  $K^-$ .