

CHARMED, STRANGE MESONS

($C = \pm 1, S = \pm 1$)

(including possibly non- $q\bar{q}$ states)

$$D_s^+ = c\bar{s}, D_s^- = \bar{c}s, \quad \text{similarly for } D_s^{*'}s$$

$D_s^\pm$

$$I(J^P) = 0(0^-)$$

$$\text{Mass } m = 1968.35 \pm 0.07 \text{ MeV}$$

$$m_{D_s^\pm} - m_{D^\pm} = 98.69 \pm 0.05 \text{ MeV}$$

$$\text{Mean life } \tau = (501.2 \pm 2.2) \times 10^{-15} \text{ s} \quad (S = 1.3)$$

$$c\tau = 150.3 \text{ } \mu\text{m}$$

CP-violating decay-rate asymmetries

$$A_{CP}(\mu^\pm \nu) = (-0.2 \pm 2.5)\%$$

$$A_{CP}(\tau^\pm \nu) \text{ in } D_s^+ \rightarrow \tau^+ \nu_\tau, D_s^- \rightarrow \tau^- \bar{\nu}_\tau = (3 \pm 5)\%$$

$$A_{CP}(K^\pm K_S^0) = (0.07 \pm 0.24)\%$$

$$A_{CP}(K^\pm K_L^0) \text{ in } D_s^\pm \rightarrow K^\pm K_L^0 = (-1.1 \pm 2.7) \times 10^{-2}$$

$$A_{CP}(K^+ K^- \pi^\pm) = (0.35 \pm 0.34)\%$$

$$A_{CP}(\phi \pi^\pm) = (-0.38 \pm 0.27)\%$$

$$A_{CP}(K^\pm K_S^0 \pi^0) = (-0.9 \pm 1.9)\%$$

$$A_{CP}(2K_S^0 \pi^\pm) = (1.3 \pm 1.6)\%$$

$$A_{CP}(K^+ K^- \pi^\pm \pi^0) = (-0.6 \pm 0.9)\%$$

$$A_{CP}(K^\pm K_S^0 \pi^+ \pi^-) = (0.7 \pm 2.9)\% \quad (S = 1.3)$$

$$A_{CP}(K_S^0 K^\mp 2\pi^\pm) = (0.7 \pm 1.8)\% \quad (S = 1.3)$$

$$A_{CP}(\pi^+ \pi^- \pi^\pm) = (-0.9 \pm 1.1)\%$$

$$A_{CP}(\pi^\pm \eta) = (0.24 \pm 0.29)\%$$

$$A_{CP}(\pi^\pm \pi^+ \pi^- \eta) = (2.4 \pm 3.0)\%$$

$$A_{CP}(\pi^\pm \eta') = (-0.08 \pm 0.17)\% \quad (S = 1.2)$$

$$A_{CP}(\eta \pi^\pm \pi^0) = (0.9 \pm 1.5)\%$$

$$A_{CP}(\eta' \pi^\pm \pi^0) = (-1.5 \pm 2.5)\%$$

$$A_{CP}(K^\pm \pi^0) = (2 \pm 4)\% \quad (S = 1.2)$$

$$A_{CP}(\bar{K}^0 / K^0 \pi^\pm) = (0.4 \pm 0.5)\%$$

$$A_{CP}(K_S^0 \pi^\pm) = (0.20 \pm 0.18)\%$$

$$A_{CP}(K^\pm \pi^+ \pi^-) = (2.2 \pm 1.9)\%$$

$$A_{CP}(K_S^0 \pi^+ \pi^0) \text{ in } D_s^\pm \rightarrow K_S^0 \pi^\pm \pi^0 = (-2 \pm 5)\%$$

$$A_{CP}(K^\pm \pi^+ \pi^- \pi^0) \text{ in } D_s^\pm \rightarrow K^\pm \pi^+ \pi^- \pi^0 = (7 \pm 5) \times 10^{-2}$$

$$A_{CP}(K^\pm \eta) = (1.8 \pm 1.9)\%$$

$$A_{CP}(K^\pm \eta'(958)) = (6 \pm 19)\%$$

***CP* violating asymmetries of *P*-odd (*T*-odd) moments**

$$A_T(K_S^0 K^\pm \pi^+ \pi^-) = (-8 \pm 6) \times 10^{-3} [a]$$

***D*_s⁺ → *φ* *ℓ*⁺ *ν*_ℓ form factors**

$$r_2 = 0.83 \pm 0.08 \quad (S = 1.8)$$

$$r_v = 1.76 \pm 0.07 \quad (S = 1.1)$$

$$\Gamma_L/\Gamma_T = 0.72 \pm 0.18$$

$$f_+(0) |V_{cs}| \text{ in } D_s^+ \rightarrow \eta e^+ \nu_e = 0.449 \pm 0.009$$

$$f_+(0) |V_{cs}| \text{ in } D_s^+ \rightarrow \eta \mu^+ \nu_\mu = 0.452 \pm 0.012$$

$$r_1 \equiv a_1/a_0 \text{ in } D_s^+ \rightarrow \eta \mu^+ \nu_\mu = -2.9 \pm 0.6$$

$$\langle A_{FB}^\eta \rangle \text{ in } D_s^+ \rightarrow \eta \mu^+ \nu_\mu = (-5.9 \pm 3.1) \times 10^{-2}$$

$$f_+(0) |V_{cs}| \text{ in } D_s^+ \rightarrow \eta' e^+ \nu_e = 0.527 \pm 0.024$$

$$r_1 \equiv a_1/a_0 \text{ in } D_s^+ \rightarrow \eta' \mu^+ \nu_\mu = -11 \pm 5$$

$$\langle A_{FB}^{\eta'} \rangle \text{ in } D_s^+ \rightarrow \eta' \mu^+ \nu_\mu = (-6 \pm 8) \times 10^{-2}$$

$$f_+(0) |V_{cs}| \text{ in } D_s^+ \rightarrow \eta' \mu^+ \nu_\mu = 0.50 \pm 0.04$$

$$f_+(0) |V_{cs}| \text{ in } D_s^+ \rightarrow f_0(980) e^+ \nu_e = 0.50 \pm 0.04$$

$$f_+(0) |V_{cd}| \text{ in } D_s^+ \rightarrow K^0 e^+ \nu_e = 0.145 \pm 0.010$$

$$r_v \equiv V(0)/A_1(0) \text{ in } D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e = 1.7 \pm 0.4$$

$$r_2 \equiv A_2(0)/A_1(0) \text{ in } D_s^+ \rightarrow K^*(892)^0 e^+ \nu_e = 0.77 \pm 0.29$$

$$f_{D_s^+} |V_{cs}| \text{ in } D_s^+ \rightarrow \mu^+ \nu_\mu = 242.7 \pm 3.0 \text{ MeV}$$

$$f_{D_s^+} |V_{cs}| \text{ in } D_s^+ \rightarrow \tau^+ \nu_\tau = 247.5 \pm 2.3 \text{ MeV}$$

Unless otherwise noted, the branching fractions for modes with a resonance in the final state include all the decay modes of the resonance. D_S^- modes are charge conjugates of the modes below.

D_S^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
Inclusive modes			
e^+ semileptonic	[b] (6.33 $\pm$ 0.15) %		—
π^+ anything	(119.3 $\pm$ 1.4) %		—
π^- anything	(43.2 $\pm$ 0.9) %		—
π^0 anything	(123 $\pm$ 7) %		—
K^- anything	(18.7 $\pm$ 0.5) %		—
K^+ anything	(28.9 $\pm$ 0.7) %		—
K_S^0 anything	(19.0 $\pm$ 1.1) %		—
η anything	[c] (29.9 $\pm$ 2.8) %		—
ω anything	(6.1 $\pm$ 1.4) %		—
η' anything	[d] (10.3 $\pm$ 1.4) %	S=1.1	—
$f_0(980)$ anything, $f_0 \rightarrow \pi^+\pi^-$	< 1.3 %	CL=90%	—
ϕ anything	(15.7 $\pm$ 1.0) %		—
K^+K^- anything	(15.8 $\pm$ 0.7) %		—
$K_S^0K^+$ anything	(5.8 $\pm$ 0.5) %		—
$K_S^0K^-$ anything	(1.9 $\pm$ 0.4) %		—
$2K_S^0$ anything	(1.70 $\pm$ 0.32) %		—
$2K^+$ anything	< 2.6 $\times 10^{-3}$	CL=90%	—
$2K^-$ anything	< 6 $\times 10^{-4}$	CL=90%	—
$2\pi^+\pi^- +$ anything	(32.8 $\pm$ 0.7) %		—
Leptonic and semileptonic modes			
$e^+\nu_e$	< 8.3 $\times 10^{-5}$	CL=90%	984
$\mu^+\nu_\mu$	(5.37 $\pm$ 0.11) $\times 10^{-3}$		981
$\tau^+\nu_\tau$	(5.39 $\pm$ 0.09) %		182
$\gamma e^+\nu_e$	< 1.3 $\times 10^{-4}$	CL=90%	984
$K^+K^-e^+\nu_e$	—		851
$K_S^0K_S^0e^+\nu_e$	< 3.8 $\times 10^{-4}$	CL=90%	849
$\phi e^+\nu_e$	[e] (2.34 $\pm$ 0.12) %	S=1.2	720
$K_1(1270)^0e^+\nu_e$	< 4.1 $\times 10^{-4}$	CL=90%	585
$b_1(1235)^0e^+\nu_e$, $b_1^0 \rightarrow \omega\pi^0$	< 6.4 $\times 10^{-4}$	CL=90%	—
$\phi\mu^+\nu_\mu$	(2.24 $\pm$ 0.11) %		715
$\eta e^+\nu_e$	[e] (2.27 $\pm$ 0.06) %		908
$\eta'(958)e^+\nu_e$	[e] (8.1 $\pm$ 0.4) $\times 10^{-3}$		751
$\eta\mu^+\nu_\mu$	(2.24 $\pm$ 0.07) %		905
$\eta'(958)\mu^+\nu_\mu$	(8.0 $\pm$ 0.6) $\times 10^{-3}$		747
$\omega e^+\nu_e$	[f] < 2.0 $\times 10^{-3}$	CL=90%	829

$K^0 e^+ \nu_e$	(2.88 $\pm$ 0.26) $\times 10^{-3}$	S=1.2	921
$K^*(892)^0 e^+ \nu_e$	[e] (2.05 $\pm$ 0.20) $\times 10^{-3}$		782
$f_0(500) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0$	< 7.3 $\times 10^{-4}$	CL=90%	—
$f_0(500) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-$	< 3.3 $\times 10^{-4}$	CL=90%	—
$f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^0 \pi^0$	(7.9 $\pm$ 1.5) $\times 10^{-4}$		—
$f_0(980) e^+ \nu_e, f_0 \rightarrow \pi^+ \pi^-$	(1.64 $\pm$ 0.13) $\times 10^{-3}$		—
$f_0(980) \mu^+ \nu_\mu, f_0 \rightarrow K^+ K^-$	< 5.45 $\times 10^{-4}$	CL=90%	—
$a_0(980)^0 e^+ \nu_e, a_0^0 \rightarrow \pi^0 \eta$	< 1.2 $\times 10^{-4}$	CL=90%	—
$\pi^0 e^+ \nu_e$	< 6.4 $\times 10^{-5}$	CL=90%	980

Hadronic modes with a $K\bar{K}$ pair

$K^+ K_S^0$	(1.500 $\pm$ 0.014) %		850
$K^+ K_L^0$	(1.49 $\pm$ 0.06) %		850
$K^+ \bar{K}^0$	(2.95 $\pm$ 0.14) %		850
$K^+ K^- \pi^+$	[g] (5.45 $\pm$ 0.08) %	S=1.3	805
$\phi \pi^+$	[e,h] (4.5 $\pm$ 0.4) %		712
$\phi \pi^+, \phi \rightarrow K^+ K^-$	[h] (2.25 $\pm$ 0.05) %		712
$K^+ \bar{K}^*(892)^0$	(12.7 $^{+4.0}_{-3.1}$) %		685
$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow$	(2.61 $\pm$ 0.05) %		416
$K^+ \bar{K}^*(892)^0, \bar{K}^{*0} \rightarrow$	(4.8 $\pm$ 0.4) $\times 10^{-3}$		—
$K_S^0 \pi^0$			
$f_0(980) \pi^+, f_0 \rightarrow K^+ K^-$	(1.12 $\pm$ 0.19) %		732
$f_0(1370) \pi^+, f_0 \rightarrow K^+ K^-$	(7.2 $\pm$ 3.0) $\times 10^{-4}$		—
$f_0(1710) \pi^+, f_0 \rightarrow K^+ K^-$	(6.8 $\pm$ 2.8) $\times 10^{-4}$		198
$a_0(980)^+ \pi^0, a_0^+ \rightarrow K^+ K_S^0$	(1.1 $\pm$ 0.4) $\times 10^{-3}$		—
$a_0(1710)^+ \pi^0, a_0^+ \rightarrow$	(3.5 $\pm$ 0.6) $\times 10^{-3}$		—
$K^+ K_S^0$			
$K^+ \bar{K}_0^*(1430)^0, \bar{K}_0^* \rightarrow$	(1.79 $\pm$ 0.26) $\times 10^{-3}$		218
$K^+ \bar{K}_0^*(1410)^0, \bar{K}_0^* \rightarrow$	(8.8 $\pm$ 2.8) $\times 10^{-4}$		—
$K_S^0 \pi^0$			
$K^+ K_S^0 \pi^0$	(1.471 $\pm$ 0.028) %		805
$K^*(892)^+ K_S^0, K^{*+} \rightarrow$	(2.04 $\pm$ 0.32) $\times 10^{-3}$		—
$K^+ \pi^0$			
$2K_S^0 \pi^+$	(7.32 $\pm$ 0.14) $\times 10^{-3}$		802
$f_0(980) \pi^+, f_0 \rightarrow K_S^0 K_S^0$	< 1.8 $\times 10^{-4}$	CL=90%	—
$f_0(1710) \pi^+, f_0 \rightarrow K_S^0 K_S^0$	(3.39 $\pm$ 0.31) $\times 10^{-3}$		—
$K^*(892)^+ K_S^0, K^{*+} \rightarrow$	(3.19 $\pm$ 0.29) $\times 10^{-3}$		683
$K_S^0 \pi^+$			
$K^0 \bar{K}^0 \pi^+$	—		802
$K^*(892)^+ \bar{K}^0$	[e] (5.4 $\pm$ 1.2) %		683
$K^+ K^- \pi^+ \pi^0$	(5.53 $\pm$ 0.15) %	S=1.3	748

$\phi \rho^+$	[e] (5.59 $\pm$ 0.34) %	401
$\bar{K}_1(1270)^0 K^+$,	(5.7 $\pm$ 0.6) $\times 10^{-3}$	—
$\bar{K}_1(1270)^0 \rightarrow K^- \rho^+$		
$\bar{K}_1(1270)^0 K^+$,	(1.31 $\pm$ 0.25) %	—
$\bar{K}_1(1270)^0 \rightarrow K^*(892) \pi$		
$\bar{K}_1(1400)^0 K^+$,	(2.0 $\pm$ 0.4) %	—
$\bar{K}_1(1400)^0 \rightarrow K^*(892) \pi$		
$a_0(980)^0 \rho^+$, $a_0^0 \rightarrow K^+ K^-$	(1.9 $\pm$ 0.4) $\times 10^{-3}$	—
$f_1(1420)^0 \pi^+$, $f_1(1420)^0 \rightarrow$	(3.9 $\pm$ 0.7) $\times 10^{-3}$	—
$K^*(892)^\mp K^\pm$		
$f_1(1420)^0 \pi^+$, $f_1(1420)^0 \rightarrow$	(4.0 $\pm$ 1.4) $\times 10^{-4}$	—
$a_0(980)^0 \pi^0$, $a_0(980)^0 \rightarrow$		
$K^+ K^-$		
$\eta(1475) \pi^+$, $\eta(1475) \rightarrow$	(7.0 $\pm$ 2.8) $\times 10^{-4}$	—
$a_0(980)^0 \pi^0$, $a_0(980)^0 \rightarrow$		
$K^+ K^-$		
$K_S^0 K^- 2\pi^+$	(1.569 $\pm$ 0.028) %	744
$K^+ K^- K_S^0 \pi^+$	(1.27 $\pm$ 0.15) $\times 10^{-4}$	527
$K^*(892)^+ \bar{K}^*(892)^0$	[e] (5.64 $\pm$ 0.35) %	417
$\eta(1475) K_S^0$, $\eta \rightarrow$	(3.5 $\pm$ 1.0) $\times 10^{-4}$	—
$K^*(892)^0 \pi^+$, $K^{*0} \rightarrow$		
$K^- \pi^+$		
$\eta(1475) \pi^+$, $\eta \rightarrow$	(3.5 $\pm$ 1.0) $\times 10^{-4}$	—
$\bar{K}^*(892)^+ K^-$, $\bar{K}^{*+} \rightarrow$		
$K_S^0 \pi^+$		
$\eta(1475) \pi^+$, $\eta \rightarrow$	(1.7 $\pm$ 0.9) $\times 10^{-3}$	—
$a_0(980)^- \pi^+$, $a_0^- \rightarrow$		
$K_S^0 K^-$		
$f_1(1285) \pi^+$, $f_1 \rightarrow$	(3.5 $\pm$ 0.8) $\times 10^{-4}$	—
$a_0(980)^- \pi^+$, $a_0^- \rightarrow$		
$K_S^0 K^-$		
$K^+ K_S^0 \pi^+ \pi^-$	(9.34 $\pm$ 0.22) $\times 10^{-3}$	744
$K^+ K^- 2\pi^+ \pi^-$	(6.6 $\pm$ 0.6) $\times 10^{-3}$	673
$\phi 2\pi^+ \pi^-$	[e] (1.21 $\pm$ 0.16) %	640
$\phi \rho^0 \pi^+$, $\phi \rightarrow K^+ K^-$	(4.9 $\pm$ 0.7) $\times 10^{-3}$	181
$\phi a_1(1260)^+$, $\phi \rightarrow$	(7.5 $\pm$ 1.2) $\times 10^{-3}$	†
$K^+ K^-$, $a_1^+ \rightarrow$		
$\rho^0 \pi^+$		
$\phi 2\pi^+ \pi^-$ non- ρ , $\phi \rightarrow$	(1.4 $\pm$ 0.5) $\times 10^{-3}$	—
$K^+ K^-$		
$K^+ K^- \rho^0 \pi^+$ non- ϕ	< 2.0 $\times 10^{-4}$ CL=90%	249
$K^+ K^- 2\pi^+ \pi^-$ nonresonant	(1.0 $\pm$ 0.4) $\times 10^{-3}$	673
$2K_S^0 2\pi^+ \pi^-$	(8.0 $\pm$ 3.3) $\times 10^{-4}$	669

Hadronic modes without K 's

$\pi^+\pi^0$	$< 1.2 \times 10^{-4}$	CL=90%	975
$2\pi^+\pi^-$	$(1.090 \pm 0.014) \%$		959
$\rho^0\pi^+$	$(1.14 \pm 0.16) \times 10^{-4}$		825
$\omega\pi^+, \omega \rightarrow \pi^+\pi^-$	$(3.9 \pm 0.4) \times 10^{-5}$		—
$\pi^+(\pi^+\pi^-)_{S\text{-wave}}$	[i] $(9.23 \pm 0.13) \times 10^{-3}$		959
$f_2(1270)\pi^+, f_2 \rightarrow \pi^+\pi^-$	$(1.42 \pm 0.10) \times 10^{-3}$		559
$f_2'(1525)^0\pi^+, f_2' \rightarrow \pi^+\pi^-$	$(5.8 \pm 2.0) \times 10^{-6}$		—
$\rho(1450)^0\pi^+, \rho^0 \rightarrow \pi^+\pi^-$	$(1.8 \pm 0.6) \times 10^{-4}$		421
$\rho(1700)^0\pi^+, \rho^0 \rightarrow \pi^+\pi^-$	$(4 \pm 4) \times 10^{-5}$		—
$\pi^+2\pi^0$	$(5.2 \pm 0.5) \times 10^{-3}$	S=1.1	961
$f_0(980)\pi^+, f_0 \rightarrow \pi^0\pi^0$	$(2.9 \pm 0.6) \times 10^{-3}$		—
$f_0(1370)\pi^+, f_0 \rightarrow \pi^0\pi^0$	$(1.3 \pm 0.6) \times 10^{-3}$		—
$f_2(1270)\pi^+, f_2 \rightarrow \pi^0\pi^0$	$(5.0 \pm 3.5) \times 10^{-4}$		—
$2\pi^+\pi^-\pi^0$	—		935
$\omega\pi^+$	[e] $(1.93 \pm 0.30) \times 10^{-3}$		822
$\eta\pi^+$	[e] $(1.686 \pm 0.027) \%$		902
$(2\pi^+\pi^-\pi^0)_{\text{non-}\eta}$	$(2.04 \pm 0.09) \%$		—
$f_0(1370)^0\rho^+, f_0(1370)^0\rho^+ \rightarrow 2\pi^+\pi^-\pi^0$	$(5.1 \pm 0.9) \times 10^{-3}$		—
$f_0(980)^0\rho^+, f_0(980)^0\rho^+ \rightarrow 2\pi^+\pi^-\pi^0$	$(2.6 \pm 0.5) \times 10^{-3}$		—
$f_2(1270)^0\rho^+, f_2(1270)^0\rho^+ \rightarrow 2\pi^+\pi^-\pi^0$	$(1.9 \pm 0.4) \times 10^{-3}$		—
$(\rho^+\rho^0)_{S\text{-wave}} \rightarrow 2\pi^+\pi^-\pi^0$	$(7.1 \pm 2.8) \times 10^{-4}$		—
$(\rho(1450)^+\rho^0)_{S\text{-wave}} \rightarrow 2\pi^+\pi^-\pi^0$	$(9.4 \pm 3.1) \times 10^{-4}$		—
$(\rho^+\rho(1450)^0)_{P\text{-wave}} \rightarrow 2\pi^+\pi^-\pi^0$	$(1.75 \pm 0.29) \times 10^{-3}$		—
$\phi\pi^+, \phi \rightarrow \rho\pi$	$(5.08 \pm 0.35) \times 10^{-3}$		—
$\omega\pi^+, \omega \rightarrow \rho\pi$	$(1.41 \pm 0.19) \times 10^{-3}$		—
$a_1(1260)^+\pi^0, a_1^+ \rightarrow (\rho^0\pi^+)_{S\text{-wave}}$	$(2.6 \pm 0.4) \times 10^{-3}$		—
$a_1(1260)^0\pi^+, a_1^0 \rightarrow (\rho\pi)_{S\text{-wave}}$	$(1.3 \pm 0.5) \times 10^{-3}$		—
$\pi(1300)^0\pi^+, \pi(1300)^0 \rightarrow (\rho\pi)_{P\text{-wave}}$	$(2.4 \pm 0.7) \times 10^{-3}$		—
$3\pi^+2\pi^-$	$(8.0 \pm 0.8) \times 10^{-3}$		899
$2\pi^+\pi^-2\pi^0$	—		902
$\eta\rho^+$	[e] $(8.9 \pm 0.8) \%$		724
$\eta\pi^+\pi^0$	$(9.10 \pm 0.17) \%$		885

$\eta(\pi^+\pi^0)_{P\text{-wave}}$	(4.9 $\pm$ 3.0) $\times 10^{-3}$	885
$a_0(980)^{+0}\pi^{0+},$	(2.1 $\pm$ 0.4) %	—
$a_0(980)^{+0} \rightarrow \eta\pi^{+0}$		
$\omega\pi^+\pi^0$	[e] (2.8 $\pm$ 0.7) %	802
$2\pi^+\pi^-\eta$	(3.08 $\pm$ 0.08) %	855
$a_1(1260)^+\eta, a_1^+ \rightarrow$	(1.71 $\pm$ 0.14) %	—
$\rho(770)^0\pi^+, \rho^0 \rightarrow$		
$\pi^+\pi^-$		
$a_1(1260)^+\eta, a_1^+ \rightarrow$	(2.5 $\pm$ 0.9) $\times 10^{-3}$	—
$f_0(500)\pi^+, f_0 \rightarrow \pi^+\pi^-$		
$a_0(980)^+\rho(770)^0, a_0^+ \rightarrow$	(2.1 $\pm$ 0.9) $\times 10^{-3}$	—
$\eta\pi^+$		
$\eta(1405)\pi^+, \eta(1405) \rightarrow$	(2.2 $\pm$ 0.7) $\times 10^{-4}$	—
$a_0(980)^-\pi^+, a_0^- \rightarrow$		
$\eta\pi^-$		
$\eta(1405)\pi^+, \eta(1405) \rightarrow$	(2.2 $\pm$ 0.7) $\times 10^{-4}$	—
$a_0(980)^+\pi^-, a_0^+ \rightarrow$		
$\eta\pi^+$		
$f_1(1420)\pi^+, f_1 \rightarrow$	(5.9 $\pm$ 1.8) $\times 10^{-4}$	—
$a_0(980)^-\pi^+, a_0^- \rightarrow$		
$\eta\pi^-$		
$f_1(1420)\pi^+, f_1 \rightarrow$	(5.2 $\pm$ 1.8) $\times 10^{-4}$	—
$a_0(980)^+\pi^-, a_0^+ \rightarrow$		
$\eta\pi^+$		
$3\pi^+2\pi^-\pi^0$	(4.9 $\pm$ 3.2) %	856
$\omega 2\pi^+\pi^-$	[e] (1.6 $\pm$ 0.5) %	766
$\eta'(958)\pi^+$	[d,e] (3.95 $\pm$ 0.08) %	743
$3\pi^+2\pi^-\pi^0$	—	803
$\omega\eta\pi^+$	[e] (5.4 $\pm$ 1.3) $\times 10^{-3}$	654
$\eta'(958)\rho^+$	[d,e] (5.8 $\pm$ 1.5) %	465
$\eta'(958)\pi^+\pi^0$	(6.14 $\pm$ 0.18) %	720
$\eta'(958)\pi^+\pi^0_{\text{nonresonant}}$	< 5.1 % CL=90%	720

Modes with one or three K's

$K^+\pi^0$	(7.5 $\pm$ 0.5) $\times 10^{-4}$	917
$K_S^0\pi^+$	(1.22 $\pm$ 0.04) $\times 10^{-3}$	916
$K^+\eta$	[e] (1.76 $\pm$ 0.08) $\times 10^{-3}$	835
$K^+\omega$	[e] (9.9 $\pm$ 1.5) $\times 10^{-4}$	741
$K^+\eta'(958)$	[e] (2.68 $\pm$ 0.24) $\times 10^{-3}$	646
$K^+\pi^+\pi^-$	(6.23 $\pm$ 0.10) $\times 10^{-3}$	900
$K^+\rho^0$	(2.18 $\pm$ 0.25) $\times 10^{-3}$	745
$K^+\rho(1450)^0, \rho^0 \rightarrow \pi^+\pi^-$	(7.3 $\pm$ 1.7) $\times 10^{-4}$	—
$K^+f_0(500), f_0 \rightarrow \pi^+\pi^-$	(4.5 $\pm$ 3.0) $\times 10^{-4}$	—

$K^+ f_0(980), f_0 \rightarrow \pi^+ \pi^-$	$(2.8 \pm 1.1) \times 10^{-4}$	—
$K^+ f_0(1370), f_0 \rightarrow \pi^+ \pi^-$	$(1.2 \pm 0.6) \times 10^{-3}$	—
$K^*(892)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(1.68 \pm 0.26) \times 10^{-3}$	775
$K^*(1410)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(6 \pm 4) \times 10^{-4}$	—
$K^*(1430)^0 \pi^+, K^{*0} \rightarrow K^+ \pi^-$	$(9.4 \pm 3.2) \times 10^{-4}$	—
$K^+ \pi^+ \pi^-$ nonresonant	$(9.9 \pm 3.2) \times 10^{-4}$	900
$K_S^0 \pi^+ \pi^0$	$(5.09 \pm 0.22) \times 10^{-3}$	899
$K_S^0 \rho(770)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(2.6 \pm 0.4) \times 10^{-3}$	—
$K_S^0 \rho(1450)^+, \rho^+ \rightarrow \pi^+ \pi^0$	$(1.04 \pm 0.32) \times 10^{-3}$	—
$K^*(892)^0 \pi^+, K^{*0} \rightarrow K_S^0 \pi^0$	$(4.3 \pm 1.2) \times 10^{-4}$	—
$K^*(892)^+ \pi^0, K^{*+} \rightarrow K_S^0 \pi^+$	$(2.3 \pm 0.7) \times 10^{-4}$	—
$K^*(1410)^0 \pi^+, K^{*0} \rightarrow K_S^0 \pi^0$	$(1.7 \pm 0.9) \times 10^{-4}$	—
$K_S^0 2\pi^+ \pi^-$	$(2.8 \pm 1.0) \times 10^{-3}$	870
$K^+ \pi^+ \pi^- \pi^0$	$(9.7 \pm 0.6) \times 10^{-3}$	873
$K^*(892)^0 \rho^+, K^{*0} \rightarrow K^+ \pi^-$	$(3.9 \pm 0.4) \times 10^{-3}$	—
$K^*(892)^+ \rho^0, K^{*+} \rightarrow K^+ \pi^0$	$(4.2 \pm 1.2) \times 10^{-4}$	—
$K_1(1270)^0 \pi^+, K_1^0 \rightarrow K^+ \rho^-$	$(3.9 \pm 1.3) \times 10^{-4}$	†
$K_1(1400)^0 \pi^+, K_1^0 \rightarrow K^+ \pi^0$	$(5.4 \pm 0.9) \times 10^{-4}$	—
$K^*(890)^+ \pi^-, K^{*+} \rightarrow K^+ \pi^0$		
$K_1(1400)^0 \pi^+, K_1^0 \rightarrow K^*(890)^0 \pi^0, K^{*0} \rightarrow K^+ \pi^-$	$(5.9 \pm 1.0) \times 10^{-4}$	—
$K^+ a_1(1260)^0, a_1 \rightarrow \rho^+ \pi^-$	$(1.8 \pm 1.1) \times 10^{-4}$	—
$K^+ a_1(1260)^0, a_1 \rightarrow \rho^- \pi^+$	$(1.8 \pm 1.1) \times 10^{-4}$	—
$K^+ \pi^+ \pi^- \pi^0$ nonresonant	$(9.2 \pm 2.4) \times 10^{-4}$	873
$(K^+ \pi^0)_{P\text{-wave}} \rho^0$	$(1.01 \pm 0.21) \times 10^{-3}$	688
$K^+ \omega \pi^0$	$[e] < 8.2 \times 10^{-3} \text{CL}=90\%$	684
$K^+ \omega \pi^+ \pi^-$	$[e] < 5.4 \times 10^{-3} \text{CL}=90\%$	603
$K^+ \omega \eta$	$[e] < 7.9 \times 10^{-3} \text{CL}=90\%$	366
$2K^+ K^-$	$(2.18 \pm 0.20) \times 10^{-4}$	628
$\phi K^+, \phi \rightarrow K^+ K^-$	$(8.9 \pm 2.0) \times 10^{-5}$	—

Radiative decays

$\rho(770)^+ \gamma$	$< 6.1 \times 10^{-4} \text{CL}=90\%$	—
----------------------	---------------------------------------	---

Doubly Cabibbo-suppressed modes


$2K^+\pi^-$	$(1.293 \pm 0.027) \times 10^{-4}$	$S=1.1$	805
$K^+K^*(892)^0, K^{*0} \rightarrow$	$(6.1 \pm 3.4) \times 10^{-5}$		—
$2K^+\pi^-\pi^0$	$< 1.7 \times 10^{-4}$	CL=90%	748

Baryon-antibaryon mode

$p\bar{n}$	$(1.22 \pm 0.11) \times 10^{-3}$		295
$p\bar{p}e^+\nu_e$	$< 2.0 \times 10^{-4}$	CL=90%	296

**$\Delta C = 1$ weak neutral current (C1) modes,
Lepton family number (LF), or
Lepton number (L) violating modes**

$\pi^+e^+e^-$	[j]	$< 5.5 \times 10^{-6}$	CL=90%	979
$\pi^+\phi, \phi \rightarrow e^+e^-$	[k]	$(1.17 \pm 0.22) \times 10^{-5}$		—
$\pi^+\pi^0e^+e^-$		$< 7.0 \times 10^{-5}$	CL=90%	974
$\rho^+\phi, \phi \rightarrow e^+e^-$		$(2.4 \pm 0.7) \times 10^{-5}$		—
$\pi^+\mu^+\mu^-$	[j]	$< 1.8 \times 10^{-7}$	CL=90%	968
$K^+e^+e^-$	C1	$< 3.7 \times 10^{-6}$	CL=90%	922
$K^+\pi^0e^+e^-$		$< 7.1 \times 10^{-5}$	CL=90%	917
$K_S^0\pi^+e^+e^-$		$< 8.1 \times 10^{-5}$	CL=90%	916
$K^+\mu^+\mu^-$	C1	$< 1.4 \times 10^{-7}$	CL=90%	909
$K^*(892)^+\mu^+\mu^-$	C1	$< 1.4 \times 10^{-3}$	CL=90%	765
$\pi^+e^+\mu^-$	LF	$< 1.1 \times 10^{-6}$	CL=90%	976
$\pi^+e^-\mu^+$	LF	$< 9.4 \times 10^{-7}$	CL=90%	976
$K^+e^+\mu^-$	LF	$< 7.9 \times 10^{-7}$	CL=90%	919
$K^+e^-\mu^+$	LF	$< 5.6 \times 10^{-7}$	CL=90%	919
π^-2e^+	L	$< 1.4 \times 10^{-6}$	CL=90%	979
$\pi^-2\mu^+$	L	$< 8.6 \times 10^{-8}$	CL=90%	968
$\pi^-e^+\mu^+$	L	$< 6.3 \times 10^{-7}$	CL=90%	976
K^-2e^+	L	$< 7.7 \times 10^{-7}$	CL=90%	922
$K^-2\mu^+$	L	$< 2.6 \times 10^{-8}$	CL=90%	909
$K^-e^+\mu^+$	L	$< 2.6 \times 10^{-7}$	CL=90%	919
$K^*(892)^-2\mu^+$	L	$< 1.4 \times 10^{-3}$	CL=90%	765



$$I(J^P) = 0(1^-)$$

Mass $m = 2112.2 \pm 0.4$ MeV

$$m_{D_s^{*\pm}} - m_{D_s^\pm} = 143.8 \pm 0.4 \text{ MeV}$$

Full width $\Gamma < 1.9$ MeV, CL = 90%

D_s^{*-} modes are charge conjugates of the modes below.

D_s^{*+} DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D_s^+ \gamma$	(93.6 $\pm$ 0.4) %	139
$D_s^+ \pi^0$	(5.77 $\pm$ 0.35) %	48
$D_s^+ e^+ e^-$	(6.7 $\pm$ 1.6) $\times 10^{-3}$	139
$e^+ \nu_e$	(2.1 $^{+1.2}_{-0.9}$) $\times 10^{-5}$	1056

$D_{s0}^*(2317)^\pm$

$I(J^P) = 0(0^+)$
 J, P need confirmation.
 J^P is natural, low mass consistent with 0^+ .

See the review on "Heavy Non- $q\bar{q}$ Mesons."

Mass $m = 2317.8 \pm 0.5$ MeV

$m_{D_{s0}^*(2317)^\pm} - m_{D_s^\pm} = 349.4 \pm 0.5$ MeV

Full width $\Gamma < 3.8$ MeV, CL = 95%

$D_{s0}^*(2317)^-$ modes are charge conjugates of modes below.

$D_{s0}^*(2317)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$D_s^+ \pi^0$	(100 $^{+0}_{-20}$) %		298
$D_s^+ \gamma$	< 5 %	90%	323
$D_{s0}^*(2112)^+ \gamma$	< 6 %	90%	—
$D_s^+ \gamma \gamma$	< 18 %	95%	323
$D_{s0}^*(2112)^+ \pi^0$	< 11 %	90%	—
$D_s^+ \pi^+ \pi^-$	< 4 $\times 10^{-3}$	90%	194
$D_s^+ \pi^0 \pi^0$	not seen		205

$D_{s1}(2460)^\pm$

$I(J^P) = 0(1^+)$

See the review on "Heavy Non- $q\bar{q}$ Mesons."

Mass $m = 2459.5 \pm 0.6$ MeV ($S = 1.1$)

$m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}} = 347.3 \pm 0.7$ MeV ($S = 1.2$)

$m_{D_{s1}(2460)^\pm} - m_{D_s^\pm} = 491.1 \pm 0.6$ MeV ($S = 1.1$)

Full width $\Gamma < 3.5$ MeV, CL = 95%

$D_{s1}(2460)^-$ modes are charge conjugates of the modes below.

$D_{s1}(2460)^+$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$D_s^{*+} \pi^0$	(48 $\pm$ 11) %		297
$D_s^+ \gamma$	(18 $\pm$ 4) %		442
$D_s^+ \pi^+ \pi^-$	(4.3 $\pm$ 1.3) %	S=1.1	363
$D_s^{*+} \gamma$	< 8 %	CL=90%	323
$D_{s0}^*(2317)^+ \gamma$	(3.7 $^{+5.0}_{-2.4}$) %		138

$D_{s1}(2536)^\pm$

$$I(J^P) = 0(1^+)$$

J, P need confirmation.

Mass $m = 2535.11 \pm 0.06$ MeV

$$m_{D_{s1}(2536)^\pm} - m_{D_s^*(2111)} = 422.9 \pm 0.4 \text{ MeV}$$

$$m_{D_{s1}(2536)^\pm} - m_{D^*(2010)^\pm} = 524.85 \pm 0.04 \text{ MeV}$$

$$m_{D_{s1}(2536)^\pm} - m_{D^*(2007)^0} = 528.26 \pm 0.05 \text{ MeV} \quad (S = 1.1)$$

$$\text{Full width } \Gamma = 0.92 \pm 0.05 \text{ MeV}$$

Branching fractions are given relative to the one **DEFINED AS 1**.

$D_{s1}(2536)^-$ modes are charge conjugates of the modes below.

$D_{s1}(2536)^+$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$D^*(2010)^+ K^0$	(31 $\pm$ 7) %		149
$(D^*(2010)^+ K^0)_{S\text{-wave}}$	(22 $\pm$ 5) %		149
$K_S^0 D^*(2010)^+$	(17 $\pm$ 4) %		149
$D^+ \pi^- K^+$	$(10.0 \pm 2.5) \times 10^{-3}$		176
$D^*(2007)^0 K^+$	(36 $\pm$ 6) %		167
$D^+ K^0$	< 12 %	90%	381
$D^0 K^+$	< 4 %	90%	391
$D_s^{*+} \gamma$	possibly seen		388
$D_s^+ \pi^+ \pi^-$	seen		437

$D_{s2}^*(2573)$

$$I(J^P) = 0(2^+)$$

Mass $m = 2569.1 \pm 0.8$ MeV ($S = 2.4$)

$$m_{D_{s2}^*(2573)} - m_{D^0} = 704 \pm 3.2 \text{ MeV}$$

$$\text{Full width } \Gamma = 16.9 \pm 0.7 \text{ MeV}$$

$D_{s2}^*(2573)^-$ modes are charge conjugates of the modes below.

$D_{s2}^*(2573)^+$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 K^+$	$(37 \pm 6) \%$	431
$D^*(2007)^0 K^+$	not seen	238
$D^+ K_S^0$	seen	422
$D^{*+} K_S^0$	seen	225

$D_{s1}^*(2700)^\pm$

$I(J^P) = 0(1^-)$

Mass $m = 2714 \pm 5$ MeV ($S = 1.5$)
Full width $\Gamma = 122 \pm 10$ MeV

$D_{s1}^*(2700)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 K^+$	seen	579
$D^+ K_S^0$	seen	573
$D^{*0} K^+$	seen	438
$D^{*+} K_S^0$	seen	431

$D_{s3}^*(2860)^\pm$

$I(J^P) = 0(3^-)$

Mass $m = 2860 \pm 7$ MeV
Full width $\Gamma = 53 \pm 10$ MeV

$D_{s3}^*(2860)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$D^0 K^+$	seen	710
$D^+ K_S^0$	seen	704
$D^{*0} K^+$	seen	589
$D^{*+} K_S^0$	seen	584

NOTES

- [a] See the Particle Listings for the (complicated) definition of this quantity.
- [b] This is the purely e^+ semileptonic branching fraction: the e^+ fraction from τ^+ decays has been subtracted off. The sum of our (non- τ) e^+ exclusive fractions — an $e^+ \nu_e$ with an η , η' , ϕ , K^0 , or K^{*0} — is 5.99 ± 0.31 %.
- [c] This fraction includes η from η' decays.
- [d] The sum of our exclusive η' fractions — $\eta' e^+ \nu_e$, $\eta' \mu^+ \nu_\mu$, $\eta' \pi^+$, $\eta' \rho^+$, and $\eta' K^+$ — is 11.8 ± 1.6 %.
- [e] This branching fraction includes all the decay modes of the final-state resonance.
- [f] A test for $u\bar{u}$ or $d\bar{d}$ content in the D_s^+ . Neither Cabibbo-favored nor Cabibbo-suppressed decays can contribute, and ω – ϕ mixing is an unlikely explanation for any fraction above about 2×10^{-4} .
- [g] The branching fraction for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Particle Listings.
- [h] We decouple the $D_s^+ \rightarrow \phi \pi^+$ branching fraction obtained from mass projections (and used to get some of the other branching fractions) from the $D_s^+ \rightarrow \phi \pi^+$, $\phi \rightarrow K^+ K^-$ branching fraction obtained from the Dalitz-plot analysis of $D_s^+ \rightarrow K^+ K^- \pi^+$. That is, the ratio of these two branching fractions is not exactly the $\phi \rightarrow K^+ K^-$ branching fraction 0.491.
- [i] This is the average of a model-independent and a K -matrix parametrization of the $\pi^+ \pi^-$ S -wave and is a sum over several f_0 mesons.
- [j] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.
- [k] This is *not* a test for the $\Delta C=1$ weak neutral current, but leads to the $\pi^+ \ell^+ \ell^-$ final state.