

$B_2^*(5747)$

$$I(J^P) = \frac{1}{2}(2^+)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

 $B_2^*(5747)$ MASS **$B_2^*(5747)^+$ mass**OUR FIT uses m_{B^0} and $m_{B_2^{*+}} - m_{B^0}$ to determine $m_{B_2^*(5747)^+}$.

VALUE (MeV)	DOCUMENT ID
5737.3 ± 0.7 OUR FIT	

 $m_{B_2^{*+}} - m_{B^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
457.5 ± 0.7 OUR FIT				
457.5 ± 0.7 OUR AVERAGE				
457.62 ± 0.72 ± 0.40	4k	¹ AAIJ	15AB	LHCB $p\bar{p}$ at 7, 8 TeV
457.3 ± 1.3 $\begin{smallmatrix} +0.3 \\ -0.9 \end{smallmatrix}$		² AALTONEN	14l	CDF $p\bar{p}$ at 1.96 TeV

¹ AAIJ 15AB reports $[m_{B_2^{*+}} - m_{B^0}] - m_{\pi^+} = 318.1 \pm 0.7 \pm 0.4$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14l reports $m_{B_2^*(5747)^+} - m_{B^0} - m_{\pi^+} = 317.7 \pm 1.2 \begin{smallmatrix} +0.3 \\ -0.9 \end{smallmatrix}$ MeV which we adjusted by the π^+ mass.

 $B_2^*(5747)^0$ massOUR FIT uses m_{B^+} , $m_{B_1^0} - m_{B^+}$, and mass differences below to determine $m_{B_2^*(5747)^0}$. The -0.659 correlation between statistical uncertainties of $m_{B_1^0} - m_{B^+}$ and $m_{B_2^{*0}} - m_{B_1^0}$ measurements reported by ABAZOV 07T is taken into account.

VALUE (MeV)	DOCUMENT ID
5739.6 ± 0.7 OUR FIT	Error includes scale factor of 1.4.

 $m_{B_2^{*0}} - m_{B_1^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
13.5 ± 1.4 OUR FIT	Error includes scale factor of 1.3.		
26.2 ± 3.1 ± 0.9	¹ ABAZOV	07T D0	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
14.9 $\begin{smallmatrix} +2.2+1.2 \\ -2.5-1.4 \end{smallmatrix}$	¹ AALTONEN	09D	CDF Repl. by AALTONEN 14l

¹ Observed in $B_2^{*0} \rightarrow B^{*+} \pi^-$ and $B_2^{*0} \rightarrow B^+ \pi^-$.

 $m_{B_2^{*0}} - m_{B^+}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
460.2 ± 0.6 OUR FIT	Error includes scale factor of 1.4.			
459.9 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.8.			
460.18 ± 0.37 ± 0.33	17k	¹ AAIJ	15AB	LHCB $p\bar{p}$ at 7, 8 TeV
457.5 ± 1.2 $\begin{smallmatrix} +0.8 \\ -0.9 \end{smallmatrix}$		² AALTONEN	14l	CDF $p\bar{p}$ at 1.96 TeV

- ¹ AAIJ 15AB reports $[m_{B_2^{*0}} - m_{B^+}] - m_{\pi^-} = 320.6 \pm 0.4 \pm 0.3$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event.
- ² AALTONEN 14I reports $m_{B_2^*(5747)^0} - m_{B^+} - m_{\pi^-} = 317.9 \pm 1.2^{+0.8}_{-0.9}$ MeV which we adjusted by the π^- mass.

$B_2^*(5747)$ WIDTH

$B_2^*(5747)^+$ width

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20 ± 5 OUR AVERAGE		Error includes scale factor of 2.2.		
23.6 ± 2.0 ± 2.1	4k	AAIJ	15AB LHCB	pp at 7, 8 TeV
11 $^{+4}_{-3}$ $^{+3}_{-4}$		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

$B_2^*(5747)^0$ width

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
24.2 ± 1.7 OUR AVERAGE				
24.5 ± 1.0 ± 1.5	17k	AAIJ	15AB LHCB	pp at 7, 8 TeV
22 $^{+3}_{-2}$ $^{+4}_{-5}$		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
22.7 $^{+3.8}_{-3.2}$ $^{+3.2}_{-10.2}$		AALTONEN	09D CDF	Repl. by AALTONEN 14I

$B_2^*(5747)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B\pi$	seen
Γ_2 $B^*\pi$	seen

$B_2^*(5747)$ BRANCHING RATIOS

$\Gamma(B\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	4k,17k	AAIJ	15AB LHCB	±0	pp at 7, 8 TeV	
seen		AALTONEN	14I CDF	±	$p\bar{p}$ at 1.96 TeV	
seen		AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV	
seen		ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV	

$\Gamma(B^*\pi)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	4k,17k	AAIJ	15AB LHCB	±0	pp at 7, 8 TeV	
seen		AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV	
seen		ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV	

$\Gamma(B^*\pi)/\Gamma(B\pi)$						Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
0.84 ± 0.27 OUR AVERAGE						
0.71 ± 0.14 ± 0.30	17k	AAIJ	15AB LHCB	0	pp at 7, 8 TeV	

$1.0 \pm 0.5 \pm 0.8$ 4k AAIJ 15AB LHCb \pm pp at 7, 8 TeV
 $1.10 \pm 0.42 \pm 0.31$ ¹ ABAZOV 07T D0 0 $p\bar{p}$ at 1.96 TeV
¹ Converted from measured ratio of $R = B(B_2^{*0} \rightarrow B^{*+} \pi^-) / B(B_2^{*0} \rightarrow B^{(*)+} \pi^-)$
 $= 0.475 \pm 0.095 \pm 0.069$.

$B_2^*(5747)$ REFERENCES

AAIJ	15AB	JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09D	PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07T	PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)