

$B_J(5840)$

$$I(J^P) = \frac{1}{2}(??)$$

I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

 $B_J(5840)$ MASS **$B_J(5840)^+$ MASS**OUR FIT uses m_{B^0} and $m_{B_J(5840)^+} - m_{B^0}$ to determine $m_{B_J(5840)^+}$.

| VALUE (MeV) | DOCUMENT ID |
|--------------------------|-------------|
| 5851 ± 19 OUR FIT | |

 $m_{B_J(5840)^+} - m_{B^0}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------------------|------|-------------|------|---------|
| 571 ± 19 OUR FIT | | | | |

| | | | | |
|----------------------|----|-------------------|-----------|------------------|
| 571 ± 13 ± 14 | 7k | ¹ AAIJ | 15AB LHCB | pp at 7, 8 TeV |
|----------------------|----|-------------------|-----------|------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------|----|-------------------|-----------|------------------|
| 595 ± 26 ± 14 | 7k | ² AAIJ | 15AB LHCB | pp at 7, 8 TeV |
|---------------|----|-------------------|-----------|------------------|

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 431 \pm 13 \pm 14$ MeV which we adjust bythe π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.² AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 455 \pm 26 \pm 14$ MeV which we adjust bythe π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$m_{B_J(5840)^+} - m_{B^{*0}}$**

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|----------------------|------|-------------------|-----------|------------------|
| 565 ± 15 ± 14 | 7k | ¹ AAIJ | 15AB LHCB | pp at 7, 8 TeV |

• • • We do not use the following data for averages, fits, limits, etc. • • •

| | | | | |
|---------------|----|-------------------|-----------|------------------|
| 565 ± 15 ± 14 | 7k | ¹ AAIJ | 15AB LHCB | pp at 7, 8 TeV |
|---------------|----|-------------------|-----------|------------------|

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 425 \pm 15 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$B_J(5840)^0$ MASS**OUR FIT uses m_{B^+} and $m_{B_J(5840)^0} - m_{B^+}$ to determine $m_{B_J(5840)^0}$.

| VALUE (MeV) | DOCUMENT ID |
|-------------------------|-------------|
| 5863 ± 9 OUR FIT | |

 $m_{B_J(5840)^0} - m_{B^+}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------|------|-------------|------|---------|
| 584 ± 9 OUR FIT | | | | |

| | | | | |
|--------------------|-----|-------------------|-----------|------------------|
| 584 ± 5 ± 7 | 12k | ¹ AAIJ | 15AB LHCB | pp at 7, 8 TeV |
|--------------------|-----|-------------------|-----------|------------------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

610 ± 22 ± 7 12k ² AAIJ 15AB LHCB *pp* at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust by

the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 471 \pm 22 \pm 7$ MeV which we adjust by

the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$m_{B_J(5840)^0} - m_{B^{*+}}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

• • • We do not use the following data for averages, fits, limits, etc. • • •

584 ± 5 ± 7 12k ¹ AAIJ 15AB LHCB *pp* at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV

which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5840)$ WIDTH

$B_J(5840)^+$ WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

224 ± 24 ± 80 7k ¹ AAIJ 15AB LHCB *pp* at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

215 ± 27 ± 80 7k ² AAIJ 15AB LHCB *pp* at 7, 8 TeV

229 ± 27 ± 80 7k ³ AAIJ 15AB LHCB *pp* at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5840)^0$ WIDTH

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|-------------|------|-------------|------|---------|
|-------------|------|-------------|------|---------|

127 ± 17 ± 34 12k ¹ AAIJ 15AB LHCB *pp* at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

107 ± 20 ± 34 12k ² AAIJ 15AB LHCB *pp* at 7, 8 TeV

119 ± 17 ± 34 12k ³ AAIJ 15AB LHCB *pp* at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5840)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|----------------------|--------------------------------|
| Γ_1 $B^* \pi$ | seen |
| Γ_2 $B \pi$ | possibly seen |

 $B_J(5840)$ BRANCHING RATIOS

| $\Gamma(B^* \pi)/\Gamma_{\text{total}}$ | | | | | | Γ_1/Γ |
|---|-------------|--------------------|-------------|------------|------------------|-------------------|
| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | |
| seen | 7k | AAIJ | 15AB LHCB | \pm | pp at 7, 8 TeV | |
| seen | 12k | AAIJ | 15AB LHCB | 0 | pp at 7, 8 TeV | |

| $\Gamma(B \pi)/\Gamma_{\text{total}}$ | | | | | | Γ_2/Γ |
|---------------------------------------|-------------|--------------------|-------------|------------|------------------|-------------------|
| <u>VALUE</u> | <u>EVTS</u> | <u>DOCUMENT ID</u> | <u>TECN</u> | <u>CHG</u> | <u>COMMENT</u> | |
| possibly seen | 7k | ¹ AAIJ | 15AB LHCB | \pm | pp at 7, 8 TeV | |
| possibly seen | | ¹ AAIJ | 15AB LHCB | 0 | pp at 7, 8 TeV | |

¹A $B \pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^* \pi$ is allowed.

 $B_J(5840)$ REFERENCES

| | | | |
|------|--------------------|-----------------------|----------------|
| AAIJ | 15AB JHEP 1504 024 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
|------|--------------------|-----------------------|----------------|