

$J/\psi(1S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $J/\psi(1S)$ MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3096.900±0.006 OUR AVERAGE				
3096.900±0.002±0.006		¹ ANASHIN 15	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ±0.09	502	² ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ±0.03 ±0.01		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ±0.1 ±0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3096.66 ±0.19 ±0.02	6.1k	⁴ AAIJ 15BI	LHCB	$pp \rightarrow J/\psi X$
3096.917±0.010±0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ±0.3		GRIBUSHIN 96	FMPS	$515 \pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ±2.0	38k	LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
3096.93 ±0.09	502	⁵ ZHOLENTZ 80	REDE	e^+e^-
3097.0 ±1		⁶ BRANDELIK 79C	DASP	e^+e^-

¹ Supersedes AULCHENKO 03.² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁴ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.⁵ Superseded by ARTAMONOV 00.⁶ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$. **$J/\psi(1S)$ WIDTH**

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
92.6 ± 1.7 OUR AVERAGE				
Error includes scale factor of 1.1.				
92.45± 1.40±1.48		¹ ANASHIN 20	KEDR	e^+e^-
96.1 ± 3.2	13k	² ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4 ± 8.9		BAI 95B	BES	e^+e^-
91 ±11 ±6		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 $\begin{smallmatrix} + 6.1 \\ - 5.8 \end{smallmatrix}$		⁴ HSUEH 92	RVUE	See Υ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
92.94± 1.83		^{5,6} ANASHIN 18A	KEDR	e^+e^-
94.1 ± 2.7		⁷ ANASHIN 10	KEDR	$3.097 e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7 ± 3.5	7.8k	² AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there.² Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(e^+e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.³ The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

⁴ Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.

⁵ Using $\Gamma(e^+e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ from PDG 16.

⁶ Superseded by ANASHIN 20 that is based on the same dataset.

⁷ Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

$J/\psi(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.46 \pm 0.07) %	
Γ_3 $g g g$	(64.1 \pm 1.0) %	
Γ_4 $\gamma g g$	(8.8 \pm 1.1) %	
Γ_5 e^+e^-	(5.971 \pm 0.032) %	
Γ_6 $e^+e^- \gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	
Γ_7 $\mu^+\mu^-$	(5.961 \pm 0.033) %	
Γ_8 $e^+e^-e^+e^-$	(5.5 \pm 0.5) $\times 10^{-5}$	
Γ_9 $e^+e^-\mu^+\mu^-$	(3.53 \pm 0.26) $\times 10^{-5}$	
Γ_{10} $\mu^+\mu^-\mu^+\mu^-$	(1.11 \pm 0.11) $\times 10^{-6}$	

Decays involving hadronic resonances

Γ_{11} $\rho\pi$	(1.88 \pm 0.12) %	S=2.6
Γ_{12} $\rho^0\pi^0$	(6.2 \pm 0.6) $\times 10^{-3}$	
Γ_{13} $a_2(1320)^0\pi^+\pi^- \rightarrow$ $2(\pi^+\pi^-)\pi^0$	(2.8 \pm 0.6) $\times 10^{-3}$	
Γ_{14} $a_2(1320)^+\pi^-\pi^0 + \text{c.c.} \rightarrow$ $2(\pi^+\pi^-)\pi^0$	(3.7 \pm 0.7) $\times 10^{-3}$	
Γ_{15} $a_2(1320)\rho$	(1.09 \pm 0.22) %	
Γ_{16} $\eta\pi^+\pi^-$	(3.8 \pm 0.7) $\times 10^{-4}$	
Γ_{17} $\eta\rho$	(1.93 \pm 0.23) $\times 10^{-4}$	
Γ_{18} $\eta\pi^+\pi^-\pi^0$	(1.17 \pm 0.20) %	
Γ_{19} $\eta\pi^+\pi^-3\pi^0$	(4.9 \pm 1.0) $\times 10^{-3}$	
Γ_{20} $\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow$ $\eta\phi\pi^+\pi^-$	(1.2 \pm 0.4) $\times 10^{-4}$	
Γ_{21} $\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0 \bar{K}^*(892)^0$	< 2.52 $\times 10^{-4}$	CL=90%
Γ_{22} $\eta K^+ K^-$	(8.6 \pm 3.0) $\times 10^{-4}$	
Γ_{23} $\eta K^\pm K_S^0 \pi^\mp$	[b] (2.2 \pm 0.4) $\times 10^{-3}$	
Γ_{24} $\eta K^*(892)^0 \bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$	
Γ_{25} $\rho\eta'(958)$	(8.1 \pm 0.8) $\times 10^{-5}$	S=1.6
Γ_{26} $\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0$	(2.8 \pm 0.8) %	
Γ_{27} $\rho^+ \rho^- \pi^+ \pi^- \pi^0$	(6 \pm 4) $\times 10^{-3}$	
Γ_{28} $\rho^+ K^+ K^- \pi^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^+ \pi^- \pi^0$	(3.5 \pm 0.8) $\times 10^{-3}$	

Γ_{29}	$\rho^\mp K^\pm K_S^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{30}	$h_1(1415)\eta' \rightarrow \gamma\eta\eta'$		
Γ_{31}	$h_1(1595)\eta' \rightarrow \gamma\eta\eta'$		
Γ_{32}	$\rho(1450)\pi$	seen	
Γ_{33}	$\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	$(2.2 \pm 1.1) \times 10^{-4}$	
Γ_{34}	$\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp$	$(3.3 \pm 0.6) \times 10^{-4}$	
Γ_{35}	$\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0$	$(2.7 \pm 0.6) \times 10^{-4}$	
Γ_{36}	$\rho(1450)\eta'(958) \rightarrow$ $\pi^+\pi^-\eta'(958)$	$(3.3 \pm 0.7) \times 10^{-6}$	
Γ_{37}	$\rho(1700)\pi$	seen	
Γ_{38}	$\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	$(1.6 \pm 1.1) \times 10^{-4}$	
Γ_{39}	$\rho(2150)\pi$	seen	
Γ_{40}	$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	$(10 \pm 40) \times 10^{-6}$	
Γ_{41}	$\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$		
Γ_{42}	$\omega\pi^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4
Γ_{43}	$\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0$	$(1.6 \pm 0.7) \times 10^{-5}$	
Γ_{44}	$\omega\pi^+\pi^-$	$(8.5 \pm 1.0) \times 10^{-3}$	S=1.3
Γ_{45}	$\omega\pi^0\pi^0$	$(3.4 \pm 0.8) \times 10^{-3}$	
Γ_{46}	$\omega 3\pi^0$	$(1.9 \pm 0.6) \times 10^{-3}$	
Γ_{47}	$\omega f_2(1270)$	$(4.3 \pm 0.6) \times 10^{-3}$	
Γ_{48}	$\omega\eta$	$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6
Γ_{49}	$\omega\pi^+\pi^-\pi^0$	$(4.0 \pm 0.7) \times 10^{-3}$	
Γ_{50}	$\omega\pi^0\eta$	$(3.4 \pm 1.7) \times 10^{-4}$	
Γ_{51}	$\omega\pi^+\pi^+\pi^-\pi^-$	$(8.5 \pm 3.4) \times 10^{-3}$	
Γ_{52}	$\omega\pi^+\pi^-2\pi^0$	$(3.3 \pm 0.5) \%$	
Γ_{53}	$\omega\eta'\pi^+\pi^-$	$(1.12 \pm 0.13) \times 10^{-3}$	
Γ_{54}	$\omega\eta'(958)$	$(1.89 \pm 0.18) \times 10^{-4}$	
Γ_{55}	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{56}	$\omega f_0(1710) \rightarrow \omega K\bar{K}$	$(4.8 \pm 1.1) \times 10^{-4}$	
Γ_{57}	$\omega f_1(1420)$	$(6.8 \pm 2.4) \times 10^{-4}$	
Γ_{58}	$\omega f_2'(1525)$	$< 2.2 \times 10^{-4}$	CL=90%
Γ_{59}	$\omega X(1835) \rightarrow \omega p\bar{p}$	$< 3.9 \times 10^{-6}$	CL=95%
Γ_{60}	$\omega K^+K^-\eta$	$(3.33 \pm 0.12) \times 10^{-4}$	
Γ_{61}	$\omega X(1835), X \rightarrow \eta'\pi^+\pi^-$	$< 6.2 \times 10^{-5}$	
Γ_{62}	ωK^+K^-	$(1.52 \pm 0.31) \times 10^{-3}$	
Γ_{63}	$\omega K^\pm K_S^0\pi^\mp$	[b] $(3.4 \pm 0.5) \times 10^{-3}$	
Γ_{64}	$\omega K\bar{K}$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{65}	$\omega K^*(892)\bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$	
Γ_{66}	$\eta' K^{*\pm} K^\mp$	$(1.48 \pm 0.13) \times 10^{-3}$	
Γ_{67}	$\eta' K^{*0}\bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$	
Γ_{68}	$\eta' h_1(1415) \rightarrow \eta' K^*\bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$	
Γ_{69}	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$	
Γ_{70}	$\eta' h_1(1415) \rightarrow \gamma\eta'\eta'$	$(4.7 \pm \frac{1.1}{2.0}) \times 10^{-7}$	

Γ_{71}	$\bar{K} K^*(892) + \text{c.c.}$	seen	
Γ_{72}	$\bar{K} K^*(892) + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(4.8 \pm 0.5) \times 10^{-3}$	
Γ_{73}	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm_{-1.0}^{0.8}) \times 10^{-3}$	S=2.9
Γ_{74}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm_{-0.20}^{0.13}) \times 10^{-3}$	
Γ_{75}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$	
Γ_{76}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$	
Γ_{77}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$	
Γ_{78}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(5.7 \pm 0.8) \times 10^{-3}$	
Γ_{79}	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$	
Γ_{80}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$	
Γ_{81}	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow$ $K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$	
Γ_{82}	$K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow$ $K^+ K^- \pi^+ \pi^-$	$(3.8 \pm 0.5) \times 10^{-3}$	
Γ_{83}	$K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm_{-0.5}^{0.6}) \times 10^{-6}$	
Γ_{84}	$K^*(892)^0 K_S^0 \pi^0$	$(7 \pm 4) \times 10^{-4}$	
Γ_{85}	$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \pm_{-0.6}^{1.0}) \times 10^{-3}$	
Γ_{86}	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$	
Γ_{87}	$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \pm_{-0.40}^{0.22}) \times 10^{-3}$	
Γ_{88}	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$	
Γ_{89}	$K^*(1410) \bar{K} + \text{c.c.}$	seen	
Γ_{90}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$	
Γ_{91}	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(8 \pm 5) \times 10^{-5}$	
Γ_{92}	$K_2^*(1430) \bar{K} + \text{c.c.}$	seen	
Γ_{93}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$	
Γ_{94}	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{95}	$\bar{K}_2^*(1430) K + \text{c.c.}$	$< 4.0 \times 10^{-3}$	CL=90%
Γ_{96}	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm_{-0.19}^{0.25}) \times 10^{-4}$	
Γ_{97}	$K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow$ $K^+ K^- \pi^+ \pi^-$	$(2.6 \pm 0.9) \times 10^{-3}$	
Γ_{98}	$K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$	
Γ_{99}	$\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$	

Γ_{100}	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$	
Γ_{101}	$K_2^*(1430)^- K^*(892)^+ +$ $\text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- +$ c.c.	$(4 \pm 4) \times 10^{-4}$	
Γ_{102}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	< 2.9	$\times 10^{-3}$ CL=90%
Γ_{103}	$\bar{K}_2^*(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow$ $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$	
Γ_{104}	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(1.10 \pm_{-0.14}^{0.60}) \times 10^{-5}$	
Γ_{105}	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(6.2 \pm_{-1.6}^{2.9}) \times 10^{-6}$	
Γ_{106}	$K_1(1270)^\pm K^\mp$	< 3.0	$\times 10^{-3}$ CL=90%
Γ_{107}	$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$	
Γ_{108}	$a_2(1320)^\pm \pi^\mp$	$[b] < 4.3$	$\times 10^{-3}$ CL=90%
Γ_{109}	$\phi \pi^0$	3×10^{-6} or 1×10^{-7}	
Γ_{110}	$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4}$	S=1.7
Γ_{111}	$\phi \pi^0 \pi^0$	$(4.9 \pm 1.0) \times 10^{-4}$	
Γ_{112}	$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$	
Γ_{113}	$\phi \eta$	$(7.4 \pm 0.6) \times 10^{-4}$	S=1.2
Γ_{114}	$\phi \eta'(958)$	$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2
Γ_{115}	$\phi \eta \eta'$	$(2.32 \pm 0.17) \times 10^{-4}$	
Γ_{116}	$\phi f_0(980)$	$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9
Γ_{117}	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	$(2.57 \pm 0.34) \times 10^{-4}$	
Γ_{118}	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$	$(1.7 \pm 0.5) \times 10^{-4}$	
Γ_{119}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$	
Γ_{120}	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$	
Γ_{121}	$\phi f_0(980) \eta \rightarrow \eta \phi \pi^+ \pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$	
Γ_{122}	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$		
Γ_{123}	$\phi(1680)^0 \pi^0 \rightarrow \phi \eta \pi^0$	$(6.7 \pm 1.1) \times 10^{-6}$	
Γ_{124}	$X(2000)^0 \pi^0 \rightarrow \phi \eta \pi^0$	$(1.70 \pm_{-0.23}^{0.50}) \times 10^{-6}$	
Γ_{125}	$h_1(1900)^0 \pi^0 \rightarrow \phi \eta \pi^0$	$(8.4 \pm_{-1.3}^{1.4}) \times 10^{-6}$	
Γ_{126}	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$	
Γ_{127}	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$	
Γ_{128}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi \pi^0 \pi^+ \pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$	
Γ_{129}	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow$ $\phi 3\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$	
Γ_{130}	$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$	
Γ_{131}	$\phi f_2'(1525)$	$(8 \pm 4) \times 10^{-4}$	S=2.7
Γ_{132}	$\phi X(1835) \rightarrow \phi p \bar{p}$	< 2.1	$\times 10^{-7}$ CL=90%
Γ_{133}	$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$	< 2.8	$\times 10^{-4}$ CL=90%
Γ_{134}	$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$	< 6.13	$\times 10^{-5}$ CL=90%

Γ_{135}	$\phi K \bar{K}$		$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3
Γ_{136}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$		$(3.6 \pm 0.6) \times 10^{-4}$	
Γ_{137}	$\phi K^+ K^-$		$(8.2 \pm 1.1) \times 10^{-4}$	
Γ_{138}	$\phi K_S^0 K_S^0$		$(5.8 \pm 1.5) \times 10^{-4}$	
Γ_{139}	$\phi K^\pm K_S^0 \pi^\mp$	[b]	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{140}	$\phi K^*(892) \bar{K} + \text{c.c.}$		$(2.18 \pm 0.23) \times 10^{-3}$	
Γ_{141}	$b_1(1235)^\pm \pi^\mp$	[b]	$(3.0 \pm 0.5) \times 10^{-3}$	
Γ_{142}	$b_1(1235)^0 \pi^0$		$(2.3 \pm 0.6) \times 10^{-3}$	
Γ_{143}	$f_2'(1525) K^+ K^-$		$(1.04 \pm 0.35) \times 10^{-3}$	
Γ_{144}	$\Delta(1232)^+ \bar{p}$		$< 1 \times 10^{-4}$	CL=90%
Γ_{145}	$\Delta(1232)^{++} \bar{p} \pi^-$		$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{146}	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$		$(1.10 \pm 0.29) \times 10^{-3}$	
Γ_{147}	$\bar{\Sigma}(1385)^0 p K^-$		$(5.1 \pm 3.2) \times 10^{-4}$	
Γ_{148}	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$		$< 8.2 \times 10^{-6}$	CL=90%
Γ_{149}	$\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.}$	[b]	$(3.0 \pm 0.7) \times 10^{-4}$	
Γ_{150}	$\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.}$		$(3.3 \pm 0.8) \times 10^{-4}$	
Γ_{151}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.}$	[b]	$(1.08 \pm 0.06) \times 10^{-3}$	
Γ_{152}	$\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.}$		$(1.25 \pm 0.07) \times 10^{-3}$	
Γ_{153}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$		$(1.07 \pm 0.08) \times 10^{-3}$	
Γ_{154}	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$		$< 4.1 \times 10^{-6}$	CL=90%
Γ_{155}	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$		$< 1.80 \times 10^{-3}$	CL=90%
Γ_{156}	$\Xi^0 \Xi^0$		$(1.17 \pm 0.04) \times 10^{-3}$	
Γ_{157}	$\Xi(1530)^- \Xi^+ + \text{c.c.}$		$(3.18 \pm 0.08) \times 10^{-4}$	
Γ_{158}	$\Xi(1530)^0 \Xi^0$		$(3.2 \pm 1.4) \times 10^{-4}$	
Γ_{159}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	[c]	$< 1.1 \times 10^{-5}$	CL=90%
Γ_{160}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	[c]	$< 2.1 \times 10^{-5}$	CL=90%
Γ_{161}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	[c]	$< 1.6 \times 10^{-5}$	CL=90%
Γ_{162}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	[c]	$< 5.6 \times 10^{-5}$	CL=90%
Γ_{163}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	[c]	$< 1.1 \times 10^{-5}$	CL=90%

Decays into stable hadrons

Γ_{164}	$2(\pi^+ \pi^-) \pi^0$		$(4.2 \pm 0.4) \%$	S=2.1
Γ_{165}	$3(\pi^+ \pi^-) \pi^0$		$(2.9 \pm 0.6) \%$	
Γ_{166}	$\pi^+ \pi^- 3\pi^0$		$(1.9 \pm 0.9) \%$	
Γ_{167}	$\rho^\pm \pi^\mp \pi^0 \pi^0$		$(1.41 \pm 0.22) \%$	
Γ_{168}	$\rho^+ \rho^- \pi^0$		$(6.0 \pm 1.1) \times 10^{-3}$	
Γ_{169}	$\pi^+ \pi^- 4\pi^0$		$(6.5 \pm 1.3) \times 10^{-3}$	
Γ_{170}	$\pi^+ \pi^- \pi^0$		$(2.00 \pm 0.07) \%$	S=2.0
Γ_{171}	$2(\pi^+ \pi^- \pi^0)$		$(1.61 \pm 0.20) \%$	
Γ_{172}	$\pi^+ \pi^- \pi^0 K^+ K^-$		$(1.52 \pm 0.27) \%$	S=1.4
Γ_{173}	$\pi^+ \pi^-$		$(1.47 \pm 0.14) \times 10^{-4}$	
Γ_{174}	$2(\pi^+ \pi^-)$		$(3.20 \pm 0.25) \times 10^{-3}$	S=1.2
Γ_{175}	$3(\pi^+ \pi^-)$		$(4.3 \pm 0.4) \times 10^{-3}$	

Γ_{176}	$2(\pi^+\pi^-)3\pi^0$	$(6.2 \pm 0.9)\%$	
Γ_{177}	$4(\pi^+\pi^-)\pi^0$	$(9.0 \pm 3.0) \times 10^{-3}$	
Γ_{178}	$2(\pi^+\pi^-)\eta$	$(2.29 \pm 0.28) \times 10^{-3}$	
Γ_{179}	$3(\pi^+\pi^-)\eta$	$(7.2 \pm 1.5) \times 10^{-4}$	
Γ_{180}	$2(\pi^+\pi^-\pi^0)\eta$	$(1.6 \pm 0.5) \times 10^{-3}$	
Γ_{181}	$\pi^+\pi^-\pi^0\pi^0\eta$	$(2.4 \pm 0.5) \times 10^{-3}$	
Γ_{182}	$\rho^\pm\pi^\mp\pi^0\eta$	$(1.9 \pm 0.8) \times 10^{-3}$	
Γ_{183}	K^+K^-	$(3.06 \pm 0.05) \times 10^{-4}$	
Γ_{184}	$K_S^0K_L^0$	$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4
Γ_{185}	$K_S^0K_S^0$	$< 4.7 \times 10^{-9}$	CL=90%
Γ_{186}	$K\bar{K}\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{187}	$K^+K^-\pi^0$	$(2.88 \pm 0.12) \times 10^{-3}$	
Γ_{188}	$K_S^0K^\pm\pi^\mp$	$(5.3 \pm 0.5) \times 10^{-3}$	
Γ_{189}	$K_S^0K_L^0\pi^0$	$(2.06 \pm 0.26) \times 10^{-3}$	
Γ_{190}	$K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K_L^0\pi^0$	$(1.21 \pm 0.18) \times 10^{-3}$	
Γ_{191}	$K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K_L^0\pi^0$	$(4.3 \pm 1.3) \times 10^{-4}$	
Γ_{192}	$K^+K^-\pi^+\pi^-$	$(7.0 \pm 1.0) \times 10^{-3}$	
Γ_{193}	$K^+K^-\pi^0\pi^0$	$(2.13 \pm 0.22) \times 10^{-3}$	
Γ_{194}	$K^+K^-\pi^0\pi^0\pi^0$	$(1.61 \pm 0.29) \times 10^{-3}$	
Γ_{195}	$K_S^0K^\pm\pi^\mp\pi^0\pi^0$	$(5.3 \pm 0.7) \times 10^{-3}$	
Γ_{196}	$K_S^0K^\pm\pi^\mp\pi^+\pi^-$	$(6.3 \pm 0.4) \times 10^{-3}$	
Γ_{197}	$K_S^0K^\pm\rho(770)^\pm\pi^0$	$(2.9 \pm 0.8) \times 10^{-3}$	
Γ_{198}	$K_S^0K_L^0\pi^+\pi^-$	$(3.8 \pm 0.6) \times 10^{-3}$	
Γ_{199}	$K_S^0K_L^0\pi^0\pi^0$	$(1.9 \pm 0.4) \times 10^{-3}$	
Γ_{200}	$K_S^0K_L^0\eta$	$(1.45 \pm 0.33) \times 10^{-3}$	
Γ_{201}	$K_S^0K_S^0\pi^+\pi^-$	$(1.68 \pm 0.19) \times 10^{-3}$	
Γ_{202}	$K^\mp K_S^0\pi^\pm\pi^0$	$(5.7 \pm 0.5) \times 10^{-3}$	
Γ_{203}	$K_S^0K^\pm\pi^\mp\rho(770)^0$	$(3.1 \pm 0.5) \times 10^{-3}$	
Γ_{204}	$K^+K^-2(\pi^+\pi^-)$	$(3.1 \pm 1.3) \times 10^{-3}$	
Γ_{205}	$K^+K^-\pi^+\pi^-\eta$	$(4.7 \pm 0.7) \times 10^{-3}$	
Γ_{206}	$2(K^+K^-)$	$(7.2 \pm 0.8) \times 10^{-4}$	
Γ_{207}	$K^+K^-K_S^0K_S^0$	$(4.2 \pm 0.7) \times 10^{-4}$	
Γ_{208}	$K_S^0K^*(892)^0\pi^+\pi^-$	$(1.7 \pm 0.6) \times 10^{-3}$	
Γ_{209}	$K_S^0K^*(892)^0\pi^0\pi^0$	$(1.01 \pm 0.18) \times 10^{-3}$	
Γ_{210}	$K^\mp K^*(892)^\pm\pi^+\pi^-$	$(3.4 \pm 1.2) \times 10^{-3}$	
Γ_{211}	$K^*(892)^\pm K^*(892)^0\pi^\mp$	$(4.8 \pm 1.0) \times 10^{-3}$	
Γ_{212}	$K^\mp K^*(892)^\pm\pi^0\pi^0$	$(1.57 \pm 0.32) \times 10^{-3}$	
Γ_{213}	$K^*(892)^+K^*(892)^-\pi^0$	$(1.12 \pm 0.23)\%$	
Γ_{214}	$\rho\bar{\rho}$	$(2.121 \pm 0.029) \times 10^{-3}$	
Γ_{215}	$\rho\bar{\rho}\pi^0$	$(1.19 \pm 0.08) \times 10^{-3}$	S=1.1

Γ_{216}	$p\bar{p}\pi^+\pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{217}	$p\bar{p}\pi^+\pi^-\pi^0$	[d] $(2.3 \pm 0.9) \times 10^{-3}$	S=1.9
Γ_{218}	$p\bar{p}\eta$	$(1.513 \pm 0.023) \times 10^{-3}$	
Γ_{219}	$p\bar{p}\rho$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{220}	$p\bar{p}\omega$	$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3
Γ_{221}	$p\bar{p}\eta'(958)$	$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0
Γ_{222}	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$	$(6.8 \pm 1.8) \times 10^{-5}$	
Γ_{223}	$p\bar{p}\phi$	$(5.19 \pm 0.33) \times 10^{-5}$	
Γ_{224}	$p\bar{n}\pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$	
Γ_{225}	$n\bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$	
Γ_{226}	$n\bar{n}\pi^+\pi^-$	$(4 \pm 4) \times 10^{-3}$	
Γ_{227}	$nN(1440)$	seen	
Γ_{228}	$nN(1520)$	seen	
Γ_{229}	$nN(1535)$	seen	
Γ_{230}	$\Lambda\bar{\Lambda}$	$(1.88 \pm 0.08) \times 10^{-3}$	S=2.6
Γ_{231}	$\Lambda\bar{\Lambda}\pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$	
Γ_{232}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$	
Γ_{233}	$\Lambda\bar{\Lambda}\eta$	$(1.62 \pm 0.17) \times 10^{-4}$	
Γ_{234}	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	[b] $(1.26 \pm 0.05) \times 10^{-3}$	S=1.2
Γ_{235}	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$(1.21 \pm 0.07) \times 10^{-3}$	S=1.8
Γ_{236}	$pK^-\bar{\Lambda} + \text{c.c.}$	$(8.6 \pm 1.1) \times 10^{-4}$	
Γ_{237}	$pK^-\bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{238}	$pK_S^0\bar{\Sigma}^- + \text{c.c.}$	$(2.73 \pm 0.05) \times 10^{-4}$	
Γ_{239}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$	
Γ_{240}	$\Lambda\bar{\Sigma} + \text{c.c.}$	$(2.83 \pm 0.23) \times 10^{-5}$	
Γ_{241}	$\Sigma^+\bar{\Sigma}^-$	$(1.07 \pm 0.04) \times 10^{-3}$	
Γ_{242}	$\Sigma^0\bar{\Sigma}^0$	$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4
Γ_{243}	$\Sigma^+\bar{\Sigma}^-\eta$	$(6.3 \pm 0.4) \times 10^{-5}$	
Γ_{244}	$\Xi^-\bar{\Xi}^+$	$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4
Γ_{245}	$\Xi^0\bar{\Sigma}^-K^+ + \text{c.c.}$	$(5.64 \pm 0.33) \times 10^{-5}$	
Γ_{246}	$\Xi^0\bar{\Sigma}^0K_S^0 + \text{c.c.}$	$(2.2 \pm 0.4) \times 10^{-5}$	
Γ_{247}	$\Xi^0\bar{\Lambda}K_S^0 + \text{c.c.}$	$(3.76 \pm 0.27) \times 10^{-5}$	

Radiative decays

Γ_{248}	$\gamma\eta_c(1S)$	$(1.82 \pm 0.15) \%$	S=1.6
Γ_{249}	$\gamma\eta_c(1S) \rightarrow 3\gamma$	seen	
Γ_{250}	$\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta'$	seen	
Γ_{251}	3γ	$(1.16 \pm 0.22) \times 10^{-5}$	
Γ_{252}	4γ	$< 9 \times 10^{-6}$	CL=90%
Γ_{253}	5γ	$< 1.5 \times 10^{-5}$	CL=90%
Γ_{254}	$\gamma\pi^0$	$(3.39 \pm 0.08) \times 10^{-5}$	
Γ_{255}	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$	
Γ_{256}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
Γ_{257}	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$	

Γ ₂₅₈	$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$	
Γ ₂₅₉	$\gamma \pi^+ \pi^- 2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	
Γ ₂₆₀	$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$	
Γ ₂₆₁	$\gamma (K \bar{K} \pi)$ [$J^{PC} = 0^{-+}$]	$(7 \pm 4) \times 10^{-4}$	S=2.1
Γ ₂₆₂	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	
Γ ₂₆₃	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$	
Γ ₂₆₄	$\gamma \eta$	$(1.090 \pm 0.013) \times 10^{-3}$	
Γ ₂₆₅	$\gamma \eta \pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$	
Γ ₂₆₆	$\gamma f_0(500) \rightarrow \gamma \pi \pi$		
Γ ₂₆₇	$\gamma f_0(500) \rightarrow \gamma K \bar{K}$		
Γ ₂₆₈	$\gamma f_0(500) \rightarrow \gamma \eta \eta$		
Γ ₂₆₉	$\gamma a_0(980)^0 \rightarrow \gamma \eta \pi^0$	< 2.5	$\times 10^{-6}$ CL=95%
Γ ₂₇₀	$\gamma a_2(1320)^0 \rightarrow \gamma \eta \pi^0$	< 6.6	$\times 10^{-6}$ CL=95%
Γ ₂₇₁	$\gamma \eta \pi \pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ ₂₇₂	$\gamma \eta_2(1870) \rightarrow \gamma \eta \pi^+ \pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$	
Γ ₂₇₃	$\gamma \eta'(958)$	$(5.28 \pm 0.06) \times 10^{-3}$	S=1.3
Γ ₂₇₄	$\gamma f_0(980) \rightarrow \gamma \pi \pi$		
Γ ₂₇₅	$\gamma f_0(980) \rightarrow \gamma K \bar{K}$		
Γ ₂₇₆	$\gamma \rho \rho$	$(4.5 \pm 0.8) \times 10^{-3}$	
Γ ₂₇₇	$\gamma \rho \omega$	< 5.4	$\times 10^{-4}$ CL=90%
Γ ₂₇₈	$\gamma \rho \phi$	< 8.8	$\times 10^{-5}$ CL=90%
Γ ₂₇₉	$\gamma \omega \omega$	$(1.61 \pm 0.33) \times 10^{-3}$	
Γ ₂₈₀	$\gamma \phi \phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1
Γ ₂₈₁	$\gamma \eta(1405/1475) \rightarrow \gamma K \bar{K} \pi$	$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6
Γ ₂₈₂	$\gamma \eta(1405/1475) \rightarrow \gamma \gamma \rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8
Γ ₂₈₃	$\gamma \eta(1405/1475) \rightarrow \gamma \eta \pi^+ \pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$	
Γ ₂₈₄	$\gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3
Γ ₂₈₅	$\gamma \eta(1405/1475) \rightarrow \gamma \gamma \phi$	$(3.6 \pm 0.6) \times 10^{-6}$	
Γ ₂₈₆	$\gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma \pi^+ \pi^- \pi^0$	$(1.50 \pm 0.16) \times 10^{-5}$	
Γ ₂₈₇	$\gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma \pi^0 \pi^0 \pi^0$	$(6.64 \pm_{-0.23}^{+1.10}) \times 10^{-6}$	
Γ ₂₈₈	$\gamma \eta(1405) \rightarrow \gamma \gamma \gamma$	< 2.63	$\times 10^{-6}$ CL=90%
Γ ₂₈₉	$\gamma \eta(1475) \rightarrow \gamma \gamma \gamma$	< 1.86	$\times 10^{-6}$ CL=90%
Γ ₂₉₀	$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$	
Γ ₂₉₁	$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	$(1.98 \pm 0.33) \times 10^{-3}$	
Γ ₂₉₂	$\gamma \eta(1760) \rightarrow \gamma \gamma \gamma$	< 4.80	$\times 10^{-6}$ CL=90%
Γ ₂₉₃	$\gamma \eta(2225)$	$(3.14 \pm_{-0.19}^{+0.50}) \times 10^{-4}$	
Γ ₂₉₄	$\gamma f_2(1270)$	$(1.63 \pm 0.12) \times 10^{-3}$	S=1.3
Γ ₂₉₅	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm_{-0.22}^{+0.60}) \times 10^{-5}$	
Γ ₂₉₆	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$	

Γ_{297}	$\gamma f_1(1285) \rightarrow \gamma\gamma\phi$	$(2.9 \pm 1.1) \times 10^{-7}$	
Γ_{298}	$\gamma f_1(1285) \rightarrow \gamma\pi^0 f_0(980) \rightarrow$ $\gamma\pi^0\pi^0\pi^0$	$(5.6 \pm 0.9) \times 10^{-7}$	
Γ_{299}	$\gamma f_1(1285) \rightarrow \gamma a_0(980)^0\pi^0 \rightarrow$ $\gamma K_S^0 K_S^0\pi^0$	$(8.6 \pm 3.4) \times 10^{-6}$	
Γ_{300}	$\gamma f_0(1370) \rightarrow \gamma\pi\pi$		
Γ_{301}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$	
Γ_{302}	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$	
Γ_{303}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta$		
Γ_{304}	$\gamma f_0(1370) \rightarrow \gamma\eta\eta'$		
Γ_{305}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$	
Γ_{306}	$\gamma f_1(1420) \rightarrow \gamma K^*(892)^0 K_S^0 \rightarrow$ $\gamma K_S^0 K_S^0\pi^0$	$(7.2 \pm 0.7) \times 10^{-5}$	
Γ_{307}	$\gamma f_1(1420) \rightarrow \gamma a_0(980)^0\pi^0 \rightarrow$ $\gamma K_S^0 K_S^0\pi^0$	$(4.6 \pm 2.4) \times 10^{-6}$	
Γ_{308}	$\gamma f_1(1420) \rightarrow \gamma\gamma\phi$	$(5.5 \pm 1.9) \times 10^{-7}$	
Γ_{309}	$\gamma f_1(1420) \rightarrow \gamma\pi^0 f_0(980) \rightarrow$ $\gamma\pi^0\pi^0\pi^0$	$(2.23 \pm 0.26) \times 10^{-6}$	
Γ_{310}	$\gamma f_0(1500) \rightarrow \gamma\pi\pi$	$(1.09 \pm 0.24) \times 10^{-4}$	
Γ_{311}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta$	$(1.7 \pm 0.6) \times 10^{-5}$	
Γ_{312}	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm 0.24) \times 10^{-5}$	
Γ_{313}	$\gamma f_0(1500) \rightarrow \gamma\eta\eta'$		
Γ_{314}	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$	
Γ_{315}	$\gamma f_1(1510) \rightarrow \gamma\gamma\phi$	$(7.8 \pm 3.5) \times 10^{-7}$	
Γ_{316}	$\gamma f_1(1510) \rightarrow \gamma\pi^0 f_0(980) \rightarrow$ $\gamma\pi^0\pi^0\pi^0$		
Γ_{317}	$\gamma f_2'(1525)$	$(5.7 \pm 0.8) \times 10^{-4}$	S=1.5
Γ_{318}	$\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm 0.7) \times 10^{-5}$	
Γ_{319}	$\gamma f_2'(1525) \rightarrow$ $\gamma K^*(892)^0 K_S^0 \rightarrow$ $\gamma K_S^0 K_S^0\pi^0$	$(9.5 \pm 1.6) \times 10^{-6}$	
Γ_{320}	$\gamma f_2'(1525) \rightarrow \gamma\eta\eta$	$(3.4 \pm 1.4) \times 10^{-5}$	
Γ_{321}	$\gamma f_2(1565) \rightarrow \gamma\eta\eta'$		
Γ_{322}	$\gamma f_2'(1525) \rightarrow \gamma\gamma\phi$	$(2.8 \pm 0.9) \times 10^{-6}$	
Γ_{323}	$\gamma f_2(1640) \rightarrow \gamma\omega\omega$	$(2.8 \pm 1.8) \times 10^{-4}$	
Γ_{324}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$	
Γ_{325}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(9.5 \pm 0.9) \times 10^{-4}$	S=1.4

Γ_{326}	$\gamma f_0(1710) \rightarrow \gamma \omega \omega$	$(3.1 \pm 1.0) \times 10^{-4}$	
Γ_{327}	$\gamma f_0(1710) \rightarrow \gamma \eta \eta$	$(2.4 \begin{smallmatrix} + 1.2 \\ - 0.7 \end{smallmatrix}) \times 10^{-4}$	
Γ_{328}	$\gamma f_0(1710) \rightarrow \gamma \eta \eta'$		
Γ_{329}	$\gamma f_0(1710) \rightarrow \gamma \omega \phi$	$(2.5 \pm 0.6) \times 10^{-4}$	
Γ_{330}	$\gamma f_0(1810) \rightarrow \gamma \pi \pi$		
Γ_{331}	$\gamma f_0(1810) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \begin{smallmatrix} + 0.20 \\ - 0.33 \end{smallmatrix}) \times 10^{-5}$	
Γ_{332}	$\gamma f_0(1810) \rightarrow \gamma K \bar{K}$		
Γ_{333}	$\gamma f_0(1810) \rightarrow \gamma \eta \eta$		
Γ_{334}	$\gamma f_0(1810) \rightarrow \gamma \eta \eta'$		
Γ_{335}	$\gamma f_0(1810) \rightarrow \gamma \omega \phi$		
Γ_{336}	$\gamma f_2(1810) \rightarrow \gamma \eta \eta$	$(5.4 \begin{smallmatrix} + 3.5 \\ - 2.4 \end{smallmatrix}) \times 10^{-5}$	
Γ_{337}	$\gamma \eta_1(1855) \rightarrow \gamma \eta \eta'$	$(2.7 \begin{smallmatrix} + 0.4 \\ - 0.5 \end{smallmatrix}) \times 10^{-6}$	
Γ_{338}	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	$(2.0 \pm 1.4) \times 10^{-4}$	
Γ_{339}	$\gamma f_2(1950) \rightarrow$ $\gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$	
Γ_{340}	$\gamma f_2(1950) \rightarrow \gamma \gamma \phi$	$(10.0 \begin{smallmatrix} + 3.5 \\ - 2.2 \end{smallmatrix}) \times 10^{-6}$	
Γ_{341}	$\gamma f_2(2010) \rightarrow \gamma \eta \eta'$		
Γ_{342}	$\gamma f_2(2010) \rightarrow \gamma \gamma \phi$	$(4.6 \pm 1.5) \times 10^{-6}$	
Γ_{343}	$\gamma f_0(2020) \rightarrow \gamma \pi \pi$		
Γ_{344}	$\gamma f_0(2020) \rightarrow \gamma K \bar{K}$		
Γ_{345}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta$		
Γ_{346}	$\gamma f_0(2020) \rightarrow \gamma \eta' \eta'$	$(2.63 \begin{smallmatrix} + 0.32 \\ - 0.50 \end{smallmatrix}) \times 10^{-4}$	
Γ_{347}	$\gamma f_0(2020) \rightarrow \gamma \eta \eta'$		
Γ_{348}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$	
Γ_{349}	$\gamma f_4(2050) \rightarrow \gamma \eta \eta'$		
Γ_{350}	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \begin{smallmatrix} + 0.60 \\ - 0.30 \end{smallmatrix}) \times 10^{-4}$	
Γ_{351}	$\gamma f_0(2100) \rightarrow \gamma K \bar{K}$		
Γ_{352}	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$	
Γ_{353}	$\gamma f_0(2200)$	seen	
Γ_{354}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$	
Γ_{355}	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \begin{smallmatrix} + 0.19 \\ - 0.50 \end{smallmatrix}) \times 10^{-4}$	
Γ_{356}	$\gamma f_0(2200) \rightarrow \gamma \pi \pi$		
Γ_{357}	$\gamma f_0(2200) \rightarrow \gamma \eta \eta$		
Γ_{358}	$\gamma f_0(2200) \rightarrow \gamma \gamma \phi$	$(2.0 \begin{smallmatrix} + 0.6 \\ - 0.8 \end{smallmatrix}) \times 10^{-7}$	
Γ_{359}	$\gamma f_J(2220)$	seen	
Γ_{360}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 3.9 \times 10^{-5}$	CL=90%
Γ_{361}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 4.1 \times 10^{-5}$	CL=90%
Γ_{362}	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$	
Γ_{363}	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$	

Γ_{364}	$\gamma f_0(2330) \rightarrow \gamma \pi \pi$		
Γ_{365}	$\gamma f_0(2330) \rightarrow \gamma \eta \eta$		
Γ_{366}	$\gamma f_0(2330) \rightarrow \gamma \eta' \eta'$	$(6.1 \pm 4.0) \times 10^{-6}$	
Γ_{367}	$\gamma f_0(2330) \rightarrow \gamma \eta \eta'$		
Γ_{368}	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \pm 2.4) \times 10^{-5}$	
Γ_{369}	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \pm 4.0) \times 10^{-5}$	
Γ_{370}	$\gamma f_2(2340) \rightarrow \gamma \eta' \eta'$	$(8.7 \pm 0.9) \times 10^{-6}$	
Γ_{371}	$\gamma f_0(2470) \rightarrow \gamma \eta' \eta'$	$(8.2 \pm 4.0) \times 10^{-7}$	
Γ_{372}	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.7 \pm 0.6) \times 10^{-4}$	S=1.6
Γ_{373}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(7.7 \pm 1.5) \times 10^{-5}$	
Γ_{374}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm 2.0) \times 10^{-5}$	
Γ_{375}	$\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)$	$(3.4 \pm 0.8) \times 10^{-6}$	
Γ_{376}	$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	< 3.56	$\times 10^{-6}$ CL=90%
Γ_{377}	$\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \pm 0.7) \times 10^{-5}$	
Γ_{378}	$\gamma \eta(2370) \rightarrow \gamma K^+ K^- \eta'$	$(1.8 \pm 0.7) \times 10^{-5}$	
Γ_{379}	$\gamma \eta(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	$(1.2 \pm 0.5) \times 10^{-5}$	
Γ_{380}	$\gamma \eta(2370) \rightarrow \gamma \eta \eta \eta'$	< 9.2	$\times 10^{-6}$ CL=90%
Γ_{381}	$\gamma D^0 + c.c.$	< 9.1	$\times 10^{-8}$ CL=90%
Γ_{382}	$\gamma p \bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$	
Γ_{383}	$\gamma p \bar{p} \pi^+ \pi^-$	< 7.9	$\times 10^{-4}$ CL=90%
Γ_{384}	$\gamma \Lambda \bar{\Lambda}$	< 1.3	$\times 10^{-4}$ CL=90%
Γ_{385}	$\gamma A^0 \rightarrow \gamma \text{invisible}$	[e] < 1.7	$\times 10^{-6}$ CL=90%
Γ_{386}	$\gamma A^0 \rightarrow \gamma \gamma \gamma$	< 4.9	$\times 10^{-7}$ CL=95%
Γ_{387}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	[f] < 7.8	$\times 10^{-7}$ CL=90%

Dalitz decays

Γ_{388}	$\pi^0 e^+ e^-$	$(8.1 \pm 0.5) \times 10^{-7}$	
Γ_{389}	$\eta e^+ e^-$	$(1.42 \pm 0.08) \times 10^{-5}$	
Γ_{390}	$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$	
Γ_{391}	$\eta(1405) e^+ e^- \rightarrow$ $f_0(980) \pi^0 e^+ e^- \rightarrow$ $\pi^+ \pi^- \pi^0 e^+ e^-$	$(2.04 \pm 0.22) \times 10^{-7}$	
Γ_{392}	$X(1835) e^+ e^-$, $X \rightarrow \pi^+ \pi^- \eta'$	$(3.58 \pm 0.25) \times 10^{-6}$	
Γ_{393}	$X(2120) e^+ e^-$, $X \rightarrow \pi^+ \pi^- \eta'$	$(8.2 \pm 1.3) \times 10^{-7}$	
Γ_{394}	$\eta(2370) e^+ e^-$, $\eta \rightarrow \pi^+ \pi^- \eta'$	$(1.08 \pm 0.17) \times 10^{-6}$	
Γ_{395}	$\eta U \rightarrow \eta e^+ e^-$	[g] < 9.11	$\times 10^{-7}$ CL=90%
Γ_{396}	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	[g] < 2.0	$\times 10^{-7}$ CL=90%
Γ_{397}	$\phi e^+ e^-$	< 1.2	$\times 10^{-7}$ CL=90%

Weak decays

Γ_{398}	$D^- e^+ \nu_e + \text{c.c.}$	< 7.1	$\times 10^{-8}$	CL=90%
Γ_{399}	$D^- \mu^+ \nu_\mu + \text{c.c.}$	< 5.6	$\times 10^{-7}$	CL=90%
Γ_{400}	$\bar{D}^0 e^+ e^- + \text{c.c.}$	< 8.5	$\times 10^{-8}$	CL=90%
Γ_{401}	$D^0 \mu^+ \mu^- + \text{c.c.}$	< 1.1	$\times 10^{-7}$	CL=90%
Γ_{402}	$D_s^- e^+ \nu_e + \text{c.c.}$	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{403}	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{404}	$D^- \pi^+ + \text{c.c.}$	< 7.0	$\times 10^{-8}$	CL=90%
Γ_{405}	$D^- \rho^+ + \text{c.c.}$	< 6.0	$\times 10^{-7}$	CL=90%
Γ_{406}	$\bar{D}^0 \pi^0 + \text{c.c.}$	< 4.7	$\times 10^{-7}$	CL=90%
Γ_{407}	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	< 1.7	$\times 10^{-4}$	CL=90%
Γ_{408}	$\bar{D}^0 \bar{K}^{*0} + \text{c.c.}$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{409}	$\bar{D}^0 \eta + \text{c.c.}$	< 6.8	$\times 10^{-7}$	CL=90%
Γ_{410}	$\bar{D}^0 \rho^0 + \text{c.c.}$	< 5.2	$\times 10^{-7}$	CL=90%
Γ_{411}	$D_s^- \pi^+ + \text{c.c.}$	< 4.1	$\times 10^{-7}$	CL=90%
Γ_{412}	$D_s^- \rho^+ + \text{c.c.}$	< 8.0	$\times 10^{-7}$	CL=90%

Charge conjugation (C), Parity (P), Lepton Family number (LF) violating modes

Γ_{413}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$	CL=90%
Γ_{414}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{415}	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$	CL=90%
Γ_{416}	$e^\pm \tau^\mp$	LF	< 7.5	$\times 10^{-8}$	CL=90%
Γ_{417}	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{418}	$K^+ K^+ e^- e^- + \text{c.c.}$	L	< 2.1	$\times 10^{-9}$	CL=90%
Γ_{419}	$p e^- + \text{c.c.}$	L,B	< 3.1	$\times 10^{-8}$	CL=90%
Γ_{420}	$\Lambda_c^+ e^- + \text{c.c.}$		< 6.9	$\times 10^{-8}$	CL=90%

Other decays

Γ_{421}	invisible		< 7	$\times 10^{-4}$	CL=90%
Γ_{422}	$\mu^+ \mu^- X^0 \rightarrow \mu^+ \mu^- + \text{invisible}$				
Γ_{423}	$\phi + \text{invisible } X$				

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] $\Theta(1540)$ is a hypothetical pentaquark state of $1.54 \text{ GeV}/c^2$ mass and a width of less than $25 \text{ MeV}/c^2$.

[d] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta, p\bar{p}\omega, p\bar{p}\eta'$.

[e] For a narrow state A with mass less than 960 MeV.

[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

[g] For a dark photon U with mass between 100 and 2100 MeV.

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 115 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 215.4$ for 96 degrees of freedom.

$J/\psi(1S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

Γ_1

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
81.37 ± 1.36 ± 1.30	¹ ANASHIN	20	KEDR e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
74.1 ± 8.1	BAI	95B	BES e^+e^-
59 ± 24	BALDINI-...	75	FRAG e^+e^-
59 ± 14	BOYARSKI	75	MRK1 e^+e^-
50 ± 25	ESPOSITO	75B	FRAM e^+e^-

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there.

$\Gamma(e^+e^-)$

Γ_5

<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.53 ± 0.10 OUR AVERAGE				
5.550 ± 0.056 ± 0.089		^{1,2} ANASHIN	18A	KEDR e^+e^-
5.36 ^{+0.29} _{-0.28}		³ HSUEH	92	RVUE See Υ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.58 ± 0.05 ± 0.08		⁴ ABLIKIM	16Q	BES3 $3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.71 ± 0.16	13k	⁵ ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57 ± 0.19	7.8k	⁵ AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14 ± 0.39		BAI	95B	BES e^+e^-
4.72 ± 0.35		ALEXANDER	89	RVUE See Υ mini-review
4.4 ± 0.6		³ BRANDELIK	79C	DASP e^+e^-
4.6 ± 0.8		⁶ BALDINI-...	75	FRAG e^+e^-
4.8 ± 0.6		BOYARSKI	75	MRK1 e^+e^-
4.6 ± 1.0		ESPOSITO	75B	FRAM e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow \text{hadrons}$ near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

⁴ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.

⁵ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

⁶ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

$\Gamma(\mu^+\mu^-)$

Γ_7

<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.13 ± 0.52	BAI	95B	BES e^+e^-

4.8 ± 0.6	BOYARSKI	75	MRK1	e^+e^-
5 ± 1	ESPOSITO	75B	FRAM	e^+e^-

$\Gamma(\gamma\gamma)$

Γ_{413}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C	DASP e^+e^-

$J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(l) in the e^+e^- annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_1\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.884 ± 0.048 ± 0.078	^{1,2} ANASHIN	18A	KEDR e^+e^-
4 ± 0.8	³ BALDINI-...	75	FRAG e^+e^-
3.9 ± 0.8	³ ESPOSITO	75B	FRAM e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow \text{hadrons}$ near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_5\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
333.1 ± 6.6 ± 4.0	^{1,2} ANASHIN	18A	KEDR e^+e^-
332.3 ± 6.4 ± 4.8	ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow e^+e^-$
350 ± 20	BRANDELIK	79C	DASP e^+e^-
320 ± 70	³ BALDINI-...	75	FRAG e^+e^-
340 ± 90	³ ESPOSITO	75B	FRAM e^+e^-
360 ± 100	³ FORD	75	SPEC e^+e^-

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow \text{hadrons}$ near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_7\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
333 ± 4 OUR AVERAGE				
333.4 ± 2.5 ± 4.4		ABLIKIM	16Q	BES3 $3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$
331.8 ± 5.2 ± 6.3		ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow \mu^+\mu^-$
338.4 ± 5.8 ± 7.1	13k	ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 90	DASP	75	DASP	e^+e^-
380 ± 50	¹ ESPOSITO	75B	FRAM	e^+e^-

¹ Data redundant with branching ratios or partial widths above.

$$\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.3 ± 0.4 OUR AVERAGE

2.34 ± 0.43 ± 0.16	49	LEES	18	BABR $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$
2.22 ± 0.96 ± 0.02	9	¹ AUBERT	07AU	BABR 10.6 $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 0.51 \pm 0.22 \pm 0.03$ eV which we divide by our best (shown rounded) value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

64.8 ± 11.1 ± 0.4 200 ¹LEES 21c BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

¹LEES 21c reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 21.1 \pm 1.7 \pm 3.2$ eV which we divide by our best (shown rounded) value $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\eta\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{19}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

26.9 ± 5.7 ± 0.1 101 ¹LEES 21c BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-3\pi^0\gamma\gamma)$

¹LEES 21c reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.6 \pm 1.6 \pm 1.6$ eV which we divide by our best (shown rounded) value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\eta K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{22}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

4.76 ± 1.64 ± 0.03 ¹LEES 23 BABR $e^+e^- \rightarrow \gamma_{ISR}$ hadrons

¹LEES 23 reports $[\Gamma(J/\psi(1S) \rightarrow \eta K^+K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 1.55 \pm 0.51 \pm 0.16$ eV which we divide by our best (shown rounded) value $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{23}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

7.3 ± 1.4 ± 0.4 44 LEES 17D BABR $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$$\Gamma(\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{26}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

155 ± 26 ± 36 14k LEES 21 BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(\rho^+ \rho^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{27}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

32 ± 13 ± 15 14k LEES 21 BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(\rho^\mp K^\pm K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{29} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.4±1.0±1.9	130	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$$\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{44} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
53.6±5.0±0.4	788	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\omega \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{45} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.8±3.5±0.2	398	¹ LEES	18E BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 24.8 \pm 1.8 \pm 2.5$ eV which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\omega 3\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{46} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±3.1±0.1	89	¹ LEES	21C BABR	$e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 4\pi^0)$

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 9.4 \pm 2.3 \pm 1.5$ eV which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\omega \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{48} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
16.9±7.6±0.2	¹ LEES	21C BABR	$e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 4\pi^0)$

¹ Different final state as in AUBERT 06. LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 4.9 \pm 2.1 \pm 0.7$ eV which we divide by our best (shown rounded) values $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$, $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) values.

$$\Gamma(\omega \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{49} \Gamma_5 / \Gamma$$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.3±0.2	170	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \pi^0 \gamma$

$$\Gamma(\omega \pi^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{50} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.96±0.01	27	¹ LEES	18E BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3$ eV which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\omega\pi^+\pi^-2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{52}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
185±30±1	14k	¹ LEES	21 BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-2\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 165 \pm 9 \pm 25$ eV which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\omega K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{64}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.70±1.98±0.03	24	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ωK ⁺ K ⁻ γ

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 3.3 \pm 1.3 \pm 1.2$ eV which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(K^+K^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{73}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
29.0±1.7±1.3	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁺ K [*] (892) ⁻ γ

$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{74}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10.96±0.85±0.70	155	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁰ γ

$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{75}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.76±1.70±1.00	89	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ

$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{76}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
26.6±2.5±1.5	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁰ $\bar{K}^*(892)^0$ γ

$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0K^\pm\pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{77}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
17.70±1.70±1.00	94	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ

$\Gamma(\bar{K}^*(892)^0K^+\pi^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{78}\Gamma_5/\Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
42.6±4.8±7.2	99	¹ LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ

¹ Dividing by 1/6 to account for $B(K^*(892)^0 \rightarrow K_S^0\pi^0)=1/6$.

$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{79}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
22.8±2.8±6.8	80	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 1/4 to account for $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$.

 $\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{80}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.0±2.8 OUR AVERAGE				
9.2±1.2±3.2	64	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
14.8±4.8±1.2	53	² LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/2 to take into account $B(K^*(892)^\pm \rightarrow K^\pm \pi^\mp) = 1/2$.

² Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

 $\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{81}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.7±1.2±0.3	53	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

 $\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{84}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.60±0.75±2.25	34	¹ LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

¹ Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$.

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{86}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.28±0.34±0.07	47±12	¹ LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.28±0.40±0.11 25±8 ^{1,2} AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$ $B(K^{*0} \rightarrow K \pi)$.

² Superseded by LEES 12F.

 $\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{87}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80±0.48±0.32	1±5	¹ LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by $(1/4)^2$ to take twice into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

 $\Gamma(K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{98}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.1±9.8±0.5	35	^{1,2} LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4$ $B(K^*(1430) \rightarrow K \pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8$ eV which we divide by our best (shown rounded) value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{99} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

25.8 ± 1.4 ± 0.6	710	1,2,3 LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
-------------------------	-----	------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1	317	2,4 AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
------------	-----	------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41$ eV which we divide by our best (shown rounded) value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

² Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4$ eV which we divide by our best (shown rounded) value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{100} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

18.6 ± 16.1 ± 0.4	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$
--------------------------	-------	----------	----------	--

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K\pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best (shown rounded) value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{101} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.32 ± 2.00 ± 0.08	8 ± 8	¹ LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$
---------------------------	-------	-------------------	----------	--

¹ Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

$\Gamma(\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{103} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.8 ± 0.4 ± 0.3	110 ± 14	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
------------------------	----------	---------------------	-----------	---

¹ Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{110} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.39 ± 0.34 OUR AVERAGE

4.37 ± 0.48 ± 0.05	181	¹ LEES	12F BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
--------------------	-----	-------------------	----------	---

$$4.42 \pm 0.47 \pm 0.05 \quad 254 \pm 23 \quad {}^2 \text{ SHEN} \quad 09 \text{ BELL} \quad 10.6 \text{ e}^+ \text{e}^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$5.2 \pm 0.7 \pm 0.1 \quad 103 \quad {}^3 \text{ AUBERT, BE 06D BABR} \quad 10.6 \text{ e}^+ \text{e}^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$$

¹LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

²SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

³Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{111} \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.71 ± 0.56 ± 0.03	45	¹ LEES	12F BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁰ π ⁰ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$3.07 \pm 0.86 \pm 0.03 \quad 23 \quad {}^2 \text{ AUBERT, BE 06D BABR} \quad 10.6 \text{ e}^+ \text{e}^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$$

¹LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

²Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\phi 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{112} \Gamma_5 / \Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.94 ± 0.19 ± 0.01	35	¹ AUBERT	06D BABR	10.6 e ⁺ e ⁻ → φ2(π ⁺ π ⁻)γ

¹AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+ \pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{113}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.6±1.4 OUR AVERAGE

4.1±1.6±0.4		¹ LEES	23 BABR	$e^+e^- \rightarrow \gamma_{ISR} \text{hadrons}$
6.1±2.7±0.4	6	² AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$

¹ LEES 23 quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi^0) = 0.64 \pm 0.26 \pm 0.06$ eV.

² AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.

$$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{117}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.42±0.18 OUR AVERAGE

1.38±0.24±0.02	57 ± 9	¹ LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
1.48±0.27±0.09	60 ± 11	² SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.00±0.23±0.01	20 ± 5	³ AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
----------------	--------	---------------------	-----------	--

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)] / \Gamma_{\text{total}} \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.69 \pm 0.11 \pm 0.05$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

² Multiplied by 2/3 to take into account the $\phi\pi^+\pi^-$ mode only. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.

³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)] / \Gamma_{\text{total}} \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \qquad \Gamma_{118}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.96±0.26±0.01	16 ± 4	¹ LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$
-----------------------	--------	-------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.94±0.39±0.01	7.0 ± 2.8	² AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$
----------------	-----------	---------------------	-----------	--

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)] / \Gamma_{\text{total}} \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.48 \pm 0.12 \pm 0.05$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)] / \Gamma_{\text{total}} \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.47 \pm 0.19 \pm 0.05$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{126} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$1.79 \pm 0.32^{+0.02}_{-0.06}$	61	1,2,3 LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
---------------------------------	----	------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.08 \pm 0.73^{+0.05}_{-0.14}$	44	2,4 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
---------------------------------	----	------------	-----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 1.51 \pm 0.25 \pm 0.10$ eV which we divide by our best (shown rounded) value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

² Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

³ Using $\pi^+ \pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28$ eV which we divide by our best (shown rounded) value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{131} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$8.1 \pm 3.2 \pm 0.2$	11	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$
-----------------------	----	----------	----------	--

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K \bar{K})$ and using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = 7.2 \pm 2.8 \pm 0.3$ eV which we divide by our best (shown rounded) value $B(f'_2(1525) \rightarrow K \bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{137} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$4.51 \pm 0.61 \pm 0.05$	163	¹ LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$
--------------------------	-----	-------------------	----------	---

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{138} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$3.19 \pm 0.82 \pm 0.04$	29	¹ LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$
--------------------------	----	-------------------	----------	--

¹ LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1$ eV which we divide by our best (shown rounded) value $B(\phi(1020) \rightarrow K^+ K^-) = (50.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(f'_2(1525)K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{143}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_S^0 K^+K^-\gamma$

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow f'_2(1525)K^+K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20$ eV which we divide by our best (shown rounded) value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{164}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
303±5±18	4990	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{166}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
100 ±50 OUR AVERAGE		Error includes scale factor of 4.3.		

55 ±16 ±1	14k	¹ LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
150.0±4.0±15.0	2.3k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] = 19.2 \pm 4.5 \pm 3.2$ eV which we divide by our best (shown rounded) value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}} = 0.3478 \pm 0.0033$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{169}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.8±4.4±5.4	340	LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-\pi^0)$

$\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{167}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.0±9.0±8.0	1.2k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

$\Gamma(\rho^+\rho^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{168}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.0±5.0±3.3	529	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{170}\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.1248±0.0019±0.0026	LEES	21B BABR	$10.5 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.122 ±0.005 ±0.008	AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
---------------------	----------	----------	---

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{171}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9±0.5±1.0	761	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

$$\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{172}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0±4.3±6.4	768	AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{174}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4±0.9±0.4		LEES	12E BABR	10.6 $e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5±1.4±1.3	270	¹ AUBERT	05D BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$
--------------	-----	---------------------	----------	---

¹ Superseded by LEES 12E.

$$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{175}\Gamma_5/\Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.16±0.14	496	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$

$$\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{176}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
345±10±50	14k	LEES	21 BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

$$\Gamma(2(\pi^+\pi^-\eta)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{178}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\eta)\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-\eta)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$ eV which we divide by our best (shown rounded) value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(2(\pi^+\pi^-\pi^0\eta)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{180}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±2.6±1.4	14k	LEES	21 BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0\eta)\gamma$

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{181}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1± 2.7 OUR AVERAGE				

26.1±17.9±0.3	14k	¹ LEES	21 BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0\eta)\gamma$
12.8± 1.8±2.0	203	LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 6 \pm 4 \pm 1$ eV which we divide by our best (shown rounded) value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{182}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±4.1±1.6	168	LEES	18E BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{183}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.78±0.11±0.05	462	¹ LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.94 \pm 0.11 \pm 0.05$	462	² LEES	15J	BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
$1.42 \pm 0.23 \pm 0.08$	51	³ LEES	13Q	BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
¹ $\sin\phi > 0$.					
² $\sin\phi < 0$.					
³ Interference with non-resonant $K^+ K^-$ production not taken into account.					

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{189} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$11.4 \pm 1.3 \pm 0.6$	182	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{190} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$6.7 \pm 0.9 \pm 0.4$	106	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{191} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 0.7 \pm 0.1$	37	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

$$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{192} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$37.94 \pm 0.81 \pm 1.10$	3.1k	LEES	12F	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$36.3 \pm 1.3 \pm 2.1$ 1.5k ¹ AUBERT 07AK BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

$33.6 \pm 2.7 \pm 2.7$ 233 ² AUBERT 05D BABR $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{193} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$11.75 \pm 0.81 \pm 0.90$	388	LEES	12F	BABR $10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$13.6 \pm 1.1 \pm 1.3$ 203 ¹ AUBERT 07AK BABR $10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

¹ Superseded by LEES 12F.

$$\Gamma(K^+ K^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{194} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$8.9 \pm 1.3 \pm 0.9$	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{195} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$29.3 \pm 2.6 \pm 2.9$	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{196} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$34.6 \pm 1.4 \pm 1.8$	LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$

$\Gamma(K_S^0 K^\pm \rho(770)^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{197}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
16.0±4.1±1.6		LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR} \text{hadrons}$

$\Gamma(K_S^0 K_L^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{198}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
20.8±2.3±2.1	248	LEES	14H	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$

$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{199}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10.3±2.3±0.5	47	LEES	17A	BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$

$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{200}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
8.0±1.8±0.4	45	LEES	17A	BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{201}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
9.3±0.9±0.5	133	LEES	14H	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

$\Gamma(K^\mp K_S^0 \pi^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{202}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
31.7±1.9±1.8	393	LEES	17D	BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

$\Gamma(K_S^0 K^\pm \pi^\mp \rho(770)^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{203}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
17.3±2.1±1.7		LEES	23	BABR	$e^+ e^- \rightarrow \gamma_{ISR} \text{hadrons}$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{204}\Gamma_5/\Gamma$
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.75±0.23±0.17	205	AUBERT	06D	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{205}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
25.9±3.9±0.1	73	¹ AUBERT	07AU	BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

¹AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8$ eV which we divide by our best (shown rounded) value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(2(K^+ K^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{206}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.00±0.33±0.29	287 ± 24	LEES	12F	BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.11±0.39±0.30	156 ± 15	¹ AUBERT	07AK	BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
4.0 ± 0.7 ± 0.6	38	² AUBERT	05D	BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

$\Gamma(K^+ K^- K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{207}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.3±0.4±0.1	29	LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$	

$\Gamma(K_S^0 K^*(892)^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{208}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
9.45±3.15±0.90		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K_S^0 K^*(892)^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{209}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
5.59±0.79±0.55		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K^\mp K^*(892)^\pm \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{210}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
18.6±6.3±1.8		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K^*(892)^\pm K^*(892)^0 \pi^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{211}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
26.6±4.5±2.7		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K^\mp K^*(892)^\pm \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{212}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
8.67±1.56±0.84		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(K^*(892)^+ K^*(892)^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{213}\Gamma_5/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
62.1±10.8±6.30		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	

$\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{214}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
11.9±0.6 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.			

11.3±0.4±0.3 821 ¹ LEES 130 BABR $e^+ e^- \rightarrow p\bar{p}\gamma$

12.9±0.4±0.4 918 ² LEES 13Y BABR $e^+ e^- \rightarrow p\bar{p}\gamma$

9.7±1.7 ³ ARMSTRONG 93B E760 $\bar{p}p \rightarrow e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.0±0.6±0.5 438 ⁴ AUBERT 06B BABR $e^+ e^- \rightarrow p\bar{p}\gamma$


¹ ISR photon reconstructed in the detector

² ISR photon undetected

³ Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.

⁴ Superseded by LEES 130

WEIGHTED AVERAGE
 11.9 ± 0.6 (Error scaled by 1.8)



$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{230}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$10.7 \pm 0.9 \pm 0.7$	AUBERT	07BD BABR	$10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma(\Sigma^+\bar{\Sigma}^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{241}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$6.8 \pm 1.5 \pm 0.8$	GONG	23 BELL	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{242}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$5.2 \pm 1.5 \pm 0.6$	GONG	23 BELL	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
$6.4 \pm 1.2 \pm 0.6$	AUBERT	07BD BABR	$10.6 e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$

$J/\psi(1S)$ BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ above.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.877 ± 0.005 OUR AVERAGE			
0.878 ± 0.005	BAI	95B BES	e^+e^-
0.86 ± 0.02	BOYARSKI	75 MRK1	e^+e^-

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$ **Γ_2/Γ**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.1346 ± 0.0007	¹ LIAO	23	RVUE e^+e^-
0.135 ± 0.003	^{2,3} SETH	04	RVUE e^+e^-
0.17 ± 0.02	² BOYARSKI	75	MRK1 e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (5.967 \pm 0.023)\%$ and $R = 2.26 \pm 0.01$ determined by a fit to data from Mark-I, DM2, BESII, KEDR, and BESIII.

² Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

³ Using $B(J/\psi \rightarrow \ell^+ \ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. Superseded by LIAO 23.

$\Gamma(g g g)/\Gamma_{\text{total}}$ **Γ_3/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
64.1 ± 1.0	6 M	¹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma g g)/\Gamma(g g g) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+ \ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\gamma \eta_c)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ **Γ_4/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.79 ± 1.05	200 k	¹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- \gamma + \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma g g)/\Gamma(g g g) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(g g g)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ measurement of BESSON 08.

$\Gamma(\gamma g g)/\Gamma(g g g)$ **Γ_4/Γ_3**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
13.7 ± 0.1 ± 0.7	6 M	BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ **Γ_5/Γ**

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.971 ± 0.032 OUR AVERAGE				
5.983 ± 0.007 ± 0.037	720k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945 ± 0.067 ± 0.042	15k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ± 0.05 ± 0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ± 0.33		BAI	95B	BES $e^+ e^-$
5.92 ± 0.15 ± 0.20		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75	MRK1 $e^+ e^-$

$\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$ **Γ_6/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
8.8 ± 1.3 ± 0.4	¹ ARMSTRONG	96	E760 $\bar{p} p \rightarrow e^+ e^- \gamma$

¹ For $E_\gamma > 100$ MeV.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.961±0.033 OUR AVERAGE				
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960±0.065±0.050	17k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ±0.06 ±0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ±0.33		BAI	95B	BES $e^+ e^-$
5.90 ±0.15 ±0.19		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1 $e^+ e^-$

 $\Gamma(e^+e^-)/\Gamma(\mu^+\mu^-)$ Γ_5/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
1.0016±0.0031 OUR AVERAGE			
1.0022±0.0044±0.0048	¹ AULCHENKO 14	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017±0.0017±0.0033	² ABLIKIM 13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 ±0.021 ±0.013	³ ANASHIN 10	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 ±0.012 ±0.006	LI 05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.011 ±0.013 ±0.016	BAI 98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 ±0.07	BAI 95B	BES	$e^+ e^-$
1.00 ±0.05	BOYARSKI 75	MRK1	$e^+ e^-$
0.91 ±0.15	ESPOSITO 75B	FRAM	$e^+ e^-$
0.93 ±0.10	FORD 75	SPEC	$e^+ e^-$

¹ From 235.3k $J/\psi \rightarrow e^+ e^-$ and 156.6k $J/\psi \rightarrow \mu^+ \mu^-$ observed events.

² Not independent of the corresponding measurements of $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$.

³ Not independent of the corresponding measurements of $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$.

 $\Gamma(e^+e^-e^+e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.48±0.31±0.45	700	¹ ABLIKIM 24L	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

¹ $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

 $\Gamma(e^+e^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.53±0.22±0.13	354	¹ ABLIKIM 24L	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

¹ $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.

 $\Gamma(\mu^+\mu^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
11.3±1.1±0.1		452	¹ AAIJ 24AE	LHCB	$J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$
<16	90	3.4	² ABLIKIM 24L	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
10.1 ^{+3.3} _{-2.8} ±0.1		12	³ HAYRAPETY...24A	CMS	$J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$

- ¹ AAIJ 24AE reports $[\Gamma(J/\psi(1S) \rightarrow \mu^+ \mu^- \mu^+ \mu^-) / \Gamma_{\text{total}}] / [B(J/\psi(1S) \rightarrow \mu^+ \mu^-)] = (1.89 \pm 0.17 \pm 0.09) \times 10^{-5}$ which we multiply by our best (shown rounded) value $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.
- ² Measured with $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$, $J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$. $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) = (34.68 \pm 0.30)\%$ from PDG 20 was used.
- ³ HAYRAPETYAN 24A reports $[\Gamma(J/\psi(1S) \rightarrow \mu^+ \mu^- \mu^+ \mu^-) / \Gamma_{\text{total}}] / [B(J/\psi(1S) \rightarrow \mu^+ \mu^-)] = (16.9_{-4.6}^{+5.5} \pm 0.6) \times 10^{-6}$ which we multiply by our best (shown rounded) value $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

$\Gamma(\mu^+ \mu^- \mu^+ \mu^-) / \Gamma(\mu^+ \mu^-)$		Γ_{10} / Γ_7		
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
18.7 ± 1.8 OUR AVERAGE				
18.9 ± 1.7 ± 0.9	452	¹ AAIJ	24AE LHCB	$J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$
16.9 _{-4.6} ^{+5.5} ± 0.6	12	HAYRAPETY...24A	CMS	$J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$

¹ Includes prompt production and inclusive decays of b -hadrons.

HADRONIC DECAYS

$\Gamma(\rho\pi) / \Gamma_{\text{total}}$		Γ_{11} / Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.88 ± 0.12 OUR AVERAGE				
2.072 ± 0.017 ± 0.062	19.8k	¹ ANASHIN	23 KEDR	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
2.18 ± 0.19		^{2,3} AUBERT,B	04N BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
2.184 ± 0.005 ± 0.201	220k	^{3,4} BAI	04H BES	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
2.091 ± 0.021 ± 0.116		^{3,5} BAI	04H BES	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
1.21 ± 0.20		BAI	96D BES	$e^+ e^- \rightarrow \rho\pi$
1.42 ± 0.01 ± 0.19		COFFMAN	88 MRK3	$e^+ e^-$
1.3 ± 0.3	150	FRANKLIN	83 MRK2	$e^+ e^-$
1.6 ± 0.4	183	ALEXANDER	78 PLUT	$e^+ e^-$
1.33 ± 0.21		BRANDELIK	78B DASP	$e^+ e^-$
1.0 ± 0.2	543	BARTEL	76 CNTR	$e^+ e^-$
1.3 ± 0.3	153	JEAN-MARIE	76 MRK1	$e^+ e^-$

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0 \pi^0$, $J/\psi \rightarrow \rho^+ \pi^-$, $J/\psi \rightarrow \rho^- \pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.


² From the ratio of $\Gamma(e^+ e^-) B(\pi^+ \pi^- \pi^0)$ and $\Gamma(e^+ e^-) B(\mu^+ \mu^-)$ (AUBERT 04).

³ Not independent of their $B(\pi^+ \pi^- \pi^0)$.

⁴ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.

⁵ Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.

WEIGHTED AVERAGE
 1.88 ± 0.12 (Error scaled by 2.6)



$\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{11}/Γ_{170}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.142 \pm 0.011 \pm 0.026$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.331 ± 0.033	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.
² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$ Γ_{12}/Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.328 \pm 0.005 \pm 0.027$	COFFMAN 88	MRK3	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.35 ± 0.08	ALEXANDER 78	PLUT	e^+e^-
0.32 ± 0.08	BRANDELIK 78B	DASP	e^+e^-
0.39 ± 0.11	BARTEL 76	CNTR	e^+e^-
0.37 ± 0.09	JEAN-MARIE 76	MRK1	e^+e^-

$\Gamma(a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{total}$ Γ_{13}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.84 \pm 0.08 \pm 0.60$	1317	ANASHIN 22	KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(a_2(1320)^+\pi^-\pi^0 + c.c \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{total}$ Γ_{14}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.67 \pm 0.09 \pm 0.73$	1628	ANASHIN 22	KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9±2.2 OUR AVERAGE				
11.7±0.7±2.5	7584	AUGUSTIN 89	DM2	$J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$
8.4±4.5	36	VANNUCCI 77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.68	471	¹ ABLIKIM 19Q	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$

¹From an energy scan of $e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$ assuming PDG 16 values for $\Gamma(e^+e^-)$, $\Gamma(\mu^+\mu^-)$, and $\Gamma(\text{total})$.

 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.193±0.023 OUR AVERAGE				
0.194±0.017±0.029	299	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
0.193±0.013±0.029		COFFMAN 88	MRK3	$e^+e^- \rightarrow \pi^+\pi^-\eta$

 $\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.20±0.14±0.37	471	ABLIKIM 15H	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

 $\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.52 × 10⁻⁴	90	ABLIKIM 10C	BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

 $\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.8±2.2±3.4	232 ± 23	ABLIKIM 08E	BES2	$e^+e^- \rightarrow J/\psi$

 $\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.15±0.13±0.22	209	ABLIKIM 10C	BES2	$J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

 $\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.1 ±0.8 OUR AVERAGE		Error includes scale factor of 1.6.		
7.90±0.19±0.49	3476	¹ ABLIKIM 17AK	BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
8.3 ±3.0 ±1.2	19	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
11.4 ±1.4 ±1.6		COFFMAN 88	MRK3	$J/\psi \rightarrow \pi^+\pi^-\eta'$

¹From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

 $\Gamma(\rho^+ K^+ K^- \pi^- + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.53±0.16±0.81	485	ANASHIN 22	KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

$\Gamma(h_1(1415)\eta' \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.08 \pm 0.01^{+0.01}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave. $\Gamma(h_1(1595)\eta' \rightarrow \gamma\eta\eta')/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.16 \pm 0.02^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave. $\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$2.2 \pm 0.2 \pm 1.1$	19.8k	¹ ANASHIN	23 KEDR	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
-----------------------	-------	----------------------	---------	---

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered. $\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{33}/Γ_{170}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.9 \pm 1.7 \pm 2.7$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
0.80 ± 0.27	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.² From a Dalitz plot analysis in a Veneziano model. $\Gamma(\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp)/\Gamma(K_S^0 K^\pm\pi^\mp)$ Γ_{34}/Γ_{188}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$6.3 \pm 0.8 \pm 0.6$	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm\pi^\mp$
-----------------------	----	-------------------	----------	---

¹ From a Dalitz plot analysis in an isobar model. $\Gamma(\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0)/\Gamma(K^+K^-\pi^0)$ Γ_{35}/Γ_{187}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$9.3 \pm 2.0 \pm 0.6$	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+K^-\pi^0$
-----------------------	----	-------------------	----------	----------------------------------

¹ From a Dalitz plot analysis in an isobar model. $\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958))/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$3.28 \pm 0.55 \pm 0.44$	119	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$
--------------------------	-----	----------------------	-----------	--------------------------------------

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

$\Gamma(\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{38}/Γ_{170}

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8 \pm 2 \pm 5$	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22 ± 6	20k	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.
² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{40}/Γ_{170}

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4 \pm 1 \pm 20$	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
600 ± 250	20k	² LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model.
² From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{41}/Γ_{170}

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.0 ± 0.8	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$


¹ From a Dalitz plot analysis in a Veneziano model.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.45 ± 0.05 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
$0.538 \pm 0.012 \pm 0.065$	2090	¹ ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\pi^0$
$0.360 \pm 0.028 \pm 0.054$	222	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
$0.482 \pm 0.019 \pm 0.064$		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

WEIGHTED AVERAGE
 0.45 ± 0.05 (Error scaled by 1.4)



$\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{43}/Γ_{170}

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8 \pm 3 \pm 2$	20k	¹ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$


¹ From a Dalitz plot analysis in an isobar model and significance 4.9σ .

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{total}$ Γ_{44}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5 ± 1.0 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
$10.6 \pm 1.2 \pm 0.1$	3531	¹ ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.0 ± 1.6	18058	AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8 ± 1.6	215	BURMESTER	77D	PLUT e^+e^-
6.8 ± 1.9	348	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-)/\Gamma_{total}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.946 \pm 0.016 \pm 0.108) \times 10^{-2}$ which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

WEIGHTED AVERAGE
 8.5 ± 1.0 (Error scaled by 1.3)



$\Gamma(\omega\pi^0\pi^0)/\Gamma_{\text{total}}$ **Γ_{45}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.4 \pm 0.3 \pm 0.7$	509	AUGUSTIN	89	DM2	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ **Γ_{47}/Γ**


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.3 ± 0.6 OUR AVERAGE					
$4.3 \pm 0.2 \pm 0.6$	5860	AUGUSTIN	89	DM2	e^+e^-
4.0 ± 1.6	70	BURMESTER	77D	PLUT	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.9 ± 0.8	81	VANNUCCI	77	MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$ **Γ_{48}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.74 ± 0.20 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.			
2.352 ± 0.273	5k	¹ ABLIKIM	06F	BES2	$J/\psi \rightarrow \omega\eta$
$1.44 \pm 0.40 \pm 0.14$	13	² AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
$1.43 \pm 0.10 \pm 0.21$	378	JOUSSET	90	DM2	$J/\psi \rightarrow \text{hadrons}$
$1.71 \pm 0.08 \pm 0.20$		COFFMAN	88	MRK3	$e^+e^- \rightarrow 3\pi\eta$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.
² Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

WEIGHTED AVERAGE
 1.74 ± 0.20 (Error scaled by 1.6)



$\Gamma(\omega \pi^+ \pi^+ \pi^- \pi^-) / \Gamma_{\text{total}}$

Γ_{51} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
85 ± 34	140	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow 3(\pi^+ \pi^-) \pi^0$

$\Gamma(\omega \eta' \pi^+ \pi^-) / \Gamma_{\text{total}}$

Γ_{53} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.12 \pm 0.02 \pm 0.13$	14k	¹ ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega \eta' \pi^+ \pi^-$

¹ Using the decays $\omega \rightarrow \pi^+ \pi^- \pi^0$ and $\eta' \rightarrow \eta \pi^+ \pi^-$.

$\Gamma(\omega \eta'(958)) / \Gamma_{\text{total}}$

Γ_{54} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89 ± 0.18 OUR AVERAGE				
$2.08 \pm 0.30 \pm 0.14$	137	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$
2.26 ± 0.43	218	² ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \eta'$
$1.8 \begin{smallmatrix} +1.0 \\ -0.8 \end{smallmatrix} \pm 0.3$	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$1.66 \pm 0.17 \pm 0.19$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

² Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+ \pi^- \gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

$\Gamma(\omega f_0(980)) / \Gamma_{\text{total}}$

Γ_{55} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$1.41 \pm 0.27 \pm 0.47$	¹ AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\omega f_0(1710) \rightarrow \omega K \bar{K})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-4})		DOCUMENT ID	TECN	COMMENT
$4.8 \pm 1.1 \pm 0.3$	^{1,2}	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios. $\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.8^{+1.9}_{-1.6} \pm 1.7$	111^{+31}_{-26}	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.2 \times 10^{-4}$	90	¹ VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.8 \times 10^{-4}$	90	¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
------------------------	----	----------------------	--------	-------------------------------------

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K \bar{K}) = 0.713$. $\Gamma(\omega X(1835) \rightarrow \omega p \bar{p})/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-6}$	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma \pi^0 p \bar{p}$

 $\Gamma(\omega X(1835), X \rightarrow \eta' \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$< 6.2 \times 10^{-5}$	¹ ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega \eta' \pi^+ \pi^-$

¹ Using the decays $\omega \rightarrow \pi^+ \pi^- \pi^0$ and $\eta' \rightarrow \eta \pi^+ \pi^-$. $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.52 \pm 0.30 \pm 0.01$	276	¹ ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = (0.136 \pm 0.008 \pm 0.026) \times 10^{-2}$ which we divide by our best (shown rounded) value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value. $\Gamma(\omega K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$3.33 \pm 0.02 \pm 0.12$	ABLIKIM	24BQ BES3	$e^+ e^- \rightarrow J/\psi(1S)$

 $\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
34 ± 5 OUR AVERAGE				
$37.7 \pm 0.8 \pm 5.8$	1972 ± 41	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
$29.5 \pm 1.4 \pm 7.0$	879 ± 41	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega K \bar{K})/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19 ± 4 OUR AVERAGE				
19.8 ± 2.1 ± 3.9		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	$e^+ e^-$

¹ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

 $\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 9 OUR AVERAGE				
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
53 ± 14 ± 14	530 ± 140	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

 $\Gamma(\eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.48 ± 0.13 OUR AVERAGE			
1.50 ± 0.02 ± 0.19	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$
1.47 ± 0.03 ± 0.17	² ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.

² From $\eta' K_S^0 K^\pm \pi^\mp$.

 $\Gamma(\eta' K^{*0} \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.66 ± 0.03 ± 0.21	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

 $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.16 ± 0.12 ± 0.29	1.1k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

 $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.51 ± 0.09 ± 0.21	1.0k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.

 $\Gamma(\eta' h_1(1415) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
4.69 ± 0.80^{+0.74}_{-1.82}	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\bar{K} K^*(892) + c.c. \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp)$ $\Gamma_{72} / \Gamma_{188}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
90.5 ± 0.9 ± 3.8	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹From a Dalitz plot analysis in an isobar model.

$\Gamma(K^+ K^*(892)^- + c.c.) / \Gamma_{total}$ Γ_{73} / Γ


VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ^{+0.8}_{-1.0} OUR AVERAGE				Error includes scale factor of 2.9. See the ideogram below.

8.07 ± 0.04 ^{+0.38} _{-0.61}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
4.57 ± 0.17 ± 0.70	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
5.26 ± 0.13 ± 0.53		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp,$ $K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$

WEIGHTED AVERAGE
6.0+0.8-1.0 (Error scaled by 2.9)



$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma_{total}$ Γ_{74} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
2.69 ± 0.01 ^{+0.13}_{-0.20}	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$ $\Gamma_{74} / \Gamma_{187}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
92.4 ± 1.5 ± 3.4	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{76} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE				
3.96 ± 0.15 ± 0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
4.33 ± 0.12 ± 0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 0.6	45	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
-----------	----	----------	---------	--

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{78} / Γ

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
5.73 ± 0.14 ± 0.82	¹ ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ² ABLIKIM 06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

¹ Obtained from $J/\psi \rightarrow K^*(892) K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$ taking the value 2/3 for the probability of the $K^*(892)^0 \rightarrow K^+ \pi^-$ decay.

² A $K_0^*(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{82} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
3.81 ± 0.10 ± 0.54	1559	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

$\Gamma(K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{83} / Γ

VALUE (units 10 ⁻⁶)	DOCUMENT ID	TECN	COMMENT
6.28^{+0.16+0.59}_{-0.17-0.52}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(K^*(892)^\pm K^*(700)^\mp) / \Gamma_{\text{total}}$ Γ_{85} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.18^{+0.94}_{-0.54}	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}}$ Γ_{86} / Γ

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5 90 VANNUCCI 77 MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(K^*(892)^\pm K^*(892)^\mp) / \Gamma_{\text{total}}$ Γ_{87} / Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.00 ± 0.19^{+0.11}_{-0.32}	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

$$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}} \quad \Gamma_{88}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.8 \pm 0.8 \pm 1.2$	1	BAI	99C	BES $e^+ e^-$

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0) \quad \Gamma_{90}/\Gamma_{187}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.3 \pm 1.1 \pm 0.7$	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(K^*(1410)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp) \quad \Gamma_{91}/\Gamma_{188}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.9$	4k	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0) \quad \Gamma_{93}/\Gamma_{187}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 1.3 \pm 0.9$	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp) \quad \Gamma_{94}/\Gamma_{188}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 1.2$	4k	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

$$\Gamma(\bar{K}_2^*(1430)K + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{95}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 40 \times 10^{-4}$	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 66 \times 10^{-4}$	90	BRAUNSCH...	76	DASP $e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$
-----------------------	----	-------------	----	---

$$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{96}/\Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.04 \pm_{-0.19}^{+0.25}$	183k	ABLIKIM	19AQ	BES $J/\psi \rightarrow K^+ K^- \pi^0$

$$\Gamma(K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{97}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.65 \pm 0.80 \pm 0.44$	1094	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

$$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{99}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7 ± 2.6	40	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<29 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

 $\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.1^{+0.6}_{-0.1}$	183k	ABLIKIM 19AQ	BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 $\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 0.7^{+2.8}_{-1.4}$	183k	ABLIKIM 19AQ	BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	¹ BAI 99C	BES	$e^+ e^-$

¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

 $\Gamma(K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$8.54^{+1.07+2.35}_{-1.20-2.13}$	ABLIKIM 18AA	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

 $\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+ e^-$

 $\Gamma(\phi \pi^0)/\Gamma_{\text{total}}$ Γ_{109}/Γ

The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	¹ ABLIKIM 15K	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$
$0.124 \pm 0.033 \pm 0.030$		35 ± 9	² ABLIKIM 15K	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.4	90	³ ABLIKIM 05B	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \gamma \gamma$
<6.8	90	COFFMAN 88	MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

³ Superseded by ABLIKIM 15K.

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.94±0.15 OUR AVERAGE Error includes scale factor of 1.7.				
1.09±0.02±0.13		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
0.78±0.03±0.12		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ±0.9	23	FELDMAN 77	MRK1	e^+e^-

$\Gamma(\phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{112}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
16.0±1.0±3.0			
	FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{113}/Γ


<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.74 ±0.06 OUR AVERAGE Error includes scale factor of 1.2.				
0.71 ±0.10 ±0.05	99±14	¹ ZHU	23	BELL $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
0.898±0.024±0.089		ABLIKIM 05B	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadr}$
0.64 ±0.04 ±0.11	346	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
0.661±0.045±0.078		COFFMAN 88	MRK3	$e^+e^- \rightarrow K^+K^-\eta$

¹ From a fit to the combined $\phi\eta$ invariant mass spectrum with a Gaussian function for the J/ψ signals and a second-order polynomial function for the backgrounds.

$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$ Γ_{114}/Γ

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.6 ±0.5 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.					
5.10±0.03±0.32		31k	ABLIKIM 16M	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
5.46±0.31±0.56			ABLIKIM 05B	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
4.1 ±0.3 ±0.8		167	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
3.08±0.34±0.36			COFFMAN 88	MRK3	$e^+e^- \rightarrow K^+K^-\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 13		90	VANNUCCI 77	MRK1	e^+e^-

WEIGHTED AVERAGE
 4.6 ± 0.5 (Error scaled by 2.2)



$\Gamma(\phi \eta \eta') / \Gamma_{\text{total}}$

Γ_{115} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.32 \pm 0.06 \pm 0.16$	2.2k	¹ ABLIKIM	19AN BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

¹Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi \eta'$ with $J^P = 1^+ \text{ or } 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi \eta') \approx 10^{-4}$.

$\Gamma(\phi f_0(980)) / \Gamma_{\text{total}}$

Γ_{116} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.9.
$4.6 \pm 0.4 \pm 0.8$		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6 ± 0.6	50	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹Assuming $B(f_0(980) \rightarrow \pi \pi) = 0.78$.

$\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$

Γ_{119} / Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.50 \pm 0.80 \pm 0.61$	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0) / \Gamma_{\text{total}}$

Γ_{120} / Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.67 \pm 0.50 \pm 0.24$	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

$\Gamma(\phi f_0(980)\eta \rightarrow \eta \phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.23 \pm 0.75 \pm 0.73$	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$

 $\Gamma(\phi a_0(980)^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.24 \pm 0.20^{+0.52}_{-0.22}$	¹ ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$
$2.74 \pm 0.13^{+0.15}_{-0.16}$	² ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$
4.37 ± 1.35	^{3,4} ABLIKIM	18D BES3	$J/\psi \rightarrow \phi \eta \pi^0$
$5.0 \pm 2.7 \pm 2.5$	⁵ ABLIKIM	11D BES3	$J/\psi \rightarrow \phi \eta \pi^0$

¹ $J/\psi \rightarrow \phi a_0(980)$ electromagnetic decay.

² $J/\psi \rightarrow \phi f_0(980)$, $\phi a_0(980)$ mixing.

³ Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.

⁴ Superseded by ABLIKIM 24CB.

⁵ Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops.

 $\Gamma(\phi(1680)^0 \pi^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.66 \pm 0.26^{+1.1}_{-1.0}$	ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$

 $\Gamma(X(2000)^0 \pi^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$1.70 \pm 0.19^{+0.48}_{-0.13}$	ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$

 $\Gamma(h_1(1900)^0 \pi^0 \rightarrow \phi \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$8.44 \pm 0.35^{+1.4}_{-1.2}$	ABLIKIM	24CB BES3	$J/\psi \rightarrow \phi \eta \pi^0$

 $\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.45	90	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
< 0.37	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

 $\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.6 ± 0.5 OUR AVERAGE

$3.4 \pm 1.8 \pm 1.5$	1.1k	¹ ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
$3.2 \pm 0.6 \pm 0.4$		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$
$2.1 \pm 0.5 \pm 0.4$	25	² JOUSSET	90 DM2	$J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.6 \pm 0.2 \pm 0.1$	16	BECKER	87 MRK3	$J/\psi \rightarrow \phi K \bar{K} \pi$
-----------------------	----	--------	---------	---

¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best (shown rounded) value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+\pi^-\eta$ invariant mass distribution at 1297 MeV.

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
9.36±2.31±1.54	78	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+K^-\pi$

$\Gamma(\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi 3\pi^0)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
2.08±1.63±1.47	9	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+K^-\pi$

$\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.01±0.58±0.82		172	¹ ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17	90	² FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$
------	----	----------------------	----	-----	-------------------------------------

¹ With 3.6 σ significance.

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$.

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4 OUR AVERAGE		Error includes scale factor of 2.7.		
12.3±0.6±2.0		^{1,2} FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$
4.8±1.8	46	¹ GIDAL	81	MRK2 $J/\psi \rightarrow K^+K^-K^+K^-$

¹ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

² Including interference with $f_0(1710)$.

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.1 × 10⁻⁷	90	¹ ABLIKIM	16K	BES3 $J/\psi \rightarrow p\bar{p}K_S^0K_L^0, p\bar{p}K^+K^-$

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

$\Gamma(\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.8 × 10⁻⁴	90	ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{134}/Γ


VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.13 × 10⁻⁵	90	ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

$\Gamma(\phi K \bar{K})/\Gamma_{\text{total}}$ Γ_{135}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17.7 ± 1.6 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
16.6 ± 1.9 ± 1.2	163 ± 19	LEES	12F BABR	10.6 $e^+e^- \rightarrow 2(K^+K^-)\gamma$
21.4 ± 0.4 ± 2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
48 $^{+20}_{-16}$ ± 6	9.0 $^{+3.7}_{-3.0}$	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6 ± 0.8 ± 2.1		3 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	e^+e^-

¹ We have multiplied $K^+ K^-$ measurement by 2 to obtain $K \bar{K}$.
² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.
³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

WEIGHTED AVERAGE
 17.7 ± 1.6 (Error scaled by 1.3)



$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K})/\Gamma_{\text{total}}$ Γ_{136}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.6 ± 0.2 ± 0.6	1,2 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

¹ Including interference with $f'_2(1525)$.
² Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.

$\Gamma(\phi K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{139}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 0.8 OUR AVERAGE				
7.4 ± 0.6 ± 1.4	227 ± 19	ABLIKIM	08E BES2	$e^+e^- \rightarrow J/\psi$
7.4 ± 0.9 ± 1.1		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
7 ± 0.6 ± 1.0	163 ± 15	BECKER	87 MRK3	$e^+e^- \rightarrow \text{hadrons}$

$$\Gamma(\phi K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{140}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
21.8±2.3 OUR AVERAGE				
20.8±2.7±3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6±3.7±4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7±2.4±3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}} \quad \Gamma_{141}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
30±5 OUR AVERAGE				
31±6	4600	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
29±7	87	BURMESTER	77D PLUT	$e^+ e^-$

$$\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{142}/\Gamma$$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
23±3±5				
	229	AUGUSTIN	89 DM2	$e^+ e^-$

$$\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}} \quad \Gamma_{144}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.1 × 10⁻³				
	90	HENRARD	87 DM2	$e^+ e^-$

$$\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{145}/\Gamma$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.58±0.23±0.40				
	332	EATON	84 MRK2	$e^+ e^-$

$$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}} \quad \Gamma_{146}/\Gamma$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.10±0.09±0.28				
	233	EATON	84 MRK2	$e^+ e^-$

$$\Gamma(\bar{\Sigma}(1385)^0 p K^-)/\Gamma_{\text{total}} \quad \Gamma_{147}/\Gamma$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.51±0.26±0.18				
	89	EATON	84 MRK2	$e^+ e^-$

$$\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{148}/\Gamma$$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.82 × 10⁻⁵				
	90	ABLIKIM	13F BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.2 × 10 ⁻³	90	HENRARD	87 DM2	$e^+ e^-$
-------------------------	----	---------	--------	-----------

$$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{149}/\Gamma$$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.30±0.07 OUR AVERAGE				
0.30±0.03±0.08	74 ± 8	HENRARD	87 DM2	$e^+ e^-$
0.29±0.11±0.10	26	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\Sigma(1385)^+\bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{150}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.33±0.08 OUR AVERAGE				
0.34±0.04±0.08	77	HENRARD	87	DM2 e^+e^-
0.31±0.11±0.11	28	EATON	84	MRK2 e^+e^-

 $\Gamma(\Sigma(1385)^-\bar{\Sigma}(1385)^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{151}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.08 ±0.06 OUR AVERAGE				
1.096±0.012±0.071	43k	ABLIKIM	16L	BES3 e^+e^-
1.23 ±0.07 ±0.30	0.8k	ABLIKIM	12P	BES2 e^+e^-
1.00 ±0.04 ±0.21	0.6k	HENRARD	87	DM2 e^+e^-
0.86 ±0.18 ±0.22	56	EATON	84	MRK2 e^+e^-

 $\Gamma(\Sigma(1385)^+\bar{\Sigma}(1385)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{152}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.25 ±0.07 OUR AVERAGE				
1.258±0.014±0.078	53k	ABLIKIM	16L	BES3 e^+e^-
1.50 ±0.08 ±0.38	1k	ABLIKIM	12P	BES2 e^+e^-
1.19 ±0.04 ±0.25	0.7k	HENRARD	87	DM2 e^+e^-
1.03 ±0.24 ±0.25	68	EATON	84	MRK2 e^+e^-

 $\Gamma(\Sigma(1385)^0\bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{153}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.071±0.009±0.082	103k	ABLIKIM	17E	BES3 $e^+e^- \rightarrow J/\psi \rightarrow$ hadrons

 $\Gamma(\Lambda(1520)\bar{\Lambda} + \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{154}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.1 × 10⁻⁶	90	ABLIKIM	12B	BES3 $J/\psi \rightarrow \Lambda\bar{\Lambda}\gamma$

 $\Gamma(\bar{\Lambda}(1520)\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{155}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.80 × 10⁻³	90	LU	19	BELL $B^+ \rightarrow \bar{p}\Lambda K^+ K^+$

 $\Gamma(\Xi^0\Xi^0)/\Gamma_{\text{total}}$ Γ_{156}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.17 ±0.04 OUR AVERAGE				
1.165±0.004±0.043	135k	ABLIKIM	17E	BES3 $e^+e^- \rightarrow J/\psi \rightarrow$ hadrons
1.20 ±0.12 ±0.21	206	ABLIKIM	080	BES2 $e^+e^- \rightarrow J/\psi$

 $\Gamma(\Xi(1530)^-\Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{157}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.318±0.008 OUR AVERAGE				
0.317±0.002±0.008	70k	ABLIKIM	20	BES3 $e^+e^- \rightarrow J/\psi$
0.59 ±0.09 ±0.12	75	HENRARD	87	DM2 e^+e^-

 $\Gamma(\Xi(1530)^0\Xi^0)/\Gamma_{\text{total}}$ Γ_{158}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.32±0.12±0.07	24 ± 9	HENRARD	87	DM2 e^+e^-

$$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{159}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90	BAI	04G	BES2 $e^+ e^-$

$$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}} \quad \Gamma_{160}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-5}$	90	BAI	04G	BES2 $e^+ e^-$

$$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}} \quad \Gamma_{161}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-5}$	90	BAI	04G	BES2 $e^+ e^-$

$$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}} \quad \Gamma_{162}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-5}$	90	BAI	04G	BES2 $e^+ e^-$

$$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}} \quad \Gamma_{163}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90	BAI	04G	BES2 $e^+ e^-$

————— STABLE HADRONS —————


$$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{164}/\Gamma$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4	OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.		
$5.44 \pm 0.07 \pm 0.33$	23k	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
$3.5 \pm 0.8 \pm 0.1$	14k	¹ LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-)3\pi^0 \gamma$
4.73 ± 0.44	228k	² ABLIKIM	19Q	BES3 $J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
3.25 ± 0.49	46055	AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
3.17 ± 0.42	147	FRANKLIN	83	MRK2 $e^+ e^- \rightarrow \text{hadrons}$
3.64 ± 0.52	1500	BURMESTER	77D	PLUT $e^+ e^-$
4 ± 1	675	JEAN-MARIE	76	MRK1 $e^+ e^-$

¹LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0)] = (14.8 \pm 2.6 \pm 2.2) \times 10^{-3}$ keV which we divide by our best (shown rounded) values $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) = (18.3 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) values.

²From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$, assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.

WEIGHTED AVERAGE
 4.2 ± 0.4 (Error scaled by 2.1)



$\Gamma(3(\pi^+ \pi^-) \pi^0) / \Gamma_{\text{total}}$

Γ_{165} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.029 ± 0.006	OUR AVERAGE			
0.028 ± 0.009	11	FRANKLIN 83	MRK2	$e^+ e^- \rightarrow \text{hadrons}$
0.029 ± 0.007	181	JEAN-MARIE 76	MRK1	$e^+ e^-$

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$

Γ_{170} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
20.0 ± 0.7	OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.		
$18.78 \pm 0.13 \pm 0.51$	19.8k	¹ ANASHIN 23	KEDR	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
$21.37 \pm 0.04^{+0.64}_{-0.62}$	1.8M	² ABLIKIM 12H	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
$23.0 \pm 2.0 \pm 0.4$	256	³ AUBERT 07AU	BABR 10.6	$e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
$21.84 \pm 0.05 \pm 2.01$	220k	^{4,5} BAI 04H	BES	$e^+ e^-$
$20.91 \pm 0.21 \pm 1.16$		^{5,6} BAI 04H	BES	$e^+ e^-$
15 ± 2	168	FRANKLIN 83	MRK2	$e^+ e^-$

¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0 \pi^0$, $J/\psi \rightarrow \rho^+ \pi^-$, $J/\psi \rightarrow \rho^- \pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.

² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.


³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best (shown rounded) value $\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-) / \Gamma_{\text{total}} = 0.809 \pm 0.014$ keV. Our first error is their experiment's

error and our second error is the systematic error from using our best (shown rounded) value.

4 From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.

5 Mostly $\rho\pi$, see also $\rho\pi$ subsection.

6 Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.



$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-) / \Gamma_{\text{total}}$ Γ_{172} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.52±0.27 OUR AVERAGE		Error includes scale factor of 1.4.		
1.74±0.08±0.24	2616	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.2 ±0.3	309	VANNUCCI	77	MRK1 $e^+ e^-$

$\Gamma(\pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{173} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE				
1.47±0.13±0.13	140	¹ METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+ \pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	$e^+ e^-$
1.0 ±0.5	5	BRANDELIK	78B	DASP $e^+ e^-$
1.6 ±1.6	1	VANNUCCI	77	MRK1 $e^+ e^-$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{174} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.20±0.25 OUR AVERAGE		Error includes scale factor of 1.2.		
2.88±0.14±0.24	2654	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+ \pi^-)$

$3.53 \pm 0.12 \pm 0.29$	1107	¹ ABLIKIM	05H	BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow$ $J/\psi \pi^+ \pi^-, J/\psi \rightarrow$ $2(\pi^+ \pi^-)$
4.0 ± 1.0	76	JEAN-MARIE	76	MRK1	e^+e^-

¹ Computed using $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ **Γ_{175}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
40 ± 20	32	JEAN-MARIE	76	MRK1 e^+e^-

$\Gamma(4(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ **Γ_{177}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
90 ± 30	13	JEAN-MARIE	76	MRK1 e^+e^-

$\Gamma(2(\pi^+ \pi^-) \eta)/\Gamma_{\text{total}}$ **Γ_{178}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.29 ± 0.28 OUR AVERAGE				
$3.1 \pm 1.5 \pm 0.1$	14k	¹ LEES	21	BABR $10.6 e^+e^- \rightarrow$ $2(\pi^+ \pi^-) 3\pi^0 \gamma$
$2.26 \pm 0.08 \pm 0.27$	4.8k	ABLIKIM	05C	BES2 $e^+e^- \rightarrow 2(\pi^+ \pi^-) \eta$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-) \eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] \times [B(\eta \rightarrow 3\pi^0)] = (5.6 \pm 2.6 \pm 0.8) \times 10^{-3}$ keV which we divide by our best (shown rounded) values $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.53 \pm 0.10$ keV, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) values.

$\Gamma(3(\pi^+ \pi^-) \eta)/\Gamma_{\text{total}}$ **Γ_{179}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.24 \pm 0.96 \pm 1.11$	616	ABLIKIM	05C	BES2 $e^+e^- \rightarrow 3(\pi^+ \pi^-) \eta$

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ **Γ_{183}/Γ**

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.06 ± 0.05 OUR AVERAGE				
$3.072 \pm 0.023 \pm 0.050$	1.8 k	¹ ABLIKIM	24AB	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-$
$2.86 \pm 0.09 \pm 0.19$	1k	² METREVELI	12	$\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2.39 \pm 0.24 \pm 0.22$	107	³ BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ± 0.9	6	³ BRANDELIK	79C	DASP e^+e^-

¹ Using $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.916 \pm 0.033)\%$.

² Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

³ Interference with non-resonant $K^+ K^-$ production not taken into account.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ **Γ_{184}/Γ**


<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.95 ± 0.11 OUR AVERAGE				
$1.93 \pm 0.01 \pm 0.05$	110k	ABLIKIM	17AH	BES3 $J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$

2.62±0.15±0.14 0.3k ¹ METREVELI 12 $\psi(2S) \rightarrow \pi^+\pi^-K_S^0K_L^0$
 1.82±0.04±0.13 2.1k ² BAI 04A BES2 $J/\psi \rightarrow K_S^0K_L^0 \rightarrow \pi^+\pi^-X$

••• We do not use the following data for averages, fits, limits, etc. •••

1.18±0.12±0.18 JOUSSET 90 DM2 $J/\psi \rightarrow$ hadrons
 1.01±0.16±0.09 74 BALTRUSAIT..85D MRK3 e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$.



$\Gamma(K_S^0 K_S^0)/\Gamma_{total}$ **Γ_{185}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.7 × 10⁻⁹	90	ABLIKIM	25CC BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<1.4 × 10 ⁻⁸	95	¹ ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$
<1 × 10 ⁻⁶	95	¹ BAI	04D BES	$e^+e^- \rightarrow J/\psi \rightarrow K_S^0 K_S^0$
<5.2 × 10 ⁻⁶	90	¹ BALTRUSAIT..85C	MRK3	e^+e^-

¹ Forbidden by CP.

$\Gamma(K\bar{K}\pi)/\Gamma_{total}$ **Γ_{186}/Γ**

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 10 OUR AVERAGE				
55.2±12.0	25	FRANKLIN	83	MRK2 $e^+e^- \rightarrow K^+K^-\pi^0$
78.0±21.0	126	VANNUCCI	77	MRK1 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{187}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.88±0.01±0.12	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

$\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{187}/\Gamma_{170}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0±0.3±0.9	23k	LEES	17C BABR	$J/\psi \rightarrow h^+ h^- \pi^0$

$\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_{188}/\Gamma_{170}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
26.5±0.5±2.1	24k	LEES	17C BABR	$J/\psi \rightarrow h^0 h^+ h^-$

$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{192}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.04±0.26±0.92	2671	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.2 ±2.3	205	VANNUCCI	77 MRK1	$e^+ e^-$

$\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{204}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
31±13	30	VANNUCCI	77 MRK1	$e^+ e^-$

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{206}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.4^{+0.5}_{-0.4} \pm 0.2$	$11.0^{+4.3}_{-3.5}$	¹ HUANG	03 BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$
0.7±0.3		VANNUCCI	77 MRK1	$e^+ e^-$
¹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.				

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{214}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.121±0.029 OUR AVERAGE				
2.112±0.004±0.031	314k	ABLIKIM	12C BES3	$e^+ e^-$
2.17 ±0.16 ±0.04	317	¹ WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
2.26 ±0.01 ±0.14	63316	BAI	04E BES2	$e^+ e^- \rightarrow J/\psi$
1.97 ±0.22	99	BALDINI	98 FENI	$e^+ e^-$
1.91 ±0.04 ±0.30		PALLIN	87 DM2	$e^+ e^-$
2.16 ±0.07 ±0.15	1420	EATON	84 MRK2	$e^+ e^-$
2.5 ±0.4	133	BRANDELIK	79C DASP	$e^+ e^-$
2.0 ±0.5		BESCH	78 BONA	$e^+ e^-$
2.2 ±0.2	331	² PERUZZI	78 MRK1	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.0 ±0.3	48	ANTONELLI	93 SPEC	$e^+ e^-$

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best (shown rounded) value $B(B^+ \rightarrow J/\psi(1S) K^+) = (1.019 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

² Assuming angular distribution $(1+\cos^2\theta)$.


$\Gamma(\rho\bar{p}\pi^0)/\Gamma_{\text{total}}$ **Γ_{215}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.1.		
1.33 ± 0.02 ± 0.11	11k	ABLIKIM	09B	BES2 e^+e^-
1.13 ± 0.09 ± 0.09	685	EATON	84	MRK2 e^+e^-
1.4 ± 0.4		BRANDELIK	79C	DASP e^+e^-
1.00 ± 0.15	109	PERUZZI	78	MRK1 e^+e^-

$\Gamma(\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{216}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
6.46 ± 0.17 ± 0.43	1435	EATON	84	MRK2 e^+e^-
3.8 ± 1.6	48	BESCH	81	BONA e^+e^-
5.5 ± 0.6	533	PERUZZI	78	MRK1 e^+e^-

WEIGHTED AVERAGE
6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(\rho\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ **Γ_{217}/Γ**

Including $\rho\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
3.36 ± 0.65 ± 0.28	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

$\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$ **Γ_{218}/Γ**


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.513 ± 0.023 OUR AVERAGE				
1.496 ± 0.001 ± 0.023	3.2M	¹ ABLIKIM	24BV	BES3 e^+e^-

1.91 ± 0.02 ± 0.17	13k	¹ ABLIKIM	09	BES2	e ⁺ e ⁻
2.03 ± 0.13 ± 0.15	826	EATON	84	MRK2	e ⁺ e ⁻
2.5 ± 1.2		BRANDELIK	79c	DASP	e ⁺ e ⁻
2.3 ± 0.4	197	PERUZZI	78	MRK1	e ⁺ e ⁻

¹ Combining the $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.

$\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$					Γ_{219}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.31 × 10⁻³	90	EATON	84	MRK2	e ⁺ e ⁻ → hadronsγ

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$					Γ_{220}/Γ
VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.98 ± 0.10 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
0.90 ± 0.02 ± 0.09	2670	ABLIKIM	13P	BES3	e ⁺ e ⁻
0.98 ± 0.03 ± 0.14	2449	ABLIKIM	08	BES2	e ⁺ e ⁻
1.10 ± 0.17 ± 0.18	486	EATON	84	MRK2	e ⁺ e ⁻
1.6 ± 0.3	77	PERUZZI	78	MRK1	e ⁺ e ⁻



$\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$					Γ_{221}/Γ
VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.129 ± 0.014 OUR AVERAGE		Error includes scale factor of 2.0.			
0.126 ± 0.002 ± 0.007	16k	¹ ABLIKIM	19N	BES3	e ⁺ e ⁻
0.200 ± 0.023 ± 0.028	265 ± 31	² ABLIKIM	09	BES2	e ⁺ e ⁻
0.68 ± 0.23 ± 0.17	19	EATON	84	MRK2	e ⁺ e ⁻
1.8 ± 0.6	19	PERUZZI	78	MRK1	e ⁺ e ⁻

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.

² From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels.

$\Gamma(p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{222}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8±1.2±1.3	ABLIKIM	14N	BES3 $e^+e^- \rightarrow J/\psi$

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{223}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.519±0.033 OUR AVERAGE				
0.523±0.006±0.033	14k	ABLIKIM	16K	BES3 $J/\psi \rightarrow p\bar{p}K_S^0 K_L^0,$ $p\bar{p}K^+ K^-$
0.45 ±0.13 ±0.07		FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{224}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.12±0.09 OUR AVERAGE				
2.36±0.02±0.21	59k	ABLIKIM	06K	BES2 $J/\psi \rightarrow p\pi^-\bar{n}$
2.47±0.02±0.24	55k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \bar{p}\pi^+n$
2.02±0.07±0.16	1288	EATON	84	MRK2 $e^+e^- \rightarrow p\pi^-$
1.93±0.07±0.16	1191	EATON	84	MRK2 $e^+e^- \rightarrow \bar{p}\pi^+$
1.7 ±0.7	32	BESCH	81	BONA $e^+e^- \rightarrow p\pi^-$
1.6 ±1.2	5	BESCH	81	BONA $e^+e^- \rightarrow \bar{p}\pi^+$
2.16±0.29	194	PERUZZI	78	MRK1 $e^+e^- \rightarrow p\pi^-$
2.04±0.27	204	PERUZZI	78	MRK1 $e^+e^- \rightarrow \bar{p}\pi^+$

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{225}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.09±0.16 OUR AVERAGE				
2.07±0.01±0.17	36k	ABLIKIM	12C	BES3 e^+e^-
2.31±0.49	79	BALDINI	98	FENI e^+e^-
1.8 ±0.9		BESCH	78	BONA e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.90±0.55	40	ANTONELLI	93	SPEC e^+e^-

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{226}/Γ


<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.8±3.6	5	BESCH	81	BONA e^+e^-

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{230}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.88 ±0.08 OUR AVERAGE				Error includes scale factor of 2.6. See the ideogram below.
1.64 ±0.12 ±0.09		ABLIKIM	23S	BES3 $e^+e^- \rightarrow \gamma\Lambda\bar{\Lambda}$
1.943±0.003±0.033	441k	ABLIKIM	17L	BES3 e^+e^-
2.03 ±0.03 ±0.15	8887	ABLIKIM	06	BES2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$
1.96 ^{+0.47} _{-0.44} ±0.04	46	¹ WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08 ±0.06 ±0.24	631	BAI	98G	BES e^+e^-

1.38 ±0.05 ±0.20	1847	PALLIN	87	DM2	e^+e^-
1.58 ±0.08 ±0.19	365	EATON	84	MRK2	e^+e^-
2.6 ±1.6	5	BESCH	81	BONA	e^+e^-
1.1 ±0.2	196	PERUZZI	78	MRK1	e^+e^-

¹WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best (shown rounded) value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.019 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{231}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

< 6.4	90		² ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$
23 ±7 ±8		11	BAI	98G BES	e^+e^-
22 ±5 ±5		19	HENRARD	87 DM2	e^+e^-

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{232}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30±0.13±0.99	2.4k	ABLIKIM	12P BES2	J/ψ

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{233}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2±1.7 OUR AVERAGE				
15.7±0.80±1.54	454	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2±6.0 ±4.4	44	² ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

 $\Gamma(\Lambda\bar{\Sigma}^-\pi^++\text{c.c.})/\Gamma_{\text{total}}$ Γ_{234}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.26 ±0.05 OUR AVERAGE Error includes scale factor of 1.2.				
1.244±0.002±0.045	2.6M	ABLIKIM	23BU BES3	e^+e^-
1.52 ±0.08 ±0.16	589	¹ ABLIKIM	07H BES2	e^+e^-
1.11 ±0.06 ±0.20	342 ± 18	HENRARD	87 DM2	e^+e^-
1.38 ±0.21 ±0.35	118	EATON	84 MRK2	e^+e^-

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

 $\Gamma(\Lambda\bar{\Sigma}^+\pi^-\text{c.c.})/\Gamma_{\text{total}}$ Γ_{235}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 ±0.07 OUR AVERAGE Error includes scale factor of 1.8.				
1.221±0.002±0.038	2.7M	ABLIKIM	23BU BES3	e^+e^-
0.90 ±0.06 ±0.16	225	HENRARD	87 DM2	e^+e^-
1.53 ±0.17 ±0.38	135	EATON	84 MRK2	e^+e^-

 $\Gamma(pK^-\bar{\Lambda}+\text{c.c.})/\Gamma_{\text{total}}$ Γ_{236}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.11 OUR AVERAGE				
0.84 ^{+0.17} _{-0.15} ±0.02	45	¹ LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
0.89±0.07±0.14	307	EATON	84 MRK2	e^+e^-

¹ LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow pK^-\bar{\Lambda}+\text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best (shown rounded) value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.019 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

 $\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{237}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.06±0.05				
	90	EATON	84 MRK2	e^+e^-

 $\Gamma(pK_S^0\bar{\Sigma}^-+\text{c.c.})/\Gamma_{\text{total}}$ Γ_{238}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.725±0.009±0.050				
	120k	¹ ABLIKIM	24H BES3	$e^+e^- \rightarrow J/\psi$

¹ The branching fractions for the charge-conjugate channels are measured separately as $(1.361 \pm 0.006 \pm 0.025) \times 10^{-4}$ for $\bar{p}K_S^0\Sigma^+$ and $(1.352 \pm 0.006 \pm 0.025) \times 10^{-4}$ for $pK_S^0\bar{\Sigma}^-$.

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{239}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46 ± 0.20 ± 1.07	1058	¹ ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

 $\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{240}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83 ± 0.23 OUR AVERAGE					
2.74 ± 0.24 ± 0.22		234 ± 21	¹ ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92 ± 0.22 ± 0.24		308 ± 24	² ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18			² HENRARD	87 DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
<15	90		PERUZZI	78 MRK1	$e^+e^- \rightarrow \Lambda X$

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{241}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.04 OUR AVERAGE				
1.061 ± 0.004 ± 0.036	87k	ABLIKIM	21AT BES3	$J/\psi \rightarrow p\pi^0\bar{p}\pi^0$
1.50 ± 0.10 ± 0.22	399	ABLIKIM	08O BES2	$e^+e^- \rightarrow J/\psi$

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{242}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.172 ± 0.032 OUR AVERAGE				Error includes scale factor of 1.4.
1.164 ± 0.004 ± 0.023	111k	ABLIKIM	17L BES3	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.33 ± 0.04 ± 0.11	1.7k	ABLIKIM	06 BES2	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 ± 0.04 ± 0.23	884	PALLIN	87 DM2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 ± 0.16 ± 0.25	90	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •


2.4 ± 2.6	3	BESCH	81 BONA	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$
-----------	---	-------	---------	---

 $\Gamma(\Sigma^+\bar{\Sigma}^- \eta)/\Gamma_{\text{total}}$ Γ_{243}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.34 ± 0.21 ± 0.37	1821	ABLIKIM	22AY BES3	$J/\psi \rightarrow \Sigma^+\bar{\Sigma}^- \eta$

 $\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{244}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.08 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
1.040 ± 0.006 ± 0.074	43k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Xi^-\bar{\Xi}^+$
0.90 ± 0.03 ± 0.18	961	ABLIKIM	12P BES2	$J/\psi \rightarrow \Xi^-\bar{\Xi}^+$
0.70 ± 0.06 ± 0.12	132	HENRARD	87 DM2	$e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$
1.14 ± 0.08 ± 0.20	194	EATON	84 MRK2	$e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$
1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$



$\Gamma(\Xi^0 \bar{\Sigma}^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{245}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.64 \pm 0.17 \pm 0.28$	1284	ABLIKIM	25BWBES3	$J/\psi \rightarrow \Xi^0 \bar{\Sigma}^- K^+$

$\Gamma(\Xi^0 \bar{\Sigma}^0 K_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{246}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.24 \pm 0.32 \pm 0.31$	91	ABLIKIM	25BWBES3	$J/\psi \rightarrow \Xi^0 \bar{\Sigma}^0 K_S^0$

$\Gamma(\Xi^0 \bar{\Lambda} K_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{247}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.76 \pm 0.14 \pm 0.23$	982	ABLIKIM	25BWBES3	$J/\psi \rightarrow \Xi^0 \bar{\Lambda} K_S^0$

————— RADIATIVE DECAYS —————

$\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{248}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.82 ± 0.15 OUR FIT Error includes scale factor of 1.6.

1.7 ± 0.4 OUR AVERAGE Error includes scale factor of 1.5.

$1.99 \pm 0.31 \pm 0.02$ ¹ MITCHELL 09 CLEO $e^+e^- \rightarrow \gamma X$

1.27 ± 0.36 GAISER 86 CBAL $J/\psi \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.40 ± 0.33 ² ANASHIN 14 KEDR $J/\psi \rightarrow \gamma \eta_c$

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming

$B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best (shown rounded) value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.78 \pm 0.33) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

²Statistical uncertainty only.

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ **Γ_{251}/Γ**

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.6±2.2 OUR AVERAGE					
11.3±1.8±2.0		113 ± 18	ABLIKIM	13I BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ± 3 ± 2		24.2 ^{+7.2} _{-6.0}	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	PARTRIDGE	80	CBAL	e^+e^-
-----	----	-----------	----	------	----------

$\Gamma(4\gamma)/\Gamma_{\text{total}}$ **Γ_{252}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<9 × 10⁻⁶	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(5\gamma)/\Gamma_{\text{total}}$ **Γ_{253}/Γ**

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<15 × 10⁻⁶	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ **Γ_{254}/Γ**

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.39±0.08 OUR AVERAGE				
3.34±0.02±0.09	176k	ABLIKIM	23BD BES3	$J/\psi \rightarrow \pi^0\gamma$
3.58±0.20±0.03	1.6k	¹ ABLIKIM	18O BES3	$\psi(2S) \rightarrow \pi^+\pi^- \gamma\gamma\gamma$
3.63±0.36±0.13		PEDLAR	09 CLE3	$J/\psi \rightarrow \pi^0\gamma$
3.13 ^{+0.65} _{-0.47}	586	ABLIKIM	06E BES2	$J/\psi \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.6 ± 1.1 ± 0.7		BLOOM	83	CBAL	e^+e^-
7.3 ± 4.7	10	BRANDELIK	79C	DASP	e^+e^-

¹ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best (shown rounded) values $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.78 \pm 0.33) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) values.

$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ **Γ_{255}/Γ**

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.15±0.05	¹ ABLIKIM	15AE BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$

¹The uncertainty is systematic as statistical is negligible.


$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

Γ_{256}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
4.32 ± 0.14 ± 0.73	¹ BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
2.08 ± 0.13 ± 0.35	² BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
3.05 ± 0.08 ± 0.45	² BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85 ± 0.45 ± 1.20	³ BURKE	82	MRK2 e^+e^-

- ¹ 4π mass less than 3.0 GeV.
- ² 4π mass less than 2.0 GeV.
- ³ 4π mass less than 2.5 GeV.

WEIGHTED AVERAGE
2.8±0.5 (Error scaled by 1.9)



$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$

Γ_{257}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.5 ± 0.7 ± 1.6	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant}))/\Gamma_{\text{total}}$

Γ_{258}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.2 ± 0.8 ± 1.7	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

- ¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

$\Gamma(\gamma \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$

Γ_{259}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.3 ± 0.2 ± 3.1	¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

- ¹ 4π mass less than 2.0 GeV.

$$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}} \quad \Gamma_{260}/\Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.1 ± 0.4	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$$\Gamma(\gamma(K\bar{K}\pi)[J^{PC} = 0^{-+}])/\Gamma_{\text{total}} \quad \Gamma_{261}/\Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
0.58 ± 0.03 ± 0.20	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
2.1 ± 0.1 ± 0.7	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

¹ For a broad structure around 1800 MeV.

² For a broad structure around 2040 MeV.

$$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{262}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1 ± 0.1 ± 0.6	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$$\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}} \quad \Gamma_{263}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.3 ± 1.3	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

¹ Summed over all charges.

$$\Gamma(\gamma f_1(1420) \rightarrow \gamma K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0 \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{306}/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
7.25 ± 0.12^{+0.73}_{-1.25}	126k	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$

$$\Gamma(\gamma f_1(1420) \rightarrow \gamma a_0(980)^0 \pi^0 \rightarrow \gamma K_S^0 K_S^0 \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{307}/\Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.62 ± 0.36^{+2.36}_{-1.94}	126k	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$

$$\Gamma(\gamma\eta)/\Gamma_{\text{total}} \quad \Gamma_{264}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.090 ± 0.013 OUR AVERAGE				
1.096 ± 0.001 ± 0.019	2.2M	ABLIKIM	23BD BES3	$J/\psi \rightarrow \eta\gamma$
1.067 ± 0.005 ± 0.023	87.9k	ABLIKIM	21AMBES3	$e^+e^- \rightarrow J/\psi$
1.11 ± 0.05 ± 0.01	18.6k	¹ ABLIKIM	18O BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma$
1.101 ± 0.029 ± 0.022		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta\gamma$
1.123 ± 0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.08 ± 0.11		BLOOM	83 CBAL	e^+e^-
0.82 ± 0.10		BRANDELIK	79C DASP	e^+e^-
1.3 ± 0.4	21	BARTEL	77 CNTR	e^+e^-

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best (shown rounded) values $B(\eta \rightarrow 2\gamma) =$

$(39.36 \pm 0.18) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.78 \pm 0.33) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) values.

$\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{265}/Γ

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
21.4±1.8±2.5	596	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma f_0(500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{266}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.5±2.0	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{267}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5±5	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{268}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4±3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{269}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.5 × 10⁻⁶	95	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{270}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6.6 × 10⁻⁶	95	ABLIKIM	16P	BES3 $J/\psi \rightarrow 5\gamma$

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$ Γ_{271}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.1 ± 1.0 OUR AVERAGE			
5.85±0.3±1.05	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta\pi^+\pi^-$
7.8 ± 1.2±2.4	¹ EDWARDS	83B	CBAL $J/\psi \rightarrow \eta 2\pi^0$
¹ Broad enhancement at 1700 MeV.			

$\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{272}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.2±2.2±0.9	BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{273}/Γ


<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.28±0.06 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
5.40±0.01±0.11	638k	ABLIKIM	23BD	BES3 $J/\psi \rightarrow \gamma\eta'$
5.27±0.03±0.05	36k	ABLIKIM	19T	BES $J/\psi \rightarrow \gamma\eta'$
5.42±0.23±0.09	5.0k	¹ ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$4.76 \pm 0.22 \pm 0.06$		² ABLIKIM	11	BES3	$J/\psi \rightarrow \eta' \gamma$
$5.24 \pm 0.12 \pm 0.11$		PEDLAR	09	CLE3	$J/\psi \rightarrow \eta' \gamma$
5.55 ± 0.44	35k	ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta' \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$4.50 \pm 0.14 \pm 0.53$		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta, \eta \rightarrow \gamma \gamma$
$4.30 \pm 0.31 \pm 0.71$		BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta, \eta \rightarrow \pi^+ \pi^- \pi^0$
$4.04 \pm 0.16 \pm 0.85$	622	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$4.39 \pm 0.09 \pm 0.66$	2420	AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
$4.1 \pm 0.3 \pm 0.6$		BLOOM	83	CBAL	$e^+ e^- \rightarrow 3\gamma + \text{hadrons}$
2.9 ± 1.1	6	BRANDELIK	79C	DASP	$e^+ e^- \rightarrow 3\gamma$
2.4 ± 0.7	57	BARTEL	76	CNTR	$e^+ e^- \rightarrow 2\gamma \rho$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma \gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma \gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best (shown rounded) values $B(\eta'(958) \rightarrow \gamma \gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.78 \pm 0.33) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) values.

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+ \pi^- \eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+ \pi^- \eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best (shown rounded) values $B(\eta'(958) \rightarrow \pi^+ \pi^- \eta) = (42.5 \pm 0.5) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) values.

WEIGHTED AVERAGE
 5.28 ± 0.06 (Error scaled by 1.3)



$\Gamma(\gamma f_0(980) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{274} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.2	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	--

$\Gamma(\gamma f_0(980) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{275} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.8 ± 0.3	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	--

$\Gamma(\gamma \rho \rho) / \Gamma_{\text{total}}$ Γ_{276} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

4.5 ± 0.8 OUR AVERAGE

$4.7 \pm 0.3 \pm 0.9$		¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
-----------------------	--	------------------------------	------	---------------------------------

$3.75 \pm 1.05 \pm 1.20$		² BURKE	82	MRK2 $J/\psi \rightarrow 4\pi\gamma$
--------------------------	--	--------------------	----	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.09	90	³ BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
----------	----	----------------------	-----	---------------------------------

¹ 4π mass less than 2.0 GeV.

² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .

³ 4π mass in the range 2.0–25 GeV.

$\Gamma(\gamma \rho \omega) / \Gamma_{\text{total}}$ Γ_{277} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 5.4 \times 10^{-4}$	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$
------------------------	----	---------	----------	-----------------------------

$\Gamma(\gamma \rho \phi) / \Gamma_{\text{total}}$ Γ_{278} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 8.8 \times 10^{-5}$	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$
------------------------	----	---------	----------	-----------------------------

$\Gamma(\gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{279} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.61 ± 0.33 OUR AVERAGE

$6.0 \pm 4.8 \pm 1.8$		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma \omega \pi^+ \pi^-$
-----------------------	--	---------	----------	--

$1.41 \pm 0.2 \pm 0.42$	120 ± 17	BISELLO	87 SPEC	e^+e^- , hadrons γ
-------------------------	--------------	---------	---------	-----------------------------

$1.76 \pm 0.09 \pm 0.45$		BALTRUSAIT..85C	MRK3	$e^+e^- \rightarrow$ hadrons γ
--------------------------	--	-----------------	------	---------------------------------------

$\Gamma(\gamma \phi \phi) / \Gamma_{\text{total}}$ Γ_{280} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------


4.0 ± 1.2 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.

$7.5 \pm 0.6 \pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
-----------------------	-----	-----	----------	--------------------------------

$3.4 \pm 0.8 \pm 0.6$	33 ± 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
-----------------------	------------	----------------------	--------	---

$3.1 \pm 0.7 \pm 0.4$		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
-----------------------	--	----------------------	---------	---

¹ $\phi\phi$ mass less than 2.9 GeV, η_c excluded.



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{281}/Γ

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
$1.66 \pm 0.1 \pm 0.58$	^{1,2} BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
$3.8 \pm 0.3 \pm 0.6$	³ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$4.0 \pm 0.7 \pm 1.0$	³ EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	^{3,4} SCHARRE	80 MRK2	$e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1.78 \pm 0.21 \pm 0.33$	^{3,5,6} AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.83 \pm 0.13 \pm 0.18$	^{3,7,8} AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
$0.66^{+0.17+0.24}_{-0.16-0.15}$	^{3,6,9} BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
$1.03^{+0.21+0.26}_{-0.18-0.19}$	^{3,8,10} BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K \bar{K} \pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

³ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.


⁵ From fit to the $a_0(980) \pi 0^-+$ partial wave.

⁶ $a_0(980) \pi$ mode.

⁷ From fit to the $K^*(892) K 0^-+$ partial wave.

⁸ $K^* K$ mode.

- ⁹ From $a_0(980)\pi$ final state.
¹⁰ From $K^*(890)K$ final state.



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0) / \Gamma_{\text{total}}$ Γ_{282} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.20 OUR AVERAGE	Error includes scale factor of 1.8.		
$1.07 \pm 0.17 \pm 0.11$	¹ BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
$0.64 \pm 0.12 \pm 0.07$	¹ COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-) / \Gamma_{\text{total}}$ Γ_{283} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ± 0.5 OUR AVERAGE				
$2.6 \pm 0.7 \pm 0.4$		BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$3.38 \pm 0.33 \pm 0.64$		¹ BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$7.0 \pm 0.6 \pm 1.1$	261	² AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Via $a_0(980)\pi$.

² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0) / \Gamma_{\text{total}}$ Γ_{284} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.3.		
2.1 ± 0.4	BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1.36 ± 0.38	^{1,2} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.² Includes unknown branching fraction to $\rho^0 \rho^0$. $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{285}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$3.57 \pm 0.18^{+0.59}_{-0.61}$			¹ ABLIKIM	25P BES3	$J/\psi \rightarrow \gamma\gamma\phi$
---------------------------------	--	--	----------------------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.03 \pm 0.92 \pm 0.91$	1.3k		² ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
$10.36 \pm 1.51 \pm 1.54$	1.9k		³ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
<82	95		BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma K^+ K^-$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\gamma\phi$ with significance 18.9σ .² Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.³ Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass. $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{286}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.5 \pm 0.11 \pm 0.11$	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma\eta(1405)$
-------------------------	-----	---------	----------	---------------------------------------

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^0\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{287}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.64^{+1.10}_{-0.23} OUR AVERAGE

$4.62 \pm 0.15^{+5.08}_{-0.18}$		¹ ABLIKIM	25AB BES3	$J/\psi \rightarrow \gamma\pi^0 f_0(980)$ $\rightarrow \gamma\pi^0\pi^0\pi^0$
---------------------------------	--	----------------------	-----------	--

$7.10 \pm 0.82 \pm 0.72$	198	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma\eta(1405)$
--------------------------	-----	---------	----------	---------------------------------------

¹ ABLIKIM 25AB measured $J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma\pi^0 f_0(980) / 0^{++} PHSP \rightarrow \gamma\pi^0\pi^0\pi^0$. Here $\gamma\pi^0 f_0(980) / 0^{++} PHSP$ indicates an intermediate state which could be either $\gamma\pi^0 f_0(980)$ or $\gamma\pi^0 0^{++}$ phase space. $\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{288}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.63 \times 10^{-6}$	90	ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
------------------------	----	---------	----------	---

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{289}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.86 \times 10^{-6}$	90	ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
------------------------	----	---------	----------	---

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{290}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.13 ± 0.09	^{1,2} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
-----------------	------------------------	---------	---------------------------------

¹ Estimated by us from various fits.² Includes unknown branching fraction to $\rho^0 \rho^0$. $\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{291}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
--------------------------	------	---------	---------	---

$\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{292}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.80 \times 10^{-6}$	90	ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$					Γ_{293}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.14^{+0.50}_{-0.19} OUR AVERAGE					
2.40 ± 0.10 ^{+2.47} _{-0.18}		1,2 ABLIKIM	16N	BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
4.4 ± 0.4 ± 0.8	196	2 ABLIKIM	08I	BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
3.3 ± 0.8 ± 0.5		2 BAI	90B	MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
2.7 ± 0.6 ± 0.6		2 BAI	90B	MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
2.4 ^{+1.5} _{-1.0}		3,4 BISELLO	89B	DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

² Includes unknown branching fraction to $\phi\phi$.

³ Estimated by us from various fits.

⁴ Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$					Γ_{294}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.63 ± 0.12 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.					
2.07 ± 0.16 ^{+0.02} _{-0.07}	2.4k	1,2 DOBBS	15		$J/\psi \rightarrow \gamma\pi\pi$
1.63 ± 0.26 ^{+0.02} _{-0.05}		3 ABLIKIM	06V	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1.42 ± 0.21 ^{+0.02} _{-0.05}		4 ABLIKIM	06V	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
1.33 ± 0.05 ± 0.20		5 AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.36 ± 0.09 ± 0.23		5 BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma\pi^+\pi^-$
1.48 ± 0.25 ± 0.30	178	EDWARDS	82B	CBAL	$e^+e^- \rightarrow 2\pi^0\gamma$
2.0 ± 0.7	35	ALEXANDER	78	PLUT	e^+e^-
1.2 ± 0.6	30	6 BRANDELIK	78B	DASP	$e^+e^- \rightarrow \pi^+\pi^-\gamma$


¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best (shown rounded) value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

³ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best (shown rounded) value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best (shown rounded) value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

- ⁵ Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.
⁶ Restated by us to take account of spread of E1, M2, E3 transitions.



$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$	Γ_{295} / Γ
VALUE (units 10^{-5})	DOCUMENT ID TECN COMMENT
$2.58^{+0.08+0.59}_{-0.09-0.20}$	ABLIKIM 18AA BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_1(1285)) / \Gamma_{\text{total}}$	Γ_{296} / Γ
VALUE (units 10^{-3})	DOCUMENT ID TECN COMMENT
0.61 ± 0.08 OUR AVERAGE	
0.69 ± 0.16 ± 0.20	1 BAI 04J BES2 $J/\psi \rightarrow \gamma \gamma \rho^0$
0.61 ± 0.04 ± 0.21	2 BAI 00D BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.45 ± 0.09 ± 0.17	3 BAI 99 BES $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.625 ± 0.063 ± 0.103	4 BOLTON 92 MRK3 $J/\psi \rightarrow \gamma f_1(1285)$
0.70 ± 0.08 ± 0.16	5 BOLTON 92B MRK3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

- ¹ Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.
² Assuming $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}} = 0.090 \pm 0.004$.
³ Assuming $\Gamma(f_1(1285) \rightarrow \eta \pi \pi) / \Gamma_{\text{total}} = 0.5 \pm 0.18$.
⁴ Obtained summing the sequential decay channels
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow K\bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.

⁵ Using $B(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.

$\Gamma(\gamma f_1(1285) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{297}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.29 \pm 0.03^{+0.11}_{-0.09}$	¹ ABLIKIM	25P BES3	$J/\psi \rightarrow \gamma\gamma\phi$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\gamma\phi$ with significance 17.3σ .

$\Gamma(\gamma f_1(1285) \rightarrow \gamma\pi^0 f_0(980) \rightarrow \gamma\pi^0\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{298}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$5.64 \pm 0.45^{+0.74}_{-3.05}$	¹ ABLIKIM	25AB BES3	

¹ ABLIKIM 25AB measured $J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\pi^0 f_0(980) / 0^{++} PHSP \rightarrow \gamma\pi^0\pi^0\pi^0$. Here $\gamma\pi^0 f_0(980) / 0^{++} PHSP$ indicates an intermediate state which could be either $\gamma\pi^0 f_0(980)$ or $\gamma\pi^0 0^{++}$ phase space.

$\Gamma(\gamma f_1(1285) \rightarrow \gamma a_0(980)^0\pi^0 \rightarrow \gamma K_S^0 K_S^0\pi^0)/\Gamma_{\text{total}}$ Γ_{299}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.55 \pm 0.41^{+3.42}_{-1.04}$	126k	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0\pi^0$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{300}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

38 ± 10 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{301}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$4.19 \pm 0.73 \pm 1.34$ 478 ¹ DOBBS 15 $J/\psi \rightarrow \gamma K\bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.4 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{302}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.07^{+0.08+0.36}_{-0.07-0.34}$ ABLIKIM 18AA BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{303}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.5 ± 1.0 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(1370) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{304} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9 ± 0.3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	--------------	------	---

 $\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{305} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.79 ± 0.13 OUR AVERAGE

0.68 ± 0.04 ± 0.24	BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
--------------------	-----	-----	---

0.76 ± 0.15 ± 0.21	^{1,2} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
--------------------	-------------------------	----	---

0.87 ± 0.14 ^{+0.14} _{-0.11}	¹ BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
---	------------------	-----	--

¹ Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.

² From fit to the $K^*(892) K 1^{++}$ partial wave.

 $\Gamma(\gamma f_1(1420) \rightarrow \gamma \gamma \phi) / \Gamma_{\text{total}}$ Γ_{308} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.55 ± 0.07 ^{+0.18}_{-0.17}	¹ ABLIKIM	25P	BES3 $J/\psi \rightarrow \gamma \gamma \phi$
---	----------------------	-----	--

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma \gamma \phi$ with significance 9.0σ .

 $\Gamma(\gamma f_1(1420) \rightarrow \gamma \pi^0 f_0(980) \rightarrow \gamma \pi^0 \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{309} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

2.23 ± 0.16 ^{+0.20}_{-1.20}	¹ ABLIKIM	25AB	BES3
---	----------------------	------	------

¹ ABLIKIM 25AB measured $J/\psi \rightarrow \gamma f_1(1420) \rightarrow \gamma \pi^0 f_0(980) / 0^{++} PHSP \rightarrow \gamma \pi^0 \pi^0 \pi^0$. Here $\gamma \pi^0 f_0(980) / 0^{++} PHSP$ indicates an intermediate state which could be either $\gamma \pi^0 f_0(980)$ or $\gamma \pi^0 0^{++}$ phase space.

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{310} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.09 ± 0.24 OUR AVERAGE

1.21 ± 0.29 ± 0.24	174	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
--------------------	-----	--------------------	----	-------------------------------------

1.00 ± 0.03 ± 0.45		² ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
--------------------	--	----------------------	-----	--

1.02 ± 0.09 ± 0.45		² ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
--------------------	--	----------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.90 ± 0.17		SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	--	--------------	------	---

5.7 ± 0.8		^{3,4} BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
-----------	--	---------------------	----	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.

⁴ Assuming that $f_0(1500)$ decays only to two S-wave dipions.

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{311} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.65 ^{+0.26 +0.51}_{-0.31 -1.40}	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$
--	------	----------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.1 ± 0.4 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{312} / \Gamma$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.16^{+0.18}_{-0.56}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.7 ± 0.3 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}} \quad \Gamma_{313} / \Gamma$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$18.1 \pm 1.1^{+1.9}_{-1.3}$ ¹ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
 12 ± 5 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ *P*-wave.

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{314} / \Gamma$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.0 \pm 0.7$	BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \gamma \phi) / \Gamma_{\text{total}} \quad \Gamma_{315} / \Gamma$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.78 \pm 0.09^{+0.34}_{-0.30}$	¹ ABLIKIM	25P BES3	$J/\psi \rightarrow \gamma \gamma \phi$

¹From a partial wave analysis of $J/\psi \rightarrow \gamma \gamma \phi$ with significance 5.3σ .

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \pi^0 f_0(980) \rightarrow \gamma \pi^0 \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{316} / \Gamma$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.91 \pm 1.20^{+0.74}_{-3.83}$ ¹ABLIKIM 25AB BES3

¹ABLIKIM 25AB measured $J/\psi \rightarrow \gamma f_1(1510) \rightarrow \gamma \pi^0 f_0(980) / 0^{++} PHSP \rightarrow \gamma \pi^0 \pi^0 \pi^0$. Here $\gamma \pi^0 f_0(980) / 0^{++} PHSP$ indicates an intermediate state which could be either $\gamma \pi^0 f_0(980)$ or $\gamma \pi^0 0^{++}$ phase space. Evidence for the $f_1(1510)$ is seen with a 5σ significance in the partial wave analysis.

$\Gamma(\gamma f_2'(1525)) / \Gamma_{\text{total}} \quad \Gamma_{317} / \Gamma$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

5.7 $^{+0.8}_{-0.5}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

8.0 ± 0.9 ± 0.2 750 ^{1,2}DOBBS 15 $J/\psi \rightarrow \gamma K\bar{K}$


$3.85 \pm 0.17^{+1.91}_{-0.73}$		³ BAI	03G	BES	$J/\psi \rightarrow \gamma K \bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		³ BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		³ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		³ BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<3.4	90	4	⁴ BRANDELIK	79C	DASP $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78	PLUT $e^+ e^- \rightarrow K^+ K^- \gamma$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best (shown rounded) value $B(f'_2(1525) \rightarrow K \bar{K}) = (88.8 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best (shown rounded) value.

³ Using $B(f'_2(1525) \rightarrow K \bar{K}) = 0.888$.

⁴ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.



$\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{318}/Γ		
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.99^{+0.03+0.69}_{-0.04-0.50}	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{319}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$9.47 \pm 0.43^{+1.51}_{-0.66}$	126k	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$	

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$					Γ_{320}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$	

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f'_2(1525) \rightarrow \gamma \gamma \phi)/\Gamma_{\text{total}}$					Γ_{322}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.76 \pm 0.18^{+0.90}_{-0.61}$		¹ ABLIKIM	25P BES3	$J/\psi \rightarrow \gamma \gamma \phi$	

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma \gamma \phi$ with significance 16.4σ .

$\Gamma(\gamma f_2(1565) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$					Γ_{321}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.32 \pm 0.05^{+0.12}_{-0.02}$		¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ *P*-wave.

$\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$					Γ_{323}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$	

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$					Γ_{324}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.8 ± 0.5 OUR AVERAGE					
$3.72 \pm 0.30 \pm 0.43$	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$	
$3.96 \pm 0.06 \pm 1.12$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$	
$3.99 \pm 0.15 \pm 2.64$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.2		³ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$	
$2.5 \pm 1.6 \pm 0.8$		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi \pi$.

³ There is a further $(2.4 \pm 0.8) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$					Γ_{325}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.5 $^{+0.9}_{-0.5}$ OUR AVERAGE					Error includes scale factor of 1.4. See the ideogram below.
$8.00^{+0.12+1.24}_{-0.08-0.40}$		¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	

below.

$8.00^{+0.12+1.24}_{-0.08-0.40}$		¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	
----------------------------------	--	----------------------	-----------	---	--

$11.76 \pm 0.54 \pm 0.94$	1.2k	² DOBBS	15		$J/\psi \rightarrow \gamma K \bar{K}$
$9.62 \pm 0.29^{+3.51}_{-1.86}$		³ BAI	03G	BES	$J/\psi \rightarrow \gamma K \bar{K}$
$5.0 \pm 0.8^{+1.8}_{-0.4}$		^{1,4} BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
$9.2 \pm 1.4 \pm 1.4$		¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$10.4 \pm 1.2 \pm 1.6$		¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$9.6 \pm 1.2 \pm 1.8$		¹ BALTRUSAIT...87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2.3 ± 0.8		⁵ SARANTSEV	21	RVUE	$J/\psi(15) \rightarrow \gamma$ ($\pi\pi, K\bar{K}, \eta\eta, \omega\phi$)
$1.6 \pm 0.2^{+0.6}_{-0.2}$		^{1,6} BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	90	⁷ BISELLO	89B		$J/\psi \rightarrow 4\pi\gamma$
$1.6 \pm 0.4 \pm 0.3$		⁸ BALTRUSAIT...87	MRK3		$J/\psi \rightarrow \gamma\pi^+\pi^-$
3.8 ± 1.6		⁹ EDWARDS	82D	CBAL	$e^+e^- \rightarrow \eta\eta\gamma$

¹ Includes unknown branching fraction to K^+K^- or $K_S^0 K_S^0$. We have multiplied K^+K^- measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Includes unknown branching ratio to K^+K^- or $K_S^0 K_S^0$.

⁴ Assuming $J^P = 2^+$ for $f_0(1710)$.

⁵ There is a further $(6 \pm 2) \times 10^{-4}$ scalar contribution at 1765 MeV.


⁶ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁷ Includes unknown branching fraction to $\rho^0\rho^0$.

⁸ Includes unknown branching fraction to $\pi^+\pi^-$.

⁹ Includes unknown branching fraction to $\eta\eta$.

WEIGHTED AVERAGE
 $9.5 \pm 0.9 - 0.5$ (Error scaled by 1.4)



$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{326} / Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H	BES3 $J/\psi \rightarrow \gamma \omega \omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{327} / Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2 ± 0.4 ² SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² There is a further $(0.7 \pm 0.1) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{328} / Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.5 ± 2.5 ¹ SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ There is a further $(2.5 \pm 1.1) \times 10^{-5}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \phi) / \Gamma_{\text{total}}$ Γ_{329} / Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.5 ± 0.6 OUR AVERAGE				

2.00 $\pm 0.08^{+1.38}_{-1.64}$ 1.3k ABLIKIM 13J BES3 $J/\psi \rightarrow \gamma \omega \phi$

2.61 $\pm 0.27 \pm 0.65$ 95 ABLIKIM 06J BES2 $J/\psi \rightarrow \gamma \omega \phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1 ± 0.1 ¹ SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ There is a further $(2.2 \pm 0.4) \times 10^{-4}$ scalar contribution at 1765 MeV.

$\Gamma(\gamma f_0(1810) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{330} / Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.4 ± 0.8 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi)$

$\Gamma(\gamma f_0(1810) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{331} / Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.11 $\pm 0.06^{+0.19}_{-0.32}$ ABLIKIM 18AA BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(1810) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{332} / Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6 ± 2 SARANTSEV 21 RVUE $J/\psi \rightarrow \gamma(K\bar{K})$

$\Gamma(\gamma f_0(1810) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{333} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.7 ± 0.1	SARANTSEV	21	RVUE $J/\psi \rightarrow \gamma(\eta\eta)$
---------------	-----------	----	--

 $\Gamma(\gamma f_0(1810) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{334} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.11 \pm 0.01^{+0.04}_{-0.03}$	¹ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	------	---

2.5 ± 1.1	SARANTSEV	21	RVUE $J/\psi \rightarrow \gamma(\eta\eta')$
---------------	-----------	----	---

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ *P*-wave. $\Gamma(\gamma f_0(1810) \rightarrow \gamma \omega \phi) / \Gamma_{\text{total}}$ Γ_{335} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.2 ± 0.4	SARANTSEV	21	RVUE $J/\psi \rightarrow \gamma(\omega\phi)$
---------------	-----------	----	--

 $\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{336} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
--------------------------	------	-------------	---------

$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N $J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	----------------------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances. $\Gamma(\gamma \eta_1(1855) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{337} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.70 \pm 0.41^{+0.16}_{-0.35}$	¹ ABLIKIM	22AI	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	------	---

¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ *P*-wave. For analysis details see ABLIKIM 22AS. $\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{338} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma \omega \omega$
--------------------------	-----	---------	-----	---

 $\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)) / \Gamma_{\text{total}}$ Γ_{339} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$0.7 \pm 0.1 \pm 0.2$	BAI	00B	BES $J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
-----------------------	-----	-----	---

 $\Gamma(\gamma f_2(1950) \rightarrow \gamma \gamma \phi) / \Gamma_{\text{total}}$ Γ_{340} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$9.96 \pm 0.60^{+3.44}_{-2.13}$	¹ ABLIKIM	25P	BES3 $J/\psi \rightarrow \gamma \gamma \phi$
---------------------------------	----------------------	-----	--

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma \gamma \phi$ with significance 13.1σ .

$\Gamma(\gamma f_2(2010) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{341} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.71 \pm 0.06^{+0.10}_{-0.06}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ *P*-wave.

 $\Gamma(\gamma f_2(2010) \rightarrow \gamma \gamma \phi) / \Gamma_{\text{total}}$ Γ_{342} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$4.63 \pm 0.43^{+1.42}_{-1.46}$	¹ ABLIKIM	25P BES3	$J/\psi \rightarrow \gamma \gamma \phi$
---------------------------------	----------------------	----------	---

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma \gamma \phi$ with significance 11.3σ .

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{343} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

42 ± 10	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	--------------	------	---

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{344} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

55 ± 25	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	--------------	------	---

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{345} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10 ± 10	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	--------------	------	---

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}}$ Γ_{346} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.63 \pm 0.06^{+0.31}_{-0.46}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
---------------------------------	----------------------	----------	---

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{347} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.28 \pm 0.12^{+0.29}_{-0.20}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ *P*-wave.

 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{348} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.7 \pm 0.5 \pm 0.5$	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
-----------------------	-----------------------------	------	---

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi \pi / \text{total} = 0.167$.

$\Gamma(\gamma f_4(2050) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{349} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.06 \pm 0.01^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{350} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	----------------------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 1.5	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	-----------	----	------	--

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_0(2100) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{351} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

32 ± 20	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
-------------	-----------	----	------	--

$\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{352} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
--------------------------	-----	--------------------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.8	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	-----------	----	------	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200)) / \Gamma_{\text{total}}$ Γ_{353} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	¹ AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	-----------------------	----	-----	---

¹ Includes unknown branching fraction to $K_S^0 K_S^0$.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{354} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$5.86 \pm 0.49 \pm 1.20$	490	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
--------------------------	-----	--------------------	----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.5 ± 0.5	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	-----------	----	------	--

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{355}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.72^{+0.08+0.17}_{-0.06-0.47}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_0(2200) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{356}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
5 ± 2	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2200) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{357}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
0.7 ± 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(2200) \rightarrow \gamma \gamma \phi)/\Gamma_{\text{total}}$ Γ_{358}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$0.20 \pm 0.04^{+0.05}_{-0.07}$	¹ ABLIKIM	25P BES3	$J/\psi \rightarrow \gamma \gamma \phi$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma \gamma \phi$ with significance 6.3σ .

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{359}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
>300			¹ BAI	96B BES	$e^+e^- \rightarrow \gamma \bar{p}p, K\bar{K}$
>250	99.9		² HASAN	96 SPEC	$\bar{p}p \rightarrow \pi^+\pi^-$
< 2.3	95		³ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+K^-$
< 1.6	95		³ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	³ BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+K^-$

¹ Using BARNES 93.

² Using BAI 96B.

³ Includes unknown branching fraction to K^+K^- or $K_S^0 K_S^0$.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{360}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.9	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$14 \pm 8 \pm 4$	BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
$8.4 \pm 2.6 \pm 3.0$	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+\pi^-$ and $\pi^0\pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{361}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 3.6		³ DEL-AMO-SA...100	BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
< 2.9		³ DEL-AMO-SA...100	BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.6 \pm 2.9 \pm 2.4$		BAI	96B	BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
$10.8 \pm 4.0 \pm 3.2$		BAI	96B	BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{362}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.6 \pm 0.5$	BAI	96B	BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \rho \bar{\rho}$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{363}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.95 \pm 0.21 \pm 0.66$ -0.72	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.1 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{364}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4 ± 2	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{365}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.5 ± 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ Γ_{366}/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$6.09 \pm 0.64 \pm 4.00$ -1.68	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ Γ_{367}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.10 \pm 0.02 \pm 0.01$ -0.02	¹ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{368}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.60^{+0.62+2.37}_{-0.65-2.07}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{369}/Γ

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$5.54^{+0.34+3.82}_{-0.40-1.49}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$\Gamma(\gamma f_2(2340) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{370}/Γ

<u>VALUE (units 10^{-6})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.67 \pm 0.70^{+0.61}_{-1.67}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta' X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma f_0(2470) \rightarrow \gamma\eta'\eta')/\Gamma_{\text{total}}$ Γ_{371}/Γ

<u>VALUE (units 10^{-7})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.18 \pm 1.77^{+3.73}_{-2.23}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta' X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

$\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{372}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7^{+0.6}_{-0.8}$		OUR AVERAGE		Error includes scale factor of 1.6.

$3.93 \pm 0.38^{+0.31}_{-0.84}$		¹ ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
---------------------------------	--	----------------------	----------	--

$2.2 \pm 0.4 \pm 0.4$	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
-----------------------	-----	---------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.87 \pm 0.09^{+0.49}_{-0.52}$	4265	² ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
---------------------------------	------	----------------------	----------	--

¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma\eta(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

$\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{373}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.77^{+0.15}_{-0.09} OUR AVERAGE

0.90 ^{+0.04+0.27} _{-0.11-0.55}		1 ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p \bar{p}$
1.14 ^{+0.43+0.42} _{-0.30-0.26}	231	2 ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p \bar{p}$
0.70 \pm 0.04 ^{+0.19} _{-0.08}		BAI	03F BES2	$J/\psi \rightarrow \gamma p \bar{p}$

¹ From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A.

² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

 $\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$ Γ_{374}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

3.31^{+0.33+1.96}_{-0.30-1.29}	ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
---	---------	----------	--

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \phi(1020))/\Gamma_{\text{total}}$ Γ_{375}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.37\pm0.19^{+0.78}_{-1.10}		1 ABLIKIM	25P BES3	$J/\psi \rightarrow \gamma \gamma \phi$
---	--	-----------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.77 \pm 0.35 \pm 0.25	305	2 ABLIKIM	18i BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
8.09 \pm 1.99 \pm 1.36	1.3k	3 ABLIKIM	18i BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma \gamma \phi$ with significance 15.3σ .

² Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma \phi$ invariant mass.

³ Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma \phi$ invariant mass.

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \gamma)/\Gamma_{\text{total}}$ Γ_{376}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<3.56 $\times 10^{-6}$	90	ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
---	----	---------	----------	---

 $\Gamma(\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{377}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.44\pm0.36^{+0.60}_{-0.74}	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$
---	------	---------	----------	--

 $\Gamma(\gamma \eta(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$ Γ_{378}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.79\pm0.23\pm0.65	ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K^+ K^- \eta'$
---	---------	----------	---

 $\Gamma(\gamma \eta(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$ Γ_{379}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.18\pm0.32\pm0.39	ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$
---	---------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.31 \pm 0.22 ^{+2.85} _{-0.84}	1 ABLIKIM	24 BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$
---	-----------	---------	---

¹ Decaying via the intermediate $f_0(980)\eta'$, fitted together with $X(1835)$, a 600 MeV broad structure around 2.8 GeV, and the tail of the $\eta_c(1S)$.

$\Gamma(\gamma\eta(2370) \rightarrow \gamma\eta\eta\eta')/\Gamma_{\text{total}}$						Γ_{380}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<9.2 \times 10^{-6}$	90	ABLIKIM	21C	BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta\eta'$	

$\Gamma(\gamma D^0 + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{381}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<9.1 \times 10^{-8}$	90	ABLIKIM	24BZ	BES3	$e^+e^- \rightarrow J/\psi(1S)$	

$\Gamma(\gamma\rho\bar{\rho})/\Gamma_{\text{total}}$						Γ_{382}/Γ
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.38 \pm 0.07 \pm 0.07$		49	EATON	84	MRK2 e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.11	90		PERUZZI	78	MRK1 e^+e^-	

$\Gamma(\gamma\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{383}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<0.79 \times 10^{-3}$	90	EATON	84	MRK2	e^+e^-	

$\Gamma(\gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$						Γ_{384}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<0.16 \times 10^{-3}$	90	BAI	98G	BES	e^+e^-	

$\Gamma(\gamma A^0 \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$						Γ_{385}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
$<1.7 \times 10^{-6}$	90	88M	¹ ABLIKIM	20K	BES3 $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<6.3 \times 10^{-6}$	90	3.7M	² INSLER	10	CLEO $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$	

¹ For a narrow state, A^0 , with mass $m_{A^0} < 1.2$ GeV. The limit varies with m_{A^0} , reaching its largest value of 1.7×10^{-6} at 1.2 GeV and being 7.0×10^{-7} for $m_{A^0} = 0$.

² The limit varies with mass m_{A^0} of a narrow state A^0 and is 4.3×10^{-6} for $m_{A^0} = 0$, reaches its largest value of 6.3×10^{-6} at $m_{A^0} = 500$ MeV, and is 3.6×10^{-6} at $m_{A^0} = 960$ MeV.

$\Gamma(\gamma A^0 \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$						Γ_{386}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<4.9 \times 10^{-7}$	95	¹ ABLIKIM	24AD	BES3	$J/\psi \rightarrow \gamma\gamma\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<1.8 \times 10^{-6}$	95	² ABLIKIM	23E	BES3	$J/\psi \rightarrow \gamma\gamma\gamma$	

¹ For a light pseudoscalar axion-like particle, A^0 , with a mass in the range 0.18–2.85 GeV. The measured 95% CL limit as a function of m_{A^0} ranges from 3.7×10^{-8} to 4.85×10^{-7} .

² For a light pseudoscalar axion-like particle, A^0 , with a mass in the range 0.165–2.84 GeV. The measured 95% CL limit as a function of m_{A^0} ranges from 8.3×10^{-8} to 1.8×10^{-6} .

$$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-) / \Gamma_{\text{total}} \quad \Gamma_{387} / \Gamma$$

(narrow state A^0 with $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.8 \times 10^{-7}$	90	¹ ABLIKIM	22H BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
$< 0.5 \times 10^{-5}$	90	² ABLIKIM	16E BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
$< 2.1 \times 10^{-5}$	90	³ ABLIKIM	12 BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3.0 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(1.2\text{--}778.0) \times 10^{-9}$.

² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

³ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

———— DALITZ DECAYS ————

$$\Gamma(\pi^0 e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{388} / \Gamma$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.06 \pm 0.31 \pm 0.38$		¹ ABLIKIM	25AD BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM	14I BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Includes the $\rho - \omega$ resonant contribution.

$$\Gamma(\eta e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{389} / \Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.04 \pm 0.07$	2.47k	^{1,2} ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+ e^-$
$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM	14I BES3	$J/\psi \rightarrow \eta e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

² Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.56 \pm 0.04 \pm 0.03$ GeV. Supersedes ABLIKIM 14I.

$$\Gamma(\eta'(958) e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{390} / \Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.59 \pm 0.07 \pm 0.17$	8.9k	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$
$5.81 \pm 0.16 \pm 0.31$	1.4k	^{1,2} ABLIKIM	14I BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using both $\eta' \rightarrow \gamma \pi^+ \pi^-$ and $\eta' \rightarrow \pi^+ \pi^- \eta$ decays.

² Superseded by ABLIKIM 19H.

$$\Gamma(\eta(1405) e^+ e^- \rightarrow f_0(980) \pi^0 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{391} / \Gamma$$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.04 \pm 0.20 \pm 0.08$	203	¹ ABLIKIM	24I BES3	$J/\psi \rightarrow e^+ e^- \eta(1405)$

¹ With a significance of 9.8σ .

$\Gamma(X(1835)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{392}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.58 \pm 0.19 \pm 0.16$	1364	¹ ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta' e^+e^-$

¹ Assuming constructive interference. Destructive interference gives a value of $(4.43 \pm 0.23 \pm 0.19) \times 10^{-6}$ for this branching fraction.

$\Gamma(X(2120)e^+e^-, X \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{393}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.82 \pm 0.12 \pm 0.06$	310	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta' e^+e^-$

$\Gamma(\eta(2370)e^+e^-, \eta \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{394}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.08 \pm 0.14 \pm 0.10$	397	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+\pi^-\eta' e^+e^-$

$\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{395}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.11 \times 10^{-7}$	90	¹ ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+e^-$

¹ For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .

$\Gamma(\eta'(958)U \rightarrow \eta'(958)e^+e^-)/\Gamma_{\text{total}}$ Γ_{396}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.0 \times 10^{-7}$	90	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+e^-$

¹ For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .

$\Gamma(\phi e^+e^-)/\Gamma_{\text{total}}$ Γ_{397}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	¹ ABLIKIM	19AB BES3	$J/\psi \rightarrow \phi e^+e^-$

¹ Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.49 \pm 0.30)\%$.

———— WEAK DECAYS ————

$\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{398}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.1 \times 10^{-8}$	90	ABLIKIM	21Q BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.2 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+e^- \rightarrow J/\psi$
------------------------	----	---------	----------	-----------------------------

$\Gamma(D^- \mu^+ \nu_\mu + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{399}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-7}$	90	¹ ABLIKIM	24AMBES3	$e^+e^- \rightarrow J/\psi$

¹ Using $B(D^- \rightarrow K^+\pi^-\pi^-) = 9.38 \pm 0.16 \%$.

$\Gamma(\overline{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{400}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<8.5 \times 10^{-8}$	90	¹ ABLIKIM	17AF BES3	$e^+ e^- \rightarrow J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<1.1 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$
¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.				

$\Gamma(D^0 \mu^+ \mu^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{401}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.1 \times 10^{-7}$	90	¹ ABLIKIM	25AH BES3	$e^+ e^- \rightarrow J/\psi$
¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.				

$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{402}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.3 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<3.6 \times 10^{-5}$	90	¹ ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$
¹ Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5 \%$.				

$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{403}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.8 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{404}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<7.0 \times 10^{-8}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$<7.5 \times 10^{-5}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(D^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{405}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.0 \times 10^{-7}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\overline{D}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{406}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<4.7 \times 10^{-7}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\overline{D}^0 \overline{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{407}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<1.7 \times 10^{-4}$	90	ABLIKIM	08J BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\overline{D}^0 \overline{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{408}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.5 \times 10^{-6}$	90	ABLIKIM	14K BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\overline{D}^0 \eta + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{409}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<6.8 \times 10^{-7}$	90	ABLIKIM	24BI BES3	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\bar{D}^0 \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{410}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-7}$	90	ABLIKIM 24BI	BES3	$e^+e^- \rightarrow J/\psi$

$\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{411}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.1 \times 10^{-7}$	90	ABLIKIM 25CG	BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.3 \times 10^{-4}$	90	¹ ABLIKIM 08J	BES2	$e^+e^- \rightarrow J/\psi$
-----------------------	----	--------------------------	------	-----------------------------

¹ Superseded by ABLIKIM 25CG.

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{412}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.0 \times 10^{-7}$	90	ABLIKIM 25CG	BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.3 \times 10^{-5}$	90	ABLIKIM 14K	BES3	$e^+e^- \rightarrow J/\psi$
-----------------------	----	-------------	------	-----------------------------

————— CHARGE CONJUGATION (C), PARITY (P), —————
 ————— LEPTON FAMILY NUMBER (LF) VIOLATING MODES —————

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{413}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7 \times 10^{-7}$	90	ABLIKIM 14Q	BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.5 \times 10^{-5}$	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
-----------------------	----	----------	------	--

$<1.6 \times 10^{-4}$	90	¹ WICHT 08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
-----------------------	----	-----------------------	------	--

$<2.2 \times 10^{-5}$	90	ABLIKIM 07J	BES2	$\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
-----------------------	----	-------------	------	--

$<50 \times 10^{-5}$	90	BARTEL 77	CNTR	e^+e^-
----------------------	----	-----------	------	----------

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] < 0.16 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(B^+ \rightarrow J/\psi(1S)K^+) = 1.019 \times 10^{-3}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ Γ_{414}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	ABLIKIM 14Q	BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{415}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-7}$	90	ABLIKIM 13L	BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-6}$	90	BAI 03D	BES	$e^+e^- \rightarrow J/\psi$
-----------------------	----	---------	-----	-----------------------------

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{416}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-8}$	90	ABLIKIM 21M	BES3	$e^+e^- \rightarrow J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8.3 \times 10^{-6}$	90	¹ ABLIKIM 04	BES	$e^+e^- \rightarrow J/\psi$
-----------------------	----	-------------------------	-----	-----------------------------

¹ Superseded by ABLIKIM 21M.

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$					Γ_{417}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	ABLIKIM 04	BES	$e^+ e^- \rightarrow J/\psi$	

$\Gamma(K^+ K^+ e^- e^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{418}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.1 \times 10^{-9}$	90	ABLIKIM 26	BES3	$e^+ e^- \rightarrow J/\psi$	

$\Gamma(pe^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{419}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.1 \times 10^{-8}$	90	¹ ABLIKIM 25AA	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi(1S)$	

¹ Using $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi(1S)$ branching fraction from PDG 24.

$\Gamma(\Lambda_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{420}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.9 \times 10^{-8}$	90	ABLIKIM 19AF	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow pK^- \pi^+ e^- (+ \text{c.c.})$	

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$					Γ_{421}/Γ_5
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.6 \times 10^{-2}$	90	LEES 13I	BABR	$B \rightarrow K^{(*)} J/\psi$	

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$					Γ_{421}/Γ_7
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-2}$	90	ABLIKIM 08G	BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

$\Gamma(\mu^+ \mu^- X^0 \rightarrow \mu^+ \mu^- + \text{invisible})/\Gamma_{\text{total}}$					Γ_{422}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.5 \times 10^{-7}$	90	¹ ABLIKIM 24G	BES3	$J/\psi \rightarrow \mu^+ \mu^- X^0$ (scalar) $\rightarrow \mu^+ \mu^- + \text{invisible}$	
$<9.6 \times 10^{-7}$	90	² ABLIKIM 24G	BES3	$J/\psi \rightarrow \mu^+ \mu^- X^0$ (vector) $\rightarrow \mu^+ \mu^- + \text{invisible}$	

¹ For a light scalar, X^0 , with a mass in the range 1–1000 MeV. The measured limit at the 90% credibility level as a function of m_{X^0} ranges from 6.2×10^{-9} to 5.5×10^{-7} .

² For a light vector, X^0 , with a mass in the range 1–1000 MeV. The measured limit at the 90% credibility level as a function of m_{X^0} ranges from 4.5×10^{-9} to 9.6×10^{-7} .

$\Gamma(\phi + \text{invisible } X)/\Gamma_{\text{total}}$					Γ_{423}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<7.0 \times 10^{-8}$	90	¹ ABLIKIM 25BS	BES3	m_X range: 0–0.96 GeV	
-----------------------	----	---------------------------	------	-------------------------	--

¹ For an invisible particle X with a mass in the interval 0–0.96 GeV, the measured 90% CL limit as a function of the mass ranges from 4×10^{-9} to 4×10^{-8} .

$J/\psi(1S)$ REFERENCES

ABLIKIM	26	CP C50 013107	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25AA	PR D111 112010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25AB	PR D112 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25AD	PR D112 L011101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25AH	JHEP 2504 061	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25BS	PRL 135 151804	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25BW	PR D112 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25CC	PR D112 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25CG	JHEP 2512 077	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25P	PR D111 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	24AE	JHEP 2412 062	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	24	PRL 132 181901	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AB	PR D110 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AD	PR D110 L031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AM	JHEP 2401 126	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BI	PR D110 032020	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BQ	PR D110 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BV	PR D110 072021	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D111 099902 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BZ	PR D110 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CB	PR D110 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24G	PR D109 L031102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24H	PR D109 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24I	PR D109 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24L	PR D109 052006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAYRAPETY... PDG	24A	PR D109 L111101	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)
	24	PR D110 030001	S. Navas <i>et al.</i>	(PDG Collab.)
ABLIKIM	23BD	PR D108 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BU	PR D108 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23E	PL B838 137698	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23M	JHEP 2303 121	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23S	PR D107 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	23	JHEP 2306 196	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
GONG	23	PR D107 072008	G. Gong <i>et al.</i>	(BELLE Collab.)
LEES	23	PR D107 072001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LIAO	23	PR D107 112007	L. Liao <i>et al.</i>	
ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	22AI	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22B	PRL 129 022002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22H	PR D105 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	22	EPJ C82 938	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
ABLIKIM	21AM	PR D104 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21M	PR D103 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Q	JHEP 2106 157	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20K	PR D101 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	20	JHEP 2007 112	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D104 099901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13L	PR D87 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT, B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
SETH	04	PR D69 097503	K.K. Seth	
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)

BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)
BALTRUSAIT...	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)
		Translated from YAF 41 733.		
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
	Also	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)
	Also	SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)
		Translated from YAF 34 1471.		

BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG
BALDINI-...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)
