

$T_{c\bar{c}1}(3900)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

was $Z_c(3900)$, $X(3900)$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Charged $T_{c\bar{c}1}(3900)^\pm$ seen as a peak in the invariant mass distribution of the $J/\psi\pi^\pm$ system by BES III (ABLIKIM 13T) in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at c.m. energy of 4.26 GeV and by radiative return from e^+e^- collisions at \sqrt{s} from 9.46 to 10.86 GeV at Belle (LIU 13B). Partial wave analysis of ABLIKIM 17J determines $J^P = 1^+$ with more than 7σ significance. Neutral $T_{c\bar{c}1}(3900)^0$ seen in the $J/\psi\pi^0$ invariant mass distribution in $e^+e^- \rightarrow \pi^0\pi^0J/\psi$ at c.m. energies of 4.23, 4.26, and 4.36 GeV by BES III (ABLIKIM 15U) and at 4.17 GeV by XIAO 13A. Peaks in $(D\bar{D}^*)^{0,\pm}$ reported by BES III (ABLIKIM 14A, ABLIKIM 15AB) are assumed to be related.

$T_{c\bar{c}1}(3900)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3886.7 ± 2.3	OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.			
$3884.6 \pm 0.7 \pm 3.3$	3.3	¹ ABLIKIM	25BU BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
$3893.1 \pm 2.2 \pm 3.0$	3.0	¹ ABLIKIM	20N BES3	0	$e^+e^- \rightarrow \pi^0\pi^0J/\psi$
$3902.6^{+5.2+3.3}_{-5.0-1.4}$	3.3	^{2,3} ABAZOV	19 D0	\pm	1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
$3885.7^{+4.3}_{-5.7} \pm 8.4$	8.4	^{2,4} ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
$3881.7 \pm 1.6 \pm 1.6$	1.2k	^{2,4} ABLIKIM	15AC BES3	\pm	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
$3883.9 \pm 1.5 \pm 4.2$	1.2k	^{2,4} ABLIKIM	14A BES3	\pm	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
$3894.5 \pm 6.6 \pm 4.5$	159	² LIU	13B BELL	\pm	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
$3886 \pm 4 \pm 2$	81	^{2,5} XIAO	13A	\pm	4.17 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$
$3904 \pm 9 \pm 5$	25	^{2,5} XIAO	13A	0	4.17 $e^+e^- \rightarrow \pi^0\pi^0J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3895.0 \pm 5.2^{+4.0}_{-2.7}$	502	^{2,6} ABAZOV	18B D0	\pm	1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
$3881.2 \pm 4.2 \pm 52.7$	6k	^{4,7} ABLIKIM	17J BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
$3894.8 \pm 2.3 \pm 3.2$	356	^{2,8} ABLIKIM	15U BES3	0	$e^+e^- \rightarrow \pi^0\pi^0J/\psi$
$3899.0 \pm 3.6 \pm 4.9$	307	^{2,9} ABLIKIM	13T BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$

¹ Pole mass obtained from a fit to a relativistic Breit-Wigner.

² Neglecting interference between the $T_{c\bar{c}1}(3900)$ and other processes.


³ Measured in weak decays of b -flavored hadrons (nonprompt).

⁴ Pole mass obtained from a fit to a Flatte-like formula.

⁵ For $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

⁶ The signal of the $T_{c\bar{c}1}(3900)$ is correlated with a parent $J/\psi\pi^+\pi^-$ system in the invariant mass range 4.2–4.7 GeV. Superseded by ABAZOV 19.

- ⁷ Superseded by ABLIKIM 25BU.
- ⁸ Superseded by ABLIKIM 20N.
- ⁹ Superseded by ABLIKIM 17J.



$T_{c\bar{c}1}(3900)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
29.5± 2.4 OUR AVERAGE					
37.2± 1.3± 6.6		¹ ABLIKIM 25BU BES3	±		$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
44.4± 5.2± 14.0		¹ ABLIKIM 20N BES3	0		$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
32 ⁺²⁸ ₋₂₁ ⁺²⁶ ₋₇		^{2,3} ABAZOV 19 D0	±		1.96 TeV $p\bar{p} \rightarrow \pi^+\pi^- J/\psi X$ (non-prompt)
35 ⁺¹¹ ₋₁₂ ± 15		^{2,4} ABLIKIM 15AB BES3	0		$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
26.6± 2.0± 2.1	1248	^{2,4} ABLIKIM 15AC BES3	±		$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
24.8± 3.3± 11.0	1212	^{2,4} ABLIKIM 14A BES3	±		$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
63 ± 24 ± 26	159	² LIU 13B BELL	±		$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
37 ± 4 ± 8	81	^{2,5} XIAO 13A	±		4.17 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
51.8± 4.6± 36.0	6 k	^{4,6} ABLIKIM 17J BES3	±		$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
29.6± 8.2± 8.2	356	^{2,7} ABLIKIM 15U BES3	0		$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
46 ± 10 ± 20	307	^{2,8} ABLIKIM 13T BES3	±		$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ Pole width obtained from a fit to a relativistic Breit-Wigner.
² Neglecting interference between the $T_{c\bar{c}1}(3900)$ and other processes.

³ Measured in weak decays of b -flavored hadrons (nonprompt).

⁴ Pole width obtained from a fit to a Flatte-like formula.

⁵ For $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

⁶ Superseded by ABLIKIM 25BU.

⁷ Superseded by ABLIKIM 20N.

⁸ Superseded by ABLIKIM 17J.

$T_{c\bar{c}1}(3900)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\pi$	seen
Γ_2 $h_c\pi^\pm$	not seen
Γ_3 $\eta_c\pi^+\pi^-$	not seen
Γ_4 $\eta_c(1S)\rho(770)^\pm$	seen
Γ_5 $(DD^*)^\pm$	seen
Γ_6 $D^0D^{*-} + \text{c.c.}$	seen
Γ_7 $D^-D^{*0} + \text{c.c.}$	seen
Γ_8 $\omega\pi^\pm$	not seen
Γ_9 $J/\psi\eta$	not seen
Γ_{10} $D^+D^{*-} + \text{c.c.}$	seen
Γ_{11} $D^0\bar{D}^{*0} + \text{c.c.}$	seen

$T_{c\bar{c}1}(3900)$ BRANCHING RATIOS

$\Gamma(J/\psi\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	20N	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
seen		¹ ABAZOV	19	D0	\pm	$1.96 \text{ TeV } \rho\bar{p} \rightarrow \pi^+\pi^- J/\psi X$ (prompt)
seen		ABLIKIM	17J	BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
seen	356	ABLIKIM	15U	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
not seen		² ADOLPH	15D	COMP	\pm	$\gamma N \rightarrow J/\psi\pi^\pm N$
seen	307	ABLIKIM	13T	BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
seen	25	³ XIAO	13A		0	$4.17 e^+e^- \rightarrow \pi^0\pi^0 J/\psi$

¹ But not seen in the "prompt" sample (no b -hadron enhancement).

² ADOLPH 15D measure $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi\pi^\pm) \sigma(\gamma N \rightarrow T_{c\bar{c}1}(3900)^\pm N)/\sigma(\gamma N \rightarrow J/\psi N) < 3.7 \times 10^{-3}$ at 90% CL.

³ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(h_c\pi^\pm)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	13X	BES3	\pm	$e^+e^- \rightarrow h_c\pi^+\pi^-$

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_3/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		¹ VINOKUROVA	15	BELL	0	$B^+ \rightarrow K^+\eta_c\pi^+\pi^-$

¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ Z_c(3900)^0) \times B(X \rightarrow \eta_c \pi^+ \pi^-) < 4.7 \times 10^{-5}$ at 90% CL.

$\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(J/\psi\pi)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.3±0.8	332	¹ ABLIKIM	19BC BES3		$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c(1S)$

¹ Using $e^+e^- \rightarrow \pi^\mp(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi\pi^\pm)$ cross section at 4.23 and 4.26 GeV from ABLIKIM 17J.

$\Gamma((D\bar{D}^*)^\pm)/\Gamma(J/\psi\pi)$ Γ_5/Γ_1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
6.2±1.1±2.7	¹ ABLIKIM	14A BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$

¹ Assuming the same origin of the $(D\bar{D}^*)^\pm$ and $\pi^\pm J/\psi$ decay modes.

$\Gamma(D^0 D^{*-} + c.c.)/\Gamma_{total}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AC BES3	±	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + c.c.$
seen	ABLIKIM	14A BES3	±	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + c.c.$

$\Gamma(D^- D^{*0} + c.c.)/\Gamma_{total}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AC BES3	±	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + c.c.$
seen	ABLIKIM	14A BES3	±	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + c.c.$

$\Gamma(\omega\pi^\pm)/\Gamma_{total}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
not seen	ABLIKIM	15R BES3	±	$e^+e^- \rightarrow \omega\pi^+\pi^-$

$\Gamma(J/\psi\eta)/\Gamma_{total}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
not seen	ABLIKIM	15Q BES3	0	4.0–4.6 $e^+e^- \rightarrow J/\psi\eta\pi^0$

$\Gamma(J/\psi\eta)/\Gamma(J/\psi\pi)$ Γ_9/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.15	90	ABLIKIM	15Q BES3	0	4.226 $e^+e^- \rightarrow J/\psi\eta\pi^0$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.65	90	ABLIKIM	15Q BES3	0	4.257 $e^+e^- \rightarrow J/\psi\eta\pi^0$
-------	----	---------	----------	---	--

$\Gamma(D^+ D^{*-} + c.c.)/\Gamma_{total}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

$\Gamma(D^0\bar{D}^{*0} + c.c.)/\Gamma_{total}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

$\Gamma(D^+ D^{*-} + c.c.)/\Gamma(D^0\bar{D}^{*0} + c.c.)$ Γ_{10}/Γ_{11}

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.96±0.18±0.12	ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

$T_{c\bar{c}1}(3900)$ REFERENCES

ABLIKIM	25BU	PR D112 092013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20N	PR D102 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP
ABAZOV	19	PR D100 012005	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	19BC	PR D100 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABAZOV	18B	PR D98 052010	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	17J	PRL 119 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP
ABLIKIM	15AB	PRL 115 222002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15R	PR D92 032009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15U	PRL 115 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADOLPH	15D	PL B742 330	C. Adolph <i>et al.</i>	(COMPASS Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also		JHEP 1702 088 (errat.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)
ABLIKIM	14A	PRL 112 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
XIAO	13A	PL B727 366	T. Xiao <i>et al.</i>	(NWES)