

$\eta_c(2S)$

$$J^{PC} = 0^+(0^-+)$$

Quantum numbers are quark model predictions.

 $\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3637.8 ± 0.6	OUR AVERAGE	Error includes scale factor of 1.1.		
3637.8 ± 0.8 ± 0.2	1.6k	ABLIKIM	24J BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$
3637.90 ± 0.54 ± 1.40	3.7k	AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$
3643.4 ± 2.3 ± 4.4	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
3635.1 ± 3.7 ± 2.9	106	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
3633.6 ± 1.7 ± 0.6	106	¹ AAIJ	17AD LHCb	$pp \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$
3636.4 ± 4.1 ± 0.7	365	² AAIJ	17BB LHCb	$pp \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-) X$
3637.0 ± 5.7 ± 3.4	178	^{3,4} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^0$
3635.1 ± 5.8 ± 2.1	47	^{3,5} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \eta$
3646.9 ± 1.6 ± 3.6	57	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6 ± 2.9 ± 1.6	127	⁶ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
3638.5 ± 1.5 ± 0.8	624	³ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5 ± 3.2 ± 2.5	1201	³ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 ^{+3.9} _{-4.2} ^{+0.7} _{-2.0}	128	⁷ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	⁸ ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c \bar{c})$
3645.0 ± 5.5 ^{+4.9} _{-7.8}	121	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

3639 ± 7	98	⁹ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c \bar{c}}$
3630.8 ± 3.4 ± 1.0	112	¹⁰ AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
3654 ± 6 ± 8	39	¹¹ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		¹² EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ AAIJ 17AD report $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$ MeV. We use the current value $m_{\psi(2S)} = 3686.097 \pm 0.025$ MeV to obtain the quoted mass.

² From a fit of the $\phi\phi$ invariant mass with the width of $\eta_c(2S)$ fixed to the PDG 16 value.

³ Ignoring possible interference with continuum.

⁴ With a width fixed to 11.3 MeV.

⁵ With a width fixed to 11.3 MeV. Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

⁷ Accounts for interference with non-resonant continuum.

⁸ From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

⁹ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁰ Superseded by DEL-AMO-SANCHEZ 11M.

¹¹ Superseded by VINOKUROVA 11.

¹² Assuming mass of $\psi(2S) = 3686$ MeV.

$\eta_c(2S)$ WIDTH

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
11.6 ± 1.4	OUR AVERAGE			
10.5 ± 1.7 ± 3.5	1.6k	ABLIKIM	24J BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$
10.77 ± 1.62 ± 1.08	3.7k	AAIJ	23AH LHCb	$B^+ \rightarrow K^+ (K_S^0 K \pi)$
19.8 ± 3.9 ± 3.1	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
9.9 ± 4.8 ± 2.9	57	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 ± 6.4 ± 4.8	127	¹ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
13.4 ± 4.6 ± 3.2	624	² DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
6.6 + 8.4 + 2.6 - 5.1 - 0.9	128	³ VINOKUROVA 11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
6.3 ± 12.4 ± 4.0	61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta_c' \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90 98	⁴ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
22 ± 14	121	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$
17.0 ± 8.3 ± 2.5	112	⁵ AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
< 55	90 39	⁶ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
< 8.0	95	⁷ EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

¹ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

² Ignoring possible interference with continuum.

³ Accounts for interference with non-resonant continuum.

⁴ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

⁵ Superseded by DEL-AMO-SANCHEZ 11M.

⁶ For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11.

⁷ For a mass value of 3594 ± 5 MeV

$\eta_c(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $K \bar{K} \pi$	$(1.9_{-1.0}^{+1.2}) \%$	
Γ_3 $K \bar{K} \eta$	$(7_{-4}^{+5}) \times 10^{-3}$	
Γ_4 $K \bar{K} \eta'$	$(2.1 \pm 1.6) \times 10^{-3}$	
Γ_5 $2\pi^+ 2\pi^-$	$< 2.6 \times 10^{-3}$	90%
Γ_6 $a_0(1450)\pi$	seen	
Γ_7 $a_2(1700)\pi$	seen	

Γ_8	$a_0(1710)\pi$	seen	
Γ_9	$\rho^0\rho^0$	< 2.3	$\times 10^{-3}$ 90%
Γ_{10}	$3\pi^+3\pi^-$	$(1.7^{+0.9}_{-1.1})$	%
Γ_{11}	$K^+K^-\pi^+\pi^-$	< 2.5	$\times 10^{-3}$ 90%
Γ_{12}	$K^{*0}\bar{K}^{*0}$	< 4	$\times 10^{-3}$ 90%
Γ_{13}	$K^+K^-\pi^+\pi^-\pi^0$	$(1.5^{+1.0}_{-0.9})$	%
Γ_{14}	$K^+K^-\pi^+\pi^-$	< 1.8	% 90%
Γ_{15}	$K_S^0K^-\pi^+\pi^- + \text{c.c.}$	$(1.4^{+0.8}_{-1.0})$	%
Γ_{16}	$2K^+2K^-$	< 1.4	$\times 10^{-3}$ 90%
Γ_{17}	$K_2^*(1430)\bar{K} + \text{c.c.}$	seen	
Γ_{18}	$K_0^*(1950)\bar{K} + \text{c.c.}$	seen	
Γ_{19}	$K_0^*(2600)\bar{K} + \text{c.c.}$	seen	
Γ_{20}	$\phi\phi$	< 1.4	$\times 10^{-3}$ 90%
Γ_{21}	$\omega\omega$	< 1.9	$\times 10^{-3}$ 90%
Γ_{22}	$\omega\phi$	< 3.4	$\times 10^{-4}$ 90%
Γ_{23}	$\rho\bar{\rho}$	< 3.2	$\times 10^{-4}$ 90%
Γ_{24}	$\rho\bar{\rho}\pi^+\pi^-$	seen	
Γ_{25}	$\rho\bar{\rho}K^+K^-$	(3.6 ± 2.9)	$\times 10^{-4}$
Γ_{26}	$\gamma\gamma$	$(1.8^{+1.0}_{-1.1})$	$\times 10^{-4}$
Γ_{27}	$\gamma J/\psi(1S)$	< 1.8	% 90%
Γ_{28}	$\pi^+\pi^-\eta$	$(5.5^{+3.3}_{-4.0})$	$\times 10^{-3}$
Γ_{29}	$2\pi^+2\pi^-\eta$	$(1.6^{+0.9}_{-1.0})$	%
Γ_{30}	$\pi^+\pi^-\eta'$	$(2.7^{+2.0}_{-1.8})$	$\times 10^{-3}$
Γ_{31}	$\pi^+\pi^-\eta_c(1S)$	< 4	% 90%

FIT INFORMATION

A multiparticle fit to $\eta_c(2S)$ and $\psi(2S)$ with 4 branching ratios uses 5 measurements to determine 3 parameters. The overall fit has a $\chi^2 = 2.6$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

$$x_3 \begin{vmatrix} & 92 \\ & x_2 \end{vmatrix}$$

$\eta_c(2S)$ PARTIAL WIDTHS $\Gamma(\gamma\gamma)$ Γ_{26}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.44 ± 0.14	106	¹ XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
1.3 ± 0.6		² ASNER	04	CLEO $\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

¹ Assuming that the branching fraction into $\eta'\pi^+\pi^-$ is the same as for $\eta_c(1S)$.

² They measure $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.

 $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2 \Gamma_{26}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
39 ± 6				OUR AVERAGE
$41 \pm 4 \pm 6$	624	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
$33.6 \pm 7.2 \pm 8.1$		¹ NAKAZAWA	08	BELL $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

¹ NAKAZAWA 08 reports $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp) \times \Gamma(\gamma\gamma) = 11.2 \pm 2.4 \pm 2.7$ eV which we multiplied by 3 to account for isospin symmetry.

 $\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5 \Gamma_{26}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.5	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

 $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{11} \Gamma_{26}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{13} \Gamma_{26}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$30 \pm 6 \pm 5$	1201	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

 $\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{16} \Gamma_{26}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+K^-)$

 $\Gamma(\pi^+\pi^-\eta') \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{30} \Gamma_{26}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$5.6^{+1.2}_{-1.1} \pm 1.1$	106	XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

 $\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{31} \Gamma_{26}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<133	90	LEES	12AE	BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(\bar{p}p)/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{23}/\Gamma \times \Gamma_{26}/\Gamma$

VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90	^{1,2,3} AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	90	^{1,2,4} AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
< 12.0	90	^{2,4} AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

¹ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

² For a total width $\Gamma=5$ MeV.

³ For the resonance mass region 3589–3599 MeV/ c^2 .

⁴ For the resonance mass region 3575–3660 MeV/ c^2 .

$\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ABREU	98O DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
seen	¹ EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

¹ For a mass value of 3594 ± 5 MeV

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.9^{+1.2}_{-1.0}$ OUR FIT				
$1.9 \pm 0.4 \pm 1.1$	59 ± 12	¹ AUBERT	08AB BABR	$B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	127 ± 18	ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$
seen	39 ± 11	² CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$

¹ Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$.

² For a mass value of 3654 ± 6 MeV

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$ Γ_3/Γ_2

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
39 ± 10 OUR FIT Error includes scale factor of 1.3.				
$27.3 \pm 7.0 \pm 9.0$	225	¹ LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$

¹ LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$, which we divide by 3 to account for isospin symmetry.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

$\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

$\Gamma(a_2(1700)\pi)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

$\Gamma(a_0(1710)\pi)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
----------	---------	----------	---

$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	---------	---------------------------------------

$\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
----------	---------	----------	---

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$ Γ_{13}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.73 \pm 0.17 \pm 0.17$	1201	¹ DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
--------------------------	------	-----------------------------------	--

¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M. We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K \bar{K} \pi)$.

$\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
------	-------------	---------	----------	---

$\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	---------	---------------------------------------

$\Gamma(K_2^*(1430)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

 $\Gamma(K_0^*(1950)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

 $\Gamma(K_0^*(2600)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + \text{c.c.}$.

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ABLIKIM 11H BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 106 ¹ AAIJ 17AD LHCb $pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$

¹ AAIJ 17AD report a 6.4 standard deviation signal, with $B(B^+ \rightarrow \eta_c(2S)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$.

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 110 ¹ CHILIKIN 19 BELL $e^+e^- \rightarrow \Upsilon(4S)$

¹ CHILIKIN 19 reports signals in $B^+ \rightarrow \eta_c(2S)K^+$ and $B^0 \rightarrow \eta_c(2S)K_S^0$ with 12.3 and 5.9 standard deviations, respectively.

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4 $\times 10^{-4}$ 90 ¹ WICHT 08 BELL $B^\pm \rightarrow K^\pm \gamma\gamma$

not seen AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$

<0.01 90 LEE 85 CBAL $\psi' \rightarrow \text{photons}$

¹ WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$ which we divide by our best (shown rounded) value $B(B^+ \rightarrow \eta_c(2S)K^+) = 4.4 \times 10^{-4}$.

 $\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma(K\bar{K}\pi)$ Γ_{31}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.33 90 ¹ LEES 12AE BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

¹ We divided the reported limit by 3 to take into account isospin relations.

$\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

$$\Gamma(\eta_c(2S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_2 / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma \psi(2S)$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.03 ± 0.12 OUR FIT				Error includes scale factor of 1.2.
1.00 ± 0.10 OUR AVERAGE				
0.97 ± 0.06 ± 0.09	1.6k	ABLIKIM	24J BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$
1.30 ± 0.20 ± 0.30	127	ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma \eta_c \rightarrow \gamma K \bar{K} \pi$

$$\Gamma(\eta_c(2S) \rightarrow K \bar{K} \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_3 / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma \psi(2S)$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.0 OUR FIT					Error includes scale factor of 1.3.
4.78 ± 0.64 ± 0.68		362	¹ ABLIKIM	24BWBES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$
• • •					We do not use the following data for averages, fits, limits, etc. • • •
<11.8	90		² CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$
¹ ABLIKIM 24BW reports a value of $(2.39 \pm 0.32 \pm 0.34) \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 to account for isospin symmetry.					
² CRONIN-HENNESSY 10 reports a limit of $< 5.9 \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 account for isospin symmetry. It assumes $\Gamma(\eta_c(2S)) = 14$ MeV. It also gives the analytic dependence of limits on width.					

$$\Gamma(\eta_c(2S) \rightarrow K \bar{K} \eta') / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_4 / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma \psi(2S)$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
11.56 ± 4.86 ± 1.9	¹ ABLIKIM	25I BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'$
¹ ABLIKIM 25I reports a value of $(5.78 \pm 2.43 \pm 0.95) \times 10^{-7}$ for $K^+ K^- \eta'$ which we multiply by 2 to account for isospin symmetry.			

$$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_5 / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma \psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.43 × 10⁻⁶	90	ABLIKIM	24BT BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
<14.6 × 10 ⁻⁶	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.				

$$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_9 / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma \psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 12.7 × 10⁻⁷	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{10} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.2±1.0±1.2		569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13.2	90		¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
-------	----	--	-----------------------------	------	---

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{11} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.6 × 10 ⁻⁶	90		¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
-------------------------	----	--	-----------------------------	------	---

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{12} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<19.6 × 10 ⁻⁷	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{13} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<43.0 × 10 ⁻⁶	90		¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$
--------------------------	----	--	-----------------------------	------	---

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{14} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<9.7 × 10 ⁻⁶	90		¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{15} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.9 ± 1.8 OUR AVERAGE					
9.31 ± 0.72 ± 2.77		3140	ABLIKIM	24Q BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
7.03 ± 2.10 ± 0.7	60		ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.2 90 ¹ CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + c.c.$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \qquad \Gamma_{20}/\Gamma \times \Gamma_{200}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 7.8 × 10⁻⁷	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \qquad \Gamma_{23}/\Gamma \times \Gamma_{200}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.4 × 10⁻⁷	90	ABLIKIM	25H BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.4 × 10⁻⁶ 90 ¹ ABLIKIM 13V BES3 $\psi(2S) \rightarrow \gamma p\bar{p}$

¹ Superseded by ABLIKIM 25H.

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p}K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \qquad \Gamma_{25}/\Gamma \times \Gamma_{200}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10 ⁻⁷)	EVTS	DOCUMENT ID	TECN	COMMENT
1.98 ± 0.41 ± 0.99	84	ABLIKIM	25V BES3	$e^+e^- \rightarrow \psi(2S)$

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \qquad \Gamma_{27}/\Gamma \times \Gamma_{200}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 9.7 × 10⁻⁶	90	33	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \qquad \Gamma_{28}/\Gamma \times \Gamma_{200}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10 ⁻⁶)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.97 ± 0.81 ± 0.26		106	ABLIKIM	23Q BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.3 90 ¹ CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow 2\pi^+2\pi^-\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}} \qquad \Gamma_{29}/\Gamma \times \Gamma_{200}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
0.85 ± 0.04 ± 0.13	565	ABLIKIM	25R BES3	$e^+e^- \rightarrow \psi(2S)$

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta') / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{30} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<14.2 \times 10^{-6}$ 90 ¹ CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta'$

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{31} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<2.21 \times 10^{-5}$ 90 ¹ ABLIKIM 24Q BES3 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.7 \times 10^{-4}$ 90 ² CRONIN-HEN..10 CLEO $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$

¹ $\eta_c(1S)$ reconstructed in the final states $K^+ K^- \pi^0$ and $K_S^0 K^\pm \pi^\mp$.

² Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \omega \omega) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{21} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.04 \times 10^{-6}$ 90 ABLIKIM 25K BES3 $\psi(2S) \rightarrow \gamma 2(\pi^+ \pi^- \pi^0)$

$$\Gamma(\eta_c(2S) \rightarrow \omega \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}} \times \Gamma_{22} / \Gamma \times \Gamma_{200}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.85 \times 10^{-7}$ 90 ABLIKIM 25K BES3 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0 K^+ K^-$

$\eta_c(2S)$ REFERENCES

ABLIKIM	25H	PR D111 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25I	PR D111 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25K	PR D111 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25R	PR D111 052013	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25V	PR D111 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BT	PR D110 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BW	PR D110 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24J	PR D109 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24Q	PR D109 072017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	23AH	PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23Q	PR D107 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHILIKIN	19	PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)

DEL-AMO-SA... 11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA 11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN... 10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT 08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
NAKAZAWA 08	NPBPS 184 220	H. Nakazawa	(BELLE Collab.)
UEHARA 08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT 08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE 07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT 06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE 04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER 04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT 04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K 02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI 02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI 01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU 98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG 95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE 85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS 82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
