

$f_0(1500)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

 $f_0(1500)$  T-MATRIX POLE  $\sqrt{s}$ Note that  $\Gamma = -2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1430–1530) – <math>i</math> (40–90) OUR ESTIMATE</b>			
(1487.8 $\pm$ 7.7 $\pm$ 3.7) – $i$ (46.1 $\pm$ 8.6 $\pm$ 6.6)	ABLIKIM	25CD BES3	$e^+ e^- \rightarrow \psi(2S)$
(1450 $\pm$ 10) – $i$ (53 $\pm$ 8)	<sup>1</sup> RODAS	22 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
(1483 $\pm$ 15) – $i$ (58 $\pm$ 6)	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
(1496 $\pm$ 1.2 <sup>+4.4</sup> <sub>-26.4</sub> ) – $i$ (40.4 $\pm$ 0.3 <sup>+10.0</sup> <sub>-2.5</sub> )	<sup>2</sup> ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
(1465 $\pm$ 18) – $i$ (50 $\pm$ 9)	<sup>3</sup> ROPERTZ	18 RVUE	$\bar{B}_S^0 \rightarrow J/\psi(\pi^+\pi^-/K^+K^-)$
(1486 $\pm$ 10) – $i$ (57 $\pm$ 5)	ANISOVICH	09 RVUE	$0.0 \bar{p}p, \pi N$
(1489 <sup>+8</sup> <sub>-4</sub> ) – $i$ (51 $\pm$ 5)	<sup>4</sup> ANISOVICH	03 RVUE	
(1515 $\pm$ 12) – $i$ (55 $\pm$ 12)	BARBERIS	00A	$450 pp \rightarrow p_f(\eta\eta', \eta'\eta')p_S$
(1511 $\pm$ 9) – $i$ (51 $\pm$ 9)	<sup>5</sup> BARBERIS	00C	$450 pp \rightarrow p_f 4\pi p_S$
(1510 $\pm$ 8) – $i$ (55 $\pm$ 8)	BARBERIS	00E	$450 pp \rightarrow p_f \eta\eta p_S$
(1502 $\pm$ 12 $\pm$ 10) – $i$ (49 $\pm$ 9 $\pm$ 8)	<sup>6</sup> BARBERIS	99D OMEG	$450 pp \rightarrow K^+K^-, \pi^+\pi^-$
(1447 $\pm$ 27) – $i$ (54 $\pm$ 23)	<sup>7</sup> KAMINSKI	99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
(1499 $\pm$ 8) – $i$ (65 $\pm$ 10)	ANISOVICH	98B RVUE	Compilation.
(1510 $\pm$ 20) – $i$ (60 $\pm$ 18)	BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$
(1449 $\pm$ 20) – $i$ (57 $\pm$ 15)	BERTIN	97C OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
(1515 $\pm$ 20) – $i$ (53 $\pm$ 8)	ABELE	96B CBAR	$0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
(1500 $\pm$ 8) – $i$ (66 $\pm$ 8)	ABELE	96C RVUE	Compilation.
(1500 $\pm$ 10) – $i$ (77 $\pm$ 15)	<sup>8</sup> AMSLER	95D CBAR	$0.0 \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$
(1520 $\pm$ 25) – $i$ (74 <sup>+10</sup> <sub>-13</sub> )	<sup>9</sup> ANISOVICH	94 CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
(1505 $\pm$ 20) – $i$ (75 $\pm$ 10)	<sup>10</sup> BUGG	94 RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0\pi^0$

<sup>1</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma\pi^0\pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).

<sup>2</sup> T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ), and BINON 84C ( $\eta\eta'$ ).

<sup>3</sup> T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants.

<sup>4</sup> Pole position from combined analysis of  $\pi^-\bar{p} \rightarrow \pi^0\pi^0 n, \pi^-\bar{p} \rightarrow K\bar{K}n, \pi^+\pi^- \rightarrow \pi^+\pi^-, \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta, \pi^+\pi^-\pi^0, K^+K^-\pi^0, K_S^0 K_S^0 \pi^0, K^+K_S^0 \pi^-$  at rest,  $\bar{p}n \rightarrow \pi^-\pi^-\pi^+, K_S^0 K^-\pi^0, K_S^0 K_S^0 \pi^-$  at rest.

<sup>5</sup> Average between  $\pi^+\pi^-2\pi^0$  and  $2(\pi^+\pi^-)$ .

<sup>6</sup> Supersedes BARBERIS 99 and BARBERIS 99B.

<sup>7</sup> T-matrix pole on sheet  $--+$ .

<sup>8</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

<sup>9</sup> From a simultaneous analysis of the annihilations  $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ .

<sup>10</sup> Reanalysis of ANISOVICH 94 data.

### $f_0(1500)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1522 ± 25</b>		<sup>1</sup> BERTIN 98	OBLX	0.05–0.405 $\bar{p}p \rightarrow \pi^+\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1492.5 ± 3.6 <sup>+</sup> <sub>−20.5</sub>		<sup>2</sup> ABLIKIM 22G	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
1447 ± 16 ± 13	163	<sup>3,4</sup> DOBBS 15		$J/\psi \rightarrow \gamma\pi^+\pi^-$
1442 ± 9 ± 4	261	<sup>3,4</sup> DOBBS 15		$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1460.9 ± 2.9		<sup>5</sup> AAIJ 14BR	LHCB	$\bar{B}_S^0 \rightarrow J/\psi\pi^+\pi^-$
1468 +14 <sup>+</sup> <sub>−15</sub> +23 <sup>+</sup> <sub>−74</sub>	5.5k	<sup>6</sup> ABLIKIM 13N	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
1470 ± 60	568	<sup>7</sup> KLEMPF 08	E791	$D_S^+ \rightarrow \pi^-\pi^+\pi^+$
1470 +6 <sup>+</sup> <sub>−7</sub> +72 <sup>+</sup> <sub>−255</sub>		<sup>8</sup> UEHARA 08A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1466 ± 6 ± 20		<sup>9</sup> ABLIKIM 06V	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1495 ± 4		AMSLER 06	CBAR	0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
1539 ± 20	9.9k	AUBERT 06O	BABR	$B^+ \rightarrow K^+K^+K^-$
1473 ± 5	80k	<sup>9,10</sup> UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
1478 ± 6		VLADIMIRSK...06	SPEC	40 $\pi^-p \rightarrow K_S^0K_S^0n$
1493 ± 7		<sup>9</sup> BINON 05	GAMS	33 $\pi^-p \rightarrow \eta\eta n$
1524 ± 14	1400	<sup>11</sup> GARMASH 05	BELL	$B^+ \rightarrow K^+K^+K^-$
1490 ± 30		<sup>9</sup> ABELE 01	CBAR	0.0 $\bar{p}d \rightarrow \pi^-4\pi^0p$
1497 ± 10		<sup>9</sup> BARBERIS 99	OMEG	450 $pp \rightarrow p_S p_f K^+K^-$
1502 ± 10		<sup>9</sup> BARBERIS 99B	OMEG	450 $pp \rightarrow p_S p_f \pi^+\pi^-$
1530 ± 45		<sup>9</sup> BELLAZZINI 99	GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
1505 ± 18		<sup>9</sup> FRENCH 99		300 $pp \rightarrow p_f(K^+K^-)p_S n$
1580 ± 80		<sup>9</sup> ALDE 98	GAM4	100 $\pi^-p \rightarrow \pi^0\pi^0n$
~ 1520		REYES 98	SPEC	800 $pp \rightarrow p_S p_f K_S^0 K_S^0$
~ 1475		FRABETTI 97D	E687	$D_S^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 1505		ABELE 96	CBAR	0.0 $\bar{p}p \rightarrow 5\pi^0$
1460 ± 20	120	<sup>9</sup> AMELIN 96B	VES	37 $\pi^-A \rightarrow \eta\eta\pi^-A$
1500 ± 8		BUGG 96	RVUE	
1500 ± 15		<sup>12</sup> AMSLER 95B	CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$
1505 ± 15		<sup>13</sup> AMSLER 95C	CBAR	0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
1445 ± 5		<sup>14</sup> ANTINORI 95	OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
1497 ± 30		<sup>9</sup> ANTINORI 95	OMEG	300,450 $pp \rightarrow pp\pi^+\pi^-$

~ 1505		BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1446 ± 5		<sup>9</sup> ABATZIS	94	OMEG 450	$pp \rightarrow pp2(\pi^+ \pi^-)$
1545 ± 25		<sup>9</sup> AMSLER	94E	CBAR 0.0	$\bar{p}p \rightarrow \pi^0 \eta \eta'$
1560 ± 25		<sup>9</sup> AMSLER	92	CBAR 0.0	$\bar{p}p \rightarrow \pi^0 \eta \eta$
1550 ± 45 ± 30		<sup>9</sup> BELADIDZE	92C	VES 36	$\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
1449 ± 4		<sup>9</sup> ARMSTRONG	89E	OMEG 300	$pp \rightarrow pp2(\pi^+ \pi^-)$
1610 ± 20		<sup>9</sup> ALDE	88	GAM4 300	$\pi^- N \rightarrow \pi^- N 2\eta$
~ 1525		ASTON	88D	LASS 11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1570 ± 20	600	<sup>9</sup> ALDE	87	GAM4 100	$\pi^- p \rightarrow 4\pi^0 n$
1575 ± 45		<sup>15</sup> ALDE	86D	GAM4 100	$\pi^- p \rightarrow 2\eta n$
1568 ± 33		<sup>9</sup> BINON	84C	GAM2 38	$\pi^- p \rightarrow \eta \eta' n$
1592 ± 25		<sup>9</sup> BINON	83	GAM2 38	$\pi^- p \rightarrow 2\eta n$
1525 ± 5		<sup>9</sup> GRAY	83	DBC 0.0	$\bar{p}N \rightarrow 3\pi$

<sup>1</sup> Breit-Wigner mass.

<sup>2</sup> The  $\pi^+ \pi^-$  mass spectrum is described by a coherent sum of two Breit-Wigner resonances,  $f_0(1500)$  and a new  $X(1540)$  with mass  $1540.2 \pm 7.0^{+36.3}_{-6.1}$  MeV and width  $157 \pm 19^{+11}_{-77}$  MeV.

<sup>3</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>4</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 109$  MeV.

<sup>5</sup> Solution I, statistical error only.

<sup>6</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>7</sup> Reanalysis of AITALA 01A data. This state could also be  $f_0(1370)$ .

<sup>8</sup> Breit-Wigner mass. May also be the  $f_0(1370)$ .

<sup>9</sup> Breit-Wigner mass.

<sup>10</sup> Statistical error only.

<sup>11</sup> Breit-Wigner, solution 1, PWA ambiguous.

<sup>12</sup> T-matrix pole, supersedes ANISOVICH 94.

<sup>13</sup> T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

<sup>14</sup> Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

<sup>15</sup> From central value and spread of two solutions. Breit-Wigner mass.

### $f_0(1500)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>108 ± 33</b>		<sup>1</sup> BERTIN	98	OBLX 0.05–0.405 $\bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
107 ± $9^{+21}_{-7}$		<sup>2</sup> ABLIKIM	22G	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
124 ± 7		<sup>3</sup> AAIJ	14BR	LHCB $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$
$136^{+41+28}_{-26-100}$	5.5k	<sup>4</sup> ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
$90^{+2+50}_{-1-22}$		<sup>5</sup> UEHARA	08A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$108^{+14}_{-11} \pm 25$		<sup>6</sup> ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
121 ± 8		AMSLER	06	CBAR $0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
257 ± 33	9.9k	AUBERT	060	BABR $B^+ \rightarrow K^+ K^+ K^-$
108 ± 9	80k	<sup>6,7</sup> UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$

119 ± 10		VLADIMIRSK...06	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$
90 ± 15		<sup>6</sup> BINON	05	GAMS	33 $\pi^- p \rightarrow \eta \eta n$
136 ± 23	1400	<sup>8</sup> GARMASH	05	BELL	$B^+ \rightarrow K^+ K^+ K^-$
140 ± 40		<sup>6</sup> ABELE	01	CBAR	0.0 $\bar{p} d \rightarrow \pi^- 4\pi^0 p$
104 ± 25		<sup>6</sup> BARBERIS	99	OMEG	450 $pp \rightarrow p_s p_f K^+ K^-$
131 ± 15		<sup>6</sup> BARBERIS	99B	OMEG	450 $pp \rightarrow p_s p_f \pi^+ \pi^-$
160 ± 50		<sup>6</sup> BELLAZZINI	99	GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
100 ± 33		<sup>6</sup> FRENCH	99		300 $pp \rightarrow p_f(K^+ K^-) p_s$
280 ± 100		<sup>6</sup> ALDE	98	GAM4	100 $\pi^- p \rightarrow \pi^0\pi^0 n$
~ 100		FRABETTI	97D	E687	$D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 169		ABELE	96	CBAR	0.0 $\bar{p} p \rightarrow 5\pi^0$
100 ± 30	120	<sup>6</sup> AMELIN	96B	VES	37 $\pi^- A \rightarrow \eta \eta \pi^- A$
132 ± 15		BUGG	96	RVUE	
120 ± 25		<sup>9</sup> AMSLER	95B	CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$
120 ± 30		<sup>10</sup> AMSLER	95C	CBAR	0.0 $\bar{p} p \rightarrow \eta \eta \pi^0$
65 ± 10		<sup>11</sup> ANTINORI	95	OMEG	300,450 $pp \rightarrow pp2(\pi^+ \pi^-)$
199 ± 30		<sup>6</sup> ANTINORI	95	OMEG	300,450 $pp \rightarrow pp\pi^+ \pi^-$
56 ± 12		<sup>6</sup> ABATZIS	94	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
100 ± 40		<sup>6</sup> AMSLER	94E	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \eta \eta'$
245 ± 50		<sup>6</sup> AMSLER	92	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \eta \eta$
153 ± 67 ± 50		<sup>6</sup> BELADIDZE	92C	VES	36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
78 ± 18		<sup>6</sup> ARMSTRONG	89E	OMEG	300 $pp \rightarrow pp2(\pi^+ \pi^-)$
170 ± 40		<sup>6</sup> ALDE	88	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$
150 ± 20	600	<sup>6</sup> ALDE	87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
265 ± 65		<sup>12</sup> ALDE	86D	GAM4	100 $\pi^- p \rightarrow 2\eta n$
260 ± 60		<sup>6</sup> BINON	84C	GAM2	38 $\pi^- p \rightarrow \eta \eta' n$
210 ± 40		<sup>6</sup> BINON	83	GAM2	38 $\pi^- p \rightarrow 2\eta n$
101 ± 13		<sup>6</sup> GRAY	83	DBC	0.0 $\bar{p} N \rightarrow 3\pi$

<sup>1</sup> Breit-Wigner width.

<sup>2</sup> The  $\pi^+ \pi^-$  mass spectrum is described by a coherent sum of two Breit-Wigner resonances,  $f_0(1500)$  and a new  $X(1540)$  with mass  $1540.2 \pm 7.0^{+36.3}_{-6.1}$  MeV and width  $157 \pm 19^{+11}_{-77}$  MeV.

<sup>3</sup> Solution I, statistical error only.

<sup>4</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>5</sup> Breit-Wigner width. May also be the  $f_0(1370)$ .

<sup>6</sup> Breit-Wigner width.

<sup>7</sup> Statistical error only.

<sup>8</sup> Breit-Wigner, solution 1, PWA ambiguous.

<sup>9</sup> T-matrix pole, supersedes ANISOVICH 94.

<sup>10</sup> T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

<sup>11</sup> Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

<sup>12</sup> From central value and spread of two solutions. Breit-Wigner mass.

### $f_0(1500)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor
$\Gamma_1$ $\pi\pi$	$(34.5 \pm 2.2) \%$	1.2
$\Gamma_2$ $\pi^+\pi^-$	seen	
$\Gamma_3$ $2\pi^0$	seen	
$\Gamma_4$ $4\pi$	$(48.9 \pm 3.3) \%$	1.2
$\Gamma_5$ $4\pi^0$	seen	
$\Gamma_6$ $2\pi^+2\pi^-$	seen	
$\Gamma_7$ $2(\pi\pi)_{S\text{-wave}}$	seen	
$\Gamma_8$ $\rho\rho$	seen	
$\Gamma_9$ $\pi(1300)\pi$	seen	
$\Gamma_{10}$ $a_1(1260)\pi$	seen	
$\Gamma_{11}$ $\eta\eta$	$(6.0 \pm 0.9) \%$	1.1
$\Gamma_{12}$ $\eta\eta'(958)$	$(2.2 \pm 0.8) \%$	1.4
$\Gamma_{13}$ $K\bar{K}$	$(8.5 \pm 1.0) \%$	1.1
$\Gamma_{14}$ $\gamma\gamma$	not seen	

### CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 10 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 5.6$  for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_4$	-88			
$x_{11}$	27	-56		
$x_{12}$	3	-32	26	
$x_{13}$	43	-64	20	2
	$x_1$	$x_4$	$x_{11}$	$x_{12}$

### $f_0(1500) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_{14}/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$33^{+12+1809}_{-6-21}$		<sup>1</sup> UEHARA	08A BELL	$10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
not seen		ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{ee} = 91, 183\text{--}209 \text{ GeV}$
<460	95	BARATE	00E ALEP	$\gamma\gamma \rightarrow \pi^+\pi^-$
<sup>1</sup> May also be the $f_0(1370)$ . Multiplied by us by 3 to obtain the $\pi\pi$ value.				

**$f_0(1500)$  BRANCHING RATIOS** **$\Gamma(\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	--------------------	-------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.454±0.104	BUGG	96 RVUE
-------------	------	---------

 **$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

<b>seen</b>	BERTIN	98 OBLX	0.05–0.405 $\bar{p}p \rightarrow \pi^+\pi^+\pi^-$
-------------	--------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	FRABETTI	97D E687	$D_S^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
---------------	----------	----------	---

 **$\Gamma(4\pi)/\Gamma(\pi\pi)$   $\Gamma_4/\Gamma_1$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

**1.42±0.18 OUR FIT** Error includes scale factor of 1.2.**1.42±0.18 OUR AVERAGE** Error includes scale factor of 1.2.

1.37±0.16	BARBERIS	00D	450 $pp \rightarrow p_f 4\pi p_S$
-----------	----------	-----	-----------------------------------

2.1 ±0.6	<sup>1</sup> AMSLER	98	RVUE
----------	---------------------	----	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 ±0.2	<sup>2</sup> ANISOVICH	02D SPEC	Combined fit
----------	------------------------	----------	--------------

3.4 ±0.8	<sup>1</sup> ABELE	96 CBAR	0.0 $\bar{p}p \rightarrow 5\pi^0$
----------	--------------------	---------	-----------------------------------

<sup>1</sup> Excluding  $\rho\rho$  contribution to  $4\pi$ .<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K}n$ ) data. **$\Gamma(2(\pi\pi)_{\text{S-wave}})/\Gamma(\pi\pi)$   $\Gamma_7/\Gamma_1$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.42±0.26	<sup>1</sup> ABELE	01 CBAR	0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
-----------	--------------------	---------	---

<sup>1</sup> From the combined data of ABELE 96 and ABELE 96c. **$\Gamma(2(\pi\pi)_{\text{S-wave}})/\Gamma(4\pi)$   $\Gamma_7/\Gamma_4$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26±0.07	ABELE	01B CBAR	0.0 $\bar{p}d \rightarrow 5\pi p$
-----------	-------	----------	-----------------------------------

 **$\Gamma(\rho\rho)/\Gamma(4\pi)$   $\Gamma_8/\Gamma_4$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13±0.08	ABELE	01B CBAR	0.0 $\bar{p}d \rightarrow 5\pi p$
-----------	-------	----------	-----------------------------------

 **$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{\text{S-wave}})$   $\Gamma_8/\Gamma_7$** 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
--------------	--------------------	----------------

**2.87±0.34 OUR AVERAGE** Error includes scale factor of 1.1.

3.3 ±0.5	BARBERIS	00C 450 $pp \rightarrow p_f \pi^+ \pi^- 2\pi^0 p_S$
----------	----------	---

2.6 ±0.4	BARBERIS	00C 450 $pp \rightarrow p_f 2(\pi^+ \pi^-) p_S$
----------	----------	---

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$  $\Gamma_9/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.50±0.25	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$  $\Gamma_{10}/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.12±0.05	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
large	ALDE	88	GAM4 300 $\pi^- N \rightarrow \eta\eta\pi^- N$
large	BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$  $\Gamma_{11}/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.173±0.024 OUR FIT</b>	Error includes scale factor of 1.1.		
<b>0.175±0.027 OUR AVERAGE</b>			
0.18 ±0.03	BARBERIS	00E	450 $p p \rightarrow p_f \eta \eta p_s$
0.157±0.060	<sup>1</sup> AMSLER	95D	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
0.080±0.033	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
0.11 ±0.03	<sup>2</sup> ANISOVICH	02D	SPEC Combined fit
0.078±0.013	<sup>3</sup> ABELE	96C	RVUE Compilation
0.230±0.097	<sup>4</sup> AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta \eta \pi^0$

<sup>1</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$ ), GAMS ( $\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$ ), and BNL ( $\pi p \rightarrow K \bar{K} n$ ) data.<sup>3</sup>  $2\pi$  width determined to be  $60 \pm 12$  MeV.<sup>4</sup> Using AMSLER 95B ( $3\pi^0$ ). $\Gamma(4\pi^0)/\Gamma(\eta\eta)$  $\Gamma_5/\Gamma_{11}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.8±0.3	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$

 $\Gamma(\eta\eta'(958))/\Gamma(\pi\pi)$  $\Gamma_{12}/\Gamma_1$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.4±2.2 OUR FIT</b>	Error includes scale factor of 1.4.		
<b>9.5±2.6</b>	BARBERIS	00A	450 $p p \rightarrow p_f \eta \eta p_s$
16.6 <sup>+4.2</sup> <sub>-4.0</sub>	<sup>1</sup> ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
0.5±0.3	<sup>2</sup> ANISOVICH	02D	SPEC Combined fit

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$ ), GAMS ( $\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$ ), and BNL ( $\pi p \rightarrow K \bar{K} n$ ) data.

$\Gamma(\eta\eta'(958))/\Gamma(\eta\eta)$  $\Gamma_{12}/\Gamma_{11}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.37±0.13 OUR FIT</b>	Error includes scale factor of 1.5.		
<b>0.29±0.10</b>	<sup>1</sup> AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.05±0.03	<sup>2</sup> ANISOVICH	02D	SPEC Combined fit
0.84±0.23	ABELE	96C	RVUE Compilation
2.7 ±0.8	BINON	84C	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$

<sup>1</sup> Using AMSLER 94E ( $\eta\eta'\pi^0$ ).<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$ ,  $\pi^0\eta\eta$ ,  $\pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n$ ,  $\eta\eta n$ ,  $\eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K}n$ ) data. $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.044±0.021	BUGG	96	RVUE

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$  $\Gamma_{13}/\Gamma_1$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.246±0.025 OUR FIT</b>			
<b>0.236±0.026 OUR AVERAGE</b>			
0.25 ±0.03	<sup>1</sup> BARGIOTTI	03	OBLX $\bar{p}p$
0.19 ±0.07	<sup>2</sup> ABELE	98	CBAR 0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
0.20 ±0.08	<sup>3</sup> ABELE	96B	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
0.16 ±0.05	<sup>4</sup> ANISOVICH	02D	SPEC Combined fit
0.33 ±0.03 ±0.07	BARBERIS	99D	OMEG 450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$

<sup>1</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ , and  $K^\pm K_S^0 \pi^\mp$ .<sup>2</sup> Using  $\pi^0\pi^0$  from AMSLER 95B.<sup>3</sup> Using AMSLER 95B ( $3\pi^0$ ), AMSLER 94C ( $2\pi^0\eta$ ) and SU(3).<sup>4</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$ ,  $\pi^0\eta\eta$ ,  $\pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n$ ,  $\eta\eta n$ ,  $\eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K}n$ ) data. $\Gamma(K\bar{K})/\Gamma(\eta\eta)$  $\Gamma_{13}/\Gamma_{11}$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.43±0.24 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>1.85±0.41</b>		BARBERIS	00E	450 $pp \rightarrow p_f \eta\eta p_S$
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
1.5 ±0.6		<sup>1</sup> ANISOVICH	02D	SPEC Combined fit
<0.4	90	<sup>2</sup> PROKOSHKIN	91	GAM4 300 $\pi^- p \rightarrow \pi^- p \eta\eta$
<0.6		<sup>3</sup> BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$

<sup>1</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$ ,  $\pi^0\eta\eta$ ,  $\pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n$ ,  $\eta\eta n$ ,  $\eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K}n$ ) data.<sup>2</sup> Combining results of GAM4 with those of WA76 on  $K\bar{K}$  central production.<sup>3</sup> Using ETKIN 82B and COHEN 80.

**$f_0(1500)$  REFERENCES**

- ABLIKIM 25CD JHEP 2510 081 M. Ablikim *et al.* (BESIII Collab.)  
 ABLIKIM 22AS PR D106 072012 M. Ablikim *et al.* (BESIII Collab.)  
 Also PR D107 079901 (errat.) M. Ablikim *et al.* (BESIII Collab.)  
 ABLIKIM 22G PRL 129 042001 M. Ablikim *et al.* (BESIII Collab.)  
 RODAS 22 EPJ C82 80 A. Rodas *et al.* (JPAC Collab.)  
 SARANTSEV 21 PL B816 136227 A.V. Sarantsev *et al.* (BONN, PNPI)  
 ALBRECHT 20 EPJ C80 453 M. Albrecht *et al.* (Crystal Barrel Collab.)  
 ABLIKIM 18AA PR D98 072003 M. Ablikim *et al.* (BESIII Collab.)  
 ROPERTZ 18 EPJ C78 1000 S. Ropertz, C. Hanhart, B. Kubis (BONN, JULI)  
 AAIJ 17V JHEP 1708 037 R. Aaij *et al.* (LHCb Collab.)  
 ABLIKIM 15AE PR D92 052003 M. Ablikim *et al.* (BESIII Collab.)  
 DOBBS 15 PR D91 052006 S. Dobbs *et al.* (NWES)  
 AAIJ 14BR PR D89 092006 R. Aaij *et al.* (LHCb Collab.)  
 ABLIKIM 13N PR D87 092009 M. Ablikim *et al.* (BESIII Collab.)  
 ANISOVICH 09 IJMP A24 2481 V.V. Anisovich, A.V. Sarantsev (PNPI)  
 KLEMP 08 EPJ C55 39 E. Klempt, M. Matveev, A.V. Sarantsev (BONN+)  
 UEHARA 08A PR D78 052004 S. Uehara *et al.* (BELLE Collab.)  
 ABLIKIM 06V PL B642 441 M. Ablikim *et al.* (BES Collab.)  
 AMSLER 06 PL B639 165 C. Amsler *et al.* (Crystal Barrel Collab.)  
 AUBERT 06O PR D74 032003 B. Aubert *et al.* (BABAR Collab.)  
 UMAN 06 PR D73 052009 I. Uman *et al.* (FNAL E835)  
 VLADIMIRSK... 06 PAN 69 493 V.V. Vladimirovsky *et al.* (ITEP, Moscow)  
 Translated from YAF 69 515.  
 BINON 05 PAN 68 960 F. Binon *et al.*  
 Translated from YAF 68 998.  
 GARMASH 05 PR D71 092003 A. Garmash *et al.* (BELLE Collab.)  
 ANISOVICH 03 EPJ A16 229 V.V. Anisovich *et al.*  
 BARGIOTTI 03 EPJ C26 371 M. Bargiotti *et al.* (OBELIX Collab.)  
 AMSLER 02 EPJ C23 29 C. Amsler *et al.* (Crystal Barrel Collab.)  
 ANISOVICH 02D PAN 65 1545 V.V. Anisovich *et al.*  
 Translated from YAF 65 1583.  
 ABELE 01 EPJ C19 667 A. Abele *et al.* (Crystal Barrel Collab.)  
 ABELE 01B EPJ C21 261 A. Abele *et al.* (Crystal Barrel Collab.)  
 ACCIARRI 01H PL B501 173 M. Acciarri *et al.* (L3 Collab.)  
 AITALA 01A PRL 86 765 E.M. Aitala *et al.* (FNAL E791 Collab.)  
 BARATE 00E PL B472 189 R. Barate *et al.* (ALEPH Collab.)  
 BARBERIS 00A PL B471 429 D. Barberis *et al.* (WA 102 Collab.)  
 BARBERIS 00C PL B471 440 D. Barberis *et al.* (WA 102 Collab.)  
 BARBERIS 00D PL B474 423 D. Barberis *et al.* (WA 102 Collab.)  
 BARBERIS 00E PL B479 59 D. Barberis *et al.* (WA 102 Collab.)  
 BARBERIS 99 PL B453 305 D. Barberis *et al.* (Omega Expt.)  
 BARBERIS 99B PL B453 316 D. Barberis *et al.* (Omega Expt.)  
 BARBERIS 99D PL B462 462 D. Barberis *et al.* (Omega Expt.)  
 BELLAZZINI 99 PL B467 296 R. Bellazzini *et al.*  
 FRENCH 99 PL B460 213 B. French *et al.* (WA76 Collab.)  
 KAMINSKI 99 EPJ C9 141 R. Kaminski, L. Lesniak, B. Loiseau (CRAC, PARIN)  
 ABELE 98 PR D57 3860 A. Abele *et al.* (Crystal Barrel Collab.)  
 ALDE 98 EPJ A3 361 D. Alde *et al.* (GAM4 Collab.)  
 Also PAN 62 405 D. Alde *et al.* (GAMS Collab.)  
 Translated from YAF 62 446.  
 AMSLER 98 RMP 70 1293 C. Amsler  
 ANISOVICH 98B SPU 41 419 V.V. Anisovich *et al.*  
 Translated from UFN 168 481.  
 BERTIN 98 PR D57 55 A. Bertin *et al.* (OBELIX Collab.)  
 REYES 98 PRL 81 4079 M.A. Reyes *et al.*  
 BARBERIS 97B PL B413 217 D. Barberis *et al.* (WA 102 Collab.)  
 BERTIN 97C PL B408 476 A. Bertin *et al.* (OBELIX Collab.)  
 FRABETTI 97D PL B407 79 P.L. Frabetti *et al.* (FNAL E687 Collab.)  
 ABELE 96 PL B380 453 A. Abele *et al.* (Crystal Barrel Collab.)  
 ABELE 96B PL B385 425 A. Abele *et al.* (Crystal Barrel Collab.)  
 ABELE 96C NP A609 562 A. Abele *et al.* (Crystal Barrel Collab.)  
 AMELIN 96B PAN 59 976 D.V. Amelin *et al.* (SERP, TBIL)  
 Translated from YAF 59 1021.  
 BUGG 96 NP B471 59 D.V. Bugg, A.V. Sarantsev, B.S. Zou (LOQM, PNPI)  
 AMSLER 95B PL B342 433 C. Amsler *et al.* (Crystal Barrel Collab.)  
 AMSLER 95C PL B353 571 C. Amsler *et al.* (Crystal Barrel Collab.)  
 AMSLER 95D PL B355 425 C. Amsler *et al.* (Crystal Barrel Collab.)  
 ANTINORI 95 PL B353 589 F. Antinori *et al.* (ATHU, BARI, BIRM+)  
 BUGG 95 PL B353 378 D.V. Bugg *et al.* (LOQM, PNPI, WASH)

ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94E	PL B340 259	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)
		Translated from YAF 55 2748.		
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)
		Translated from DANS 316 900.		
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
ALDE	88	PL B201 160	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
Also		SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
		Translated from YAF 38 934.		
GRAY	83	PR D27 307	L. Gray <i>et al.</i>	(SYRA)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)

---