

$\psi(4415)$ [a]

$$I^G(J^{PC}) = 0^-(1^{--})$$

[a] J^{PC} known by production in e^+e^- via single photon annihilation. I^G is not known; interpretation of this state as a single resonance is unclear because of the expectation of substantial threshold effects in this energy region.

$\psi(4415)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4415 ± 5 OUR AVERAGE			
4413.6 ± 9.0 ± 0.8	¹ ABLIKIM	24D BES3	$e^+e^- \rightarrow \omega\gamma J/\psi$
4414.6 ± 3.4 ± 6.1	ABLIKIM	23BH BES3	$e^+e^- \rightarrow D_s^{*+} D_s^{*-}$
4415.1 ± 7.9	² ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4412 ± 15	³ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4411 ± 7	⁴ PAKHLOVA	08A BELL	10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
4425 ± 6	⁵ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4429 ± 9	⁶ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4417 ± 10	BRANDELIK	78C DASP	e^+e^-
4414 ± 7	SIEGRIST	76 MRK1	e^+e^-

¹ Assuming one single Breit-Wigner resonance in $\omega\chi_{c2}(1P)$ ($\chi_{c2} \rightarrow \gamma J/\psi$).

² Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

³ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.

⁴ Systematic uncertainties not estimated.


⁵ From a fit to Crystal Ball (OSTERHELD 86) data.

⁶ From a fit to BES (BAI 02C) data.

$\psi(4415)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
110 ± 13 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
110.5 ± 15.0 ± 2.9	⁷ ABLIKIM	24D BES3	$e^+e^- \rightarrow \omega\gamma J/\psi$
122.5 ± 7.5 ± 8.1	ABLIKIM	23BH BES3	$e^+e^- \rightarrow D_s^{*+} D_s^{*-}$
71.5 ± 19.0	⁸ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118 ± 32	⁹ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
77 ± 20	¹⁰ PAKHLOVA	08A BELL	10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
119 ± 16	¹¹ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
118 ± 35	¹² SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
66 ± 15	BRANDELIK	78C DASP	e^+e^-
33 ± 10	SIEGRIST	76 MRK1	e^+e^-

- ⁷ Assuming one single Breit-Wigner resonance in $\omega\chi_{c2}(1P)$ ($\chi_{c2} \rightarrow \gamma J/\psi$).
- ⁸ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.
- ⁹ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8-4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.
- ¹⁰ Systematic uncertainties not estimated.
- ¹¹ From a fit to Crystal Ball (OSTERHELD 86) data.
- ¹² From a fit to BES (BAI 02C) data.



$\psi(4415)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $D\bar{D}$	seen	
Γ_2 $D^0\bar{D}^0$	seen	
Γ_3 D^+D^-	seen	
Γ_4 $D^*\bar{D} + c.c.$	seen	
Γ_5 $D^*(2007)^0\bar{D}^0 + c.c.$	seen	
Γ_6 $D^*(2010)^+D^- + c.c.$	seen	

Γ_7	$D^* \bar{D}^*$	seen		
Γ_8	$D^*(2007)^0 \bar{D}^*(2007)^0 + \text{c.c.}$	seen		
Γ_9	$D^*(2010)^+ D^*(2010)^- + \text{c.c.}$	seen		
Γ_{10}	$D^0 D^- \pi^+ (\text{excl. } D^*(2010)^+ D^- + \text{c.c.})$	< 2.3	%	90%
Γ_{11}	$D \bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.}$	(10 ± 4)	%	
Γ_{12}	$D^0 D^{*-} \pi^+ + \text{c.c.}$	< 31	%	90%
Γ_{13}	$D_1(2420) \bar{D} + \text{c.c.}$	possibly seen		
Γ_{14}	$D_s^+ D_s^-$	not seen		
Γ_{15}	$\omega \chi_{c2}$	$(9 \pm 4) \times 10^{-3}$		
Γ_{16}	$D_s^{*+} D_s^- + \text{c.c.}$	seen		
Γ_{17}	$D_s^{*+} D_s^{*-}$	seen		
Γ_{18}	$\psi_2(3823) \pi^+ \pi^-$	possibly seen		
Γ_{19}	$\psi(3770) \pi^+ \pi^-$	possibly seen		
Γ_{20}	$J/\psi \eta$	< 1.0	%	90%
Γ_{21}	$\chi_{c1} \gamma$	< 1.3	$\times 10^{-3}$	90%
Γ_{22}	$\chi_{c2} \gamma$	< 7	$\times 10^{-3}$	90%
Γ_{23}	$\Lambda \bar{\Lambda}$	< 5	$\times 10^{-6}$	90%
Γ_{24}	$\Sigma^+ \bar{\Sigma}^-$	< 1.8	$\times 10^{-4}$	90%
Γ_{25}	$\Sigma^0 \bar{\Sigma}^0$	< 4	$\times 10^{-5}$	90%
Γ_{26}	$\Xi^0 \bar{\Xi}^0$	< 1.4	$\times 10^{-4}$	90%
Γ_{27}	$\Xi^- \bar{\Xi}^+$	< 6	$\times 10^{-5}$	90%
Γ_{28}	$p K^- \bar{\Lambda} + \text{c.c.}$	< 1.0	$\times 10^{-5}$	90%
Γ_{29}	$\Lambda \bar{\Xi}^+ K^- + \text{c.c.}$	< 4	$\times 10^{-5}$	90%
Γ_{30}	$\Sigma^0 \bar{\Xi}^+ K^- + \text{c.c.}$	< 2.5	$\times 10^{-4}$	90%
Γ_{31}	$p K^- K^- \bar{\Xi}^+ + \text{c.c.}$	< 8	$\times 10^{-5}$	90%
Γ_{32}	$\omega \pi^0$	not seen		
Γ_{33}	$\omega \eta$	not seen		
Γ_{34}	$e^+ e^-$	$(3.2 \pm 1.2) \times 10^{-6}$		
Γ_{35}	$\mu^+ \mu^-$	$(1.1 \pm 0.4) \times 10^{-5}$		

$\psi(4415)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$					Γ_{34}
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
0.35 ± 0.12	¹³ ABLIKIM	08D	BES2	$e^+ e^- \rightarrow$ hadrons	
• • •	We do not use the following data for averages, fits, limits, etc. • • •				
0.4 to 0.8	¹⁴ MO	10	RVUE	$e^+ e^- \rightarrow$ hadrons	
0.72 ± 0.11	¹⁵ SETH	05A	RVUE	$e^+ e^- \rightarrow$ hadrons	
0.64 ± 0.23	¹⁶ SETH	05A	RVUE	$e^+ e^- \rightarrow$ hadrons	
0.49 ± 0.13	BRANDELIK	78C	DASP	$e^+ e^-$	
0.44 ± 0.14	SIEGRIST	76	MRK1	$e^+ e^-$	

¹³ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

¹⁴ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e^+e^- partial widths. We quote only the range of values.

¹⁵ From a fit to Crystal Ball (OSTERHELD 86) data.

¹⁶ From a fit to BES (BAI 02C) data.

$\Gamma(\mu^+\mu^-)$ Γ_{35}

VALUE (keV)		DOCUMENT ID	TECN	COMMENT
$1.25 \pm 0.28 \pm 0.35$	17,18	ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

¹⁷ From a fit to the $e^+e^- \rightarrow \mu^+\mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$.

¹⁸ From solution 1 of 8 with equal fit quality. Other solutions range from $1.24 \pm 0.28 \pm 0.35$ to $1.27 \pm 0.41 \pm 0.36$ keV.

$\psi(4415) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\omega\chi_{c2}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_{34}/\Gamma$

VALUE (eV)		DOCUMENT ID	TECN	COMMENT
$3.17 \pm 0.39 \pm 0.24$	19	ABLIKIM	24D BES3	$e^+e^- \rightarrow \omega\gamma J/\psi$

¹⁹ Assuming one single Breit-Wigner resonance in $\omega\chi_{c2}(1P)$ ($\chi_{c2} \rightarrow \gamma J/\psi$).

$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_{34}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 3.6	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{21}\Gamma_{34}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.47	90	²⁰ HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

²⁰ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_{34}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 2.3	90	²¹ HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

²¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{23}\Gamma_{34}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.8 \times 10^{-3}$	90	²² ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(4415)$

²² From a measurement of the $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

$\Gamma(\Sigma^+\bar{\Sigma}^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_{34}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 62.1 \times 10^{-3}$	90	²³ ABLIKIM	24AH BES3	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$

²³ Interference effect between resonance and continuum amplitudes is considered. Two solutions from the fit.

$\Gamma(\Sigma^0 \bar{\Sigma}^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{25} \Gamma_{34} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<14.0 \times 10^{-3}$	90	²⁴ ABLIKIM	25U BES3	$e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$

²⁴ Interference effect between resonance and continuum amplitudes is considered. Upper limit is for the larger of the two solutions from the fit.

 $\Gamma(\Xi^0 \bar{\Xi}^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{26} \Gamma_{34} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<48.0 \times 10^{-3}$	90	²⁵ ABLIKIM	24CD BES3	$e^+ e^- \rightarrow \psi(4415)$

²⁵ From a fit to $e^+ e^- \rightarrow \Xi^0 \bar{\Xi}^0$ cross sections.

 $\Gamma(\Xi^- \bar{\Xi}^+) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{27} \Gamma_{34} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<21.7 \times 10^{-3}$	90	²⁶ ABLIKIM	23BK BES3	$e^+ e^- \rightarrow \psi(4415)$

²⁶ From a fit to $e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$ cross sections.

 $\Gamma(p K^- \bar{\Lambda} + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{28} \Gamma_{34} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-3}$	90	²⁷ ABLIKIM	23BL BES3	$e^+ e^- \rightarrow \psi(4415)$

²⁷ From a fit to $e^+ e^- \rightarrow p K^- \bar{\Lambda} + \text{c.c.}$ cross sections.

 $\Gamma(\Lambda \bar{\Xi}^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{29} \Gamma_{34} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<14.3 \times 10^{-3}$	90	²⁸ ABLIKIM	24AL BES3	$e^+ e^- \rightarrow \Lambda \bar{\Xi}^+ K^- + \text{c.c.}$

²⁸ A fit to the Born cross section of $e^+ e^- \rightarrow \Lambda \bar{\Xi}^+ K^- + \text{c.c.}$ including interference with the continuum. Two solutions from the fit.

 $\Gamma(\Sigma^0 \bar{\Xi}^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{30} \Gamma_{34} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<87.0 \times 10^{-3}$	90	²⁹ ABLIKIM	24AL BES3	$e^+ e^- \rightarrow \Sigma^0 \bar{\Xi}^+ K^- + \text{c.c.}$

²⁹ A fit to the Born cross section of $e^+ e^- \rightarrow \Sigma^0 \bar{\Xi}^+ K^- + \text{c.c.}$ including interference with the continuum. Two solutions from the fit.

 $\Gamma(e^+ e^-) \times \Gamma(p K^- K^- \bar{\Xi}^+ + \text{c.c.}) / \Gamma_{\text{total}}$ $\Gamma_{34} \Gamma_{31} / \Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<29.4 \times 10^{-3}$	90	ABLIKIM	25CF BES3	$e^+ e^- \rightarrow p K^- K^- \bar{\Xi}^+ + \text{c.c.}$

 $\psi(4415) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$ $\Gamma(D^0 D^{*-} \pi^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{12} / \Gamma \times \Gamma_{34} / \Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.99 \times 10^{-6}$	90	³⁰ PAKHLOVA	09 BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

³⁰ Using 4421 ± 4 MeV for the mass of $\psi(4415)$.

$\psi(4415)$ BRANCHING RATIOS $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$ Γ_2/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

seen	³¹ ABLIKIM	24BH	BES3 $e^+e^- \rightarrow D^0\bar{D}^0$
seen	PAKHLOVA	08	BELL $e^+e^- \rightarrow D^0\bar{D}^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^0\bar{D}^0\gamma$
----------	--------	-----	--

³¹ A precision measurement of the $e^+e^- \rightarrow D^0\bar{D}^0$ cross section shows complex structure in this mass region.

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$ Γ_3/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

seen	³² ABLIKIM	24BH	BES3 $e^+e^- \rightarrow D^+D^-$
seen	PAKHLOVA	08	BELL $e^+e^- \rightarrow D^+D^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^+D^-\gamma$
----------	--------	-----	--

³² A precision measurement of the $e^+e^- \rightarrow D^+D^-$ cross section shows complex structure in this mass region.

 $\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$
-------------	--------	-----	---

 $\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

seen	³³ ZHUKOVA	18	BELL $e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*+}D^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	PAKHLOVA	07	BELL $e^+e^- \rightarrow D^{*+}D^-\gamma$
------	----------	----	---

³³ Supersedes PAKHLOVA 07.

 $\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$ Γ_1/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.14 ± 0.12 ± 0.03	AUBERT	09M	BABR $e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$
---------------------------	--------	-----	---

 $\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$ Γ_4/Γ_7

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

0.17 ± 0.25 ± 0.03	AUBERT	09M	BABR $e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$
---------------------------	--------	-----	---

 $\Gamma(D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$
-------------	--------	-----	--

 $\Gamma(D^*(2010)^+D^*(2010)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

seen	³⁴ ZHUKOVA	18	BELL $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$
seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen PAKHLOVA 07 BELL $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$
³⁴ Supersedes PAKHLOVA 07.

$\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
10.5 ± 2.4 ± 3.8	³⁵ PAKHLOVA 08A	BELL	10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$

³⁵ Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.

$\Gamma(D^0 D^- \pi^+ (\text{excl. } D^*(2010)^+ D^- + \text{c.c.})/\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})$ Γ_{10}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	³⁶ PAKHLOVA 08A	BELL	10.6 $e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$

³⁶ Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.

$\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	³⁷ ABLIKIM 19AR	BES3	$e^+e^- \rightarrow \pi^+ \pi^- D\bar{D}$

³⁷ Evidence for $e^+e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV, not necessarily resonant.

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 11	BELL	$e^+e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA...10N	BABR	$e^+e^- \rightarrow D_s^+ D_s^- \gamma$

$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11	BELL	$e^+e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA...10N	BABR	$e^+e^- \rightarrow D_s^{*+} D_s^- \gamma$

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM 23BH	BES3	$e^+e^- \rightarrow D_s^{*+} D_s^{*-}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen PAKHLOVA 11 BELL $e^+e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
 not seen DEL-AMO-SA...10N BABR $e^+e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$

$\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
possibly seen	19	³⁸ ABLIKIM 15S	BES3	$e^+e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$

³⁸ From a fit of $e^+e^- \rightarrow \pi^+ \pi^- \psi_2(3823)$, $\psi_2(3823) \rightarrow \chi_{c1} \gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4415)$ line shape.

$\Gamma(\psi(3770)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	³⁹ ABLIKIM 19AR	BES3	$e^+e^- \rightarrow \pi^+ \pi^- D\bar{D}$

³⁹ Observe $e^+e^- \rightarrow \pi^+\pi^-\psi(3770)$ at $\sqrt{s} = 4.26, 4.36, \text{ and } 4.42$ GeV but cannot establish if continuum or resonant.

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$				Γ_{32}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	ABLIKIM	22K	BES3	$e^+e^- \rightarrow \omega\pi^0$

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$				Γ_{33}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	ABLIKIM	22K	BES3	$e^+e^- \rightarrow \omega\eta$

$\psi(4415)$ REFERENCES

ABLIKIM	25CF	JHEP 2511 111	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	25U	PR D111 L051502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AH	JHEP 2405 022	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24AL	JHEP 2407 258	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24BH	PRL 133 081901	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24CD	JHEP 2411 062	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	24D	PRL 132 161901	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BH	PRL 131 151903	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BL	JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
SIEGRIST	76	PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)