

$\psi(4660)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as $Y(4660)$; was $X(4660)$


See the reviews on the "Spectroscopy of Mesons Containing two Heavy Quarks" and on "Heavy Non-qqbar Mesons."

 $\psi(4660)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4623 ±10	OUR AVERAGE	Error includes scale factor of 3.7. See the ideogram below.		
4603.1 ± 3.9 ± 0.8		¹ ABLIKIM	24BN BES3	$e^+e^- \rightarrow D_s^+ D_{s2}^*(2573)^-$
4584 ±14 ±80		² ABLIKIM	24BN BES3	$e^+e^- \rightarrow D_s^+ D_{s1}(2536)^-$
4708 $^{+17}_{-15}$ ±21		³ ABLIKIM	23BI BES3	$e^+e^- \rightarrow K^+ K^- J/\psi$
4701.8 ±10.9 ± 2.7		⁴ ABLIKIM	23H BES3	$e^+e^- \rightarrow \phi \chi_{c2}$
4675.3 ±29.5 ± 3.5		⁵ ABLIKIM	23X BES3	$e^+e^- \rightarrow D^{*0} D^{*-} \pi^+$
4651.0 ±37.8 ± 2.1		⁶ ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4619.8 $^{+8.9}_{-8.0}$ ± 2.3	66	⁷ JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$
4625.9 $^{+6.2}_{-6.0}$ ± 0.4	89	⁸ JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
4652 ±10 ±11	279	⁹ WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4669 ±21 ± 3	37	¹⁰ LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4634 $^{+8}_{-7}$ $^{+5}_{-8}$	142	¹¹ PAKHLOVA	08B BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4647.9 ± 8.6 ± 0.8		¹² ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
4652.5 ± 3.4 ± 1.1		¹³ DAI	17 RVUE	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
4645.2 ± 9.5 ± 6.0		¹⁴ ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4646.4 ± 9.7 ± 4.8		¹⁵ ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$
4661 $^{+9}_{-8}$ ± 6	44	¹⁶ LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4664 ±11 ± 5	44	WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

¹ Extracted in a fit that employs two BW resonances. The second one at about 4720 MeV shows low statistical significance of 2.7 σ .² Extracted from a fit with two BW functions. The second one located at about 4750 MeV show a low statistical significance of 4.3 σ .³ Seen as a peak in the c.m. energy dependence of the $e^+e^- \rightarrow K^+ K^- J/\psi$ cross section using 5.85 fb⁻¹ of data at c.m. energies 4.61–4.95 GeV. Statistical significance is over 5 σ .⁴ Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term.⁵ From a cross-section measurement of $e^+e^- \rightarrow D^{*0} D^{*-} \pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. The two other resonances have masses (widths) 4209.6 ± 7.5 (81.6 ± 19.9) MeV and 4469.1 ± 26.4 (246.3 ± 37.9) MeV.


- ⁶ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.
- ⁷ Using $D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-$ decays.
- ⁸ From a fit of a Breit-Wigner convolved with a Gaussian.
- ⁹ From a two-resonance fit. Supersedes WANG 07D.
- ¹⁰ From a two-resonance fit.
- ¹¹ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.
- ¹² From a fit to the $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV.
- ¹³ The pole parameters are extracted from the speed plot.
- ¹⁴ From a three-resonance fit.
- ¹⁵ From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.
- ¹⁶ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.



$\psi(4660)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
55 ± 9	OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
57 ± 12 ± 219	1	ABLIKIM	24BN BES3	$e^+ e^- \rightarrow D_s^+ D_{s1}^-(2536)^-$
45.2 ± 5.7 ± 0.7	2	ABLIKIM	24BN BES3	$e^+ e^- \rightarrow D_s^+ D_{s2}^*(2573)^-$
126 $\begin{smallmatrix} +27 \\ -23 \end{smallmatrix}$ ± 30	3	ABLIKIM	23BI BES3	$e^+ e^- \rightarrow K^+ K^- J/\psi$
30.5 ± 22.3 ± 14.6	4	ABLIKIM	23H BES3	$e^+ e^- \rightarrow \phi \chi_{c2}$
218.3 ± 72.9 ± 9.3	5	ABLIKIM	23X BES3	$e^+ e^- \rightarrow D^{*0} D^{*-} \pi^+$

$155.4 \pm 24.8 \pm 0.8$	6	ABLIKIM	21AJ	BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$47.0^{+31.3}_{-14.8} \pm 4.6$	66	7	JIA	20	BELL $e^+e^- \rightarrow \gamma D_s^+ D_{s2}^{*-}(2573)^-$
$49.8^{+13.9}_{-11.5} \pm 4.0$	89	8	JIA	19A	BELL $e^+e^- \rightarrow \gamma D_s^+ D_{s1}^-(2536)^-$
$68 \pm 11 \pm 5$	279	9	WANG	15A	BELL $10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
$104 \pm 48 \pm 10$	37	10	LEES	14F	BABR $10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
$92^{+40}_{-24} \pm 10$	142	11	PAKHLOVA	08B	BELL $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$33.1 \pm 18.6 \pm 4.1$	12	ABLIKIM	22R	BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
$62.6 \pm 5.6 \pm 4.3$	13	DAI	17	RVUE	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
$113.8 \pm 18.1 \pm 3.4$	14	ZHANG	17B	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$103.5 \pm 15.6 \pm 4.0$	15	ZHANG	17C	RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
$42^{+17}_{-12} \pm 6$	44	16	LIU	08H	RVUE $10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
$48 \pm 15 \pm 3$	44	WANG	07D	BELL	$10.58 e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$



¹ Extracted from a fit with two BW functions. The second one located at about 4750 MeV show a low statistical significance of 4.3σ .

² Extracted in a fit that employs two BW resonances. The second one at about 4720 MeV shows low statistical significance of 2.7σ .

³ Seen as a peak in the c.m. energy dependence of the $e^+e^- \rightarrow K^+K^- J/\psi$ cross section using 5.85 fb^{-1} of data at c.m. energies 4.61–4.95 GeV. Statistical significance is over 5σ .

- ⁴ Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term.
- ⁵ From a cross-section measurement of $e^+e^- \rightarrow D^{*0}D^{*-}\pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. The two other resonances have masses (widths) 4209.6 ± 7.5 (81.6 ± 19.9) MeV and 4469.1 ± 26.4 (246.3 ± 37.9) MeV.
- ⁶ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.
- ⁷ Using $D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-$ decays.
- ⁸ From a fit of a Breit-Wigner convolved with a Gaussian.
- ⁹ From a two-resonance fit. Supersedes WANG 07D.
- ¹⁰ From a two-resonance fit.
- ¹¹ The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+\Lambda_c^-$ states are not necessarily the same.
- ¹² From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV.
- ¹³ The pole parameters are extracted from the speed plot.
- ¹⁴ From a three-resonance fit.
- ¹⁵ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.
- ¹⁶ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

$\psi(4660)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	not seen
Γ_2 $\psi(2S)\pi^+\pi^-$	seen
Γ_3 $J/\psi\eta$	not seen
Γ_4 $D^0D^{*-}\pi^+$	not seen
Γ_5 $D^{*0}D^{*-}\pi^+$	seen
Γ_6 $\psi_2(3823)\pi^+\pi^-$	seen
Γ_7 $\chi_{c0}\phi$	not seen
Γ_8 $\chi_{c1}\gamma$	not seen
Γ_9 $\chi_{c1}\phi$	not seen
Γ_{10} $\chi_{c2}\gamma$	not seen
Γ_{11} $\chi_{c2}\phi$	seen
Γ_{12} $\Lambda_c^+\Lambda_c^-$	seen
Γ_{13} $D_s^+D_{s1}(2536)^-$	seen
Γ_{14} $D_s^+D_{s2}^*(2573)^-$	seen
Γ_{15} $\omega\pi^0$	not seen
Γ_{16} $\omega\eta$	not seen
Γ_{17} $\Sigma^+\bar{\Sigma}^-$	not seen
Γ_{18} $\Sigma^0\bar{\Sigma}^0$	
Γ_{19} $\Xi^0\bar{\Xi}^0$	not seen
Γ_{20} $\Xi^-\bar{\Xi}^+$	not seen
Γ_{21} $pK^-\bar{\Lambda} + \text{c.c.}$	not seen
Γ_{22} $\Lambda\bar{\Xi}^+K^- + \text{c.c.}$	not seen
Γ_{23} $\Sigma^0\bar{\Xi}^+K^- + \text{c.c.}$	not seen

$\Gamma_{24} \quad pK^- K^- \Xi^+ + \text{c.c.}$

$\psi(4660) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

$\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_2 \Gamma_1 / \Gamma$

<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.7±3.8		¹ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
11.2±3.2		² ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4.7±4.2		³ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
11.3±3.3		⁴ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
2.0±0.3±0.2	279	⁵ WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
8.1±1.1±1.0	279	⁶ WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.7±1.3±0.5	37	⁷ LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.5±1.7±0.7	37	⁸ LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.2 ^{+0.7} _{-0.6}	44	⁹ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
5.9±1.6	44	¹⁰ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
3.0±0.9±0.3	44	⁷ WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.6±1.8±0.8	44	⁸ WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

- ¹ Solution I of four equivalent solutions in a fit using three interfering resonances.
- ² Solution II of four equivalent solutions in a fit using three interfering resonances.
- ³ Solution III of four equivalent solutions in a fit using three interfering resonances.
- ⁴ Solution IV of four equivalent solutions in a fit using three interfering resonances.
- ⁵ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.
- ⁶ Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.
- ⁷ Solution I of two equivalent solutions in a fit using two interfering resonances.
- ⁸ Solution II of two equivalent solutions in a fit using two interfering resonances.
- ⁹ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.
- ¹⁰ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

$\Gamma(J/\psi\eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_3 \Gamma_1 / \Gamma$

<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.94	90	WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

$\Gamma(D^{*0} D^{*-} \pi^+) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_5 \Gamma_1 / \Gamma$

<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
19 to 2005	¹ ABLIKIM	23X BES3	$e^+ e^- \rightarrow D^{*0} D^{*-} \pi^+$

- ¹ From a cross-section measurement of $e^+ e^- \rightarrow D^{*0} D^{*-} \pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. Depending on solutions I – VIII with same fit qualities.

$\Gamma(\chi_{c0}\phi) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_7 \Gamma_1 / \Gamma$

<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.35	90	ABLIKIM	25J BES3	$e^+ e^- \rightarrow \phi \chi_{c0}$

$$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.45	90	¹ HAN	15 BELL	10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$$\Gamma(\chi_{c1}\phi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_9\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	90	¹ ABLIKIM	23H BES3	$e^+e^- \rightarrow \phi\chi_{c1}$
-------	----	----------------------	----------	------------------------------------

¹ Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term.

$$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{10}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<2.1	90	¹ HAN	15 BELL	10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$
------	----	------------------	---------	--

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$$\Gamma(\chi_{c2}\phi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 ± 0.13		¹ ABLIKIM	23H BES3	$e^+e^- \rightarrow \phi\chi_{c2}$
-----------------	--	----------------------	----------	------------------------------------

¹ Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term. Constructive solution of the interference. Destructive solution gives 0.66 ± 0.41 eV.

$$\Gamma(D_s^+ D_{s1}(2536)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{13}\Gamma_1/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$14.3^{+2.8}_{-2.6} \pm 1.5$	89	¹ JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
------------------------------	----	------------------	----------	--

¹ Assuming $B(D_{s1}(2536)^- \rightarrow \bar{D}^{*0} K^-) = 1$.

$$\Gamma(D_s^+ D_{s2}^*(2573)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_1/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

$14.7^{+5.9}_{-4.5} \pm 3.6$	66	¹ JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$
------------------------------	----	------------------	---------	--

¹ Assuming $B(D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-) = 1$.

$$\Gamma(\Sigma^+ \bar{\Sigma}^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<49.6 \times 10^{-3}$	90	¹ ABLIKIM	24AH BES3	$e^+e^- \rightarrow \Sigma^+ \bar{\Sigma}^-$
------------------------	----	----------------------	-----------	--

¹ Interference effect between resonance and continuum amplitudes is considered. Two solutions from the fit.

$$\Gamma(\Sigma^0 \bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<17.4 \times 10^{-3}$	90	¹ ABLIKIM	25U BES3	$e^+e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$
------------------------	----	----------------------	----------	--

¹ Interference effect between resonance and continuum amplitudes is considered. Upper limit is for the larger of the two solutions from the fit.

$$\Gamma(\Xi^0 \Xi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{19} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<45.0 \times 10^{-3}$ 90 ¹ ABLIKIM 24CD BES3 $e^+ e^- \rightarrow \psi(4660)$

¹ From a fit to $e^+ e^- \rightarrow \Xi^0 \Xi^0$ cross sections.

$$\Gamma(\Xi^- \Xi^+) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{20} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<19.9 \times 10^{-3}$ 90 ¹ ABLIKIM 23BK BES3 $e^+ e^- \rightarrow \psi(4660)$

¹ From a fit to $e^+ e^- \rightarrow \Xi^- \Xi^+$ cross sections.

$$\Gamma(p K^- \bar{\Lambda} + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{21} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<2.8 \times 10^{-3}$ 90 ¹ ABLIKIM 23BL BES3 $e^+ e^- \rightarrow \psi(4660)$

¹ From a fit to $e^+ e^- \rightarrow p K^- \bar{\Lambda} + \text{c.c.}$ cross sections.

$$\Gamma(\Lambda \Xi^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{22} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<13.0 \times 10^{-3}$ 90 ¹ ABLIKIM 24AL BES3 $e^+ e^- \rightarrow \Lambda \Xi^+ K^- + \text{c.c.}$

¹ A fit to the Born cross section of $e^+ e^- \rightarrow \Lambda \Xi^+ K^- + \text{c.c.}$ including interference with the continuum. Two solutions from the fit.

$$\Gamma(\Sigma^0 \Xi^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{23} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<77.3 \times 10^{-3}$ 90 ¹ ABLIKIM 24AL BES3 $e^+ e^- \rightarrow \Sigma^0 \Xi^+ K^- + \text{c.c.}$

¹ A fit to the Born cross section of $e^+ e^- \rightarrow \Sigma^0 \Xi^+ K^- + \text{c.c.}$ including interference with the continuum. Two solutions from the fit.

$$\Gamma(e^+ e^-) \times \Gamma(p K^- K^- \Xi^+ + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_1 \Gamma_{24} / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

$<46.2 \times 10^{-3}$ 90 ABLIKIM 25CF BES3 $e^+ e^- \rightarrow p K^- K^- \Xi^+ + \text{c.c.}$

$$\psi(4660) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$$

$$\Gamma(D^0 D^{*-} \pi^+) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_4 / \Gamma \times \Gamma_1 / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<0.37 \times 10^{-6}$ 90 ¹ PAKHLOVA 09 BELL $e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

¹ Using $4664 \pm 11 \pm 5$ MeV for the mass of $\psi(4660)$.

$$\Gamma(\Lambda_c^+ \Lambda_c^-) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma \times \Gamma_1 / \Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.68^{+0.16+0.29}_{-0.15-0.30}$ 142 ¹ PAKHLOVA 08B BELL $e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$

¹ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

$\psi(4660)$ BRANCHING RATIOS $\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S)\pi^+\pi^-)$ Γ_4/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<10	90	PAKHLOVA 09	BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$

 $\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM 22R	BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$

¹ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances.

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 22K	BES3	$e^+e^- \rightarrow \omega\pi^0$

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM 22K	BES3	$e^+e^- \rightarrow \omega\eta$

 $\psi(4660)$ REFERENCES

ABLIKIM 25CF	JHEP 2511 111	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 25J	PR D111 012016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 25U	PR D111 L051502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 24AH	JHEP 2405 022	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 24AL	JHEP 2407 258	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 24BN	PRL 133 171903	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 24CD	JHEP 2411 062	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 23BI	PRL 131 211902	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 23BL	JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 23H	JHEP 2301 132	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 23X	PRL 130 121901	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 21AJ	PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
JIA 20	PR D101 091101	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA 19A	PR D100 111103	S. Jia <i>et al.</i>	(BELLE Collab.)
DAI 17	PR D96 116001	L.-Y. Dai, J. Haidenbauer, U.-G. Meissner	(JULI+)
ZHANG 17B	PR D96 054008	J. Zhang, J. Zhang	
ZHANG 17C	EPJ C77 727	J. Zhang, L. Yuan	
HAN 15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG 15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES 14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG 13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
LIU 08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
PAKHLOVA 08B	PRL 101 172001	C. Pakhlova <i>et al.</i>	(BELLE Collab.)
AUBERT 07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG 07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)