

Ω BARYONS

(S = -3, I = 0)

$\Omega^- = sss$

Ω⁻

$$I(J^P) = 0(\frac{3}{2}^+)$$

$J^P = \frac{3}{2}^+$ is the quark-model prediction; and $J = 3/2$ is fairly well established.

- Mass $m = 1672.45 \pm 0.29$ MeV
- $(m_{\Omega^-} - m_{\bar{\Omega}^+}) / m_{\Omega^-} = (-1 \pm 8) \times 10^{-5}$
- Mean life $\tau = (0.821 \pm 0.011) \times 10^{-10}$ s
- $c\tau = 2.461$ cm
- $(\tau_{\Omega^-} - \tau_{\bar{\Omega}^+}) / \tau_{\Omega^-} = 0.00 \pm 0.05$
- Magnetic moment $\mu = -2.02 \pm 0.05 \mu_N$

Decay parameters

- $\alpha(\Omega^-) \alpha_{-}(\Lambda)$ FOR $\Omega^- \rightarrow \Lambda K^- = 0.0115 \pm 0.0015$
- $\Lambda K^- \quad \alpha = 0.0154 \pm 0.0020$
- $\Lambda K^-, \bar{\Lambda} K^+ \quad (\alpha + \bar{\alpha}) / (\alpha - \bar{\alpha}) = -0.02 \pm 0.13$
- $\Xi^0 \pi^- \quad \alpha = 0.09 \pm 0.14$
- $\Xi^- \pi^0 \quad \alpha = 0.05 \pm 0.21$

Ω ⁻ DECAY MODES	Fraction (Γ _i /Γ)	Scale factor/ Confidence level	p (MeV/c)
ΛK ⁻	(67.7 ± 0.7) %		211
Ξ ⁰ π ⁻	(24.3 ± 0.7) %	S=1.5	294
Ξ ⁻ π ⁰	(8.55±0.33) %		289
Ξ ⁻ π ⁺ π ⁻	(3.7 ^{+0.7} _{-0.6}) × 10 ⁻⁴		189
Ξ(1530) ⁰ π ⁻	< 7 × 10 ⁻⁵	CL=90%	17
Ξ ⁰ e ⁻ $\bar{\nu}_e$	(5.6 ± 2.8) × 10 ⁻³		319
Ξ ⁻ γ	< 4.6 × 10 ⁻⁴	CL=90%	314
ΔS = 2 forbidden (S2) modes			
Λπ ⁻	S2 < 2.9 × 10 ⁻⁶	CL=90%	449
Σ ⁰ π ⁻	< 5.4 × 10 ⁻⁴	CL=90%	393
nK ⁻	< 2.4 × 10 ⁻⁴	CL=90%	415

$\Omega(2012)^-$

$I(J^P) = 0(?^-)$

Mass $m = 2012.9 \pm 0.4$ MeVFull width $\Gamma = 6.3 \pm 2.0$ MeV

$\Omega(2012)^-$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi^0 K^-$	$(34^{+17}_{-12})\%$	403
$\Xi^- \bar{K}^0$	$(28^{+12}_{-7})\%$	393
$\Xi^- \pi^+ K^-$	seen	225

 $\Omega(2250)^-$

$I(J^P) = 0(?^?)$

Mass $m = 2252 \pm 9$ MeVFull width $\Gamma = 55 \pm 18$ MeV

$\Omega(2250)^-$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi^- \pi^+ K^-$	seen	532
$\Xi(1530)^0 K^-$	seen	437