

BOTTOM, STRANGE MESONS

($B = \pm 1, S = \mp 1$)

$$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b, \quad \text{similarly for } B_s^{*0}$$

B_s^0

$$I(J^P) = 0(0^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

$$\text{Mass } m_{B_s^0} = 5366.93 \pm 0.10 \text{ MeV}$$

$$m_{B_s^0} - m_B = 87.37 \pm 0.12 \text{ MeV}$$

$$\text{Mean life } \tau = (1.515 \pm 0.006) \times 10^{-12} \text{ s}$$

$$c\tau = 454.2 \text{ } \mu\text{m}$$

$$\Delta\Gamma_{B_s^0} = \Gamma_{B_{sL}^0} - \Gamma_{B_{sH}^0} = (0.0783 \pm 0.0035) \times 10^{12} \text{ s}^{-1}$$

B_s^0 - \bar{B}_s^0 mixing parameters

$$\Delta m_{B_s^0} = m_{B_{sH}^0} - m_{B_{sL}^0} = (17.766 \pm 0.006) \times 10^{12} \hbar \text{ s}^{-1}$$

$$= (1.1694 \pm 0.0004) \times 10^{-8} \text{ MeV}$$

$$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0} = 26.91 \pm 0.11$$

$$\chi_s (B_s^0$$
- \bar{B}_s^0 mixing parameter) = 0.499313 ± 0.000006

CP violation parameters in B_s^0

$$\text{Re}(\epsilon_{B_s^0}) / (1 + |\epsilon_{B_s^0}|^2) = (-0.15 \pm 0.70) \times 10^{-3}$$

$$C_{KK}(B_s^0 \rightarrow K^+ K^-) = 0.162 \pm 0.035$$

$$S_{KK}(B_s^0 \rightarrow K^+ K^-) = 0.14 \pm 0.05 \quad (S = 1.3)$$

$$r_B(B_s^0 \rightarrow D_s^\mp K^\pm) = 0.318 \pm 0.034$$

$$r_B(B_s^0 \rightarrow D_s^\mp K^\pm \pi^\pm \pi^\mp) = 0.47 \pm 0.08$$

$$\delta_B(B_s^0 \rightarrow D_s^\pm K^\mp) = (348 \pm 6)^\circ$$

$$\delta_B(B_s^0 \rightarrow D_s^\pm K^\mp \pi^\pm \pi^\mp) = (-6^{+10}_{-13})^\circ$$

$$\text{CP Violation phase } \beta_s (b \rightarrow c\bar{c}s) = (2.7 \pm 0.7) \times 10^{-2} \text{ rad}$$

$$\text{CP Violation phase } \beta_s (b \rightarrow s\bar{s}s) = (3.7 \pm 3.5) \times 10^{-2} \text{ rad}$$

$$\text{CP Violation phase } \beta_s (B_s^0 \rightarrow J/\psi(1S)\phi) = (3.0 \pm 0.7) \times 10^{-2} \text{ rad}$$

$$|\lambda| (B_s^0 \rightarrow J/\psi(1S)\phi) = 0.995 \pm 0.009$$

$$|\lambda| (b \rightarrow c\bar{c}s) = 0.989 \pm 0.008$$

$$A, \text{ CP violation parameter} = -0.79 \pm 0.08$$

$$C, \text{ CP violation parameter} = 0.19 \pm 0.06$$

$$S, \text{ CP violation parameter} = 0.17 \pm 0.06$$

$$\begin{aligned}
 A_{CP}^L(B_s \rightarrow J/\psi \bar{K}^*(892)^0) &= 0.021 \pm 0.027 \\
 A_{CP}^{\parallel}(B_s \rightarrow J/\psi \bar{K}^*(892)^0) &= -0.07 \pm 0.06 \\
 A_{CP}^{\perp}(B_s \rightarrow J/\psi \bar{K}^*(892)^0) &= 0.06 \pm 0.05 \\
 \mathbf{A}_{CP}(B_s \rightarrow \pi^+ K^-) &= 0.224 \pm 0.012 \\
 A_{CP}(B_s^0 \rightarrow [K^+ K^-]_D \bar{K}^*(892)^0) &= 0.06 \pm 0.04 \\
 A_{CP}(B_s^0 \rightarrow [\pi^+ K^-]_D K^*(892)^0) &= -0.009 \pm 0.023 \\
 A_{CP}(B_s^0 \rightarrow [\pi^+ \pi^-]_D K^*(892)^0) &= 0.00 \pm 0.06 \\
 A_{CP}(B_s^0 \rightarrow [K^+ \pi^- \pi^+ \pi^-]_D \bar{K}^*(892)^0) &= -0.029 \pm 0.024 \\
 A_{CP}(B_s^0 \rightarrow [\pi^+ \pi^- \pi^+ \pi^-]_D \bar{K}^*(892)^0) &= 0.02 \pm 0.05 \\
 R_s^+ = \Gamma(B_s^0 \rightarrow [\pi^- K^+]_D \bar{K}^{*0}) / \Gamma(B_s^0 \rightarrow [\pi^+ K^-]_D \bar{K}^{*0}) &= 0.004 \pm 0.006 \\
 R_s^- = \Gamma(\bar{B}_s^0 \rightarrow [\pi^+ K^-]_D K^{*0}) / \Gamma(\bar{B}_s^0 \rightarrow [\pi^- K^+]_D K^{*0}) &= 0.004 \pm 0.006 \\
 R_s^+ = \Gamma(B_s^0 \rightarrow [\pi^- K^+ \pi^+ \pi^-]_D \bar{K}^{*0}) / \Gamma(B_s^0 \rightarrow [\pi^+ K^- \pi^+ \pi^-]_D \bar{K}^{*0}) &= \\
 &= 0.019 \pm 0.008 \\
 R_s^- = \Gamma(\bar{B}_s^0 \rightarrow [\pi^+ K^- \pi^+ \pi^-]_D K^{*0}) / \Gamma(\bar{B}_s^0 \rightarrow [\pi^- K^+ \pi^+ \pi^-]_D K^{*0}) &= \\
 &= 0.015 \pm 0.008 \\
 S(B_s^0 \rightarrow \phi \gamma) &= 0.43 \pm 0.32 \\
 C(B_s^0 \rightarrow \phi \gamma) &= 0.11 \pm 0.31 \\
 A^{\Delta}(B_s^0 \rightarrow \phi \gamma) &= -0.7 \pm 0.4 \\
 \Delta a_{\perp} &< 1.2 \times 10^{-12} \text{ GeV, CL} = 95\% \\
 \Delta a_{\parallel} &= (-0.9 \pm 1.5) \times 10^{-14} \text{ GeV} \\
 \Delta a_{\chi} &= (1.0 \pm 2.2) \times 10^{-14} \text{ GeV} \\
 \Delta a_{\gamma} &= (-3.8 \pm 2.2) \times 10^{-14} \text{ GeV} \\
 \text{Re}(\xi) &= -0.022 \pm 0.033 \\
 \text{Im}(\xi) &= 0.004 \pm 0.011
 \end{aligned}$$

These branching fractions all scale with $B(\bar{b} \rightarrow B_s^0)$.

The branching fraction $B(B_s^0 \rightarrow D_s^- \ell^+ \nu_{\ell} \text{ anything})$ is not a pure measurement since the measured product branching fraction $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_{\ell} \text{ anything})$ was used to determine $B(\bar{b} \rightarrow B_s^0)$, as described in the note on “ B^0 - \bar{B}^0 Mixing”

For inclusive branching fractions, e.g., $B \rightarrow D^{\pm} \text{ anything}$, the values usually are multiplicities, not branching fractions. They can be greater than one.

| B_s^0 DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level | p (MeV/c) |
|--------------------------|--------------------------------|-----------------------------------|----------------|
| D_s^{\pm} anything | (65 ± 5) % | | — |
| D^0/\bar{D}^0 anything | (24 ± 4) % | | — |
| D^{\pm} anything | (13 ± 5) % | | — |
| $\ell \nu_{\ell} X$ | (9.6 ± 0.8) % | | — |
| $e^+ \nu X^-$ | (9.1 ± 0.8) % | | — |

| | | | |
|--|--|-------|------|
| $\mu^+ \nu X^-$ | (10.2 \pm 1.0) % | | — |
| $D_s^- \ell^+ \nu_\ell$ anything | [a] (8.1 \pm 1.3) % | | — |
| $D_s^{*-} \ell^+ \nu_\ell$ anything | (5.4 \pm 1.1) % | | — |
| $D_s^- \mu^+ \nu_\mu$ | (2.31 \pm 0.21) % | | 2321 |
| $D_s^{*-} \mu^+ \nu_\mu$ | (5.2 \pm 0.5) % | | 2266 |
| $D_{s1}(2536)^- \mu^+ \nu_\mu, D_{s1}^- \rightarrow D_s^{*-} K_S^0$ | (2.7 \pm 0.7) $\times 10^{-3}$ | | — |
| $D_{s1}(2536)^- X \mu^+ \nu, D_{s1}^- \rightarrow \bar{D}^0 K^+$ | (4.4 \pm 1.3) $\times 10^{-3}$ | | — |
| $D_{s2}(2573)^- X \mu^+ \nu, D_{s2}^- \rightarrow \bar{D}^0 K^+$ | (2.7 \pm 1.0) $\times 10^{-3}$ | | — |
| $K^- \mu^+ \nu_\mu$ | (1.06 \pm 0.09) $\times 10^{-4}$ | | 2660 |
| $D_s^- \pi^+$ | (2.98 \pm 0.13) $\times 10^{-3}$ | | 2320 |
| $D_s^- \rho^+$ | (6.9 \pm 1.4) $\times 10^{-3}$ | | 2249 |
| $D_s^- \pi^+ \pi^+ \pi^-$ | (6.1 \pm 1.0) $\times 10^{-3}$ | | 2301 |
| $D_{s1}(2536)^- \pi^+, D_{s1}^- \rightarrow D_s^- \pi^+ \pi^-$ | (2.4 \pm 0.8) $\times 10^{-5}$ | | — |
| $D_s^\mp K^\pm$ | (2.25 \pm 0.12) $\times 10^{-4}$ | | 2293 |
| $D_{s1}(2536)^\mp K^\pm, D_{s1}^- \rightarrow \bar{D}^*(2007)^0 K^-$ | (2.48 \pm 0.28) $\times 10^{-5}$ | | — |
| $D_s^- K^+ \pi^+ \pi^-$ | (3.2 \pm 0.6) $\times 10^{-4}$ | | 2249 |
| $D_s^+ D_s^-$ | (4.5 \pm 0.6) $\times 10^{-3}$ | S=1.3 | 1824 |
| $D_s^- D^+$ | (3.1 \pm 0.5) $\times 10^{-4}$ | | 1875 |
| $D^+ D^-$ | (2.2 \pm 0.6) $\times 10^{-4}$ | | 1925 |
| $D_s^{*+} D_s^{*-}$ | (2.14 \pm 0.32) $\times 10^{-4}$ | | 1778 |
| $D^0 \bar{D}^0$ | (1.9 \pm 0.5) $\times 10^{-4}$ | | 1930 |
| $D_s^{*-} \pi^+$ | (1.9 $^{+0.5}_{-0.4}$) $\times 10^{-3}$ | | 2265 |
| $D_s^{*\mp} K^\pm$ | (1.32 $^{+0.40}_{-0.32}$) $\times 10^{-4}$ | | — |
| $D_s^{*-} \rho^+$ | (9.5 \pm 2.0) $\times 10^{-3}$ | | 2191 |
| $D_s^{*+} D_s^- + D_s^{*-} D_s^+$ | (1.51 \pm 0.13) % | | 1742 |
| $D_s^{*+} D_s^{*-}$ | (1.58 \pm 0.20) % | S=1.3 | 1655 |
| $D_s^{(*)+} D_s^{(*)-}$ | (4.5 \pm 1.4) % | | — |
| $D_s^{*-} D_s^+$ | (4.0 \pm 0.7) $\times 10^{-4}$ | | 1801 |
| $\bar{D}^{*0} \bar{K}^0$ | (2.8 \pm 1.1) $\times 10^{-4}$ | | 2278 |
| $\bar{D}^0 \bar{K}^0$ | (4.3 \pm 0.9) $\times 10^{-4}$ | | 2330 |
| $\bar{D}^0 K^- \pi^+$ | (1.04 \pm 0.13) $\times 10^{-3}$ | | 2312 |
| $\bar{D}^*(2007)^0 K^- \pi^+$ | (7.3 \pm 2.6) $\times 10^{-4}$ | | 2259 |
| $\bar{D}^0 \bar{K}^*(892)^0$ | (4.4 \pm 0.6) $\times 10^{-4}$ | | 2264 |
| $\bar{D}^0 \bar{K}^*(1410)$ | (3.9 \pm 3.5) $\times 10^{-4}$ | | 2117 |
| $\bar{D}^0 \bar{K}_0^*(1430)$ | (3.0 \pm 0.7) $\times 10^{-4}$ | | 2113 |

| | | | |
|---|--|--------|------|
| $\bar{D}^0 \bar{K}_2^*(1430)$ | $(1.1 \pm 0.4) \times 10^{-4}$ | | 2112 |
| $\bar{D}^0 \bar{K}^*(1680)$ | $< 7.8 \times 10^{-5}$ | CL=90% | 1997 |
| $\bar{D}^0 \bar{K}_0^*(1950)$ | $< 1.1 \times 10^{-4}$ | CL=90% | 1884 |
| $\bar{D}^0 \bar{K}_3^*(1780)$ | $< 2.6 \times 10^{-5}$ | CL=90% | 1970 |
| $\bar{D}^0 \bar{K}_4^*(2045)$ | $< 3.1 \times 10^{-5}$ | CL=90% | 1835 |
| $\bar{D}^0 K^- \pi^+$ (non-resonant) | $(2.1 \pm 0.8) \times 10^{-4}$ | | 2312 |
| $[K^+ K^-]_D \bar{K}^*(892)^0$ | $(4.4 \pm 0.6) \times 10^{-4}$ | | — |
| $[\pi^+ \pi^-]_D \bar{K}^*(892)^0$ | $(4.4 \pm 0.6) \times 10^{-4}$ | | — |
| $[\pi^+ \pi^- \pi^+ \pi^-]_D \bar{K}^*(892)^0$ | $(4.4 \pm 0.6) \times 10^{-4}$ | | — |
| $D_{s2}^*(2573)^- \pi^+, D_{s2}^* \rightarrow$ | $(2.6 \pm 0.4) \times 10^{-4}$ | | — |
| $\bar{D}^0 K^-$ | | | |
| $D_{s1}^*(2700)^- \pi^+, D_{s1}^* \rightarrow$ | $(1.6 \pm 0.8) \times 10^{-5}$ | | — |
| $\bar{D}^0 K^-$ | | | |
| $D_{s1}^*(2860)^- \pi^+, D_{s1}^* \rightarrow$ | $(5 \pm 4) \times 10^{-5}$ | | — |
| $\bar{D}^0 K^-$ | | | |
| $D_{s3}^*(2860)^- \pi^+, D_{s3}^* \rightarrow$ | $(2.2 \pm 0.6) \times 10^{-5}$ | | — |
| $\bar{D}^0 K^-$ | | | |
| $\bar{D}^0 K^+ K^-$ | $(5.6 \pm 0.9) \times 10^{-5}$ | | 2243 |
| $\bar{D}^0 f_0(980)$ | $< 3.1 \times 10^{-6}$ | CL=90% | 2242 |
| $\bar{D}^0 \phi$ | $(2.30 \pm 0.25) \times 10^{-5}$ | | 2235 |
| $\bar{D}^{*0} \phi$ | $(3.2 \pm 0.4) \times 10^{-5}$ | | 2178 |
| $D^{*0} \pi^\pm$ | $< 6.1 \times 10^{-6}$ | CL=90% | — |
| $\eta_c \phi$ | $(5.0 \pm 0.9) \times 10^{-4}$ | | 1663 |
| $\eta_c \pi^+ \pi^-$ | $(1.8 \pm 0.7) \times 10^{-4}$ | | 1840 |
| $J/\psi(1S) \phi$ | $(1.01 \pm 0.04) \times 10^{-3}$ | | 1588 |
| $J/\psi(1S) \phi \phi$ | $(1.17^+_{-0.16} \pm 0.14) \times 10^{-5}$ | | 764 |
| $J/\psi(1S) \pi^0$ | $< 1.21 \times 10^{-5}$ | CL=90% | 1787 |
| $J/\psi(1S) \eta$ | $(4.45 \pm 0.25) \times 10^{-4}$ | S=1.1 | 1733 |
| $J/\psi(1S) K_S^0$ | $(1.92 \pm 0.14) \times 10^{-5}$ | | 1743 |
| $J/\psi(1S) \bar{K}^*(892)^0$ | $(3.95 \pm 0.24) \times 10^{-5}$ | | 1637 |
| $J/\psi(1S) \eta'$ | $(3.53 \pm 0.22) \times 10^{-4}$ | S=1.1 | 1612 |
| $J/\psi(1S) \pi^+ \pi^-$ | $(2.00 \pm 0.17) \times 10^{-4}$ | S=1.7 | 1775 |
| $J/\psi(1S) f_0(500), f_0 \rightarrow$ | $< 4 \times 10^{-6}$ | CL=90% | — |
| $\pi^+ \pi^-$ | | | |
| $J/\psi(1S) \rho, \rho \rightarrow \pi^+ \pi^-$ | $< 3.4 \times 10^{-6}$ | CL=90% | — |
| $J/\psi(1S) f_0(980), f_0 \rightarrow$ | $(1.23 \pm 0.15) \times 10^{-4}$ | S=2.1 | — |
| $\pi^+ \pi^-$ | | | |
| $J/\psi(1S) f_2(1270), f_2 \rightarrow$ | $(1.0 \pm 0.4) \times 10^{-6}$ | | — |
| $\pi^+ \pi^-$ | | | |
| $J/\psi(1S) f_2(1270)_0, f_2 \rightarrow$ | $(7.2 \pm 1.6) \times 10^{-7}$ | | — |
| $\pi^+ \pi^-$ | | | |
| $J/\psi(1S) f_2(1270)_\parallel, f_2 \rightarrow$ | $(1.04 \pm 0.32) \times 10^{-6}$ | | — |
| $\pi^+ \pi^-$ | | | |

| | | | |
|---|---|--------|------|
| $J/\psi(1S) f_2(1270)_{\perp}, f_2 \rightarrow \pi^+ \pi^-$ | $(1.3 \pm 0.7) \times 10^{-6}$ | — | |
| $J/\psi(1S) f_0(1370), f_0 \rightarrow \pi^+ \pi^-$ | $(4.4 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.6 \\ 4.0 \end{smallmatrix}) \times 10^{-5}$ | — | |
| $J/\psi(1S) f_0(1500), f_0 \rightarrow \pi^+ \pi^-$ | $(2.02 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.32 \\ 0.24 \end{smallmatrix}) \times 10^{-5}$ | — | |
| $J/\psi(1S) f'_2(1525)_0, f'_2 \rightarrow \pi^+ \pi^-$ | $(1.02 \pm 0.22) \times 10^{-6}$ | — | |
| $J/\psi(1S) f'_2(1525)_{\parallel}, f'_2 \rightarrow \pi^+ \pi^-$ | $(1.2 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2.6 \\ 0.8 \end{smallmatrix}) \times 10^{-7}$ | — | |
| $J/\psi(1S) f'_2(1525)_{\perp}, f'_2 \rightarrow \pi^+ \pi^-$ | $(5 \pm 4) \times 10^{-7}$ | — | |
| $J/\psi(1S) f_0(1790), f_0 \rightarrow \pi^+ \pi^-$ | $(4.8 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 10.0 \\ 1.0 \end{smallmatrix}) \times 10^{-6}$ | — | |
| $J/\psi(1S) \pi^+ \pi^-$ (nonresonant) | $(1.72 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.00 \\ 0.34 \end{smallmatrix}) \times 10^{-5}$ | | 1775 |
| $J/\psi(1S) \bar{K}^0 \pi^+ \pi^-$ | $< 4.4 \times 10^{-5}$ | CL=90% | 1675 |
| $J/\psi(1S) K^+ K^-$ | $(7.9 \pm 0.7) \times 10^{-4}$ | | 1601 |
| $J/\psi(1S) K^0 K^- \pi^+ + \text{c.c.}$ | $(9.5 \pm 1.3) \times 10^{-4}$ | | 1538 |
| $J/\psi(1S) \bar{K}^0 K^+ K^-$ | $< 1.2 \times 10^{-5}$ | CL=90% | 1333 |
| $J/\psi K^*(892)^0 \bar{K}^*(892)^0$ | $(1.07 \pm 0.09) \times 10^{-4}$ | | 1082 |
| $J/\psi(1S) f'_2(1525)$ | $(2.6 \pm 0.6) \times 10^{-4}$ | | 1310 |
| $J/\psi(1S) p \bar{p}$ | $(3.6 \pm 0.4) \times 10^{-6}$ | | 982 |
| $J/\psi(1S) \gamma$ | $< 7.3 \times 10^{-6}$ | CL=90% | 1790 |
| $J/\psi \mu^+ \mu^-, J/\psi \rightarrow \mu^+ \mu^-$ | $< 2.6 \times 10^{-9}$ | CL=95% | — |
| $J/\psi(1S) \pi^+ \pi^- \pi^+ \pi^-$ | $(7.4 \pm 0.8) \times 10^{-5}$ | | 1731 |
| $J/\psi(1S) f_1(1285)$ | $(7.2 \pm 1.4) \times 10^{-5}$ | | 1460 |
| $J/\psi(1S) \bar{D}^0$ | $< 1.0 \times 10^{-6}$ | CL=90% | 996 |
| $\psi(2S) \eta$ | $(3.7 \pm 0.8) \times 10^{-4}$ | | 1338 |
| $\psi(2S) \eta'$ | $(1.37 \pm 0.33) \times 10^{-4}$ | | 1158 |
| $\psi(2S) \pi^+ \pi^-$ | $(6.8 \pm 1.2) \times 10^{-5}$ | | 1397 |
| $\psi(2S) \phi$ | $(5.1 \pm 0.4) \times 10^{-4}$ | | 1120 |
| $\psi(2S) K^0$ | $(1.9 \pm 0.5) \times 10^{-5}$ | | 1352 |
| $\psi(2S) K^- \pi^+$ | $(3.1 \pm 0.4) \times 10^{-5}$ | | 1310 |
| $\psi(2S) \bar{K}^*(892)^0$ | $(3.3 \pm 0.5) \times 10^{-5}$ | | 1196 |
| $\chi_{c1} \phi$ | $(1.92 \pm 0.25) \times 10^{-4}$ | | 1275 |
| $\chi_{c1}(3872) \phi$ | $(9.6 \pm 3.2) \times 10^{-5}$ | | 936 |
| $\chi_{c1}(3872) (K^+ K^-)_{non-\phi}$ | $(7.6 \pm 3.0) \times 10^{-5}$ | | 961 |
| $\chi_{c1}(3872) \pi^+ \pi^-$ | $(3.7 \pm 1.5) \times 10^{-5}$ | | 1264 |
| $\pi^+ \pi^-$ | $(7.1 \pm 0.8) \times 10^{-7}$ | | 2680 |
| $\pi^0 \pi^0$ | $< 7.7 \times 10^{-6}$ | CL=90% | 2680 |
| $\eta \pi^0$ | $< 1.0 \times 10^{-3}$ | CL=90% | 2654 |
| $\eta \eta$ | $< 1.43 \times 10^{-4}$ | CL=90% | 2627 |
| $\rho^0 \rho^0$ | $< 3.20 \times 10^{-4}$ | CL=90% | 2569 |

| | | | | |
|---|--|--|--------|------|
| $\eta' K_S^0$ | | $< 8.16 \times 10^{-6}$ | CL=90% | 2573 |
| $\eta' \eta$ | | $< 6.5 \times 10^{-5}$ | CL=90% | 2568 |
| $\eta' \eta'$ | | $(3.3 \pm 0.7) \times 10^{-5}$ | | 2507 |
| $\eta' \phi$ | | $< 8.2 \times 10^{-7}$ | CL=90% | 2495 |
| $\phi f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$ | | $(1.12 \pm 0.21) \times 10^{-6}$ | | — |
| $\phi f_2(1270), f_2(1270) \rightarrow \pi^+ \pi^-$ | | $(6.1 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.8 \\ 1.5 \end{smallmatrix}) \times 10^{-7}$ | | — |
| $\phi \rho^0$ | | $(2.7 \pm 0.8) \times 10^{-7}$ | | 2526 |
| $\phi \pi^+ \pi^-$ | | $(3.5 \pm 0.5) \times 10^{-6}$ | | 2579 |
| $\phi \phi$ | | $(1.83 \pm 0.14) \times 10^{-5}$ | | 2482 |
| $\phi \phi \phi$ | | $(2.1 \pm 0.6) \times 10^{-6}$ | | 2165 |
| $\pi^+ K^-$ | | $(5.9 \pm 0.6) \times 10^{-6}$ | | 2659 |
| $K^+ K^-$ | | $(2.61 \pm 0.16) \times 10^{-5}$ | | 2638 |
| $K^0 \bar{K}^0$ | | $(1.76 \pm 0.31) \times 10^{-5}$ | | 2637 |
| $K^0 \pi^+ \pi^-$ | | $(9.5 \pm 2.1) \times 10^{-6}$ | | 2653 |
| $K^0 K^\pm \pi^\mp$ | | $(8.4 \pm 0.9) \times 10^{-5}$ | | 2622 |
| $K^*(892)^- \pi^+$ | | $(2.9 \pm 1.1) \times 10^{-6}$ | | 2607 |
| $K^*(892)^\pm K^\mp$ | | $(1.9 \pm 0.5) \times 10^{-5}$ | | 2585 |
| $K_0^*(1430)^\pm K^\mp$ | | $(3.1 \pm 2.5) \times 10^{-5}$ | | — |
| $K_2^*(1430)^\pm K^\mp$ | | $(1.0 \pm 1.7) \times 10^{-5}$ | | — |
| $K^*(892)^0 \bar{K}^0 + c.c.$ | | $(2.0 \pm 0.6) \times 10^{-5}$ | | 2585 |
| $K_0^*(1430) \bar{K}^0 + c.c.$ | | $(3.3 \pm 1.0) \times 10^{-5}$ | | 2468 |
| $K_2^*(1430)^0 \bar{K}^0 + c.c.$ | | $(1.7 \pm 2.2) \times 10^{-5}$ | | 2467 |
| $K_S^0 \bar{K}^*(892)^0 + c.c.$ | | $(1.6 \pm 0.4) \times 10^{-5}$ | | 2585 |
| $K^0 K^+ K^-$ | | $(1.3 \pm 0.6) \times 10^{-6}$ | | 2568 |
| $\bar{K}^*(892)^0 \rho^0$ | | $< 7.67 \times 10^{-4}$ | CL=90% | 2550 |
| $\bar{K}^*(892)^0 K^*(892)^0$ | | $(1.11 \pm 0.27) \times 10^{-5}$ | | 2531 |
| $\phi K^*(892)^0$ | | $(1.14 \pm 0.30) \times 10^{-6}$ | | 2507 |
| $p \bar{p}$ | | $< 4.4 \times 10^{-9}$ | CL=90% | 2514 |
| $p \bar{p} K^0$ | | $(9.1 \pm 2.0) \times 10^{-7}$ | | 2396 |
| $p \bar{p} K^+ K^-$ | | $(4.5 \pm 0.5) \times 10^{-6}$ | | 2231 |
| $p \bar{p} K^+ \pi^-$ | | $(1.39 \pm 0.26) \times 10^{-6}$ | | 2355 |
| $p \bar{p} \pi^+ \pi^-$ | | $(4.3 \pm 2.0) \times 10^{-7}$ | | 2454 |
| $p \bar{p} p \bar{p}$ | | $(2.3 \pm 1.0) \times 10^{-8}$ | | 1797 |
| $p \bar{\Lambda} K^- + c.c.$ | | $(5.5 \pm 1.0) \times 10^{-6}$ | | 2358 |
| $\Lambda_c^- \Lambda \pi^+$ | | $(3.6 \pm 1.6) \times 10^{-4}$ | | 1979 |
| $\Lambda_c^- \Lambda_c^+$ | | $< 8.0 \times 10^{-5}$ | CL=95% | 1405 |

Lepton family (LF), lepton (L), baryon (B) number violating modes or $\Delta B = 1$ weak neutral current (B1) modes

| | | | | |
|--------------------|----|--|--------|------|
| $\gamma \gamma$ | B1 | $< 3.1 \times 10^{-6}$ | CL=90% | 2683 |
| $\phi \gamma$ | B1 | $(3.4 \pm 0.4) \times 10^{-5}$ | | 2587 |
| $f_2(1270) \gamma$ | B1 | $(9 \begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 4 \\ 5 \end{smallmatrix}) \times 10^{-6}$ | | 2532 |

| | | | | |
|---|------------|--|--------|------|
| $f_2'(1525)\gamma$ | <i>B1</i> | $(6.7 \pm_{-0.8}^{+0.9}) \times 10^{-6}$ | | 2469 |
| $\phi(1680)\gamma, \phi \rightarrow K^+K^-$ | <i>B1</i> | $(9.3 \pm 2.4) \times 10^{-7}$ | | — |
| $\phi_3(1850)\gamma, \phi_3 \rightarrow K^+K^-$ | <i>B1</i> | $(7 \pm_{-5}^{+6}) \times 10^{-8}$ | | — |
| $f_2(2010)\gamma, f_2 \rightarrow K^+K^-$ | <i>B1</i> | $(1.0 \pm_{-0.5}^{+0.7}) \times 10^{-7}$ | | — |
| $\mu^+\mu^-$ | <i>B1</i> | $(3.34 \pm 0.27) \times 10^{-9}$ | | 2681 |
| e^+e^- | <i>B1</i> | $< 9.4 \times 10^{-9}$ | CL=90% | 2683 |
| $\tau^+\tau^-$ | <i>B1</i> | $< 6.8 \times 10^{-3}$ | CL=95% | 2011 |
| $\mu^+\mu^-\gamma$ | <i>B1</i> | $< 4.2 \times 10^{-8}$ | CL=95% | 2681 |
| $\mu^+\mu^-\mu^+\mu^-$ | <i>B1</i> | $< 8.6 \times 10^{-10}$ | CL=95% | 2673 |
| $SP, S \rightarrow \mu^+\mu^-,$ $P \rightarrow \mu^+\mu^-$ | <i>B1</i> | [b] $< 2.2 \times 10^{-9}$ | CL=95% | — |
| $aa, a \rightarrow \mu^+\mu^-$ | <i>B1</i> | $< 5.8 \times 10^{-10}$ | CL=95% | — |
| $\phi(1020)\mu^+\mu^-$ | <i>B1</i> | $(8.2 \pm 0.4) \times 10^{-7}$ | | 2582 |
| $f_2'(1525)\mu^+\mu^-$ | <i>B1</i> | $(1.57 \pm 0.22) \times 10^{-7}$ | | 2464 |
| $\bar{K}^*(892)^0\mu^+\mu^-$ | <i>B1</i> | $(2.9 \pm 1.1) \times 10^{-8}$ | | 2605 |
| $\pi^+\pi^-\mu^+\mu^-$ | <i>B1</i> | $(8.4 \pm 1.7) \times 10^{-8}$ | | 2670 |
| $\bar{D}^0\mu^+\mu^-$ | <i>B1</i> | $< 1.2 \times 10^{-7}$ | CL=90% | 2354 |
| $\phi\nu\bar{\nu}$ | <i>B1</i> | $< 5.4 \times 10^{-3}$ | CL=90% | 2587 |
| $e^\pm\mu^\mp$ | <i>LF</i> | [c] $< 5.4 \times 10^{-9}$ | CL=90% | 2682 |
| $e^\pm\tau^\mp$ | <i>LF</i> | $< 1.4 \times 10^{-3}$ | CL=90% | 2389 |
| $\mu^\pm\tau^\mp$ | <i>LF</i> | $< 4.2 \times 10^{-5}$ | CL=95% | 2388 |
| $\phi\mu^\pm e^\mp$ | <i>LF</i> | $< 1.6 \times 10^{-8}$ | CL=90% | 2586 |
| $\phi\mu^\pm\tau^\mp$ | <i>LF</i> | $< 1.0 \times 10^{-5}$ | CL=90% | 2241 |
| $\rho\mu^-$ | <i>L,B</i> | $< 1.21 \times 10^{-8}$ | CL=90% | 2600 |

B_s^*

$$I(J^P) = 0(1^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

$$\text{Mass } m = 5415.4 \pm 1.4 \text{ MeV} \quad (S = 2.6)$$

$$m_{B_s^*} - m_{B_s} = 48.5 \pm 1.4 \text{ MeV} \quad (S = 2.6)$$

| B_s^* DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---------------------------------------|--------------------------------|-------------|
| $B_s\gamma$ | seen | 48 |

$B_{s1}(5830)^0$

$I(J^P) = 0(1^+)$
 I, J, P need confirmation.

Mass $m = 5828.73 \pm 0.20$ MeV
 $m_{B_{s1}^0} - m_{B^{*+}} = 503.98 \pm 0.17$ MeV
 Full width $\Gamma = 0.5 \pm 0.4$ MeV

| $B_{s1}(5830)^0$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|--|--------------------------------|-------------|
| $B^{*+} K^-$ | seen | 97 |

$B_{s2}^*(5840)^0$

$I(J^P) = 0(2^+)$
 I, J, P need confirmation.

Mass $m = 5839.88 \pm 0.12$ MeV
 $m_{B_{s2}^{*0}} - m_{B^+} = 560.48 \pm 0.12$ MeV
 Full width $\Gamma = 1.49 \pm 0.27$ MeV

Branching fractions are given relative to the one **DEFINED AS 1**.

| $B_{s2}^*(5840)^0$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|--|--------------------------------|-------------|
| $B^+ K^-$ | DEFINED AS 1 | 252 |
| $B^{*+} K^-$ | 0.093 ± 0.018 | 141 |
| $B^0 K_S^0$ | 0.43 ± 0.11 | 245 |
| $B^{*0} K_S^0$ | 0.04 ± 0.04 | — |

NOTES

- [a] Not a pure measurement. See note at head of B_s^0 Decay Modes.
- [b] Here S and P are the hypothetical scalar and pseudoscalar particles with masses of $2.5 \text{ GeV}/c^2$ and $214.3 \text{ MeV}/c^2$, respectively.
- [c] The value is for the sum of the charge states or particle/antiparticle states indicated.