

$\Xi_b(5955)^-$ $J^P = \frac{3}{2}^+$

Status: ***

 $\Xi_b(5955)^-$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|--|-------------|-----------|----------------------------|
| 5955.5 ± 0.4 OUR AVERAGE | | | |
| 5955.50 ± 0.03 ± 0.41 | 1,2 AAIJ | 23AU LHCb | $p\bar{p}$ at 7, 8, 13 TeV |
| 5955.2 ± 0.1 ± 0.4 | 1,3 AAIJ | 15H LHCb | $p\bar{p}$ at 7, 8 TeV |
| ¹ Observed in $\Xi_b^0\pi^-$ channel with $\Xi_b^0 \rightarrow \Xi_c^+\pi^-$ and $\Xi_c^+ \rightarrow pK^-\pi^+$. ² AAIJ 23AU measures $m(\Xi_b(5955)^-) - m(\Xi_b^0) - m(\pi^-) = 24.27 \pm 0.03 \pm 0.01$ MeV. We have adjusted the measurement to our best values of $m(\Xi_b^0) = 5791.7 \pm 0.4$ MeV, $m(\pi^-) = 139.57039 \pm 0.00018$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values. ³ AAIJ 15H measures $m(\Xi_b(5955)^-) - m(\Xi_b^0) - m(\pi^-) = 24.96 \pm 0.12 \pm 0.06$ MeV. We have adjusted the measurement to our best values of $m(\Xi_b^0) = 5791.7 \pm 0.4$ MeV, $m(\pi^-) = 139.57039 \pm 0.00018$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values. | | | |

 $\Xi_b(5955)^-$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|-----------|----------------------------|
| 1.43 ± 0.08 ± 0.08 | | | |
| 1.43 ± 0.08 ± 0.08 | 1 AAIJ | 23AU LHCb | $p\bar{p}$ at 7, 8, 13 TeV |
| $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ 1.65 ± 0.31 ± 0.10 | | | |
| 1.65 ± 0.31 ± 0.10 | 1 AAIJ | 15H LHCb | $p\bar{p}$ at 7, 8 TeV |
| ¹ Observed in $\Xi_b^0\pi^-$ channel with $\Xi_b^0 \rightarrow \Xi_c^+\pi^-$ and $\Xi_c^+ \rightarrow pK^-\pi^+$. | | | |

 $\Xi_b(5955)^-$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| $\Gamma_1 \quad \Xi_b^0\pi^- \times B(\bar{b} \rightarrow \Xi_b^*(5955)^-)/B(\bar{b} \rightarrow \Xi_b^0)$ | (20.7 ± 3.5) % |

 $\Xi_b(5955)^-$ BRANCHING RATIOS

| $\Gamma(\Xi_b^0\pi^- \times B(\bar{b} \rightarrow \Xi_b^*(5955)^-)/B(\bar{b} \rightarrow \Xi_b^0)) / \Gamma_{\text{total}}$ | Γ_1/Γ |
|---|-------------------|
| 0.207 ± 0.032 ± 0.015 | 1 AAIJ |
| ¹ Observed in $\Xi_b^0\pi^-$ channel with $\Xi_b^0 \rightarrow \Xi_c^+\pi^-$ and $\Xi_c^+ \rightarrow pK^-\pi^+$. | |

 $\Xi_b(5955)^-$ REFERENCES

| | | | |
|------|---------------------|-----------------------|----------------|
| AAIJ | 23AU PRL 131 171901 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| AAIJ | 15H PRL 114 062004 | R. Aaij <i>et al.</i> | (LHCb Collab.) |

NODE=B170M

NODE=B170M

OCCUR=2;SYCLP=A
SYCLP=ANODE=B170M;LINKAGE=AA
NODE=B170M;LINKAGE=C

NODE=B170M;LINKAGE=E

NODE=B170W

NODE=B170W

NODE=B170W;LINKAGE=AA

NODE=B170215;NODE=B170

DESIG=1

NODE=B170220

NODE=B170R01
NODE=B170R01

NODE=B170R01;LINKAGE=AA

NODE=B170

REFID=62459
REFID=56376