

$P_{c\bar{c}}(4457)^+$ $I(J^P) = \frac{1}{2}(?)$ Status: *

OMITTED FROM SUMMARY TABLE
was $P_c(4450)$

A resonance seen in $\Lambda_b^0 \rightarrow P_c^+ K^-$, then $P_c \rightarrow J/\psi p$, with a significance of 12 standard deviations. The $J/\psi p$ quark content is $u u d c \bar{c}$, a pentaquark. See also the $P_{c\bar{c}}(4380)^+$. In the best amplitude fit, the two states have opposite parity, one having $J = 3/2$, the other $J = 5/2$.

Extraction of the pentaquark signals requires some understanding of the dominant $K^- p$ background. AAIJ 15P used a model-dependent approach. AAIJ 16AG reanalyzed the data making minimal assumptions about the $K^- p$ background, and thus confirmed the strong significance of the pentaquark signals.

 $P_{c\bar{c}}(4457)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4457.3±0.6^{+4.1}_{-1.7}	AAIJ	19W LHCb	$p\bar{p}$ at 7, 8, 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4449.8±1.7±2.5	¹ AAIJ	15P LHCb	Repl. by AAIJ 19W
1 Considering $P_{c\bar{c}}(4440)$ and $P_{c\bar{c}}(4457)$ as a single resonance.			

NODE=B172M

NODE=B172M

NODE=B172M;LINKAGE=A

NODE=B172W

NODE=B172W

NODE=B172W;LINKAGE=A

NODE=B172215;NODE=B172

 $P_{c\bar{c}}(4457)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 J/\psi p$	seen
$\Gamma_2 \Lambda_c^+ \bar{D}^0$	not seen
$\Gamma_3 \Lambda_c^+ \pi^+ D^-$	not seen
$\Gamma_4 \Sigma_c(2455)^{++} D^-$	not seen
$\Gamma_5 \Sigma_c(2520)^{++} D^-$	not seen
$\Gamma_6 \bar{\Lambda}_c^- \pi^+ D^+$	[a] not seen
$\Gamma_7 \bar{\Sigma}_c(2455)^0 D^+$	[a] not seen
$\Gamma_8 \bar{\Sigma}_c(2520)^0 D^+$	[a] not seen
$\Gamma_9 \Lambda_c^+ \pi^+ D^{*-}$	not seen
$\Gamma_{10} \bar{\Lambda}_c^- \pi^+ D^{*+}$	[a] not seen

DESIG=1

DESIG=2

DESIG=3

DESIG=6

DESIG=8

DESIG=9

DESIG=5

DESIG=7

DESIG=4

DESIG=10

LINKAGE=PCC

 $P_{c\bar{c}}(4457)^+$ BRANCHING RATIOS

$\Gamma(J/\psi p)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	¹ POPOV	21 D0	$p\bar{p}$ at 1.96 TeV	
seen	AAIJ	19W LHCb	$p\bar{p}$ at 7, 8, 13 TeV	
seen	AAIJ	15P LHCb	$p\bar{p}$ at 7, 8 TeV	

NODE=B172225

NODE=B172R01
NODE=B172R01

NODE=B172R01;LINKAGE=A

¹ Search for J/ψ inclusive production in association with a charged particle, assumed to be a proton. POPOV 21 observes a resonant signal consistent with a superposition of the $P_{c\bar{c}}(4440)^+$ and $P_{c\bar{c}}(4457)^+$, using masses and widths measured by AAIJ 19W, at significance of 3σ .

$\Gamma(\Lambda_c^+ \bar{D}^0)/\Gamma_{\text{total}}$				Γ_2/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R00 NODE=B172R00
not seen	AAIJ	24Z	LHCb	$p p, 5.7 \text{ fb}^{-1}$ at 13 TeV	
$\Gamma(\Lambda_c^+ \pi^+ D^-)/\Gamma_{\text{total}}$				Γ_3/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R02 NODE=B172R02
not seen	AAIJ	24Z	LHCb	$p p, 5.7 \text{ fb}^{-1}$ at 13 TeV	
$\Gamma(\Sigma_c(2455)^{++} D^-)/\Gamma_{\text{total}}$				Γ_4/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R08 NODE=B172R08
not seen	AAIJ	24Z	LHCb	$p p, 5.7 \text{ fb}^{-1}$ at 13 TeV	OCCUR=2
$\Gamma(\Sigma_c(2520)^{++} D^-)/\Gamma_{\text{total}}$				Γ_5/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R07 NODE=B172R07
not seen	AAIJ	24Z	LHCb	$p p, 5.7 \text{ fb}^{-1}$ at 13 TeV	OCCUR=2
$\Gamma(\bar{\Lambda}_c^- \pi^+ D^+)/\Gamma_{\text{total}}$				Γ_6/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R10 NODE=B172R10
not seen	AAIJ	24Z	LHCb	$\bar{P}_{c\bar{c}}^- \rightarrow \Lambda_c^+ \pi^- D^-$	
$\Gamma(\bar{\Sigma}_c(2455)^0 D^+)/\Gamma_{\text{total}}$				Γ_7/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R04 NODE=B172R04
not seen	AAIJ	24Z	LHCb	$\bar{P}_{c\bar{c}}^- \rightarrow \Sigma_c(2455)^0 D^-$	OCCUR=2
$\Gamma(\bar{\Sigma}_c(2520)^0 D^+)/\Gamma_{\text{total}}$				Γ_8/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R06 NODE=B172R06
not seen	AAIJ	24Z	LHCb	$\bar{P}_{c\bar{c}}^- \rightarrow \Sigma_c(2520)^0 D^-$	OCCUR=2
$\Gamma(\Lambda_c^+ \pi^+ D^{*-})/\Gamma_{\text{total}}$				Γ_9/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R03 NODE=B172R03
not seen	AAIJ	24Z	LHCb	$p p, 5.7 \text{ fb}^{-1}$ at 13 TeV	OCCUR=2
$\Gamma(\bar{\Lambda}_c^- \pi^+ D^{*+})/\Gamma_{\text{total}}$				Γ_{10}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=B172R09 NODE=B172R09
not seen	AAIJ	24Z	LHCb	$\bar{P}_{c\bar{c}}^- \rightarrow \Lambda_c^+ \pi^- D^{*-}$	

$P_{c\bar{c}}(4457)^+$ REFERENCES

AAIJ	24Z	PR D110 032001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=62930
POPOV	21	PAN 83 1383	A.V. Popov <i>et al.</i>	(D0 Collab.)	REFID=61409
AAIJ	19W	PRL 122 222001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59789
AAIJ	16AG	PRL 117 082002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57460
AAIJ	15P	PRL 115 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56488

NODE=B172