

Σ BARYONS ($S = -1, I = 1$)

$$\Sigma^+ = uus, \quad \Sigma^0 = uds, \quad \Sigma^- = dds$$

 Σ^+

$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass $m = 1189.37 \pm 0.07$ MeV ($S = 2.2$)
 Mean life $\tau = (0.8018 \pm 0.0026) \times 10^{-10}$ s
 $c\tau = 2.404$ cm
 $(\tau_{\Sigma^+} - \tau_{\Sigma^-}) / \tau_{\Sigma^+} = -0.0006 \pm 0.0012$
 Magnetic moment $\mu = 2.458 \pm 0.010 \mu_N$ ($S = 2.1$)
 $(\mu_{\Sigma^+} + \mu_{\Sigma^-}) / \mu_{\Sigma^+} = 0.014 \pm 0.015$
 $\Gamma(\Sigma^+ \rightarrow n\ell^+\nu) / \Gamma(\Sigma^- \rightarrow n\ell^-\bar{\nu}_\ell) < 0.043$

Decay parameters

$p\pi^0$	$\alpha_0 = -0.982 \pm 0.014$
$\bar{p}\pi^0$	$\bar{\alpha}_0 = 0.99 \pm 0.04$
$(\alpha_0 + \bar{\alpha}_0) / (\alpha_0 - \bar{\alpha}_0)$	$= 0.00 \pm 0.04$
$p\pi^0$	$\phi_0 = (36 \pm 34)^\circ$
"	$\gamma_0 = 0.16$ [a]
"	$\Delta_0 = (187 \pm 6)^\circ$ [a]
$n\pi^+$	$\alpha_+ = (4.89 \pm 0.26) \times 10^{-2}$
"	$\phi_+ = (167 \pm 20)^\circ$ ($S = 1.1$)
$\bar{\alpha}_-$ FOR $\bar{\Sigma}^- \rightarrow \bar{n}\pi^-$	$= (-5.7 \pm 0.5) \times 10^{-2}$
$\bar{\alpha}_- / \bar{\alpha}_0$	$= (-5.7 \pm 0.6) \times 10^{-2}$
$(\alpha_+ + \bar{\alpha}_-) / (\alpha_+ - \bar{\alpha}_-)$	$= (-8 \pm 6) \times 10^{-2}$
"	$\gamma_+ = -0.97$ [a]
"	$\Delta_+ = (-73 \pm 133)^\circ$ [a]
$p\gamma$	$\alpha_\gamma = -0.69 \pm 0.05$

Σ^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$p\pi^0$	$(51.47 \pm 0.30) \%$		189
$n\pi^+$	$(48.43 \pm 0.30) \%$		185
$p\gamma$	$(1.04 \pm 0.06) \times 10^{-3}$	$S=2.4$	225
$n\pi^+\gamma$	[b] $(4.5 \pm 0.5) \times 10^{-4}$		185
$\Lambda e^+\nu_e$	$(2.3 \pm 0.4) \times 10^{-5}$		71

 **$\Delta S = \Delta Q$ (SQ) violating modes or
 $\Delta S = 1$ weak neutral current ($S1$) modes**

$ne^+\nu_e$	SQ	$< 5 \times 10^{-6}$	$CL=90\%$	224
$n\mu^+\nu_\mu$	SQ	$< 3.0 \times 10^{-5}$	$CL=90\%$	202
pe^+e^-	$S1$	$< 7 \times 10^{-6}$		225
$p\mu^+\mu^-$	$S1$	$(2.4 \pm 1.7) \times 10^{-8}$		121
$p\gamma_{dark}$		$< 3.2 \times 10^{-5}$	$CL=90\%$	—

 Σ^0

$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass $m = 1192.642 \pm 0.024$ MeV
 $m_{\Sigma^-} - m_{\Sigma^0} = 4.807 \pm 0.035$ MeV ($S = 1.1$)
 $m_{\Sigma^0} - m_\Lambda = 76.959 \pm 0.023$ MeV
 Mean life $\tau = (7.4 \pm 0.7) \times 10^{-20}$ s
 $c\tau = 2.22 \times 10^{-11}$ m
 Transition magnetic moment $|\mu_{\Sigma^0}| = 1.61 \pm 0.08 \mu_N$
 Magnetic moment $\mu = (-1.7 \pm 2.8) \times 10^{-3}$
 Magnetic moment $\mu = (2.1 \pm 3.0) \times 10^{-3}$
 Magnetic moment $\mu = (0.4 \pm 3.2) \times 10^{-3}$

NODE=BXXX025

NODE=S019

NODE=S019M;DTYPE=M
 NODE=S019T;DTYPE=T
 NODE=S019CTA;DTYPE=C;OUR EVAL
 NODE=S019DT;DTYPE=x
 NODE=S019MM;DTYPE=m
 NODE=S019MMD;DTYPE=i
 NODE=S019R7;DTYPE=Y;OUR LIM;
 \rightarrow UNCHECKED \leftarrow
 CLUMP=D
 NODE=S019A0;DTYPE=d;CLUMP=D
 NODE=S019A1;DTYPE=d;CLUMP=D
 NODE=S019A00;DTYPE=d;CLUMP=D
 NODE=S019F0;DTYPE=d;CLUMP=D
 NODE=S019G0;DTYPE=d;CLUMP=D;OUR EVAL;
 \rightarrow UNCHECKED \leftarrow
 NODE=S019D0;DTYPE=d;CLUMP=D;OUR EVAL;
 \rightarrow UNCHECKED \leftarrow
 NODE=S019A+;DTYPE=d;CLUMP=D
 NODE=S019F+;DTYPE=d;CLUMP=D
 NODE=S019A01;DTYPE=d;CLUMP=D
 NODE=S019A02;DTYPE=d;CLUMP=D
 NODE=S019A03;DTYPE=d;CLUMP=D
 NODE=S019G+;DTYPE=d;CLUMP=D;OUR EVAL;
 \rightarrow UNCHECKED \leftarrow
 NODE=S019D+;DTYPE=d;CLUMP=D;OUR EVAL;
 \rightarrow UNCHECKED \leftarrow
 NODE=S019AG;DTYPE=d;CLUMP=D

NODE=S019220;DESIG=1

DESIG=2

DESIG=5

DESIG=3

DESIG=4

NODE=S019;CLUMP=A

DESIG=7;OUR LIM; \rightarrow UNCHECKED \leftarrow
 DESIG=6;OUR LIM; \rightarrow UNCHECKED \leftarrow
 DESIG=8
 DESIG=9
 DESIG=195

NODE=S021

NODE=S021M;DTYPE=M
 NODE=S021D1;DTYPE=D
 NODE=S021DL;DTYPE=D
 NODE=S021T;DTYPE=T;OUR EVAL;
 \rightarrow UNCHECKED \leftarrow
 NODE=S021CTA;DTYPE=C;OUR EVAL
 NODE=S021MM;DTYPE=m
 NODE=S021A00;DTYPE=m
 NODE=S021A01;DTYPE=m
 NODE=S021A02;DTYPE=m

Σ^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
$\Lambda\gamma$	100 %		74
$\Lambda\gamma\gamma$	< 3 %	90%	74
$\Lambda e^+ e^-$	[c] 5×10^{-3}		74

 Σ^-

$$I(J^P) = 1(\frac{1}{2}^+)$$

Mass $m = 1197.449 \pm 0.029$ MeV (S = 1.1)
 $m_{\Sigma^-} - m_{\Sigma^+} = 8.08 \pm 0.08$ MeV (S = 1.9)
 $m_{\Sigma^-} - m_\Lambda = 81.766 \pm 0.029$ MeV (S = 1.1)
Mean life $\tau = (1.479 \pm 0.011) \times 10^{-10}$ s (S = 1.3)
 $c\tau = 4.434$ cm
Magnetic moment $\mu = -1.160 \pm 0.025 \mu_N$ (S = 1.7)
 Σ^- charge radius = 0.78 ± 0.10 fm

Decay parameters

$n\pi^-$	$\alpha_- = -0.068 \pm 0.008$
"	$\phi_- = (10 \pm 15)^\circ$
"	$\gamma_- = 0.98$ [a]
"	$\Delta_- = (249^{+12}_{-120})^\circ$ [a]
$ne^-\bar{\nu}_e$	$g_A/g_V = 0.340 \pm 0.017$ [d]
"	$f_2(0)/f_1(0) = 0.97 \pm 0.14$
"	$D = 0.11 \pm 0.10$
$\Lambda e^-\bar{\nu}_e$	$g_V/g_A = 0.01 \pm 0.10$ [d] (S = 1.5)
"	$g_{WM}/g_A = 2.4 \pm 1.7$ [d]

Σ^- DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	P (MeV/c)
$n\pi^-$	$(99.848 \pm 0.005) \%$		193
$n\pi^-\gamma$	[b] $(4.6 \pm 0.6) \times 10^{-4}$		193
$ne^-\bar{\nu}_e$	$(1.017 \pm 0.034) \times 10^{-3}$		230
$n\mu^-\bar{\nu}_\mu$	$(4.5 \pm 0.4) \times 10^{-4}$		210
$\Lambda e^-\bar{\nu}_e$	$(5.73 \pm 0.27) \times 10^{-5}$		79
$\Sigma^+ X$	$< 1.2 \times 10^{-4}$	90%	—

Lepton number (L) violating modes

$pe^- e^-$	L	$< 6.7 \times 10^{-5}$	90%	231
------------	-----	------------------------	-----	-----

 $\Sigma(1385) 3/2^+$

$$I(J^P) = 1(\frac{3}{2}^+)$$

$\Sigma(1385)^+$ mass $m = 1382.83 \pm 0.34$ MeV (S = 1.9)
 $\Sigma(1385)^0$ mass $m = 1383.7 \pm 1.0$ MeV (S = 1.4)
 $\Sigma(1385)^-$ mass $m = 1387.2 \pm 0.5$ MeV (S = 2.2)
 $\Sigma(1385)^+$ full width $\Gamma = 36.2 \pm 0.7$ MeV
 $\Sigma(1385)^0$ full width $\Gamma = 36 \pm 5$ MeV
 $\Sigma(1385)^-$ full width $\Gamma = 39.4 \pm 2.1$ MeV (S = 1.7)
Below $\bar{K}N$ threshold

NODE=S021225;DESIG=1;OUR EVAL;
→UNCHECKED←
DESIG=3
DESIG=2;OUR EVAL;→UNCHECKED←

NODE=S020

NODE=S020M;DTYPE=M
NODE=S020D;DTYPE=D
NODE=S020DL;DTYPE=D
NODE=S020T;DTYPE=T
NODE=S020CTA;DTYPE=C;OUR EVAL
NODE=S020MM;DTYPE=m
NODE=S020CR;DTYPE=x

CLUMP=D

NODE=S020A-;DTYPE=d;CLUMP=D
NODE=S020F-;DTYPE=d;CLUMP=D
NODE=S020G-;DTYPE=d;CLUMP=D;OUR EVAL;
→UNCHECKED←
NODE=S020D-;DTYPE=d;CLUMP=D;OUR EVAL;
→UNCHECKED←
NODE=S020AV2;DTYPE=d;CLUMP=D
NODE=S020F2;DTYPE=d;CLUMP=D
NODE=S020TC;DTYPE=d;CLUMP=D
NODE=S020AV;DTYPE=d;CLUMP=D
NODE=S020WM;DTYPE=d;CLUMP=D

NODE=S020230;DESIG=1
DESIG=2
DESIG=4
DESIG=3
DESIG=5
DESIG=193

NODE=S020;CLUMP=L
DESIG=192

NODE=B043

NODE=B043M+;DTYPE=M
NODE=B043M0;DTYPE=M
NODE=B043M-;DTYPE=M
NODE=B043W+;DTYPE=G
NODE=B043W0;DTYPE=G
NODE=B043W-;DTYPE=G

$\Sigma(1385)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$\Lambda\pi$	(87.0 \pm 1.5) %	208	
$\Sigma\pi$	(11.7 \pm 1.5) %	129	
$\Lambda\gamma$	(1.25 $^{+0.13}_{-0.12}$) %	241	
$\Sigma^+\gamma$	(7.0 \pm 1.7) \times 10 $^{-3}$	180	
$\Sigma^-\gamma$	< 2.4 \times 10 $^{-4}$	90%	173

 $\Sigma(1660)$ 1/2 $^+$

$I(J^P) = 1(\frac{1}{2}^+)$

Re(pole position) = 1585 \pm 20 MeV
 $-2\text{Im}(\text{pole position}) = 290^{+140}_{-40}$ MeV
 Mass m = 1640 to 1680 (\approx 1660) MeV
 Full width Γ = 100 to 300 (\approx 200) MeV

$\Sigma(1660)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	0.05 to 0.15 (\approx 0.10)	405
$\Lambda\pi$	(35 \pm 12) %	440
$\Sigma\pi$	(37 \pm 10) %	387
$\Sigma\sigma$	(20 \pm 8) %	—
$\Lambda(1405)\pi$	(4.0 \pm 2.0) %	199

 $\Sigma(1670)$ 3/2 $^-$

$I(J^P) = 1(\frac{3}{2}^-)$

Mass m = 1665 to 1685 (\approx 1675) MeV
 Full width Γ = 40 to 100 (\approx 70) MeV

$\Sigma(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	0.06 to 0.12	419
$\Lambda\pi$	5–15 %	452
$\Sigma\pi$	30–60 %	398
$\Sigma\sigma$	(7.0 \pm 3.0) %	—

 $\Sigma(1750)$ 1/2 $^-$

$I(J^P) = 1(\frac{1}{2}^-)$

Mass m = 1700 to 1800 (\approx 1750) MeV
 Full width Γ = 100 to 200 (\approx 150) MeV

$\Sigma(1750)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$N\bar{K}$	0.06 to 0.12	486
$\Lambda\pi$	(14 \pm 5) %	507
$\Sigma\pi$	(16 \pm 4) %	456
$\Sigma\eta$	15–55 %	98
$\Sigma(1385)\pi$, D-wave	< 1 %	305
$\Lambda(1520)\pi$	(2.0 \pm 1.0) %	175
$N\bar{K}^*(892)$, S=1/2	(8 \pm 4) %	†

 $\Sigma(1775)$ 5/2 $^-$

$I(J^P) = 1(\frac{5}{2}^-)$

Mass m = 1770 to 1780 (\approx 1775) MeV
 Full width Γ = 105 to 135 (\approx 120) MeV

NODE=B043235;DESIG=1;OUR EST
 DESIG=2;OUR EST
 DESIG=3
 DESIG=6
 DESIG=5

NODE=B079

NODE=B079RE;DTYPE=i
 NODE=B079IM;DTYPE=i
 NODE=B079M;DTYPE=M;OUR EST;
 \rightarrow UNCHECKED \leftarrow
 NODE=B079W;DTYPE=G;OUR EST;
 \rightarrow UNCHECKED \leftarrow

NODE=B079215;DESIG=1
 DESIG=3
 DESIG=2
 DESIG=5
 DESIG=6

NODE=B044

NODE=B044M;DTYPE=M;OUR EST;
 \rightarrow UNCHECKED \leftarrow
 NODE=B044W;DTYPE=G;OUR EST;
 \rightarrow UNCHECKED \leftarrow

NODE=B044215;DESIG=1
 DESIG=2;OUR EST
 DESIG=3;OUR EST
 DESIG=62

NODE=B057

NODE=B057M;DTYPE=M;OUR EST;
 \rightarrow UNCHECKED \leftarrow
 NODE=B057W;DTYPE=G;OUR EST;
 \rightarrow UNCHECKED \leftarrow

NODE=B057215;DESIG=1
 DESIG=3
 DESIG=4
 DESIG=2;OUR EST
 DESIG=5
 DESIG=6
 DESIG=7

NODE=B045

NODE=B045M;DTYPE=M;OUR EST;
 \rightarrow UNCHECKED \leftarrow
 NODE=B045W;DTYPE=G;OUR EST;
 \rightarrow UNCHECKED \leftarrow

$\Sigma(1775)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)	
$N\bar{K}$	37–43%	508	NODE=B045215;DESIG=1;OUR EST
$\Lambda\pi$	14–20%	525	DESIG=2;OUR EST
$\Sigma\pi$	2–5%	475	DESIG=5;OUR EST
$\Sigma(1385)\pi$	8–12%	327	DESIG=4;OUR EST
$\Lambda(1520)\pi$, P -wave	17–23%	202	DESIG=3;OUR EST

 $\Sigma(1910) \frac{3}{2}^-$

$I(J^P) = 1(\frac{3}{2}^-)$

was $\Sigma(1940)$

Mass $m = 1870$ to 1950 (≈ 1910) MeV
 Full width $\Gamma = 150$ to 300 (≈ 220) MeV

$\Sigma(1910)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)	
$N\bar{K}$	0.01 to 0.05 (≈ 0.02)	615	NODE=B098M;DTYPE=M;OUR EST;
$\Lambda\pi$	(6 \pm 4) %	619	→ UNCHECKED ↵
$\Sigma\pi$	(86 \pm 21) %	574	NODE=B098W;DTYPE=G;OUR EST;
$\Sigma(1385)\pi$	seen	439	→ UNCHECKED ↵
$\Lambda(1520)\pi$	seen	329	
$\Delta(1232)\bar{K}$	(3.0 \pm 1.0) %	377	
$N\bar{K}^*(892)$	seen	274	
$N\bar{K}^*(892)$, $S=1/2$, D -wave	(1.0 \pm 1.0) %	274	

 $\Sigma(1915) \frac{5}{2}^+$

$I(J^P) = 1(\frac{5}{2}^+)$

Mass $m = 1900$ to 1935 (≈ 1915) MeV
 Full width $\Gamma = 80$ to 160 (≈ 120) MeV

$\Sigma(1915)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)	
$N\bar{K}$	0.05 to 0.15	618	NODE=B046215;DESIG=1
$\Lambda\pi$	(6.0 \pm 2.0) %	623	DESIG=2
$\Sigma\pi$	(10.0 \pm 2.0) %	577	DESIG=3
$\Sigma(1385)\pi$, P -wave	(2.0 \pm 2.0) %	443	DESIG=4
$\Sigma(1385)\pi$, F -wave	(4.0 \pm 2.0) %	443	DESIG=5
$\Lambda(1520)\pi$, D -wave	(8.0 \pm 2.0) %	334	DESIG=190
$N\bar{K}^*(892)$, $S=1/2$, F -wave	(5.0 \pm 3.0) %	282	DESIG=183
$N\bar{K}^*(892)$, $S=3/2$, F -wave	(5.0 \pm 2.0) %	282	DESIG=185
$\Delta\bar{K}$, P -wave	(16 \pm 5) %	383	DESIG=188
$\Delta\bar{K}$, F -wave	(5.0 \pm 3.0) %	383	DESIG=189

 $\Sigma(2030) \frac{7}{2}^+$

$I(J^P) = 1(\frac{7}{2}^+)$

Mass $m = 2025$ to 2040 (≈ 2030) MeV
 Full width $\Gamma = 150$ to 200 (≈ 180) MeV

$\Sigma(2030)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)	
$N\bar{K}$	17–23 %	702	NODE=B047215;DESIG=1;OUR EST
$\Lambda\pi$	17–23 %	700	DESIG=2;OUR EST
$\Sigma\pi$	5–10 %	657	DESIG=3;OUR EST
ΞK	<2 %	422	DESIG=4;OUR EST
$\Sigma(1385)\pi$	5–15 %	532	DESIG=10;OUR EST
$\Sigma(1385)\pi$, F -wave	(1.0 \pm 1.0) %	532	DESIG=21
$\Lambda(1520)\pi$	10–20 %	431	DESIG=181;OUR EST
$\Delta(1232)\bar{K}$	10–20 %	498	DESIG=182;OUR EST
$\Delta(1232)\bar{K}$, F -wave	(15 \pm 5) %	498	DESIG=8
$\Delta(1232)\bar{K}$, H -wave	(1.0 \pm 1.0) %	498	DESIG=9
$N\bar{K}^*(892)$, $S=3/2$, F -wave	(14 \pm 8) %	439	DESIG=12

NOTES

[a] The decay parameters γ and Δ are calculated from α and ϕ using

$$\gamma = \sqrt{1-\alpha^2} \cos\phi, \quad \tan\Delta = -\frac{1}{\alpha} \sqrt{1-\alpha^2} \sin\phi.$$

See the “Note on Baryon Decay Parameters” in the neutron Particle Listings.

[b] See the Listings for the pion momentum range used in this measurement.

[c] A theoretical value using QED.

[d] The parameters g_A , g_V , and g_{WM} for semileptonic modes are defined by $\bar{B}_f[\gamma_\lambda(g_V + g_A\gamma_5) + i(g_{WM}/m_{B_i}) \sigma_{\lambda\nu} q^\nu]B_i$, and ϕ_{AV} is defined by $g_A/g_V = |g_A/g_V|e^{i\phi_{AV}}$. See the “Note on Baryon Decay Parameters” in the neutron Particle Listings.

LINKAGE=SBE

LINKAGE=SD

LINKAGE=SU

LINKAGE=SBD