NODE=M002

$\eta'(958)$

$I^G(J^{PC}) = 0^+(0^{-+})$

$\eta'(958)$ MASS

NODE=M002M

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	COMMENT
957.78 ±0.06 OUR AVERAGE					
957.793±0.054±0.036	3.9k	LIBBY	08	CLEO	$J/\psi \to \gamma\eta'$
957.9 ±0.2 ±0.6	4800	WURZINGER	96	SPEC	1.68 $pd \to {}^3\mathrm{He}\eta'$
957.46 ±0.33		DUANE	74	MMS	$\pi^- p \to n\mathrm{MM}$
958.2 ±0.5	1414	DANBURG	73	HBC	2.2 $K^- p \to \Lambda\eta'$
958 ±1	400	JACOBS	73	HBC	2.9 $K^- p \to \Lambda\eta'$
956.1 ±1.1	3415	[1] BASILE	71	CNTR	1.6 $\pi^- p \to n\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
957.5 ±0.2		BAI	04J	BES2	$J/\psi \to \gamma\gamma\pi^+\pi^-$
959 ±1	630	[2] BELADIDZE	92C	VES	36 π^- Be $\to \pi^-\eta'\eta$ Be
958 ±1	340	[2] ARMSTRONG	91B	OMEG	300 $pp \to pp\eta\pi^+\pi^-$
958.2 ±0.4	622	[2] AUGUSTIN	90	DM2	$J/\psi \to \gamma\eta\pi^+\pi^-$
957.8 ±0.2	2420	[2] AUGUSTIN	90	DM2	$J/\psi \to \gamma\gamma\pi^+\pi^-$ OCCUR=2
956.3 ±1.0	143	[2] GIDAL	87	MRK2	$e^+e^- \to e^+e^-\eta\pi^+\pi^-$
957.4 ±1.4	535	[3] BASILE	71	CNTR	1.6 $\pi^- p \to n\eta'$ OCCUR=2
957 ±1		RITTENBERG	69	HBC	1.7–2.7 $K^- p$

[1] Using all η' decays. NODE=M002M;LINKAGE=BS

[2] Systematic uncertainty not estimated. NODE=M002M;LINKAGE=NS

[3] Using η' decays into neutrals. Not independent of the other listed BASILE 71 η' mass measurement. NODE=M002M;LINKAGE=BA

$\eta'(958)$ WIDTH

NODE=M002W

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	CHG	COMMENT
0.188±0.006 OUR FIT						
0.230±0.021 OUR AVERAGE						
0.226±0.017±0.014	2300	CZERWINSKI	10	MMS		$pp \to pp\eta'$
0.40 ±0.22	4800	WURZINGER	96	SPEC		1.68 $pd \to {}^3\mathrm{He}\eta'$
0.28 ±0.10	1000	BINNIE	79	MMS	0	$\pi^- p \to n\mathrm{MM}$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.20 ±0.04		BAI	04J	BES2		$J/\psi \to \gamma\gamma\pi^+\pi^-$

$\eta'(958)$ DECAY MODES

NODE=M002215;NODE=M002

	Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1	$\pi^+\pi^-\eta$	(42.5 ±0.5) %		DESIG=1
Γ_2	$\rho^0\gamma$(including non-resonant $\pi^+\ \pi^-\ \gamma$)	(29.5 ±0.4) %		DESIG=9
Γ_3	$\rho^0\gamma$			DESIG=213
Γ_4	$\pi^0\pi^0\eta$	(22.4 ±0.5) %		DESIG=2
Γ_5	$\omega\gamma$	(2.52 ±0.07) %		DESIG=7
Γ_6	ωe^+e^-	(2.0 ±0.4) $\times 10^{-4}$		DESIG=205
Γ_7	$\gamma\gamma$	(2.307±0.033) %		DESIG=6
Γ_8	$3\pi^0$	(2.50 ±0.17) $\times 10^{-3}$		DESIG=8
Γ_9	$\mu^+\mu^-\gamma$	(1.13 ±0.28) $\times 10^{-4}$		DESIG=20
Γ_{10}	$\pi^+\pi^-\mu^+\mu^-$	(1.9 ±0.4) $\times 10^{-5}$		DESIG=201
Γ_{11}	$\pi^+\pi^-\pi^0$	(3.61 ±0.17) $\times 10^{-3}$		DESIG=121
Γ_{12}	$(\pi^+\pi^-\pi^0)$ S-wave	(3.8 ±0.5) $\times 10^{-3}$		DESIG=211
Γ_{13}	$\pi^\mp\rho^\pm$	(7.4 ±2.3) $\times 10^{-4}$		DESIG=210
Γ_{14}	$2(\pi^+\pi^-)$	(8.3 ±0.9) $\times 10^{-5}$		DESIG=131
Γ_{15}	$\pi^+\pi^-2\pi^0$	(1.8 ±0.4) $\times 10^{-4}$		DESIG=202
Γ_{16}	$2(\pi^+\pi^-)$ neutrals	< 1 %	95%	DESIG=132

Γ_{17}	$2(\pi^+\pi^-)\pi^0$	$<$	1.8		$\times 10^{-3}$	90%	DESIG=141
Γ_{18}	$2(\pi^+\pi^-)2\pi^0$	$<$	1		%	95%	DESIG=15
Γ_{19}	$3(\pi^+\pi^-)$	$<$	3.1		$\times 10^{-5}$	90%	DESIG=203
Γ_{20}	$K^\pm\pi^\mp$	$<$	4		$\times 10^{-5}$	90%	DESIG=207
Γ_{21}	$\pi^+\pi^- e^+ e^-$	(	2.42	± 0.10)	$\times 10^{-3}$		DESIG=10
Γ_{22}	$\pi^+ e^- \nu_e +$ c.c.	$<$	2.1		$\times 10^{-4}$	90%	DESIG=204
Γ_{23}	$\gamma e^+ e^-$	(	4.91	± 0.27)	$\times 10^{-4}$		DESIG=28
Γ_{24}	$\pi^0\gamma\gamma$	(	3.20	± 0.24)	$\times 10^{-3}$		DESIG=24
Γ_{25}	$\pi^0\gamma\gamma$(non resonant)	(	6.2	± 0.9)	$\times 10^{-4}$		DESIG=212
Γ_{26}	$\eta\gamma\gamma$	$<$	1.33		$\times 10^{-4}$	90%	DESIG=214
Γ_{27}	$4\pi^0$	$<$	4.94		$\times 10^{-5}$	90%	DESIG=26
Γ_{28}	$e^+ e^-$	$<$	5.6		$\times 10^{-9}$	90%	DESIG=150
Γ_{29}	$e^+ e^- e^+ e^-$	(	4.5	± 1.1)	$\times 10^{-6}$		DESIG=215
Γ_{30}	invisible	$<$	6		$\times 10^{-4}$	90%	DESIG=200

Charge conjugation (C), Parity (P), Lepton family number (LF) violating modes NODE=M002;CLUMP=B

Γ_{31}	$\pi^+\pi^-$	P,CP		$<$	1.8	$\times 10^{-5}$	90%	DESIG=111
Γ_{32}	$\pi^0\pi^0$	P,CP		$<$	4	$\times 10^{-4}$	90%	DESIG=25
Γ_{33}	$\pi^0 e^+ e^-$	C	[a]	$<$	1.4	$\times 10^{-3}$	90%	DESIG=16
Γ_{34}	$\pi^0\rho^0$	C		$<$	4	%	90%	DESIG=18
Γ_{35}	$\eta e^+ e^-$	C	[a]	$<$	2.4	$\times 10^{-3}$	90%	DESIG=17
Γ_{36}	3γ	C		$<$	1.0	$\times 10^{-4}$	90%	DESIG=23
Γ_{37}	$\mu^+\mu^-\pi^0$	C	[a]	$<$	6.0	$\times 10^{-5}$	90%	DESIG=22
Γ_{38}	$\mu^+\mu^-\eta$	C	[a]	$<$	1.5	$\times 10^{-5}$	90%	DESIG=21
Γ_{39}	$e\mu$	LF		$<$	4.7	$\times 10^{-4}$	90%	DESIG=27

[a] C parity forbids this to occur as a single-photon process. LINKAGE=CS

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 20 branching ratios uses 52 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 69.5$ for 44 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i, including the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	−25							
x_4	−75	−43						
x_5	−7	−6	−2					
x_7	−11	−7	9	−1				
x_8	−17	−10	19	0	2			
x_{11}	−1	−1	−1	0	0	0		
x_{21}	−8	30	−14	−2	−2	−3	0	
Γ	11	−10	−1	1	−40	0	0	−3
	x_1	x_2	x_4	x_5	x_7	x_8	x_{11}	x_{21}

	Mode	Rate (MeV)	
Γ_1	$\pi^+\pi^-\eta$	0.0799 $\pm$0.0029	DESIG=1
Γ_2	$\rho^0\gamma$(including non-resonant $\pi^+\ \pi^-\ \gamma$)	0.0554 $\pm$0.0019	DESIG=9
Γ_4	$\pi^0\pi^0\eta$	0.0421 $\pm$0.0017	DESIG=2
Γ_5	$\omega\gamma$	0.00474 $\pm$0.00020	DESIG=7
Γ_7	$\gamma\gamma$	0.00434 $\pm$0.00013	DESIG=6
Γ_8	$3\pi^0$	(4.7 $\pm$0.4) $\times 10^{-4}$	DESIG=8
Γ_{11}	$\pi^+\pi^-\pi^0$	(6.8 $\pm$0.4) $\times 10^{-4}$	DESIG=121
Γ_{21}	$\pi^+\pi^- e^+ e^-$	(4.54 $\pm$0.23) $\times 10^{-4}$	DESIG=10

$\eta'(958)$ PARTIAL WIDTHS

NODE=M002220

$\Gamma(\gamma\gamma)$ Γ_7

NODE=M002W4
NODE=M002W4

VALUE (keV)	EVTS	DOCUMENT ID		TECN	COMMENT
4.34±0.14 OUR FIT					
4.28±0.19 OUR AVERAGE					
4.17±0.10±0.27	2000	[1] ACCIARRI	98Q	L3	$e^+e^- \to e^+e^-\pi^+\pi^-\gamma$
4.53±0.29±0.51	266	KARCH	92	CBAL	$e^+e^- \to e^+e^-\eta\pi^0\pi^0$
3.61±0.13±0.48		[2] BEHREND	91	CELL	$e^+e^- \to e^+e^-\eta'(958)$
4.6 ±1.1 ±0.6	23	BARU	90	MD1	$e^+e^- \to e^+e^-\pi^+\pi^-\gamma$
4.57±0.25±0.44		BUTLER	90	MRK2	$e^+e^- \to e^+e^-\eta'(958)$
5.08±0.24±0.71	547	[3] ROE	90	ASP	$e^+e^- \to e^+e^-2\gamma$
3.8 ±0.7 ±0.6	34	AIHARA	88C	TPC	$e^+e^- \to e^+e^-\eta\pi^+\pi^-$
4.9 ±0.5 ±0.5	136	[4] WILLIAMS	88	CBAL	$e^+e^- \to e^+e^-2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.7 ±0.6 ±0.9	143	[5] GIDAL	87	MRK2	$e^+e^- \to e^+e^-\eta\pi^+\pi^-$
4.0 ±0.9		[6] BARTEL	85E	JADE	$e^+e^- \to e^+e^-2\gamma$

[1] No non-resonant $\pi^+\pi^-$ contribution found. NODE=M002W4;LINKAGE=AC
[2] Reevaluated by us using B($\eta' \to \rho(770)\gamma$) = (30.2 ± 1.3)%. NODE=M002W4;LINKAGE=K1
[3] Reevaluated by us using B($\eta' \to \gamma\gamma$) = (2.11 ± 0.13)%. NODE=M002W4;LINKAGE=K2
[4] Reevaluated by us using B($\eta' \to \gamma\gamma$) = (2.11 ± 0.13)%. NODE=M002W4;LINKAGE=K3
[5] Superseded by BUTLER 90. NODE=M002W4;LINKAGE=C
[6] Systematic error not evaluated. NODE=M002W4;LINKAGE=A

$\Gamma(e^+e^-)$ Γ_{28}

NODE=M002W1
NODE=M002W1

VALUE (eV)	CL%	DOCUMENT ID		TECN	COMMENT
$<1.1\times10^{-3}$	90	[1,2] ACHASOV	15	SND	0.958 $e^+e^- \to \pi\pi\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.0\times10^{-3}$	90	[2] ACHASOV	15	SND	0.958 $e^+e^- \to \pi\pi\eta$
$<2.4\times10^{-3}$	90	[2] AKHMETSHIN	15	CMD3	0.958 $e^+e^- \to \pi^+\pi^-\eta$

OCCUR=2

[1] Combining data of ACHASOV 15 and AKHMETSHIN 15. NODE=M002W1;LINKAGE=A
[2] Using η and η' branching fractions from PDG 14. NODE=M002W1;LINKAGE=B

$\eta'(958)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(total)$

NODE=M002223

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

NODE=M002223

$\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{total}$ $\Gamma_7\Gamma_2/\Gamma$

NODE=M002G1
NODE=M002G1

VALUE (keV)	EVTS	DOCUMENT ID		TECN	COMMENT
1.28±0.04 OUR FIT					
1.26±0.07 OUR AVERAGE	Error includes scale factor of 1.2.				
1.09±0.04±0.13		BEHREND	91	CELL	$e^+e^- \to e^+e^-\rho(770)^0\gamma$
1.35±0.09±0.21		AIHARA	87	TPC	$e^+e^- \to e^+e^-\rho\gamma$
1.13±0.04±0.13	867	ALBRECHT	87B	ARG	$e^+e^- \to e^+e^-\rho\gamma$
1.53±0.09±0.21		ALTHOFF	84E	TASS	$e^+e^- \to e^+e^-\rho\gamma$
1.14±0.08±0.11	243	BERGER	84B	PLUT	$e^+e^- \to e^+e^-\rho\gamma$
1.73±0.34±0.35	95	JENNI	83	MRK2	$e^+e^- \to e^+e^-\rho\gamma$
1.49±0.13±0.027	213	BARTEL	82B	JADE	$e^+e^- \to e^+e^-\rho\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.85±0.31±0.24	43	BEHREND	82C	CELL	$e^+e^- \to e^+e^-\rho\gamma$

$\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{total}$ $\Gamma_7\Gamma_4/\Gamma$

NODE=M002G2
NODE=M002G2

VALUE (keV)	DOCUMENT ID		TECN	COMMENT
0.97±0.04 OUR FIT Error includes scale factor of 1.1.				
0.92±0.06±0.11	[1] KARCH	92	CBAL	$e^+e^- \to e^+e^-\eta\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.95±0.05±0.08	[2] KARCH	90	CBAL	$e^+e^- \to e^+e^-\eta\pi^0\pi^0$
1.00±0.08±0.10	[2,3] ANTREASYAN	87	CBAL	$e^+e^- \to e^+e^-\eta\pi^0\pi^0$

[1] Reevaluated by us using B($\eta \to \gamma\gamma$) = (39.21 ± 0.34)%. Supersedes ANTREASYAN 87 and KARCH 90. NODE=M002G2;LINKAGE=K4
[2] Superseded by KARCH 92. NODE=M002G2;LINKAGE=A
[3] Using BR($\eta \to 2\gamma$)=(38.9 ± 0.5)%. NODE=M002G2;LINKAGE=D

$\eta'(958)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma(total)$

NODE=M002224

$\Gamma(\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_1\Gamma_{28}/\Gamma$

NODE=M002G01
NODE=M002G01

VALUE (10^{-3} eV)	CL%	DOCUMENT ID		TECN	COMMENT
<1.0	90	[1] AKHMETSHIN	15	CMD3	0.958 $e^+e^- \to \pi^+\pi^-\eta$

[1] AKHMETSHIN 15 reports $[\Gamma(\eta'(958) \to \pi^+\pi^-\eta) \times \Gamma(\eta'(958) \to e^+e^-)/\Gamma_{total}] \times [B(\eta \to 2\gamma)] < 4.1 \times 10^{-4}$ eV which we divide by our best value $B(\eta \to 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002G01;LINKAGE=A

$\eta'(958)$ BRANCHING RATIOS

NODE=M002230

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{total}$ Γ_1/Γ

NODE=M002R47
NODE=M002R47

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
42.5 ±0.5 OUR FIT	Error includes scale factor of 1.1.				
41.24±0.08±1.24	312k	ABLIKIM	19T	BES	$J/\psi \to \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
42.4 ±1.1 ±0.4	1.2k	[1] PEDLAR	09	CLEO	$J/\psi \to \gamma\eta'$

[1] Not independent of other η' branching fractions and ratios in PEDLAR 09.

NODE=M002R47;LINKAGE=PE

$\Gamma(\pi^+\pi^-\eta(\text{charged decay}))/\Gamma_{total}$ $0.2804\Gamma_1/\Gamma$

NODE=M002R3
NODE=M002R3

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.1191±0.0015 OUR FIT	Error includes scale factor of 1.1.				
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.123 ±0.014	107	RITTENBERG	69	HBC	1.7–2.7 K^-p
0.10 ±0.04	10	LONDON	66	HBC	2.24 $K^-p \to \Lambda 2\pi^+ 2\pi^- \pi^0$
0.07 ±0.04	7	BADIER	65B	HBC	3 K^-p

$\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))/\Gamma_{total}$ $0.7196\Gamma_1/\Gamma$

NODE=M002R1
NODE=M002R1

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.306±0.004 OUR FIT	Error includes scale factor of 1.1.				
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.314±0.026	281	RITTENBERG	69	HBC	1.7–2.7 K^-p

$\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{total}$ Γ_2/Γ

NODE=M002R6
NODE=M002R6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
29.5 ± 0.4 OUR FIT	Error includes scale factor of 1.1.				
29.90± 0.03±0.55	913k	ABLIKIM	19T	BES	$J/\psi \to \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
28.7 ± 0.7 ±0.4	0.2k	[1] PEDLAR	09	CLEO	$J/\psi \to \gamma\eta'$
32.9 ± 3.3	298	RITTENBERG	69	HBC	1.7–2.7 K^-p
20 ±10	20	LONDON	66	HBC	2.24 $K^-p \to \Lambda\pi^+\pi^-\gamma$
34 ± 9	35	BADIER	65B	HBC	3 K^-p

[1] Not independent of other η' branching fractions and ratios in PEDLAR 09.

NODE=M002R6;LINKAGE=PE

$\Gamma(\rho^0\gamma)/\Gamma_{total}$ Γ_3/Γ

NODE=M002R66
NODE=M002R66

VALUE (%)	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
33.34±0.06±1.60	970k	[1] ABLIKIM	18C	BES3	$\eta'(958) \to \gamma\pi^+\pi^-$
34.43±0.52±1.97	970k	[2] ABLIKIM	18C	BES3	$\eta'(958) \to \gamma\pi^+\pi^-$

OCCUR=2

[1] From a fit to $\pi^+\pi^-$ mass using $\rho(770)$, $\omega(782)$, and box anomaly components.

NODE=M002R66;LINKAGE=A

[2] From a fit to $\pi^+\pi^-$ mass using $\rho(770)$, $\omega(782)$, and $\rho(1450)$ components.

NODE=M002R66;LINKAGE=B

$\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta)$ Γ_2/Γ_1

NODE=M002R43
NODE=M002R43

VALUE	DOCUMENT ID		TECN	COMMENT
0.694±0.014 OUR FIT	Error includes scale factor of 1.1.			
0.683±0.020 OUR AVERAGE				
0.677±0.024±0.011	PEDLAR	09	CLE3	$J/\psi \to \eta'\gamma$
0.69 ±0.03	ABLIKIM	06E	BES2	$J/\psi \to \eta'\gamma$

$\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))$ $\Gamma_2/0.714\Gamma_1$

NODE=M002R27
NODE=M002R27

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.972±0.020 OUR FIT Error includes scale factor of 1.1.					
0.97 ±0.09 OUR AVERAGE					
0.70 ±0.22		AMSLER	04B	CBAR	0 $\bar{p}p \to \pi^+\pi^-\eta$
1.07 ±0.17		BELADIDZE	92C	VES	36 π^- Be $\to \pi^-\eta'\eta$ Be
0.92 ±0.14	473	DANBURG	73	HBC	2.2 $K^-p \to \Lambda X^0$
1.11 ±0.18	192	JACOBS	73	HBC	2.9 $K^-p \to \Lambda X^0$

$\Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$ Γ_4/Γ

NODE=M002R48
NODE=M002R48

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
22.4 ±0.6 OUR FIT Error includes scale factor of 1.1.					
21.36±0.10±0.92	52k	ABLIKIM	19T	BES	$J/\psi \to \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
23.5 ±1.3 ±0.4	3.2k	[1] PEDLAR	09	CLEO	$J/\psi \to \gamma\eta'$

[1] Not independent of other η' branching fractions and ratios in PEDLAR 09. NODE=M002R48;LINKAGE=PE

$\Gamma(\pi^0\pi^0\eta(3\pi^0\text{decay}))/\Gamma_{\text{total}}$ $0.321\Gamma_4/\Gamma$

NODE=M002R26
NODE=M002R26

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0718±0.0018 OUR FIT Error includes scale factor of 1.1.					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.11 ±0.06	4	BENSINGER	70	DBC	2.2 $\pi^+ d$

$\Gamma(\pi^0\pi^0\eta)/\Gamma(\pi^+\pi^-\eta)$ Γ_4/Γ_1

NODE=M002R45
NODE=M002R45

VALUE	DOCUMENT ID		TECN	COMMENT
0.527±0.019 OUR FIT Error includes scale factor of 1.1.				
0.555±0.043±0.013	PEDLAR	09	CLE3	$J/\psi \to \eta'\gamma$

$\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi\pi\eta)$ $\Gamma_2/(\Gamma_1+\Gamma_4)$

NODE=M002R7
NODE=M002R7

VALUE	DOCUMENT ID		TECN	COMMENT
0.454±0.009 OUR FIT Error includes scale factor of 1.1.				
0.43 ±0.02 ±0.02	BARBERIS	98C	OMEG	450 $pp \to p_f\eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.31 ±0.15	DAVIS	68	HBC	5.5 K^-p

$\Gamma(\omega\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

NODE=M002R49
NODE=M002R49

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
2.52 ±0.07 OUR FIT					
2.50 ±0.07 OUR AVERAGE					
2.489±0.018±0.074	23k	ABLIKIM	19T	BES	$J/\psi \to \gamma\eta'$
2.55 ±0.03 ±0.16	33.2k	[1] ABLIKIM	15AD	BES3	$J/\psi \to \eta'\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.34 ±0.30 ±0.04	70	[2] PEDLAR	09	CLEO	$J/\psi \to \gamma\eta'$

[1] Using B($J/\psi \to \eta'\gamma$) = (5.15 ± 0.16)×10^{-3} and B($\omega \to \pi^+\pi^-\pi^0$) = (89.2 ± 0.7)%. NODE=M002R49;LINKAGE=A
[2] Not independent of other η' branching fractions and ratios in PEDLAR 09. NODE=M002R49;LINKAGE=PE

$\Gamma(\omega\gamma)/\Gamma(\pi^+\pi^-\eta)$ Γ_5/Γ_1

NODE=M002R17
NODE=M002R17

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0593±0.0018 OUR FIT Error includes scale factor of 1.1.					
0.055 ±0.007 ±0.001		PEDLAR	09	CLE3	$J/\psi \to \eta'\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.068 ±0.013	68	ZANFINO	77	ASPK	8.4 $\pi^- p$

$\Gamma(\omega\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_5/Γ_4

NODE=M002R33
NODE=M002R33

VALUE	DOCUMENT ID		TECN	COMMENT
0.113±0.004 OUR FIT				
0.147±0.016	ALDE	87B	GAM2	38 $\pi^- p \to n4\gamma$

$\Gamma(\omega e^+e^-)/\Gamma(\omega\gamma)$ Γ_6/Γ_5

NODE=M002R60
NODE=M002R60

VALUE (units 10^{-3})	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.71±1.34±0.54	[1] ABLIKIM	15AD	BES3	$J/\psi \to \eta'\gamma$

[1] Obtained from other ABLIKIM 15AD meausurements with common systematics taken into account. NODE=M002R60;LINKAGE=A

$\Gamma(\omega e^+ e^-)/\Gamma_{total}$ Γ_6/Γ

NODE=M002R59
NODE=M002R59

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
1.97±0.34±0.17	66	[1] ABLIKIM	15AD	BES3	$J/\psi \to \eta'\gamma$

[1] Using $B(J/\psi \to \eta'\gamma) = (5.15 \pm 0.16)\times 10^{-3}$ and $B(\omega \to \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$. NODE=M002R59;LINKAGE=A

$\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/[\Gamma(\pi^+\pi^-\eta)+\Gamma(\pi^0\pi^0\eta)+\Gamma(\omega\gamma)]$ $\Gamma_2/(\Gamma_1+\Gamma_4+\Gamma_5)$

NODE=M002R18
NODE=M002R18

VALUE	DOCUMENT ID		TECN	COMMENT
0.437±0.008 OUR FIT Error includes scale factor of 1.1.				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.25 ±0.14	DAUBER	64	HBC	1.95 $K^- p$

$[\Gamma(\pi^0\pi^0\eta(\text{charged decay}))+\Gamma(\omega(\text{charged decay})\gamma)]/\Gamma_{total}$ $(0.286\Gamma_4+0.89\Gamma_5)/\Gamma$

NODE=M002R4
NODE=M002R4

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.0864±0.0017 OUR FIT Error includes scale factor of 1.1.					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.045 ±0.029	42	RITTENBERG	69	HBC	1.7–2.7 $K^- p$

$\Gamma(\pi^+\pi^-\text{neutrals})/\Gamma_{total}$ $(0.714\Gamma_1+0.286\Gamma_4+0.89\Gamma_5)/\Gamma$

NODE=M002R2
NODE=M002R2

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.3897±0.0028 OUR FIT Error includes scale factor of 1.1.					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.4 ±0.1	39	LONDON	66	HBC	2.24 $K^- p \to \Lambda\pi^+\pi^-$ neutrals
0.35 ±0.06	33	BADIER	65B	HBC	3 $K^- p$

$\Gamma(\gamma\gamma)/\Gamma_{total}$ Γ_7/Γ

NODE=M002R19
NODE=M002R19

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
2.307±0.035 OUR FIT Error includes scale factor of 1.1.					
2.31 ±0.06 OUR AVERAGE Error includes scale factor of 1.8.					
2.331±0.012±0.035	71k	ABLIKIM	19T	BES	$J/\psi \to \gamma\eta'$
1.99 $^{+0.31}_{-0.27}$ ±0.07	114	[1] WICHT	08	BELL	$B^\pm \to K^\pm\gamma\gamma$
2.00 ±0.18		[2] STANTON	80	SPEC	8.45 $\pi^- p \to n\pi^+\pi^- 2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.25 ±0.16 ±0.03	0.3k	[3] PEDLAR	09	CLEO	$J/\psi \to \gamma\eta'$
1.8 ±0.2	6000	[4] APEL	79	NICE	15–40 $\pi^- p \to n2\gamma$
2.5 ±0.7		DUANE	74	MMS	$\pi^- p \to n$MM
1.71 ±0.33	68	DALPIAZ	72	CNTR	1.6 $\pi^- p \to nX^0$
2.0 $^{+0.8}_{-0.6}$	31	HARVEY	71	OSPK	3.65 $\pi^- p \to nX^0$

[1] WICHT 08 reports $[\Gamma(\eta'(958) \to \gamma\gamma)/\Gamma_{total}] \times [B(B^+ \to \eta' K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$ which we divide by our best value $B(B^+ \to \eta' K^+) = (7.04 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M002R19;LINKAGE=WI

[2] Includes APEL 79 result. NODE=M002R19;LINKAGE=S

[3] Not independent of other η' branching fractions and ratios in PEDLAR 09. NODE=M002R19;LINKAGE=PE

[4] Data is included in STANTON 80 evaluation. NODE=M002R19;LINKAGE=A

$\Gamma(\gamma\gamma)/\Gamma(\pi^+\pi^-\eta)$ Γ_7/Γ_1

NODE=M002R46
NODE=M002R46

VALUE	DOCUMENT ID		TECN	COMMENT
0.0543±0.0012 OUR FIT Error includes scale factor of 1.1.				
0.053 ±0.004 ±0.001	PEDLAR	09	CLE3	$J/\psi \to \eta'\gamma$

$\Gamma(\gamma\gamma)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_7/Γ_2

NODE=M002R42
NODE=M002R42

VALUE	DOCUMENT ID		TECN	COMMENT
0.0783±0.0016 OUR FIT Error includes scale factor of 1.1.				
0.080 ±0.008	ABLIKIM	06E	BES2	$J/\psi \to \eta'\gamma$

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_7/Γ_4

NODE=M002R38
NODE=M002R38

VALUE	DOCUMENT ID		TECN	COMMENT
0.1031±0.0028 OUR FIT				
0.105 ±0.010 OUR AVERAGE Error includes scale factor of 1.9.				
0.091 ±0.009	AMSLER	93	CBAR	0.0 $\bar{p}p$
0.112 ±0.002 ±0.006	ALDE	87B	GAM2	38 $\pi^- p \to n2\gamma$

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta(\text{neutral decay}))$ $\Gamma_7/0.714\Gamma_4$

NODE=M002R28
NODE=M002R28

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.144±0.004 OUR FIT					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.188±0.058	16	APEL	72	OSPK	3.8 $\pi^- p \rightarrow n X^0$

$\Gamma(\text{neutrals})/\Gamma_{\text{total}}$ $(0.714\Gamma_4+0.09\Gamma_5+\Gamma_7)/\Gamma$

NODE=M002R5
NODE=M002R5

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
0.185±0.004 OUR FIT Error includes scale factor of 1.1.					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.185±0.022	535	BASILE	71	CNTR	1.6 $\pi^- p \rightarrow n X^0$
0.189±0.026	123	RITTENBERG	69	HBC	1.7–2.7 $K^- p$

$\Gamma(3\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

NODE=M002R55
NODE=M002R55

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
2.50 ±0.17 OUR FIT					
3.57 ±0.26 OUR AVERAGE					
3.522±0.082±0.254	2015	ABLIKIM	17	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
4.79 ±0.59 ±1.14	183	[1] ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.56 ±0.22 ±0.34	309	[2] ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$

[1] We have added all systematic uncertainties in quadrature to a single value.
NODE=M002R55;LINKAGE=A
[2] Superseded by ABLIKIM 17.
NODE=M002R55;LINKAGE=B

$\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_8/Γ_4

NODE=M002R32
NODE=M002R32

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
112± 8 OUR FIT					
78±10 OUR AVERAGE					
86±19	235	BLIK	08	GAMS	32 $\pi^- p \rightarrow \eta' n$
74±15		ALDE	87B	GAM2	38 $\pi^- p \rightarrow n 6\gamma$
75±18		BINON	84	GAM2	30–40 $\pi^- p \rightarrow n 6\gamma$

$\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$ Γ_9/Γ_7

NODE=M002R29
NODE=M002R29

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
4.9±1.2	33	VIKTOROV	80	CNTR	25,33 $\pi^- p \rightarrow 2\mu\gamma$

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

NODE=M002R50
NODE=M002R50

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
1.9 ±0.4 OUR AVERAGE [(2.0 ± 0.4) × 10^{-5} OUR 2023 AVERAGE]						NEW
1.94±0.37±0.02		53	[1] ABLIKIM	21I	BES3	$J/\psi \rightarrow \gamma\eta'(958)$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 2.9	90		[2] ABLIKIM	13O	BES3	$J/\psi \rightarrow \gamma\eta'$
<24	90		[3] NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$

[1] ABLIKIM 21I reports $(1.97 \pm 0.33 \pm 0.19) \times 10^{-5}$ from a measurement of $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}] \times [\text{B}(J/\psi(1S) \rightarrow \gamma\eta'(958))]$ assuming $\text{B}(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.21 \pm 0.17) \times 10^{-3}$, which we rescale to our best value $\text{B}(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.28 \pm 0.06) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
NODE=M002R50;LINKAGE=E
[2] Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.
NODE=M002R50;LINKAGE=A
[3] Not independent of measured value of Γ_{10}/Γ_1 from NAIK 09.
NODE=M002R50;LINKAGE=NA

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_{10}/Γ_1

NODE=M002R03
NODE=M002R03

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<0.5	90	[1] NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$

[1] NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)]$ / $[\text{B}(\eta \rightarrow 2\gamma)] < 1.3 \times 10^{-3}$ which we multiply by our best value $\text{B}(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.
NODE=M002R03;LINKAGE=NA

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_{10}/Γ_2

NODE=M002R57
NODE=M002R57

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<1.0	90	ABLIKIM	13O	BES3	$J/\psi \rightarrow \gamma\eta'$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{11}/Γ

NODE=M002R21
NODE=M002R21

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
3.61 ±0.18 OUR FIT					
3.61 ±0.18 OUR AVERAGE					
3.591±0.054±0.174	6067	ABLIKIM	17	BES3	$J/\psi \to \gamma(\pi^+\pi^-\pi^0)$
4.28 ±0.49 ±1.11	78	[1] ABLIKIM	15P	BES3	$J/\psi \to K^+K^-3\pi$
3.7 $^{+1.1}_{-0.9}$ ±0.4		[2] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.83 ±0.15 ±0.39	1014	[3] ABLIKIM	12E	BES3	$J/\psi \to \gamma(\pi^+\pi^-\pi^0)$

[1] We have added all systematic uncertainties in quadrature to a single value. NODE=M002R21;LINKAGE=A
[2] Not independent of measured value of Γ_{11}/Γ_1 from NAIK 09. NODE=M002R21;LINKAGE=NA
[3] Superseded by ABLIKIM 17. NODE=M002R21;LINKAGE=B

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{11}/Γ_1

NODE=M002R01
NODE=M002R01

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
8.5 ±0.4 OUR FIT Error includes scale factor of 1.1.					
8.27 $^{+2.49}_{-2.12}$ ±0.04	20	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] NAIK 09 reports $[\Gamma(\eta'(958) \to \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \to \pi^+\pi^-\eta)]$ / $[B(\eta \to 2\gamma)] = (21^{+6}_{-5} \pm 2)\times10^{-3}$ which we multiply by our best value $B(\eta \to 2\gamma) = (39.36 \pm 0.18)\times10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M002R01;LINKAGE=NA

$\Gamma((\pi^+\pi^-\pi^0)$ S-wave$)/\Gamma_{total}$ Γ_{12}/Γ

NODE=M002R63
NODE=M002R63

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
37.63±0.77±5.00	6580	[1] ABLIKIM	17	BES3	$J/\psi \to \gamma(\pi^+\pi^-\pi^0)$

[1] We have added all systematic uncertainties in quadrature . NODE=M002R63;LINKAGE=A

$\Gamma(\pi^\mp\rho^\pm)/\Gamma_{total}$ Γ_{13}/Γ

NODE=M002R62
NODE=M002R62

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
7.44±0.60±2.23	1231	[1] ABLIKIM	17	BES3	$J/\psi \to \gamma(\pi^\mp\rho^\pm)$

[1] We have added all systematic uncertainties in quadrature . NODE=M002R62;LINKAGE=A

$\Gamma(2(\pi^+\pi^-))/\Gamma_{total}$ Γ_{14}/Γ

NODE=M002R24
NODE=M002R24

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
8.3±0.9 OUR AVERAGE [(8.4 ± 0.9) × 10^{-5} OUR 2023 AVERAGE]						NEW
8.3±0.9±0.1		199	[1] ABLIKIM	14M	BES3	$J/\psi \to \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 24	90		[2] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$
<1000	90		RITTENBERG	69	HBC	1.7–2.7 K^-p

[1] ABLIKIM 14M reports $[\Gamma(\eta'(958) \to 2(\pi^+\pi^-))/\Gamma_{total}] \times [B(J/\psi(1S) \to \gamma\eta'(958))]$ = (4.40 ± 0.35 ± 0.30) × 10^{-7} which we divide by our best value $B(J/\psi(1S) \to \gamma\eta'(958)) = (5.28 \pm 0.06)\times10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M002R24;LINKAGE=A
[2] Not independent of measured value of Γ_{14}/Γ_1 from NAIK 09. NODE=M002R24;LINKAGE=NA

$\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{14}/Γ_1

NODE=M002R04
NODE=M002R04

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<0.6	90	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] NAIK 09 reports $[\Gamma(\eta'(958) \to 2(\pi^+\pi^-))/\Gamma(\eta'(958) \to \pi^+\pi^-\eta)]$ / $[B(\eta \to 2\gamma)]$ < 1.4×10^{-3} which we multiply by our best value $B(\eta \to 2\gamma) = 39.36\times10^{-2}$. NODE=M002R04;LINKAGE=NA

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{total}$ Γ_{15}/Γ

NODE=M002R51
NODE=M002R51

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
1.78±0.38±0.02	84		[1] ABLIKIM	14M	BES3	$J/\psi \to \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<27	90		[2] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] ABLIKIM 14M reports $[\Gamma(\eta'(958) \to \pi^+\pi^-2\pi^0)/\Gamma_{total}] \times [B(J/\psi(1S) \to \gamma\eta'(958))]$ = (9.38±1.79±0.89)×10^{-7} which we divide by our best value $B(J/\psi(1S) \to \gamma\eta'(958))$ = (5.28 ± 0.06) × 10^{-3}. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M002R51;LINKAGE=A
[2] Not independent of measured value of Γ_{15}/Γ_1 from NAIK 09. NODE=M002R51;LINKAGE=NA

$\Gamma(\pi^+\pi^- 2\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{15}/Γ_1

NODE=M002R05
NODE=M002R05

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<6	90	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] NAIK 09 reports $[\Gamma(\eta'(958) \to \pi^+\pi^- 2\pi^0)/\Gamma(\eta'(958) \to \pi^+\pi^-\eta)]$ / $[B(\eta \to 2\gamma)]$ $< 15 \times 10^{-3}$ which we multiply by our best value $B(\eta \to 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R05;LINKAGE=NA

$\Gamma(2(\pi^+\pi^-)\text{ neutrals})/\Gamma_{\text{total}}$ Γ_{16}/Γ

NODE=M002R22
NODE=M002R22

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.01	95	DANBURG	73	HBC	2.2 $K^- p \to \Lambda X^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.01	90	RITTENBERG	69	HBC	1.7–2.7 $K^- p$

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

NODE=M002R23
NODE=M002R23

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.002	90	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$
<0.01	90	RITTENBERG	69	HBC	1.7–2.7 $K^- p$

[1] Not independent of measured value of Γ_{17}/Γ_1 from NAIK 09.

NODE=M002R23;LINKAGE=NA

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{17}/Γ_1

NODE=M002R06
NODE=M002R06

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<4	90	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] NAIK 09 reports $[\Gamma(\eta'(958) \to 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \to \pi^+\pi^-\eta)]$ / $[B(\eta \to 2\gamma)]$ $< 11 \times 10^{-3}$ which we multiply by our best value $B(\eta \to 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R06;LINKAGE=NA

$\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

NODE=M002R16
NODE=M002R16

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.01	95	KALBFLEISCH	64B	HBC	$K^- p \to \Lambda 2(\pi^+\pi^-)$+MM
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.01	90	LONDON	66	HBC	Compilation

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{19}/Γ

NODE=M002R07
NODE=M002R07

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
< 3.1	90	[1] ABLIKIM	13U	BES3	$J/\psi \to \gamma 3(\pi^+\pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 53	90	[2] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$
<500	95	KALBFLEISCH	64B	HBC	$K^- p \to \Lambda 2(\pi^+\pi^-)$

[1] Using $B(J/\psi \to \gamma\eta'(958)) = (5.16 \pm 0.15) \times 10^{-3}$.

NODE=M002R07;LINKAGE=A

[2] Not independent of measured value of Γ_{19}/Γ_1 from NAIK 09.

NODE=M002R07;LINKAGE=NA

$\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{19}/Γ_1

NODE=M002R08
NODE=M002R08

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
<1.2	90	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] NAIK 09 reports $[\Gamma(\eta'(958) \to 3(\pi^+\pi^-))/\Gamma(\eta'(958) \to \pi^+\pi^-\eta)]$ / $[B(\eta \to 2\gamma)]$ $< 3.0 \times 10^{-3}$ which we multiply by our best value $B(\eta \to 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R08;LINKAGE=NA

$\Gamma(K^\pm\pi^\mp)/\Gamma(\rho^0\gamma(\text{including non-resonant }\pi^+\pi^-\gamma))$ Γ_{20}/Γ_2

NODE=M002R61
NODE=M002R61

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.3 \times 10^{-4}$	90	ABLIKIM	16M	BES3	$e^+e^- \to J/\psi \to$ hadrons

$\Gamma(\pi^+\pi^- e^+ e^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

NODE=M002R12
NODE=M002R12

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID		TECN	COMMENT
2.42±0.10 OUR FIT						
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2.11±0.12±0.14		429	[1] ABLIKIM	13O	BES3	$J/\psi \to \gamma\eta'$
$2.5\ ^{+1.2}_{-0.9}\ \pm 0.5$			[2] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$
<6	90		RITTENBERG	65	HBC	2.7 $K^- p$

[1] Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.

NODE=M002R12;LINKAGE=A

[2] Not independent of measured value of Γ_{21}/Γ_1 from NAIK 09.

NODE=M002R12;LINKAGE=NA

$\Gamma(\pi^+\pi^- e^+ e^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_{21}/Γ_1

NODE=M002R02
NODE=M002R02

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
5.69±0.25 OUR FIT					
$5.51^{+3.00}_{-2.30}\pm0.03$	8	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] NAIK 09 reports $[\Gamma(\eta'(958) \to \pi^+\pi^- e^+ e^-)/\Gamma(\eta'(958) \to \pi^+\pi^-\eta)]$ / $[B(\eta \to 2\gamma)] = (14^{+7}_{-5} \pm 3) \times 10^{-3}$ which we multiply by our best value $B(\eta \to 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R02;LINKAGE=NA

$\Gamma(\pi^+\pi^- e^+ e^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_{21}/Γ_2

NODE=M002R56
NODE=M002R56

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
8.20±0.31 OUR FIT					
8.20±0.16±0.27	2584	ABLIKIM	21J	BES3	$J/\psi \to \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.2 ±0.4 ±0.5	429	[1] ABLIKIM	13O	BES3	$J/\psi \to \gamma\eta'$

[1] Superseded by ABLIKIM 21J.

NODE=M002R56;LINKAGE=A

$\Gamma(\pi^+ e^-\nu_e + \text{c.c.})/\Gamma(\pi^+\pi^-\eta)$ Γ_{22}/Γ_1

NODE=M002R54
NODE=M002R54

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<5.0	90	ABLIKIM	13G	BES3	$J/\psi \to \phi\eta'$

$\Gamma(\gamma e^+ e^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

NODE=M002R40
NODE=M002R40

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.9	90	BRIERE	00	CLEO	10.6 e^+e^-

$\Gamma(\gamma e^+ e^-)/\Gamma(\gamma\gamma)$ Γ_{23}/Γ_7

NODE=M002R00
NODE=M002R00

VALUE (units 10^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
2.13±0.09±0.07	864	ABLIKIM	15O	BES3	$J/\psi \to \gamma e^+ e^-$

$\Gamma(\pi^0\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

NODE=M002R64
NODE=M002R64

VALUE (units 10^{-3})	EVTS	DOCUMENT ID		TECN	COMMENT
3.20±0.07±0.23	3.4k	ABLIKIM	17T	BES3	$J/\psi \to \gamma\eta'$

$\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_{24}/Γ_4

NODE=M002R35
NODE=M002R35

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<37	90	ALDE	87B	GAM2	38 $\pi^- p \to n4\gamma$

$\Gamma(\pi^0\gamma\gamma(\text{non resonant}))/\Gamma_{\text{total}}$ Γ_{25}/Γ

NODE=M002R65
NODE=M002R65

VALUE (units 10^{-4})	EVTS	DOCUMENT ID		TECN	COMMENT
6.16±0.64±0.67	655	ABLIKIM	17T	BES3	$J/\psi \to \gamma\eta'$

$\Gamma(\eta\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

NODE=M002R67
NODE=M002R67

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<1.33 \times 10^{-4}$	90	ABLIKIM	19AW	BES3	$J/\psi \to \gamma\eta' \to \gamma\gamma\gamma 2\gamma$

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

NODE=M002R58
NODE=M002R58

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
$<4.94 \times 10^{-5}$	90	ABLIKIM	20E	BES3	$J/\psi \to \eta'\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<3.2 \times 10^{-4}$	90	DONSKOV	14	GAM4	32.5 $\pi^- p \to \eta' n$

$\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_{27}/Γ_4

NODE=M002R37
NODE=M002R37

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<23	90	ALDE	87B	GAM2	38 $\pi^- p \to n8\gamma$

$\Gamma(e^+e^-)/\Gamma_{total}$ Γ_{28}/Γ

NODE=M002R39
NODE=M002R39

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
< 5.6 × 10⁻⁹	90	[1] ACHASOV	15	SND	0.958 $e^+e^- \to \pi\pi\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<12 × 10⁻⁹	90	[2] AKHMETSHIN	15	CMD3	0.958 $e^+e^- \to \pi^+\pi^-\eta$
< 2.1 × 10⁻⁷	90	VOROBYEV	88	ND	$e^+e^- \to \pi^+\pi^-\eta$

[1] Combining data of ACHASOV 15 and AKHMETSHIN 15 and using $\Gamma(\eta') = 0.198 \pm 0.009$ MeV.
NODE=M002R39;LINKAGE=B

[2] Using $\Gamma_{\eta'(958)} = 198 \pm 9$ keV, B($\eta'(958) \to \pi^+\pi^-\eta$) = (42.9 ± 0.7)%, and B($\eta \to \gamma\gamma$) = (39.41 ± 0.20)%.
NODE=M002R39;LINKAGE=A

$\Gamma(e^+e^-e^+e^-)/\Gamma_{total}$ Γ_{29}/Γ

NODE=M002R68
NODE=M002R68

VALUE (units 10⁻⁶)	EVTS	DOCUMENT ID		TECN	COMMENT
4.5±1.1±0.1	30	[1] ABLIKIM	22E	BES3	$J/\psi \to \gamma\eta'$

[1] ABLIKIM 22E reports (4.5 ± 1.0 ± 0.5) × 10⁻⁶ from a measurement of [$\Gamma(\eta'(958) \to e^+e^-e^+e^-)/\Gamma_{total}$] × [B($J/\psi(1S) \to \gamma\eta'(958)$)] assuming B($J/\psi(1S) \to \gamma\eta'(958)$) = (5.25 ± 0.07) × 10⁻³, which we rescale to our best value B($J/\psi(1S) \to \gamma\eta'(958)$) = (5.28 ± 0.06) × 10⁻³. Our first error is their experiment's error and our second error is the systematic error from using our best value.
NODE=M002R68;LINKAGE=B

$\Gamma(\text{invisible})/\Gamma_{total}$ Γ_{30}/Γ

NODE=M002R52
NODE=M002R52

VALUE (units 10⁻⁴)	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<9.5	90	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] Not independent of measured value of Γ_{30}/Γ_1 from NAIK 09.
NODE=M002R52;LINKAGE=NA

$\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\eta)$ Γ_{30}/Γ_1

NODE=M002R09
NODE=M002R09

VALUE (units 10⁻³)	CL%	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.1	90	[1] NAIK	09	CLEO	$J/\psi \to \gamma\eta'$

[1] NAIK 09 reports [$\Gamma(\eta'(958) \to$ invisible)/$\Gamma(\eta'(958) \to \pi^+\pi^-\eta)$] / [B($\eta \to 2\gamma$)] < 5.4 × 10⁻³ which we multiply by our best value B($\eta \to 2\gamma$) = 39.36 × 10⁻².
NODE=M002R09;LINKAGE=NA

$\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$ Γ_{30}/Γ_7

NODE=M002R44
NODE=M002R44

VALUE (units 10⁻²)	CL%	DOCUMENT ID		TECN	COMMENT
<2.4	90	ABLIKIM	13	BES3	$J/\psi \to \phi\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.69	90	ABLIKIM	06Q	BES	$J/\psi \to \phi\eta'$

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ Γ_{31}/Γ

NODE=M002R20
NODE=M002R20

VALUE (units 10⁻⁴)	CL%	DOCUMENT ID		TECN	COMMENT
< 0.18	90	[1] AAIJ	17D	LHCB	$D^+_{(s)} \to \pi^+\pi^-\pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.5	90	[2] ABLIKIM	11G	BES3	$J/\psi \to \gamma\pi^+\pi^-$
< 29	90	[3] MORI	07A	BELL	$\gamma\gamma \to \pi^+\pi^-$
< 3.3	90	[4] MORI	07A	BELL	$\gamma\gamma \to \pi^+\pi^-$
<800	95	DANBURG	73	HBC	2.2 $K^-p \to \Lambda X^0$
<200	90	RITTENBERG	69	HBC	1.7–2.7 K^-p

OCCUR=2

[1] Using branching fractions of $D^+_{(s)}$ decays from PDG 15.
NODE=M002R20;LINKAGE=A

[2] ABLIKIM 11G reports [$\Gamma(\eta'(958) \to \pi^+\pi^-)/\Gamma_{total}$] × [B($J/\psi(1S) \to \gamma\eta'(958)$)] < 2.84 × 10⁻⁷ which we divide by our best value B($J/\psi(1S) \to \gamma\eta'(958)$) = 5.28 × 10⁻³.
NODE=M002R20;LINKAGE=AL

[3] Taking into account interference with the $\gamma\gamma \to \pi^+\pi^-$ continuum.
NODE=M002R20;LINKAGE=MO

[4] Without interference with the $\gamma\gamma \to \pi^+\pi^-$ continuum.
NODE=M002R20;LINKAGE=MR

$\Gamma(\pi^0\pi^0)/\Gamma_{total}$ Γ_{32}/Γ

NODE=M002R53
NODE=M002R53

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<4 × 10⁻⁴	90	[1] ABLIKIM	11G	BES3	$J/\psi \to \gamma\pi^0\pi^0$

[1] ABLIKIM 11G reports [$\Gamma(\eta'(958) \to \pi^+\pi^-)/\Gamma_{total}$] × [B($J/\psi(1S) \to \gamma\eta'(958)$)] < 2.84 × 10⁻⁷ which we divide by our best value B($J/\psi(1S) \to \gamma\eta'(958)$) = 5.28 × 10⁻³.
NODE=M002R53;LINKAGE=AL

$\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_{32}/Γ_4

NODE=M002R36
NODE=M002R36

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<45	90	ALDE	87B	GAM2	38 $\pi^- p \rightarrow n4\gamma$

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{total}$ Γ_{33}/Γ

NODE=M002R8
NODE=M002R8

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
< 1.4	90	BRIERE	00	CLEO	10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<13	90	RITTENBERG	65	HBC	2.7 $K^- p$

$\Gamma(\pi^0\rho^0)/\Gamma_{total}$ Γ_{34}/Γ

NODE=M002R10
NODE=M002R10

VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.04	90	RITTENBERG	65	HBC	2.7 $K^- p$

$\Gamma(\eta e^+ e^-)/\Gamma_{total}$ Γ_{35}/Γ

NODE=M002R9
NODE=M002R9

VALUE (units 10^{-3})	CL%	DOCUMENT ID		TECN	COMMENT
< 2.4	90	BRIERE	00	CLEO	10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<11	90	RITTENBERG	65	HBC	2.7 $K^- p$

$\Gamma(3\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_{36}/Γ_4

NODE=M002R34
NODE=M002R34

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<4.6	90	ALDE	87B	GAM2	38 $\pi^- p \rightarrow n3\gamma$

$\Gamma(\mu^+\mu^-\pi^0)/\Gamma_{total}$ Γ_{37}/Γ

NODE=M002R31
NODE=M002R31

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<6.0	90	DZHELYADIN	81	CNTR	30 $\pi^- p \rightarrow \eta' n$

$\Gamma(\mu^+\mu^-\eta)/\Gamma_{total}$ Γ_{38}/Γ

NODE=M002R30
NODE=M002R30

VALUE (units 10^{-5})	CL%	DOCUMENT ID		TECN	COMMENT
<1.5	90	DZHELYADIN	81	CNTR	30 $\pi^- p \rightarrow \eta' n$

$\Gamma(e\mu)/\Gamma_{total}$ Γ_{39}/Γ

NODE=M002R41
NODE=M002R41

VALUE (units 10^{-4})	CL%	DOCUMENT ID		TECN	COMMENT
<4.7	90	BRIERE	00	CLEO	10.6 $e^+ e^-$

$\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

NODE=M002225

$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha Y|^2 + CX + DX^2$

NODE=M002225

X and Y are Dalitz variables; α is complex and C, and D are real-valued. Parameters C and D are not necessarily equal to c and d, respectively, in the generalized parameterization following this one. May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays. Because of different initial assumptions and strong correlations of the parameters we do not average the parameters in the section below.

$Re(\alpha)$ decay parameter

NODE=M002A0
NODE=M002A0

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$-0.034 \pm 0.002 \pm 0.002$	351k	ABLIKIM	18	BES3	$\eta' \rightarrow \eta\pi^+\pi^-$
$-0.054 \pm 0.004 \pm 0.001$	56k	ABLIKIM	18	BES3	$\eta' \rightarrow \eta\pi^0\pi^0$
$-0.033 \pm 0.005 \pm 0.003$	44k	[1] ABLIKIM	11	BES3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$-0.072 \pm 0.012 \pm 0.006$	7k	[2] AMELIN	05A	VES	28 $\pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
$-0.021 \pm 0.018 \pm 0.017$	6.7k	[3] BRIERE	00	CLEO	10.6 $e^+ e^- \rightarrow \eta\pi^+\pi^- X$
$-0.058 \pm 0.013 \pm 0.003$	5.4k	[4] ALDE	86	GAM2	38 $\pi^- p \rightarrow n\eta\pi^0\pi^0$
-0.08 ± 0.03		[4,5] KALBFLEISCH	74	RVUE	$\eta' \rightarrow \eta\pi^+\pi^-$

OCCUR=2

[1] See ABLIKIM 11 for the full correlation matrix. NODE=M002A0;LINKAGE=AB
[2] Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below. NODE=M002A0;LINKAGE=AM
[3] Assuming $\text{Im}(\alpha) = 0$, $C = 0$, and $D = 0$. NODE=M002A0;LINKAGE=BR
[4] Assuming $C = 0$. NODE=M002A0;LINKAGE=A
[5] From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73. NODE=M002A0;LINKAGE=KA

$Im(\alpha)$ decay parameter

NODE=M002IA0
NODE=M002IA0

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.000 \pm 0.019 \pm 0.001$	351k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^+\pi^-$
$0.000 \pm 0.038 \pm 0.002$	56k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^0\pi^0$
$0.000 \pm 0.049 \pm 0.001$	44k	[1] ABLIKIM	11	BES3	$J/\psi \to \gamma\eta\pi^+\pi^-$
$0.0 \quad \pm 0.1 \quad \pm 0.0$	7k	[2] AMELIN	05A	VES	28 $\pi^- A \to \eta\pi^+\pi^-\pi^- A^*$
$-0.00 \pm 0.13 \pm 0.00$	5.4k	[3] ALDE	86	GAM2	38 $\pi^- p \to n\eta\pi^0\pi^0$
$0.0 \quad \pm 0.3$		[3,4] KALBFLEISCH	74	RVUE	$\eta' \to \eta\pi^+\pi^-$

[1] See ABLIKIM 11 for the full correlation matrix. NODE=M002IA0;LINKAGE=AB
[2] Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below. NODE=M002IA0;LINKAGE=AM
[3] Assuming $C = 0$. NODE=M002IA0;LINKAGE=A
[4] From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73. NODE=M002IA0;LINKAGE=KA

C decay parameter

NODE=M002C0
NODE=M002C0

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.0027 \pm 0.0024 \pm 0.0015$	351k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^+\pi^-$
$0.018 \pm 0.009 \pm 0.003$	44k	[1] ABLIKIM	11	BES3	$J/\psi \to \gamma\eta\pi^+\pi^-$
$0.020 \pm 0.018 \pm 0.004$	7k	[2] AMELIN	05A	VES	28 $\pi^- A \to \eta\pi^+\pi^-\pi^- A^*$

[1] See ABLIKIM 11 for the full correlation matrix. NODE=M002C0;LINKAGE=AB
[2] Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below. NODE=M002C0;LINKAGE=AM

D decay parameter

NODE=M002D0
NODE=M002D0

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$-0.053 \pm 0.004 \pm 0.004$	351k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^+\pi^-$
$-0.061 \pm 0.009 \pm 0.005$	56k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^0\pi^0$
$-0.059 \pm 0.012 \pm 0.004$	44k	[1] ABLIKIM	11	BES3	$J/\psi \to \gamma\eta\pi^+\pi^-$
$-0.066 \pm 0.030 \pm 0.015$	7k	[2] AMELIN	05A	VES	28 $\pi^- A \to \eta\pi^+\pi^-\pi^- A^*$
$0.00 \pm 0.03 \pm 0.00$	5.4k	[3] ALDE	86	GAM2	38 $\pi^- p \to n\eta\pi^0\pi^0$
0		[3,4] KALBFLEISCH	74	RVUE	$\eta' \to \eta\pi^+\pi^-$

[1] See ABLIKIM 11 for the full correlation matrix. NODE=M002D0;LINKAGE=AB
[2] Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below. NODE=M002D0;LINKAGE=AM
[3] Assuming $C = 0$. NODE=M002D0;LINKAGE=AL
[4] From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73. NODE=M002D0;LINKAGE=KA

$\eta'(958) \to \eta\pi\pi$ DECAY PARAMETERS

NODE=M002227

$|\text{MATRIX ELEMENT}|^2 \propto 1 + a\,Y + b\,Y^2 + c\,X + d\,X^2$

X and Y are Dalitz variables and a, b, c, and d are real-valued parameters. May be different for $\eta'(958) \to \eta\pi^+\pi^-$ and $\eta'(958) \to \eta\pi^0\pi^0$ decays. We do not average measurements in the section below because parameter values from each experiment are strongly correlated. NODE=M002227

a decay parameter

NODE=M002DPA
NODE=M002DPA

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$-0.077 \pm 0.003 \pm 0.001$		[1] ABLIKIM	23AH	BES3	$\eta' \to \eta\pi^0\pi^0$
$-0.056 \pm 0.004 \pm 0.002$	351k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^+\pi^-$
$-0.087 \pm 0.009 \pm 0.006$	56k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^0\pi^0$
$-0.074 \pm 0.008 \pm 0.006$	124k	ADLARSON	18A	A2MM	$\eta' \to \eta\pi^0\pi^0$
$-0.072 \pm 0.007 \pm 0.008$		[2] GONZALEZ-S..	18A	RVUE	$\eta' \to \eta\pi^0\pi^0$
$-0.047 \pm 0.011 \pm 0.003$	44k	[3] ABLIKIM	11	BES3	$J/\psi \to \gamma\eta\pi^+\pi^-$
$-0.066 \pm 0.016 \pm 0.003$	15k	[4] BLIK	09	GAM4	32.5 $\pi^- p \to \eta' n$
$-0.127 \pm 0.016 \pm 0.008$	20k	[5] DOROFEEV	07	VES	27 $\pi^- p \to \eta' n$, $\pi^- A \to \eta'\pi^- A^*$

[1] Fit IV, ignoring noncusp terms. Supersedes ABLIKIM 18. NODE=M002DPA;LINKAGE=B
[2] Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop. NODE=M002DPA;LINKAGE=A
[3] See ABLIKIM 11 for the full correlation matrix. NODE=M002DPA;LINKAGE=AB
[4] From $\eta' \to \eta\pi^0\pi^0$ decay. NODE=M002DPA;LINKAGE=BL
[5] From $\eta' \to \eta\pi^+\pi^-$ decay. NODE=M002DPA;LINKAGE=DO

b decay parameter

NODE=M002DPB
NODE=M002DPB

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$-0.066 \pm 0.006 \pm 0.001$		[1] ABLIKIM	23AH	BES3	$\eta' \to \eta\pi^0\pi^0$
$-0.049 \pm 0.006 \pm 0.006$	351k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^+\pi^-$
$-0.073 \pm 0.014 \pm 0.005$	56k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^0\pi^0$ OCCUR=2
$-0.063 \pm 0.014 \pm 0.005$	124k	ADLARSON	18A	A2MM	$\eta' \to \eta\pi^0\pi^0$
$-0.052 \pm 0.001 \pm 0.002$		[2] GONZALEZ-S..	18A	RVUE	$\eta' \to \eta\pi^0\pi^0$
$-0.069 \pm 0.019 \pm 0.009$	44k	[3] ABLIKIM	11	BES3	$J/\psi \to \gamma\eta\pi^+\pi^-$
$-0.063 \pm 0.028 \pm 0.004$	15k	[4] BLIK	09	GAM4	32.5 $\pi^- p \to \eta' n$
$-0.106 \pm 0.028 \pm 0.014$	20k	[5] DOROFEEV	07	VES	27 $\pi^- p \to \eta' n$, $\pi^- A \to \eta'\pi^- A^*$

[1] Fit IV, ignoring noncusp terms. Supersedes ABLIKIM 18. NODE=M002DPB;LINKAGE=B
[2] Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop. NODE=M002DPB;LINKAGE=A
[3] See ABLIKIM 11 for the full correlation matrix. NODE=M002DPB;LINKAGE=AB
[4] From $\eta' \to \eta\pi^0\pi^0$ decay. NODE=M002DPB;LINKAGE=BL
[5] From $\eta' \to \eta\pi^+\pi^-$ decay. NODE=M002DPB;LINKAGE=DO

c decay parameter

NODE=M002DPC
NODE=M002DPC

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.0027 \pm 0.0024 \pm 0.0018$	351k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^+\pi^-$
$0.019 \pm 0.011 \pm 0.003$	44k	[1] ABLIKIM	11	BES3	$J/\psi \to \gamma\eta\pi^+\pi^-$
$-0.107 \pm 0.096 \pm 0.003$	15k	[2] BLIK	09	GAM4	32.5 $\pi^- p \to \eta' n$
$0.015 \pm 0.011 \pm 0.014$	20k	[3] DOROFEEV	07	VES	27 $\pi^- p \to \eta' n$, $\pi^- A \to \eta'\pi^- A^*$

[1] See ABLIKIM 11 for the full correlation matrix. NODE=M002DPC;LINKAGE=AB
[2] From $\eta' \to \eta\pi^0\pi^0$ decay. NODE=M002DPC;LINKAGE=BL
[3] From $\eta' \to \eta\pi^+\pi^-$ decay. NODE=M002DPC;LINKAGE=DO

d decay parameter

NODE=M002DPD
NODE=M002DPD

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$-0.068 \pm 0.004 \pm 0.001$		[1] ABLIKIM	23AH	BES3	$\eta' \to \eta\pi^0\pi^0$
$-0.063 \pm 0.004 \pm 0.003$	351k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^+\pi^-$
$-0.074 \pm 0.009 \pm 0.004$	56k	ABLIKIM	18	BES3	$\eta' \to \eta\pi^0\pi^0$ OCCUR=2
$-0.050 \pm 0.009 \pm 0.005$	124k	ADLARSON	18A	A2MM	$\eta' \to \eta\pi^0\pi^0$
$-0.051 \pm 0.008 \pm 0.006$		[2] GONZALEZ-S..	18A	RVUE	$\eta' \to \eta\pi^0\pi^0$
$-0.073 \pm 0.012 \pm 0.003$	44k	[3] ABLIKIM	11	BES3	$J/\psi \to \gamma\eta\pi^+\pi^-$
$0.018 \pm 0.078 \pm 0.006$	15k	[4] BLIK	09	GAM4	32.5 $\pi^- p \to \eta' n$
$-0.082 \pm 0.017 \pm 0.008$	20k	[5] DOROFEEV	07	VES	27 $\pi^- p \to \eta' n$, $\pi^- A \to \eta'\pi^- A^*$

[1] Fit IV, ignoring noncusp terms. Supersedes ABLIKIM 18. NODE=M002DPD;LINKAGE=B
[2] Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop. NODE=M002DPD;LINKAGE=A
[3] See ABLIKIM 11 for the full correlation matrix. NODE=M002DPD;LINKAGE=AB
[4] From $\eta' \to \eta\pi^0\pi^0$ decay. If $c \equiv 0$ from Bose-Einstein symmetry, $d = -0.067 \pm 0.020 \pm 0.003$. NODE=M002DPD;LINKAGE=BL
[5] From $\eta' \to \eta\pi^+\pi^-$ decay. NODE=M002DPD;LINKAGE=DO

$\eta'(958)$ β PARAMETER
$|\text{MATRIX ELEMENT}|^2 = (1 + 2\beta Z)$

NODE=M002226

See the "Note on η Decay Parameters" in our 1994 edition Physical Review **D50** 1173 (1994), p. 1454. NODE=M002226

β decay parameter

NODE=M002B0
NODE=M002B0

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
-0.61 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.2.			
$-0.640 \pm 0.046 \pm 0.047$	1.8k	ABLIKIM	15G	BES3	$J/\psi \to \gamma(\pi^0\pi^0\pi^0)$
-0.59 ± 0.18	235	BLIK	08	GAMS	32 $\pi^- p \to \eta' n$
-0.1 ± 0.3		ALDE	87B	GAM2	38 $\pi^- p \to n3\pi^0$

$\eta'(958)$ C-NONCONSERVING DECAY PARAMETER

NODE=M002235

See the note on η decay parameters in the Stable Particle Particle Listings for definition of this parameter.

NODE=M002235

DECAY ASYMMETRY PARAMETER FOR $\pi^+\pi^-\gamma$

NODE=M002A

VALUE	EVTS	DOCUMENT ID		TECN	COMMENT
-0.03 ±0.04 OUR AVERAGE					
-0.019 ± 0.056		AIHARA	87	TPC	$2\gamma \to \pi^+\pi^-\gamma$
-0.069 ± 0.078	295	GRIGORIAN	75	STRC	2.1 $\pi^- p$
0.00 ±0.10	103	KALBFLEISCH	75	HBC	2.18 $K^- p \to \Lambda\pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.07 ±0.08	152	RITTENBERG	65	HBC	2.1–2.7 $K^- p$

NODE=M002A

$\eta'(958) \to \gamma\ell^+\ell^-$ TRANSITION FORM FACTOR SLOPE

NODE=M002FFL

Related to the effective virtual meson mass Λ, via slope $\approx \Lambda^{-2}$. See e.g. LANDSBERG 85, eq. (3.8), for a detailed definition.

NODE=M002FFL

VALUE (GeV^{-2})	EVTS	DOCUMENT ID		TECN	COMMENT
1.62±0.17 OUR AVERAGE					
1.60±0.17±0.08	864	[1] ABLIKIM	15O	BES3	$J/\psi \to \gamma e^+ e^-$
1.7 ±0.4	33	[1] VIKTOROV	80		25,33 $\pi^- p \to 2\mu\gamma$

NODE=M002FFL

[1] In the single-pole Ansatz where slope $= 1/(\Lambda^2 + \gamma^2)$ with Λ, γ being a Breit-Wigner mass, width for the effective contributing vector meson.

NODE=M002FFL;LINKAGE=A

$\eta'(958)$ REFERENCES

NODE=M002

ABLIKIM	23AH	PRL 130 081901	M. Ablikim *et al.*	(BESIII Collab.)	REFID=62206
ABLIKIM	22E	PR D105 112010	M. Ablikim *et al.*	(BESIII Collab.)	REFID=61639
ABLIKIM	21I	PR D103 072006	M. Ablikim *et al.*	(BESIII Collab.)	REFID=61112
ABLIKIM	21J	PR D103 092005	M. Ablikim *et al.*	(BESIII Collab.)	REFID=61114
ABLIKIM	20E	PR D101 032001	M. Ablikim *et al.*	(BESIII Collab.)	REFID=60252
ABLIKIM	19AW	PR D100 052015	M. Ablikim *et al.*	(BESIII Collab.)	REFID=60000
ABLIKIM	19T	PRL 122 142002	M. Ablikim *et al.*	(BESIII Collab.)	REFID=59773
ABLIKIM	18	PR D97 012003	M. Ablikim *et al.*	(BESIII Collab.)	REFID=58711
ABLIKIM	18C	PRL 120 242003	M. Ablikim *et al.*	(BESIII Collab.)	REFID=58843
ADLARSON	18A	PR D98 012001	P. Adlarson *et al.*	(A2 Collab. at MAMI)	REFID=58957
GONZALEZ-S...	18A	EPJ C78 758	S. Gonzalez-Solis, E. Passemar	(BEIJ, IND+)	REFID=59316
AAIJ	17D	PL B764 233	R. Aaij *et al.*	(LHCb Collab.)	REFID=57702
ABLIKIM	17	PRL 118 012001	M. Ablikim *et al.*	(BESIII Collab.)	REFID=57650
ABLIKIM	17T	PR D96 012005	M. Ablikim *et al.*	(BESIII Collab.)	REFID=58013
ABLIKIM	16M	PR D93 072008	M. Ablikim *et al.*	(BESIII Collab.)	REFID=57511
ABLIKIM	15AD	PR D92 051101	M. Ablikim *et al.*	(BESIII Collab.)	REFID=56983
ABLIKIM	15G	PR D92 012014	M. Ablikim *et al.*	(BESIII Collab.)	REFID=56602
ABLIKIM	15O	PR D92 012001	M. Ablikim *et al.*	(BESIII Collab.)	REFID=56780
ABLIKIM	15P	PR D92 012007	M. Ablikim *et al.*	(BESIII Collab.)	REFID=56781
ACHASOV	15	PR D91 092010	M.N. Achasov *et al.*	(SND Collab.)	REFID=56788
AKHMETSHIN	15	PL B740 273	R.R. Akhmetshin *et al.*	(CMD-3 Collab.)	REFID=56386
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=1
ABLIKIM	14M	PRL 112 251801	M. Ablikim *et al.*	(BESIII Collab.)	REFID=55904
DONSKOV	14	MPL A29 1450213	S. Donskov *et al.*	(GAMS-4π Collab.)	REFID=56321
PDG	14	CP C38 070001	K. Olive *et al.*	(PDG Collab.)	REFID=55687
ABLIKIM	13	PR D87 012009	M. Ablikim *et al.*	(BESIII Collab.)	REFID=54829
ABLIKIM	13G	PR D87 032006	M. Ablikim *et al.*	(BESIII Collab.)	REFID=54952
ABLIKIM	13O	PR D87 092011	M. Ablikim *et al.*	(BESIII Collab.)	REFID=55388
ABLIKIM	13U	PR D88 091502	M. Ablikim *et al.*	(BESIII Collab.)	REFID=55582
ABLIKIM	12E	PRL 108 182001	M. Ablikim *et al.*	(BESIII Collab.)	REFID=54270
PDG	12	PR D86 010001	J. Beringer *et al.*	(PDG Collab.)	REFID=54066
ABLIKIM	11	PR D83 012003	M. Ablikim *et al.*	(BESIII Collab.)	REFID=53646
ABLIKIM	11G	PR D84 032006	M. Ablikim *et al.*	(BESIII Collab.)	REFID=53711
CZERWINSKI	10	PRL 105 122001	E. Czerwinski *et al.*	(COSY-11 Collab.)	REFID=53364
BLIK	09	PAN 72 231 Translated from YAF 72 258.	A.M. Blik *et al.*	(IHEP (Protvino))	REFID=52727
NAIK	09	PRL 102 061801	P. Naik *et al.*	(CLEO Collab.)	REFID=52678
PEDLAR	09	PR D79 111101	T.K. Pedlar *et al.*	(CLEO Collab.)	REFID=52998
BLIK	08	PAN 71 2124 Translated from YAF 71 2161.	A. Blik *et al.*	(GAMS-4π Collab.)	REFID=52663
LIBBY	08	PRL 101 182002	J. Libby *et al.*	(CLEO Collab.)	REFID=52591
WICHT	08	PL B662 323	J. Wicht *et al.*	(BELLE Collab.)	REFID=52204
DOROFEEV	07	PL B651 22	V. Dorofeev *et al.*	(VES Collab.)	REFID=51711
MORI	07A	JPSJ 76 074102	T. Mori *et al.*	(BELLE Collab.)	REFID=51691
ABLIKIM	06E	PR D73 052008	M. Ablikim *et al.*	(BES Collab.)	REFID=51057
ABLIKIM	06Q	PRL 97 202002	M. Ablikim *et al.*	(BES Collab.)	REFID=51487
AMELIN	05A	PAN 68 372 Translated from YAF 68 401.	D.V. Amelin *et al.*	(VES Collab.)	REFID=50766
AMSLER	04B	EPJ C33 23	C. Amsler *et al.*	(Crystal Barrel Collab.)	REFID=51079
BAI	04J	PL B594 47	J.Z. Bai *et al.*	(BES Collab.)	REFID=50167
BRIERE	00	PRL 84 26	R. Briere *et al.*	(CLEO Collab.)	REFID=47410
ACCIARRI	98Q	PL B418 399	M. Acciarri *et al.*	(L3 Collab.)	REFID=46316
BARBERIS	98C	PL B440 225	D. Barberis *et al.*	(WA 102 Collab.)	REFID=46346
WURZINGER	96	PL B374 283	R. Wurzinger *et al.*	(BONN, ORSAY, SACL+)	REFID=44992
PDG	94	PR D50 1173	L. Montanet *et al.*	(CERN, LBL, BOST+)	REFID=43653
AMSLER	93	ZPHY C58 175	C. Amsler *et al.*	(Crystal Barrel Collab.)	REFID=43311
BELADIDZE	92C	SJNP 55 1535 Translated from YAF 55 2748.	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)	REFID=43175

KARCH	92	ZPHY C54 33	K. Karch *et al.*	(Crystal Ball Collab.)	REFID=42170
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong *et al.*	(ATHU, BARI, BIRM+)	REFID=41862
BEHREND	91	ZPHY C49 401	H.J. Behrend *et al.*	(CELLO Collab.)	REFID=41497
AUGUSTIN	90	PR D42 10	J.E. Augustin *et al.*	(DM2 Collab.)	REFID=41352
BARU	90	ZPHY C48 581	S.E. Baru *et al.*	(MD-1 Collab.)	REFID=41366
BUTLER	90	PR D42 1368	F. Butler *et al.*	(Mark II Collab.)	REFID=41363
KARCH	90	PL B249 353	K. Karch *et al.*	(Crystal Ball Collab.)	REFID=41377
ROE	90	PR D41 17	N.A. Roe *et al.*	(ASP Collab.)	REFID=41014
AIHARA	88C	PR D38 1	H. Aihara *et al.*	(TPC-2γ Collab.)	REFID=40564
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev *et al.*	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
WILLIAMS	88	PR D38 1365	D.A. Williams *et al.*	(Crystal Ball Collab.)	REFID=40567
AIHARA	87	PR D35 2650	H. Aihara *et al.*	(TPC-2γ Collab.) JP	REFID=40009
ALBRECHT	87B	PL B199 457	H. Albrecht *et al.*	(ARGUS Collab.)	REFID=40265
ALDE	87B	ZPHY C36 603	D.M. Alde *et al.*	(LANL, BELG, SERP, LAPP)	REFID=40236
ANTREASYAN	87	PR D36 2633	D. Antreasyan *et al.*	(Crystal Ball Collab.)	REFID=40008
GIDAL	87	PRL 59 2012	G. Gidal *et al.*	(LBL, SLAC, HARV)	REFID=40223
ALDE	86	PL B177 115	D.M. Alde *et al.*	(SERP, BELG, LANL, LAPP)	REFID=20310
BARTEL	85E	PL 160B 421	W. Bartel *et al.*	(JADE Collab.)	REFID=10843
LANDSBERG	85	PRPL 128 301	L.G. Landsberg	(SERP)	REFID=10844
ALTHOFF	84E	PL 147B 487	M. Althoff *et al.*	(TASSO Collab.)	REFID=20305
BERGER	84B	PL 142B 125	C. Berger	(PLUTO Collab.)	REFID=20306
BINON	84	PL 140B 264	F.G. Binon *et al.*	(SERP, BELG, LAPP+)	REFID=20307
JENNI	83	PR D27 1031	P. Jenni *et al.*	(SLAC, LBL)	REFID=20304
BARTEL	82B	PL 113B 190	W. Bartel *et al.*	(JADE Collab.)	REFID=20300
BEHREND	82C	PL 114B 378	H.J. Behrend *et al.*	(CELLO Collab.)	REFID=20303
Also		PL 125B 518 (errat.)	H.J. Behrend *et al.*	(CELLO Collab.)	REFID=20302
DZHELYADIN	81	PL 105B 239	R.I. Dzhelyadin *et al.*	(SERP)	REFID=10836
STANTON	80	PL B92 353	N.R. Stanton *et al.*	(OSU, CARL, MCGI+)	REFID=40294
VIKTOROV	80	SJNP 32 520	V.A. Viktorov *et al.*	(SERP)	REFID=20298
		Translated from YAF 32 1005.			
APEL	79	PL 83B 131	W.D. Apel, K.H. Augenstein, E. Bertolucci	(KARLK+)	REFID=20295
BINNIE	79	PL 83B 141	D.M. Binnie *et al.*	(LOIC)	REFID=20296
ZANFINO	77	PRL 38 930	C. Zanfino *et al.*	(CARL, MCGI, OHIO+)	REFID=20293
GRIGORIAN	75	NP B91 232	A. Grigorian *et al.*	(+)	REFID=20287
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
DUANE	74	PRL 32 425	A. Duane *et al.*	(LOIC, SHMP)	REFID=20284
KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)	REFID=20286
DANBURG	73	PR D8 3744	J.S. Danburg *et al.*	(BNL, MICH) JP	REFID=20280
JACOBS	73	PR D8 18	S.M. Jacobs *et al.*	(BRAN, UMD, SYRA+) JP	REFID=20281
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez *et al.*	(BNL)	REFID=20205
APEL	72	PL 40B 680	W.D. Apel *et al.*	(KARLK, KARLE, PISA)	REFID=20275
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz *et al.*	(CERN)	REFID=20278
BASILE	71	NC 3A 371	M. Basile *et al.*	(CERN, BGNA, STRB)	REFID=20270
HARVEY	71	PRL 27 885	E.H. Harvey *et al.*	(MINN, MICH)	REFID=20272
BENSINGER	70	PL 33B 505	J.R. Bensinger *et al.*	(WISC)	REFID=20268
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL) I	REFID=20266
DAVIS	68	PL 27B 532	R. Davis *et al.*	(NWES, ANL)	REFID=20263
LONDON	66	PR 143 1034	G.W. London *et al.*	(BNL, SYRA) IJP	REFID=11774
BADIER	65B	PL 17 337	J. Badier *et al.*	(EPOL, SACL, AMST)	REFID=20253
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)	REFID=10761
DAUBER	64	PRL 13 449	P.M. Dauber *et al.*	(UCLA) JP	REFID=20247
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP	REFID=20252