

$a_2(1700)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

NODE=M162

 $a_2(1700)$ T-MATRIX POLE $\sqrt{s}$ Note that $\Gamma = -2 \operatorname{Im}(\sqrt{s})$.

NODE=M162PP

NODE=M162PP

NODE=M162PP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1630–1780) – i (60–250) OUR ESTIMATE			
$(1686 \pm 22^{+19}_{-7}) - i(211 \pm 38^{+32}_{-29})$	¹ KOPF	21	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$ and $191 \pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
$(1638.9 \pm 2.3^{+57.4}_{-0.1}) - i(112.0 \pm 1.3^{+0.9}_{-24.2})$	² ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1722 \pm 15 \pm 67) - i(124 \pm 9 \pm 32)$	³ RODAS	19	RVUE $191 \pi^- p \rightarrow \eta' \pi^- p$
$(1698 \pm 44) - i(133 \pm 28)$	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta \eta$
¹ Based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi$, $\eta'\pi$ and $K\bar{K}$ systems. ² Based on 2 poles, 2 channels ($\eta\pi$, $K\bar{K}$). ³ The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15 data.			

NODE=M162PP;LINKAGE=A

NODE=M162PP;LINKAGE=B

NODE=M162PP;LINKAGE=C

 $a_2(1700)$ MASS

NODE=M162M

NODE=M162M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1706 ± 14 OUR AVERAGE Error includes scale factor of 1.2.				
1681^{+22}_{-35}	46M	^{1,2} AGHASYAN	18B	COMP $190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$1726 \pm 12 \pm 25$		² ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
$1722 \pm 9 \pm 15$	18k	³ SCHEGELSKY	06	RVUE $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
1660 ± 40		² ABELE	99B	CBAR $1.94 \bar{p}p \rightarrow \pi^0 \eta \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1720 \pm 10 \pm 60$		⁴ JACKURA	18	RVUE $\pi^- p \rightarrow \eta \pi^- p$
1675 ± 25		ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
1702 ± 7	80k	⁵ UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
$1721 \pm 13 \pm 44$	145k	LU	05	B852 $18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
$1737 \pm 5 \pm 7$		ABE	04	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
1767 ± 14	221	⁶ ACCIARRI	01H	L3 $\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183-209 \text{ GeV}$
~ 1775		⁷ GRYGOREV	99	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1752 \pm 21 \pm 4$		ACCIARRI	97T	L3 $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
¹ Statistical error negligible. ² Breit-Wigner mass. ³ From analysis of L3 data at 183–209 GeV. ⁴ Superseded by RODAS 19. ⁵ Statistical error only. ⁶ Spin 2 dominant, isospin not determined, could also be $I=1$. ⁷ Possibly two $J^P = 2^+$ resonances with isospins 0 and 1.				

NODE=M162M;LINKAGE=B

NODE=M162M;LINKAGE=E

NODE=M162M;LINKAGE=SC

NODE=M162M;LINKAGE=D

NODE=M162M;LINKAGE=ST

NODE=M162M;LINKAGE=HA

NODE=M162M;LINKAGE=GR

 $a_2(1700)$ WIDTH

NODE=M162W

NODE=M162W


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
380^{+60}_{-50} OUR AVERAGE Error includes scale factor of 3.9. See the ideogram below.				
436^{+20}_{-16}	46M	^{1,2} AGHASYAN	18B	COMP $190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$190 \pm 18 \pm 30$		² ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
$336 \pm 20 \pm 20$	18k	³ SCHEGELSKY	06	RVUE $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
280 ± 70		² ABELE	99B	CBAR $1.94 \bar{p}p \rightarrow \pi^0 \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

280± 10±70		⁴ JACKURA	18	RVUE	$\pi^- p \rightarrow \eta \pi^- p$
270 ⁺ ₋ 50 20		ANISOVICH	09	RVUE	0.0 $\bar{p} p, \pi N$
417± 19	80k	⁵ UMAN	06	E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
279± 49±66	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
151± 22±24		ABE	04	BELL	10.6 $e^+ e^- \rightarrow$ $e^+ e^- K^+ K^-$
187± 60	221	⁶ ACCIARRI	01H L3		$\gamma \gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{see}} =$ 91, 183–209 GeV
150±110±34		ACCIARRI	97T L3		$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$

- ¹ Statistical error negligible.
² Breit-Wigner width.
³ From analysis of L3 data at 183–209 GeV.
⁴ Superseded by RODAS 19.
⁵ Statistical error only.
⁶ Spin 2 dominant, isospin not determined, could also be $I=1$.

NODE=M162W;LINKAGE=B
NODE=M162W;LINKAGE=E
NODE=M162W;LINKAGE=SC
NODE=M162W;LINKAGE=D
NODE=M162W;LINKAGE=ST
NODE=M162W;LINKAGE=HA



$a_2(1700)$ DECAY MODES

NODE=M162215;NODE=M162

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta \pi$	$(2.5 \pm 0.6) \%$
Γ_2 $\eta' \pi$	seen
Γ_3 $\gamma \gamma$	$(7.9 \pm 1.7) \times 10^{-7}$
Γ_4 $\rho \pi$	seen
Γ_5 $f_2(1270) \pi$	seen
Γ_6 $K \bar{K}$	$(1.3 \pm 0.8) \%$
Γ_7 $\omega \pi^- \pi^0$	seen
Γ_8 $\omega \rho$	seen

DESIG=4
DESIG=8;OUR EVAL;→ UNCHECKED ←
DESIG=1
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=5
DESIG=6;OUR EVAL;→ UNCHECKED ←
DESIG=7;OUR EVAL;→ UNCHECKED ←

$a_2(1700)$ PARTIAL WIDTHS

NODE=M162220

$\Gamma(\eta \pi)$				Γ_1	
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
9.5±2.0	870	¹ SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	

NODE=M162W3
NODE=M162W3

- ¹ From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

NODE=M162W3;LINKAGE=SC

$\Gamma(\gamma\gamma)$ Γ_3

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.30±0.05	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

NODE=M162W2
NODE=M162W2

NODE=M162W2;LINKAGE=SC

 $\Gamma(K\bar{K})$ Γ_6

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.0±3.0	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

NODE=M162W1
NODE=M162W1

NODE=M162W1;LINKAGE=SC

 $a_2(1700) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

NODE=M162225

$$\left[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi) \right] \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}} \quad (\Gamma_4 + \Gamma_5) \Gamma_3 / \Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.04±0.02		ACCIARRI 97T	L3	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.37^{+0.12}_{-0.08} \pm 0.10$	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
---------------------------------	-----	----------------------------	------	--

¹ From analysis of L3 data at 183–209 GeV.

NODE=M162G1
NODE=M162G1

NODE=M162G1;LINKAGE=SC

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$ $\Gamma_6 \Gamma_3 / \Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

20.6 ± 4.2 ± 4.6	¹ ABE 04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
49 ± 11 ± 13	² ACCIARRI 01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183\text{--}209 \text{ GeV}$

¹ Assuming spin 2.

² Spin 2 dominant, isospin not determined, could also be $I=1$.

NODE=M162G2
NODE=M162G2

NODE=M162G2;LINKAGE=AB
NODE=M162G;LINKAGE=HA

 $a_2(1700) \text{ BRANCHING RATIOS}$

NODE=M162235

 $\Gamma(\rho\pi) / \Gamma(f_2(1270)\pi)$ Γ_4 / Γ_5

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.4 \pm 0.4 \pm 0.1$	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
-----------------------	-----	----------------------------	------	--

¹ From analysis of L3 data at 183–209 GeV.

NODE=M162R01
NODE=M162R01

NODE=M162R01;LINKAGE=SC

 $\Gamma(K\bar{K}) / \Gamma(\eta\pi)$ Γ_6 / Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.029 \pm 0.04^{+0.011}_{-0.012}$	¹ KOPF 21	RVUE	$0.9 p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^- \text{ and } 191 \pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
$4.134 \pm 0.106^{+4.909}_{-2.988}$	² ALBRECHT 20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$

¹ From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.

² Residues from T-matrix pole, 2 poles, 2 channels ($\pi\eta, K\bar{K}$).

NODE=M162R00
NODE=M162R00

NODE=M162R00;LINKAGE=B

NODE=M162R00;LINKAGE=A

 $\Gamma(\eta'\pi) / \Gamma(\eta\pi)$ Γ_2 / Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.035 \pm 0.044^{+0.069}_{-0.012}$	¹ KOPF 21	RVUE	$0.9 p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^- \text{ and } 191 \pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
-------------------------------------	----------------------	------	--

¹ From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.

NODE=M162R02
NODE=M162R02

NODE=M162R02;LINKAGE=A

$a_2(1700)$ REFERENCES

NODE=M162

KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)	REFID=61470
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=59554
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)	REFID=59003
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57953
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46904
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>		REFID=46909
		Translated from YAF 62 513.			
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45761