

$\eta_b(1S)$

$I^G(J^P C) = 0^+(0^- +)$

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$.

NODE=M171

NODE=M171

NODE=M171M


NODE=M171M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9398.7 ± 2.0 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.			
9394.8 ± 2.7 ± 4.5	29k	FULSOM	18	BELL $\Upsilon(2S) \rightarrow \gamma X$
9400.7 ± 1.7 ± 1.6	33.1k	TAMPONI	15	BELL $e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$
9402.4 ± 1.5 ± 1.8	34k	¹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$
9391.8 ± 6.6 ± 2.0	2.3k	² BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
9394.2 ± 4.8 ± 2.0	13k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
9388.9 ± 3.1 ± 2.7	19k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9393.2 ± 3.4 ± 2.3	10	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$
9300 ± 20 ± 20		HEISTER	02D ALEP	181–209 $e^+ e^-$

¹With floating width. Not independent of the corresponding mass difference measurement.

²Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass difference measurements.

³Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.



NODE=M171M;LINKAGE=MI

NODE=M171M;LINKAGE=AU

NODE=M171M;LINKAGE=DO

$m_{\Upsilon(1S)} - m_{\eta_b}$


NODE=M171M2

NODE=M171M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62.3 ± 3.2 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
57.9 ± 1.5 ± 1.8	34k	¹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$
68.5 ± 6.6 ± 2.0	2.3 ± 0.5k	² BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
66.1 ± 4.8 ± 2.0	13 ± 5k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
71.4 ± 2.3 ± 2.7	19 ± 3k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
67.1 ± 3.4 ± 2.3	10 ± 5	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$

- ¹ With floating width. Not independent of the corresponding mass measurement.
² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass measurements.
³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M2;LINKAGE=MI
 NODE=M171M2;LINKAGE=AU
 NODE=M171M2;LINKAGE=DO



γ ENERGY IN $\gamma(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
920.6^{+2.8}_{-3.2} OUR AVERAGE				
$918.6 \pm 6.0 \pm 1.9$	$2.3 \pm 0.5k$	1 BONVICINI	10	CLEO $\gamma(3S) \rightarrow \gamma X$
$921.2 \pm 2.1 \pm 2.4$	$19 \pm 3k$	1 AUBERT	08V	BABR $\gamma(3S) \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171DM

NODE=M171DM

NODE=M171DM;LINKAGE=BO

γ ENERGY IN $\gamma(2S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
609.3^{+4.6}_{-4.5} ± 1.9				
	$13 \pm 5k$	1 AUBERT	09AQ BABR	$\gamma(2S) \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171U2S

NODE=M171U2S

NODE=M171U2S;LINKAGE=AU

$\eta_b(1S)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10⁺⁵₋₄ OUR AVERAGE				
$8 \pm 5 \pm 5$	$33.1k$	1 TAMPONI	15	BELL $e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$
$10.8 \pm 4.0 \pm 4.5$	$34k$	1 MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$

¹ With floating mass.

NODE=M171W

NODE=M171W

NODE=M171W;LINKAGE=MI

$\eta_b(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $3h^+ 3h^-$	not seen	
Γ_3 $2h^+ 2h^-$	not seen	
Γ_4 $4h^+ 4h^-$	not seen	
Γ_5 $\gamma\gamma$	not seen	
Γ_6 $\mu^+ \mu^-$	$<9 \times 10^{-3}$	90%
Γ_7 $\tau^+ \tau^-$	$<8 \%$	90%

 $\eta_b(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ **$\Gamma(3h^+ 3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_5/\Gamma$
------------	-----	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<470	95	ABDALLAH	06	DLPH	161–209 $e^+ e^-$
<132	95	HEISTER	02D	ALEP	181–209 $e^+ e^-$

 $\Gamma(2h^+ 2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_3\Gamma_5/\Gamma$
------------	-----	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<190	95	ABDALLAH	06	DLPH	161–209 $e^+ e^-$
< 48	95	HEISTER	02D	ALEP	181–209 $e^+ e^-$

 $\Gamma(4h^+ 4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_4\Gamma_5/\Gamma$
------------	-----	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<660	95	ABDALLAH	06	DLPH	161–209 $e^+ e^-$
------	----	----------	----	------	-------------------

 $\eta_b(1S)$ BRANCHING RATIOS **$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
-------	------	-------------	------	---------	-------------------

seen	34k	MIZUK	12	BELL	$e^+ e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
------	-----	-------	----	------	---

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
-------	-----	-------------	------	---------	-------------------

$<9 \times 10^{-3}$	90	1 AUBERT	09Z	BABR	$e^+ e^- \rightarrow \gamma(2S, 3S) \rightarrow \gamma\eta_b$
---------------------	----	----------	-----	------	---

1 Obtained using $B(\gamma(2S) \rightarrow \gamma\eta_b) = (4.2^{+1.1}_{-1.0} \pm 0.9) \times 10^{-4}$ and $B(\gamma(3S) \rightarrow \gamma\eta_b) = (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$. This limit is equivalent to $B(\eta_b \rightarrow \mu^+ \mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$ measurement.

 $\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
-------	-----	-------------	------	---------	-------------------

$<8 \times 10^{-2}$	90	AUBERT	09P	BABR	$e^+ e^- \rightarrow \gamma\tau^+\tau^-$
---------------------	----	--------	-----	------	--

 $\eta_b(1S)$ REFERENCES

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRC 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER	02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)

NODE=M171225;NODE=M171

DESIG=7
DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=5
DESIG=6

NODE=M171230

NODE=M171G1
NODE=M171G1

NODE=M171G2
NODE=M171G2

NODE=M171G3
NODE=M171G3

NODE=M171235

NODE=M171R03
NODE=M171R03

NODE=M171R01
NODE=M171R01

NODE=M171R01;LINKAGE=AU

NODE=M171R02
NODE=M171R02

NODE=M171

REFID=59535
REFID=56996
REFID=54288
REFID=54718
REFID=53231
REFID=53106
REFID=53062
REFID=52930
REFID=52262
REFID=51042
REFID=48577