

$\psi_2(3823)$

$I^G(J^{PC}) = 0^-(2^{--})$
 I, J, P need confirmation.

was $\psi(3823)$, $X(3823)$

Seen by BHARDWAJ 13 in $B \rightarrow \chi_{c1} \gamma K$ and ABLIKIM 15S in $e^+ e^- \rightarrow \pi^+ \pi^- \gamma \chi_{c1}$ decays as a narrow peak in the invariant mass distribution of the $\chi_{c1} \gamma$ system. Properties consistent with the $\psi_2(1^3D_2) c \bar{c}$ state.

NODE=M212

NODE=M212

$\psi_2(3823)$ MASS

NODE=M212M

NODE=M212M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3823.51 ± 0.34 OUR AVERAGE				
3824.5 ± 2.4 ± 1.0	30	¹ ABLIKIM	23J BES3	$e^+ e^- \rightarrow \pi^0 \pi^0 \chi_{c1} \gamma$
3823.12 ± 0.43 ± 0.13	120	ABLIKIM	22R BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
3824.08 ± 0.53 ± 0.14	137	² AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
3823.1 ± 1.8 ± 0.7	33 ± 10	³ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1} \gamma K^\pm$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3821.7 ± 1.3 ± 0.7	19 ± 5	⁴ ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$

¹ Using the measured $m_{\psi_2(3823)} - m_{\psi(2S)}$ and assuming $m_{\psi(2S)} = 3686.097$ MeV from PDG 22.

² Using the measured $m_{\psi_2(3823)} - m_{\psi(2S)} = 137.98 \pm 0.53 \pm 0.14$ MeV.

³ From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$ and $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$ with significance 4.0σ including systematics. Corrected for the measured $\psi(2S)$ mass using $B \rightarrow \psi(2S) K \rightarrow (\gamma \chi_{c1}) K$ decays.

⁴ From a simultaneous unbinned maximum likelihood fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at $\sqrt{s}$ values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1} \gamma$ and $\psi_2(3823) \rightarrow \chi_{c1} \gamma$ together, with floating mass scale offset for $\psi(2S)$, floating $\psi_2(3823)$ mass, and zero $\psi_2(3823)$ width, resulting in a significance of 5.9σ when including systematic uncertainties. Superseded by ABLIKIM 22R.

NODE=M212M;LINKAGE=D

NODE=M212M;LINKAGE=C

NODE=M212M;LINKAGE=A

NODE=M212M;LINKAGE=B

$m_{\psi_2(3823)} = m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
137.98 ± 0.53 ± 0.14	137	¹ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
¹ AAIJ 20S also reports $m_{\chi_{c1}(3872)} - m_{\psi_2(3823)} = 47.50 \pm 0.53 \pm 0.13$ MeV.				

NODE=M212A00
 NODE=M212A00

NODE=M212A00;LINKAGE=A

$\psi_2(3823)$ WIDTH

NODE=M212W

NODE=M212W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 2.9					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 18.8	90	30	¹ ABLIKIM	23J BES3	$e^+ e^- \rightarrow \pi^0 \pi^0 \chi_{c1} \gamma$
< 5.2	90		² AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
< 16	90		³ ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
< 24	90		⁴ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1} \gamma K^\pm$

¹ From a fit of $e^+ e^- \rightarrow \pi^0 \pi^0 \chi_{c1} \gamma$ data at $\sqrt{s}$ values from 4.23 to 4.70 GeV to a Breit-Wigner function with floating width, using the Bayesian approach.

² AAIJ 20S also provides a limit of < 6.6 MeV with 95% CL.

³ From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at $\sqrt{s}$ values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to a Breit-Wigner function with the mass fixed from the likelihood fit above, Gaussian resolution smearing, and floating width.

⁴ From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$ and $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$ with significance 4.0σ including systematics.

NODE=M212W;LINKAGE=D

NODE=M212W;LINKAGE=C

NODE=M212W;LINKAGE=B

NODE=M212W;LINKAGE=A

$\psi_2(3823)$ DECAY MODES

Branching fractions are given relative to the one **DEFINED AS 1**.

NODE=M212215;NODE=M212

NODE=M212

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $J/\psi(1S)\pi^+\pi^-$	<0.06	90%
Γ_2 $J/\psi(1S)\pi^0\pi^0$	<0.11	90%
Γ_3 $J/\psi(1S)\pi^0$	<0.030	90%
Γ_4 $J/\psi(1S)\eta$	<0.14	90%
Γ_5 $\chi_{c0}\gamma$	<0.24	90%
Γ_6 $\chi_{c1}\gamma$	DEFINED AS 1	
Γ_7 $\chi_{c2}\gamma$	0.28 $^{+0.14}_{-0.11}$	

DESIG=3

DESIG=5

DESIG=6

DESIG=7

DESIG=4

DESIG=1

DESIG=2

 $\psi_2(3823)$ BRANCHING RATIOS

NODE=M212225

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	EVTS

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	¹ ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	137 $\pm$ 26	AAIJ	20S	LHCB $B^+ \rightarrow J/\psi\pi^+\pi^-K^+$

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R00;LINKAGE=A

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(\chi_{c1}\gamma)$	Γ_1/Γ_6
VALUE	CL%

<0.06	90	¹ ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
-------	----	----------------------	-----	------	----------------------------------

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R06
NODE=M212R06

NODE=M212R06;LINKAGE=A

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(\chi_{c1}\gamma)$	Γ_2/Γ_6
VALUE	CL%

<0.11	90	¹ ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
-------	----	----------------------	-----	------	----------------------------------

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R07
NODE=M212R07

NODE=M212R07;LINKAGE=A

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(\chi_{c1}\gamma)$	Γ_3/Γ_6
VALUE	CL%

<0.03	90	¹ ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
-------	----	----------------------	-----	------	----------------------------------

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R08
NODE=M212R08

NODE=M212R08;LINKAGE=A

$\Gamma(J/\psi(1S)\eta)/\Gamma(\chi_{c1}\gamma)$	Γ_4/Γ_6
VALUE	CL%

<0.14	90	¹ ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
-------	----	----------------------	-----	------	----------------------------------

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R09
NODE=M212R09

NODE=M212R09;LINKAGE=A

$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$	Γ_4/Γ_1
VALUE	DOCUMENT ID

4.4 $^{+2.5}_{-1.9} \pm 0.9$	¹ AAIJ	22D	LHCB	$B^+ \rightarrow J/\psi(1S)\eta K^+$
------------------------------	-------------------	-----	------	--------------------------------------

¹ Using the branching ratio for $B^+ \rightarrow \psi_2(3823)K^+$ with $\psi_2(3823) \rightarrow J/\psi(1S)\pi^+\pi^-$ from AAIJ 20S.

NODE=M212R10
NODE=M212R10

NODE=M212R10;LINKAGE=A

$\Gamma(\chi_{c0}\gamma)/\Gamma_{\text{total}}$	Γ_5/Γ
VALUE	DOCUMENT ID

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	¹ ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
----------	----------------------	-----	------	----------------------------------

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R04
NODE=M212R04

NODE=M212R04;LINKAGE=A

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen	120	¹ ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
seen	63 ± 9	² ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	16 ± 5	³ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^0\pi^0X$
seen	33 ± 10	⁴ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

NODE=M212R01
 NODE=M212R01

OCCUR=2

NODE=M212R01;LINKAGE=D

NODE=M212R01;LINKAGE=B

NODE=M212R01;LINKAGE=C

NODE=M212R01;LINKAGE=A

¹ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak R with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV, which leads to $\Gamma(e^+e^-)B(R \rightarrow \pi^+\pi^-\psi_2(3823))B(\psi_2(3823) \rightarrow \chi_{c1}\gamma) = 0.57 \pm 0.08$ eV.

² From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$. Signal has a 11.8 σ significance.

³ From a fit of the invariant $\pi^0\pi^0$ recoil-mass distribution. Signal has a 4.3 σ significance.

⁴ BHARDWAJ 13 reports $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$ with statistical significance 3.8 σ .

 $\Gamma(\chi_{c0}\gamma)/\Gamma(\chi_{c1}\gamma)$ Γ_5/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M212R05
 NODE=M212R05

NODE=M212R05;LINKAGE=A

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	¹ ABLIKIM 210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$	
not seen	² ABLIKIM 15s BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$	
not seen	³ BHARDWAJ 13 BELL	$B^\pm \rightarrow \chi_{c2}\gamma K^\pm$	

NODE=M212R02
 NODE=M212R02

NODE=M212R02;LINKAGE=C

NODE=M212R02;LINKAGE=B

NODE=M212R02;LINKAGE=A

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$. Signal has a 3.2 σ significance.

² From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at $\sqrt{s}$ values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c2}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c2}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.

³ BHARDWAJ 13 reports $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c2}) < 3.6 \times 10^{-6}$ at 90% CL.

 $\Gamma(\chi_{c2}\gamma)/\Gamma(\chi_{c1}\gamma)$ Γ_7/Γ_6

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.28^{+0.14}_{-0.11} \pm 0.02$	9 ± 4		¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$

NODE=M212R03
 NODE=M212R03

NODE=M212R03;LINKAGE=B

NODE=M212R03;LINKAGE=A

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<0.42	90	² ABLIKIM	15s BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
<0.41	90	BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1/c2}\gamma K^\pm$

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

² From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1(2)}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at $\sqrt{s}$ values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1(2)}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c1(2)}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.

 $\psi_2(3823)$ REFERENCES

NODE=M212

ABLIKIM	23J	JHEP 2302 171	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	22D	JHEP 2204 046	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ABLIKIM	210	PR D103 L091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)

REFID=62052
 REFID=61647
 REFID=61664
 REFID=61634
 REFID=61121
 REFID=60526
 REFID=56784
 REFID=55412